Extremely projective normed spaces

N. T. Nemesh

Abstract: A normed space is called extremely projective if it is C-topologically projective in the category of normed spaces for every $C > 1$. Helemskii proved that the spaces $\ell_1^0(\Lambda)$ of finitely supported functions on a set Λ with the ℓ_1-norm are extremely projective and asked whether every extremely projective normed space is isometrically isomorphic to some $\ell_1^0(\Lambda)$; the same conjecture was raised by Grønbæk. We answer this question in the negative and prove that the conjecture holds in the almost isometric form: a normed space P is extremely projective if and only if it is topologically projective and its completion is isometrically isomorphic to $\ell_1(\Gamma)$ for some set Γ, if and only if the Banach–Mazur distance between P and $\ell_1^0(\Gamma)$ equals 1. For spaces of countable dimension this means that P is extremely projective if and only if its completion is isometrically isomorphic to ℓ_1 or ℓ_1^n. The proof is a transfinite back-and-forth construction. Along the way we show that extremely projective normed spaces contain no infinite dimensional complete subspaces, so that no infinite dimensional Banach space is extremely projective in the category of normed spaces. Finally, we show that a dense subspace of $\ell_1(\Gamma)$ is extremely projective if and only if it admits a continuous filtration with countable blocks, dense pieces and countable dimensional successive quotients. For $|\Gamma| = \aleph_1$ this becomes a criterion in the spirit of the theory of almost free abelian groups, requiring the countably generated pieces to have countable dimension and the set of discontinuities of their chain to be non-stationary. Using ladder systems on stationary sets we construct $2^{\aleph_1}$ pairwise non-isomorphic dense subspaces of $\ell_1(\omega_1)$ all of whose countably generated pieces are extremely projective, none of which is topologically projective.

Keywords: projective normed space, topological projectivity, extreme projectivity, Banach–Mazur distance, ℓ_1-space.

1 Introduction

Projectivity in categories of normed and Banach modules admits several non-equivalent versions. Besides the classical relative projectivity of Helemskii [1], which ignores the norms of lifting operators, there are so-called metric and topological projectivity [2, 4], where the norm of a lifting is controlled by a constant. In the metric theory this constant is 1, and in the topological theory it is some fixed $C \geq 1$. The present paper is devoted to the limiting case between these two theories: a module is called extremely projective if it is C-topologically projective for every $C > 1$. This notion was introduced and studied by Helemskii [3, 2], whose definition is formally different but equivalent to ours, see proposition 2.1. We restrict ourselves to the simplest possible algebra, namely the zero algebra. Normed modules over the zero algebra are just normed spaces, so we deal with extreme projectivity in the category **Nor** of normed spaces and bounded linear operators.

For Banach spaces the corresponding questions are classical. By a theorem of Köthe [5] topologically projective Banach spaces are exactly the Banach spaces topologically isomorphic to $\ell_1(\Lambda)$ for some set Λ, and by a theorem of Grothendieck [6] a Banach space has the metric lifting property if and only if it is isometrically isomorphic to some $\ell_1(\Lambda)$. For normed spaces the answers are different, because the category **Nor** has its own free objects. These are the spaces $\ell_1^0(\Lambda)$ of finitely supported functions on a set Λ with the ℓ_1-norm. Grønbæk [8] extended Köthe's theorem to **Nor**: a normed space E is topologically projective if and only if it is topologically isomorphic to

$\ell_1^0(\Lambda)$, where Λ is a set whose cardinality is the density character of E (for infinite dimensional E). Helemskii [2] proved the metric counterpart: a normed space is metrically projective if and only if it is isometrically isomorphic to some $\ell_1^0(\Lambda)$. For extreme projectivity the following facts were known. The spaces $\ell_1^0(\Lambda)$ are extremely projective, and a normed space is extremely projective if and only if it is a near-retract of some $\ell_1^0(\Lambda)$ [3, proposition 2.2 and the proposition of the introduction]. Every extremely projective normed space is, up to an isometric isomorphism, a dense subspace of $\ell_1(\Lambda)$ for some set Λ [3, proposition 2.3], see also [8, section 3], and every separable topologically projective normed space has at most countable algebraic dimension [3, proposition 2.4]. Both Helemskii [3, section 2], [2, remark 3.6] and Grønbæk [8, conjecture 3.1] asked whether, conversely, every extremely projective normed space is isometrically isomorphic to some $\ell_1^0(\Lambda)$; in a more detailed form, Helemskii asked which dense subspaces of $\ell_1(\Lambda)$ are extremely projective.

The main results of this paper are the following. First, the answer to the question of Helemskii and Grønbæk is negative: the subspace of ℓ_1 spanned by the finitely supported sequences and a single sequence with infinite support is extremely projective but not isometrically isomorphic to any $\ell_1^0(\Lambda)$. Second, extreme projectivity is an almost isometric property: a normed space P is extremely projective if and only if it is topologically projective and its completion is isometrically isomorphic to $\ell_1(\Gamma)$ for some set Γ, if and only if for every $\varepsilon > 0$ there is a linear bijection $T : P \to \ell_1^0(\Gamma)$ with $\|T\|\|T^{-1}\| < 1 + \varepsilon$. Thus, the conjecture of Grønbæk holds in the almost isometric form. For spaces of countable algebraic dimension the condition of topological projectivity is superfluous: such a space is extremely projective if and only if its completion is isometrically isomorphic to ℓ_1 or to ℓ_1^n, so every dense subspace of ℓ_1 of countable algebraic dimension is extremely projective. Third, we extend the dimension restriction of Helemskii to non-separable spaces: every separable subspace of a topologically projective normed space has countable algebraic dimension, and therefore topologically projective normed spaces contain no infinite dimensional subspaces which are complete with respect to a norm dominating the original one. In particular, an infinite dimensional Banach space is never extremely projective in **Nor**, although it may well be extremely projective in the category **Ban** of Banach spaces. The last fact explains the main technical difficulty of the subject: when the lifting problems are posed for all normed spaces rather than Banach spaces, the standard tools of Banach space geometry, which rely on completeness, are no longer available, and we must work with the finite dimensional pieces of our spaces directly. Our main tool is a back-and-forth construction, which in the case of countable dimension replaces the decomposition method of Pełczyński in the proofs of Köthe and Grønbæk, and in the general case takes the form of a transfinite recursion along a filtration of Γ whose perturbations are controlled on disjoint blocks of coordinates. Fourth, we describe the topologically projective dense subspaces of $\ell_1(\Gamma)$ in terms of the coordinate structure of $\ell_1(\Gamma)$ only: a dense subspace P is extremely projective if and only if Γ admits a continuous filtration with countable blocks along which the pieces $P \cap \ell_1(\Gamma_\delta)$ are dense, form a continuous chain and have countable dimensional successive quotients. Thus, for uncountable Λ the question of Helemskii on dense subspaces of $\ell_1(\Lambda)$ does not reduce to the countable case: a dense subspace of $\ell_1(\Lambda)$ is extremely projective if and only if it is topologically projective, and for uncountable Λ this is a genuine restriction. Fifth, for $|\Gamma| = \aleph_1$ we obtain a criterion in the spirit of the theory of almost free abelian groups: a dense subspace P of $\ell_1(\omega_1)$ is extremely projective if and only if its subspaces $P \cap \ell_1(\delta)$, $\delta < \omega_1$, have countable dimension and the set of limit ordinals at which the chain of these subspaces is discontinuous is not stationary. The class of this set modulo non-stationary sets is an isomorphism invariant of P. Using it we construct, from ladder systems on stationary subsets of ω_1, $2^{\aleph_1}$ pairwise non-isomorphic dense subspaces of $\ell_1(\omega_1)$ which are not topologically projective, although all their separable subspaces have countable dimension and all their countably generated pieces are extremely projective.

Let us fix the notation and recall the definitions. All normed spaces are considered over the complex field; the real case is identical. For a normed space E we denote by B_E its closed unit ball and by B_E° its open unit ball. By **Nor** we denote the category of normed spaces and bounded linear operators and by **Ban** its full subcategory of Banach spaces. For a set Λ we denote by $\ell_1(\Lambda)$ the Banach space of absolutely summable families and by $\ell_1^0(\Lambda)$ its dense subspace consisting of finitely supported families. For $\lambda \in \Lambda$ the symbol e_λ denotes the family which equals 1 at λ and 0 elsewhere; the family $(e_\lambda)_{\lambda\in\Lambda}$ is a Hamel basis of $\ell_1^0(\Lambda)$. For a normed space E and a set Λ there is a bijection between bounded families $(x_\lambda)_{\lambda\in\Lambda}$ in E and bounded linear operators $T : \ell_1^0(\Lambda) \to E$, given by $T(e_\lambda) = x_\lambda$, and $\|T\| = \sup_{\lambda\in\Lambda}\|x_\lambda\|$ [3, proposition 2.1], [8, proposition 1.10]. This is the sense in which the spaces $\ell_1^0(\Lambda)$ are free objects of **Nor**; note that no completeness of E is required here, in contrast with the case of $\ell_1(\Lambda)$.

We say that a bounded linear operator $\xi : X \to Y$ between normed spaces is c-topologically surjective, where $c \geq 1$, if $c\,\xi(B_X^\circ) \supset B_Y^\circ$, and that ξ is coisometric if it is contractive and 1-topologically surjective, that is, if $\xi(B_X^\circ) = B_Y^\circ$. Equivalently, ξ is c-topologically surjective if for every $y \in Y$ and every $\delta > 0$ there exists $x \in X$ with $\xi(x) = y$ and $\|x\| < c\|y\| + \delta$. A linear bijection $T : E \to F$ is called a c-isomorphism if $\|T\|\|T^{-1}\| \leq c$. The Banach–Mazur distance between E and F is

$$d_{BM}(E, F) = \inf\{\|T\|\|T^{-1}\| : T : E \to F \text{ is a linear bijection}\},$$

with $d_{BM}(E, F) = +\infty$ if there are no bounded linear bijections with bounded inverses between E and F. A bounded linear operator $\zeta : E \to F$ is called a c-retraction if there exists a bounded linear operator $\eta : F \to E$ with $\zeta\eta = 1_F$ and $\|\zeta\|\|\eta\| \leq c$. In this case F is called a c-retract of E.

Definition 1.1. *Let $C \geq 1$. A normed space P is called C-topologically projective if for every $c \geq 1$, every c-topologically surjective operator $\xi : X \to Y$ between normed spaces and every bounded linear operator $\phi : P \to Y$ there exists a bounded linear operator $\psi : P \to X$ such that $\xi\psi = \phi$ and $\|\psi\| \leq cC\|\phi\|$. A normed space is called topologically projective if it is C-topologically projective for some $C \geq 1$, and extremely projective if it is C-topologically projective for every $C > 1$.*

The operator ψ in definition 1.1 is called a lifting of ϕ along ξ. Extreme projectivity in the sense of definition 1.1 is precisely the extreme projectivity of Helemskii, see proposition 2.1. We emphasize that the spaces X and Y in definition 1.1 are arbitrary normed spaces. If we restrict them to Banach spaces, we obtain the notions of topological and extreme projectivity in **Ban**, which we shall mention only for comparison. The two notions differ substantially, see section 3.

The paper is organized as follows. In section 2 we collect the basic properties of extremely projective normed spaces: the spaces $\ell_1^0(\Lambda)$ are extremely projective, extreme projectivity passes to $(1+\varepsilon)$-retracts, and a normed space is extremely projective if and only if it is a $(1+\varepsilon)$-retract of a free normed space for every $\varepsilon > 0$. In section 3 we discuss the dimension restrictions on topologically projective normed spaces. Section 4 is devoted to the almost isometric characterization of extremely projective normed spaces of countable dimension, and section 5 contains the counterexample to the question of Helemskii and Grønbæk. Section 6 treats spaces of uncountable dimension: it contains the transfinite back-and-forth construction, the closing-off arguments and the main theorem. Section 7 is devoted to the case $|\Gamma| = \aleph_1$: it contains the criterion in terms of stationary sets, the isomorphism invariant and the ladder spaces.

2 Basic properties of extremely projective normed spaces

We start by comparing definition 1.1 with Helemskii's one. Our definition of extreme projectivity is equivalent to Helemskii's definition [3, definition I]: a normed space P is extremely projective in the sense of Helemskii if for every coisometric operator $\xi : X \to Y$, every bounded linear operator $\phi : P \to Y$ and every $\delta > 0$ there exists a lifting $\psi : P \to X$ of ϕ along ξ with $\|\psi\| < \|\phi\| + \delta$.

Proposition 2.1. *A normed space is extremely projective in the sense of definition 1.1 if and only if it is extremely projective in the sense of Helemskii.*

Proof. Let P be extremely projective in the sense of definition 1.1, $\xi : X \to Y$ be a coisometric operator, $\phi : P \to Y$ be a bounded linear operator and $\delta > 0$. If $\phi = 0$ we take $\psi = 0$. Otherwise we put $C = 1 + \delta\|\phi\|^{-1}/2$. Since ξ is 1-topologically surjective, C-topological projectivity of P gives a lifting ψ of ϕ along ξ with $\|\psi\| \le C\|\phi\| < \|\phi\| + \delta$.

Conversely, let P be extremely projective in the sense of Helemskii, $C > 1$, $\xi : X \to Y$ be a c-topologically surjective operator and $\phi : P \to Y$ be a non-zero bounded linear operator. Consider the operator $\xi' = c\xi$ and endow the vector space X with the norm $\|x\|' = \max(\|x\|, \|\xi'(x)\|)$, which is equivalent to the original one since $\|x\| \le \|x\|' \le \max(1, c\|\xi\|)\|x\|$. Denote the resulting normed space by X'. The operator $\xi' : X' \to Y$ is contractive by construction. If $y \in B_Y^\circ$, then by c-topological surjectivity of ξ there is $x \in B_X^\circ$ with $\xi'(x) = y$, and $\|x\|' = \max(\|x\|, \|y\|) < 1$. Thus, $\xi' : X' \to Y$ is coisometric. By Helemskii's definition there is a lifting $\psi' : P \to X'$ of ϕ along ξ' with $\|\psi'\| < \|\phi\| + (C-1)\|\phi\| = C\|\phi\|$. Regarded as an operator into X, the operator $\psi = c\psi'$ satisfies $\xi\psi = \xi'\psi' = \phi$ and $\|\psi\| \le c\|\psi'\|' \le cC\|\phi\|$. Since $C > 1$ is arbitrary, P is extremely projective in the sense of definition 1.1. □

Now we turn to the free objects of **Nor**.

Proposition 2.2. *Let Λ be a set. Then $\ell_1^0(\Lambda)$ is $(1+\varepsilon)$-topologically projective for every $\varepsilon > 0$. In particular, $\ell_1^0(\Lambda)$ is extremely projective.*

Proof. By [3, proposition 2.2] the space $\ell_1^0(\Lambda)$ is extremely projective in the sense of Helemskii, and the proof of proposition 2.1 shows that Helemskii's extreme projectivity gives exactly $(1+\varepsilon)$-topological projectivity for every $\varepsilon > 0$. □

The constant $1 + \varepsilon$ in proposition 2.2 cannot be replaced by 1. In fact, with the convention of definition 1.1, 1-topological projectivity is an empty notion in **Nor**.

Proposition 2.3. *The only 1-topologically projective normed space is the zero space.*

Proof. Let X be the space of finitely supported sequences with the sup-norm and $f : X \to \mathbb{C}$ be the functional $f(x) = s^{-1}\sum_{n=1}^\infty 2^{-n}(1-2^{-n})x_n$, where $s = \sum_{n=1}^\infty 2^{-n}(1-2^{-n})$. Clearly, $\|f\| \le 1$. The operator f is coisometric: for $z \in B_{\mathbb{C}}^\circ$ choose N with $|z| < s^{-1}\sum_{n=1}^N 2^{-n}(1-2^{-n})$, then $z = f(x)$ for the sequence $x = t\,\mathrm{sgn}(z)\sum_{n=1}^N e_n$ with a suitable $t \in [0,1)$, and $\|x\| < 1$. On the other hand, $|f(x)| = 1$ for $x \in B_X$ would force $x_n = \mathrm{sgn}(f(x))$ for all n, which is impossible for a finitely supported sequence. Thus, $f(B_X) = f(B_X^\circ) = B_{\mathbb{C}}^\circ$.

Now let P be a non-zero 1-topologically projective normed space. Take $p \in P$ with $\|p\| = 1$ and, by the Hahn–Banach theorem, a functional $g \in P^*$ with $\|g\| = g(p) = 1$. Lifting g along the coisometric operator f we get an operator $\psi : P \to X$ with $f\psi = g$ and $\|\psi\| \le 1$. Then $x = \psi(p) \in B_X$ and $f(x) = g(p) = 1$, which contradicts the previous paragraph. □

Proposition 2.3 shows why in **Nor** it is natural to consider all constants $C > 1$ at once, instead of the single constant $C = 1$ of the metric theory.

Remark 2.4. *The proof of proposition 2.3 uses only the fact that the coisometric functional f does not attain its norm. Functionals of this kind exist on Banach spaces too, for example $x \mapsto \sum_{n=1}^{\infty}(1-2^{-n})x_n$ on ℓ_1. Therefore, if c-topologically surjective operators are defined via open balls as in definition 1.1, then 1-topological projectivity is an empty notion in* **Ban** *as well, and the free Banach spaces $\ell_1(\Lambda)$ are only $(1+\varepsilon)$-topologically projective for every $\varepsilon > 0$. The metric theory of [2] avoids this problem by posing lifting problems along strictly coisometric operators, that is, operators mapping the closed unit ball onto the closed unit ball; with this convention $\ell_1(\Lambda)$ and $\ell_1^0(\Lambda)$ are 1-topologically projective in* **Ban** *and* **Nor** *respectively, and 1-topological projectivity coincides with metric projectivity in the sense of [2]. For the purposes of this paper the choice of convention is immaterial: if we call $\xi : X \to Y$ strictly c-topologically surjective whenever $c\xi(B_X) \supset B_Y$ and define strict C-topological projectivity by lifting along such operators with the same bound $\|\psi\| \le cC\|\phi\|$, then every C-topologically projective space is strictly C-topologically projective, and every strictly C-topologically projective space is $C(1+\varepsilon)$-topologically projective for every $\varepsilon > 0$. The first claim is obvious, since strictly c-topologically surjective operators are c-topologically surjective. For the second one, given a c-topologically surjective operator $\xi : X \to Y$ and $\varepsilon > 0$, endow X with the equivalent norm $\|x\|'' = \max(\|x\|, (1+\varepsilon)c\|\xi(x)\|)$; then $(1+\varepsilon)c\xi$ becomes a contractive operator mapping the closed unit ball onto the closed unit ball (for $y \in B_Y$ the c-topological surjectivity of ξ gives x with $(1+\varepsilon)c\xi(x) = y$ and $\|x\| < (1+\varepsilon)^{-1}\|y\| + \varepsilon(1+\varepsilon)^{-1} \le 1$, and then $\|x\|'' \le 1$), and a lifting ψ'' along it with $\|\psi''\|'' \le C\|\phi\|$ gives the lifting $\psi = (1+\varepsilon)c\psi''$ along ξ with $\|\psi\| \le (1+\varepsilon)cC\|\phi\|$. In particular, the two conventions lead to the same notion of extreme projectivity.*

Next we record the standard permanence property of projectivity. The proof is the same as in the Banach case, see [4, lemma 2.7]; for the second part see also [3, proposition 1.1].

Proposition 2.5. *Let P' be a C-topologically projective normed space and P be a c-retract of P'. Then P is cC-topologically projective. Consequently, a normed space which is a $(1+\varepsilon)$-retract of an extremely projective normed space for every $\varepsilon > 0$ is extremely projective.*

For a normed space P consider the operator

$$\pi_P : \ell_1^0(B_P) \to P, \quad e_x \mapsto x.$$

It is well defined and contractive, because the family $(x)_{x \in B_P}$ is bounded by 1. Moreover, π_P maps the closed unit ball of $\ell_1^0(B_P)$ onto B_P, since $\pi_P(e_x) = x$ for $x \in B_P$. Thus, π_P is strictly coisometric and, a fortiori, coisometric.

Proposition 2.6. *Let P be a normed space and $C \ge 1$. Then*

(i) if P is C-topologically projective, then π_P has a right inverse bounded linear operator of norm at most C;

(ii) if π_P has a right inverse bounded linear operator of norm at most C, then P is $C(1+\varepsilon)$-topologically projective for every $\varepsilon > 0$.

Proof. (*i*) Lift 1_P along the 1-topologically surjective operator π_P: we get $\sigma : P \to \ell_1^0(B_P)$ with $\pi_P\sigma = 1_P$ and $\|\sigma\| \le C$.

(*ii*) If $\pi_P\sigma = 1_P$ with $\|\sigma\| \le C$, then P is a C-retract of $\ell_1^0(B_P)$, which is $(1+\varepsilon)$-topologically projective by proposition 2.2. It remains to apply proposition 2.5. □

The equivalence of conditions (i) and (iii) in the following theorem is the main observation of [3, section 2] on extremely projective normed spaces: a normed space is extremely projective if and only if it is a near-retract of a free normed space, where a near-retract is a retract via a contractive operator which has right inverses of norm $< 1+\varepsilon$ for every $\varepsilon > 0$.

Theorem 2.7. *For a normed space P the following conditions are equivalent:*

(i) P is extremely projective;

(ii) for every $\varepsilon > 0$ the operator π_P has a right inverse bounded linear operator of norm at most $1+\varepsilon$;

(iii) for every $\varepsilon > 0$ there is a set Λ such that P is a $(1+\varepsilon)$-retract of $\ell_1^0(\Lambda)$;

(iv) for every $\varepsilon > 0$ there are a set Λ, a subspace $Z \subset \ell_1^0(\Lambda)$ and a projection $Q : \ell_1^0(\Lambda) \to \ell_1^0(\Lambda)$ onto Z with $\|Q\| \leq 1+\varepsilon$ such that $d_{BM}(P, Z) \leq 1+\varepsilon$.

Proof. $(i) \implies (ii)$ is proposition 2.6 (i), and $(ii) \implies (iii)$ is obvious.

$(iii) \implies (iv)$ Let $\zeta : \ell_1^0(\Lambda) \to P$ and $\eta : P \to \ell_1^0(\Lambda)$ satisfy $\zeta\eta = 1_P$ and $\|\zeta\|\|\eta\| \leq 1+\varepsilon$. Replacing ζ by $\|\zeta\|^{-1}\zeta$ and η by $\|\zeta\|\eta$ we may assume that $\|\zeta\| = 1$ and $\|\eta\| \leq 1+\varepsilon$. Put $Z = \eta(P)$ and $Q = \eta\zeta$. Then $Q^2 = \eta(\zeta\eta)\zeta = Q$, $\mathrm{Im}(Q) = Z$ and $\|Q\| \leq 1+\varepsilon$. The operator $\eta : P \to Z$ is a bijection with $\|\eta\| \leq 1+\varepsilon$ and $\eta^{-1} = \zeta|_Z$, so $\|\eta^{-1}\| \leq 1$ and $d_{BM}(P, Z) \leq 1+\varepsilon$.

$(iv) \implies (i)$ Since $d_{BM}(P, Z) \leq 1+\varepsilon$, there is a linear bijection $T : P \to Z$ with $\|T\|\|T^{-1}\| \leq 1+2\varepsilon$. Put $\eta = T$ and $\zeta = T^{-1}Q : \ell_1^0(\Lambda) \to P$. Then $\zeta\eta = T^{-1}QT = T^{-1}T = 1_P$ and $\|\zeta\|\|\eta\| \leq (1+2\varepsilon)(1+\varepsilon)$. By propositions 2.2 and 2.5 the space P is $(1+2\varepsilon)(1+\varepsilon)^2$-topologically projective. As $\varepsilon > 0$ is arbitrary, P is extremely projective. □

3 Dimension restrictions on topologically projective normed spaces

The free normed spaces $\ell_1^0(\Lambda)$ are far from being complete: for an infinite set Λ the Hamel basis $(e_\lambda)_{\lambda\in\Lambda}$ of $\ell_1^0(\Lambda)$ has the same cardinality as a dense subset of $\ell_1^0(\Lambda)$ of minimal cardinality, whereas the algebraic dimension of an infinite dimensional Banach space is uncountable by the Baire category theorem, and in fact at least continuum [12]. In this section we show that this smallness is inherited by all topologically projective normed spaces. Helemskii proved that a separable topologically projective normed space has at most countable algebraic dimension [3, proposition 2.4], and deduced that $\ell_1(\Lambda)$ is not topologically projective in **Nor** for any infinite set Λ [3, proposition 2.5]. His argument uses only the existence of a bounded injective operator into a free normed space, and we state it in this form.

Lemma 3.1. *Let Z be a normed space, S be a set and $\rho : Z \to \ell_1^0(S)$ be an injective bounded linear operator. Then the algebraic dimension of Z is at most* $\max(\mathrm{dens}(Z), \aleph_0)$, *where* $\mathrm{dens}(Z)$ *is the density character of Z, that is, the smallest cardinality of a dense subset of Z.*

Proof. Let M be a dense subset of Z of cardinality $\mathrm{dens}(Z)$ and $S_0 = \bigcup_{x\in M} \mathrm{supp}(\rho(x))$, a set of cardinality at most $\max(\mathrm{dens}(Z), \aleph_0)$. For $z \in Z$ the vector $\rho(z)$ is the limit in $\ell_1(S)$ of a sequence of elements of $\rho(M)$, which are supported in S_0, so $\rho(z)$ is supported in S_0 by the continuity of the coordinate functionals. Thus, ρ is an injective linear map of Z into $\ell_1^0(S_0)$, whose algebraic dimension is the cardinality of S_0. □

Theorem 3.2. *Let P be a topologically projective normed space and Z be a subspace of P. Then the algebraic dimension of Z is at most $\max(\operatorname{dens}(Z), \aleph_0)$. In particular, every separable subspace of P has at most countable algebraic dimension.*

Proof. By proposition 2.6 (i) there exists a bounded linear operator $\sigma : P \to \ell_1^0(B_P)$ with $\pi_P \sigma = 1_P$; in particular, σ is injective. It remains to apply lemma 3.1 to $\sigma|_Z$. □

Corollary 3.3. *Let P be a topologically projective normed space, Z be a subspace of P and $\|\cdot\|'$ be a norm on Z such that $(Z, \|\cdot\|')$ is complete and $\|z\| \le K\|z\|'$ for all $z \in Z$ and some constant $K > 0$. Then Z is finite dimensional. In particular, every complete subspace of P is finite dimensional, and no infinite dimensional Banach space is topologically projective in* **Nor** *(for the spaces $\ell_1(\Lambda)$ this is [3, proposition 2.5]).*

Proof. Assume that Z is infinite dimensional. Choose a linearly independent sequence $(z_n)_{n\in\mathbb{N}}$ in Z and let Z_0 be the closure of its linear span in $(Z, \|\cdot\|')$. Then $(Z_0, \|\cdot\|')$ is a separable infinite dimensional Banach space, so its algebraic dimension is uncountable. On the other hand, Z_0 is separable with respect to $\|\cdot\|$ because $\|\cdot\| \le K\|\cdot\|'$, so by theorem 3.2 the algebraic dimension of Z_0 is countable. This contradiction proves the first claim; the rest follows by taking $Z = P$ and $\|\cdot\|' = \|\cdot\|$. □

Let us compare theorem 3.2 with the theorem of Grønbæk [8, theorem 2.1]: a normed space E is topologically projective if and only if it is topologically isomorphic to $\ell_1^0(\Lambda)$, where the cardinality of Λ is the density character of E. Since for infinite Λ the algebraic dimension of $\ell_1^0(\Lambda)$ equals the cardinality of Λ, the algebraic dimension of an infinite dimensional topologically projective normed space coincides with its density character. Theorem 3.2 is a hereditary version of this equality: it applies to all subspaces of P, which are, of course, not topologically projective in general.

The following example shows that corollary 3.3 is sharper than its last claim: a normed space may have no infinite dimensional complete subspaces and still fail to be topologically projective. It also shows that topological projectivity of a normed space is not determined by its completion.

Example 3.4. *Let $(w_n)_{n\in\mathbb{N}}$ be a sequence of real numbers with $w_n \ge 1$ and $\lim_n w_n = +\infty$, and let*

$$P_w = \Big\{ x \in \ell_1 : \sum_{n=1}^{\infty} w_n |x_n| < +\infty \Big\}$$

with the norm inherited from ℓ_1. Then P_w is a dense subspace of ℓ_1 containing $\ell_1^0(\mathbb{N})$, so its completion is ℓ_1, and

(i) every complete subspace of P_w is finite dimensional;

(ii) P_w is not topologically projective.

Proof. Denote $\|x\|_w = \sum_{n=1}^{\infty} w_n |x_n|$ for $x \in P_w$. Then $(P_w, \|\cdot\|_w)$ is the Banach space $\ell_1(w)$ of w-weighted summable sequences and $\|x\| \le \|x\|_w$ for $x \in P_w$.

(i) Let Z be a complete subspace of P_w. If $(z_k)_{k\in\mathbb{N}}$ is a sequence in Z converging to $z \in P_w$ with respect to $\|\cdot\|_w$, then it converges to z in ℓ_1, so $z \in Z$. Thus, Z is closed in $(P_w, \|\cdot\|_w)$ and the two norms $\|\cdot\|$ and $\|\cdot\|_w$ are complete on Z. Since $\|\cdot\| \le \|\cdot\|_w$, by the open mapping theorem there is a constant K with $\|z\|_w \le K\|z\|$ for all $z \in Z$. Assume that Z is infinite dimensional and fix $N \in \mathbb{N}$. The linear map $Z \to \mathbb{C}^N$, $z \mapsto (z_1, \dots, z_N)$ has a non-zero kernel, so there exists $z \in Z$ with $\|z\| = 1$ and $z_n = 0$ for $n \le N$. Then $K \ge \|z\|_w = \sum_{n>N} w_n |z_n| \ge \inf_{n>N} w_n$. Since N is arbitrary and $\lim_n w_n = +\infty$, we get a contradiction.

(*ii*) Apply corollary 3.3 to $Z = P_w$ and $\|\cdot\|' = \|\cdot\|_w$. Alternatively, note that P_w is separable, being dense in ℓ_1, while its algebraic dimension is uncountable, since $(P_w, \|\cdot\|_w)$ is an infinite dimensional Banach space; so theorem 3.2 applies directly. □

4 Extremely projective normed spaces of countable dimension

We start with a necessary condition which holds in all dimensions. Recall that the completion $\overline{P}$ of a normed space P is a Banach space containing P as a dense subspace. Every bounded linear operator between normed spaces extends uniquely to a bounded linear operator between their completions with the same norm, and these extensions respect compositions. The following theorem is [3, proposition 2.3]; see also [8, section 3].

Theorem 4.1. *Let P be an extremely projective normed space. Then $\overline{P}$ is isometrically isomorphic to $\ell_1(\Gamma)$ for some set Γ.*

Proof. We only indicate the argument. By theorem 2.7 for every $\varepsilon > 0$ the space $\overline{P}$ is a $(1+\varepsilon)$-retract of the Banach space $\ell_1(B_P)$, so by Grothendieck's theorem [6] in the form given in [3, section 2] (see also [2, proposition 3.2]) $\overline{P}$ is isometrically isomorphic to $\ell_1(\Gamma)$ for some set Γ. Explicitly, let $\overline{\pi} : \ell_1(B_P) \to \overline{P}$ be the extension of π_P and $\overline{\sigma}_\varepsilon$ be the extensions of right inverses of π_P of norm at most $1+\varepsilon$. Taking adjoints one sees that $\overline{\pi}^*$ is an isometric weak*-continuous embedding of $\overline{P}^*$ into $\ell_\infty(B_P)$ with weak*-closed range and that $\overline{\pi}^*\overline{\sigma}_\varepsilon{}^*$ are projections onto this range of norm at most $1+\varepsilon$; a limit point of these projections in the weak* operator topology is a projection of norm 1, so $\overline{P}^*$ is a 1-complemented subspace of the 1-injective space $\ell_\infty(B_P)$. Hence $\overline{P}^*$ is 1-injective, hence isometrically isomorphic to $C(K)$ for a Stonean space K [7, chapter 3, §11, theorem 6]; a Banach space whose dual is 1-injective is isometrically isomorphic to some $L_1(\Omega, \mu)$ [7, chapter 7]; and $\overline{P}$ has the Schur property as a subspace of $\ell_1(B_P)$, which forces μ to be purely atomic. □

Theorems 3.2 and 4.1 describe extremely projective normed spaces from two opposite sides: they are dense subspaces of the spaces $\ell_1(\Gamma)$ which are small enough to have separable subspaces of countable algebraic dimension only. In this section we show that for spaces of countable dimension there is nothing in between: every dense subspace of ℓ_1 of countable algebraic dimension is extremely projective, and even almost isometric to $\ell_1^0(\mathbb{N})$. This answers the question of Helemskii [3, section 2] on dense subspaces of $\ell_1(\Lambda)$ for countable Λ. The proof is a back-and-forth construction based on the following three lemmas. For a sequence $x \in \ell_1$ and $N \in \mathbb{N} \cup \{0\}$ we denote by $x|_N$ the sequence which coincides with x at $n \le N$ and vanishes at $n > N$.

Lemma 4.2. *Let $v \in \ell_1$, $N \in \mathbb{N} \cup \{0\}$ and $0 < \delta < 1$. Then there exists a bounded linear operator $R : \ell_1 \to \ell_1$ with $\|R\| \le \delta$ such that*

(i) $R(x) = 0$ *for every* $x \in \ell_1$ *with* $x = x|_N$;

(ii) $R(\ell_1^0(\mathbb{N})) \subset \ell_1^0(\mathbb{N})$;

(iii) $(1_{\ell_1} - R)(v) \in \ell_1^0(\mathbb{N})$;

(iv) $(1_{\ell_1} - R)^{-1}(\ell_1^0(\mathbb{N})) \subset \ell_1^0(\mathbb{N}) + \mathbb{C}v$.

In particular, $T = 1_{\ell_1} - R$ is a linear bijection of ℓ_1 with $\|T\| \le 1+\delta$ and $\|T^{-1}\| \le (1-\delta)^{-1}$ such that $T(x) = x$ whenever $x = x|_N$, $T(\ell_1^0(\mathbb{N})) \subset \ell_1^0(\mathbb{N})$ and $T(\ell_1^0(\mathbb{N}) + \mathbb{C}v) = \ell_1^0(\mathbb{N})$.

Proof. Since $\|R\| \le \delta < 1$, the operator $T = 1_{\ell_1} - R$ is a linear bijection with $T^{-1} = \sum_{m=0}^{\infty} R^m$, $\|T\| \le 1+\delta$ and $\|T^{-1}\| \le (1-\delta)^{-1}$, and the properties of T listed in the last sentence follow from (i)–(iv). So we only need to construct R with the properties (i)–(iv).

If $v - v|_N$ is finitely supported, take $R = 0$. Assume that $v - v|_N$ has infinite support. For $M \ge N$ put $t_M = \|v - v|_M\| = \sum_{n>M} |v_n|$. Then the sequence $(t_M)_{M\ge N}$ is non-increasing, $t_M > 0$ for all $M \ge N$ because $v - v|_N$ has infinite support, and $\lim_M t_M = 0$. Choose integers $N = N_0 < N_1 < N_2 < \dots$ such that $t_{N_k} \le (\delta/2) t_{N_{k-1}}$ for all $k \in \mathbb{N}$. The sets $B_k = \{n \in \mathbb{N} : N_{k-1} < n \le N_k\}$, $k \in \mathbb{N}$, are finite, pairwise disjoint and cover $\{n \in \mathbb{N} : n > N\}$. Put $w_k = v|_{N_k} - v|_{N_{k-1}}$, that is, w_k is the restriction of v to B_k. Then $w_k \in \ell_1^0(\mathbb{N})$, $v - v|_N = \sum_{k=1}^{\infty} w_k$ in ℓ_1, $\operatorname{supp}(w_k) \subset B_k$, and

$$\|w_k\| = t_{N_{k-1}} - t_{N_k} \ge (1-\delta/2) t_{N_{k-1}} > 0, \qquad \|w_{k+1}\| \le t_{N_k} \le (\delta/2) t_{N_{k-1}} \le \frac{\delta/2}{1-\delta/2} \|w_k\| \le \delta \|w_k\|,$$

where the last inequality holds because $\delta \le 1$. For $k \in \mathbb{N}$ define the functional

$$g_k : \ell_1 \to \mathbb{C}, \quad g_k(x) = \sum_{n \in B_k} \overline{\operatorname{sgn}(v_n)}\, x_n,$$

where $\operatorname{sgn}(z) = z/|z|$ for $z \ne 0$ and $\operatorname{sgn}(0) = 0$. Then $|g_k(x)| \le \sum_{n\in B_k} |x_n|$ for all $x \in \ell_1$, $g_k(w_k) = \|w_k\|$ and $g_k(w_j) = 0$ for $j \ne k$, because the supports of w_j and g_k are disjoint. Now define

$$R : \ell_1 \to \ell_1, \quad R(x) = \sum_{k=1}^{\infty} \frac{g_k(x)}{\|w_k\|}\, w_{k+1}.$$

The series converges absolutely, since

$$\sum_{k=1}^{\infty} \frac{|g_k(x)|}{\|w_k\|} \|w_{k+1}\| \le \delta \sum_{k=1}^{\infty} \sum_{n\in B_k} |x_n| \le \delta \|x\|,$$

so R is a well defined bounded linear operator with $\|R\| \le \delta$. We verify the properties (i)–(iv).

(i) If $x = x|_N$, then $g_k(x) = 0$ for all k, because $\operatorname{supp}(g_k) \subset B_k \subset \{n : n > N\}$. Hence $R(x) = 0$.

(ii) If $x \in \ell_1^0(\mathbb{N})$, then $g_k(x) \ne 0$ only for the finitely many k with $B_k \cap \operatorname{supp}(x) \ne \varnothing$, so $R(x)$ is a finite linear combination of the finitely supported sequences w_{k+1}.

(iii) Since $g_k(v|_N) = 0$ and $g_k(v - v|_N) = \sum_{j=1}^{\infty} g_k(w_j) = g_k(w_k) = \|w_k\|$, we get $R(v) = \sum_{k=1}^{\infty} w_{k+1} = v - v|_N - w_1$, so $(1_{\ell_1} - R)(v) = v|_N + w_1 \in \ell_1^0(\mathbb{N})$.

(iv) Let $x \in \ell_1^0(\mathbb{N})$ and let K be the finite set of those k for which $g_k(x) \ne 0$. Put $c_k = g_k(x)/\|w_k\|$ for $k \in K$. Since $g_j(w_k) = \delta_{jk}\|w_k\|$, we have $R(w_k) = w_{k+1}$ for all k, hence $R^m(x) = \sum_{k\in K} c_k w_{k+m}$ for $m \ge 1$ and

$$(1_{\ell_1} - R)^{-1}(x) = \sum_{m=0}^{\infty} R^m(x) = x + \sum_{k\in K} c_k \sum_{m=1}^{\infty} w_{k+m} = x + \sum_{k\in K} c_k (v - v|_{N_k}) \in \ell_1^0(\mathbb{N}) + \mathbb{C}v.$$

□

Lemma 4.3. *Let Y be a finite dimensional subspace of $\ell_1^0(\mathbb{N})$ and $n \in \mathbb{N}$ be such that $e_n \notin Y$. Then there is a constant $C \geq 1$ such that for every $z \in \ell_1^0(\mathbb{N})$ with $\|z - e_n\| \leq (4C)^{-1}$ there exists a linear bijection $S : \ell_1 \to \ell_1$ with $\|S - 1_{\ell_1}\| \leq 2C\|z - e_n\| \leq 1/2$, $S(y) = y$ for all $y \in Y$ and $S(z) = e_n$.*

Proof. Since Y is finite dimensional, it is closed in ℓ_1, so $d = \inf\{\|e_n - y\| : y \in Y\} > 0$. By the Hahn–Banach theorem there is $f \in \ell_1^*$ with $f(y) = 0$ for $y \in Y$, $f(e_n) = 1$ and $\|f\| = d^{-1}$. Put $C = \max(1, d^{-1})$. Let $z \in \ell_1^0(\mathbb{N})$ satisfy $\|z - e_n\| \leq (4C)^{-1}$. Then $|f(z) - 1| = |f(z - e_n)| \leq C\|z - e_n\| \leq 1/4$, so $|f(z)| \geq 3/4$. Define

$$S : \ell_1 \to \ell_1, \quad S(x) = x + \frac{f(x)}{f(z)}(e_n - z).$$

Then $S(y) = y$ for $y \in Y$ because f vanishes on Y, and $S(z) = z + (e_n - z) = e_n$. Moreover, $\|S - 1_{\ell_1}\| \leq \|f\||f(z)|^{-1}\|e_n - z\| \leq \frac{4}{3}C\|z - e_n\| \leq 2C\|z - e_n\| \leq 1/2$, so S is a linear bijection of ℓ_1 by the Neumann series. □

Lemma 4.4. *Let X be a Banach space and $(F_m)_{m\in\mathbb{N}}$ be a sequence of bounded linear operators on X such that $\theta_m = \|F_m - 1_X\| < 1$ for all m and $\sum_{m=1}^{\infty} \theta_m < +\infty$. Put $A_j = F_j F_{j-1} \cdots F_1$ for $j \in \mathbb{N}$. Then the sequence $(A_j)_{j\in\mathbb{N}}$ converges in the operator norm to a linear bijection T of X with*

$$\|T\| \leq \prod_{m=1}^{\infty}(1 + \theta_m), \qquad \|T^{-1}\| \leq \prod_{m=1}^{\infty}(1 - \theta_m)^{-1}.$$

Proof. Both products converge because $\sum_m \theta_m < +\infty$; denote them by M and M'. Then $\|A_j\| \leq M$ and, since each F_m is invertible with $\|F_m^{-1}\| \leq (1 - \theta_m)^{-1}$ by the Neumann series, $\|A_j^{-1}\| \leq M'$ for all j. From $\|A_j - A_{j-1}\| = \|(F_j - 1_X)A_{j-1}\| \leq \theta_j M$ we see that $(A_j)_j$ is a Cauchy sequence; let T be its limit, so $\|T\| \leq M$. Also $\|A_j^{-1} - A_k^{-1}\| = \|A_j^{-1}(A_k - A_j)A_k^{-1}\| \leq M'^2\|A_k - A_j\|$, so $(A_j^{-1})_j$ is a Cauchy sequence with a limit B, $\|B\| \leq M'$. Passing to the limit in $A_j A_j^{-1} = A_j^{-1} A_j = 1_X$ we get $TB = BT = 1_X$. □

Theorem 4.5. *Let D be a dense subspace of ℓ_1 of countable algebraic dimension and $\varepsilon > 0$. Then there exists a linear bijection $T : \ell_1 \to \ell_1$ with $\|T\|\|T^{-1}\| \leq 1 + \varepsilon$ and $T(D) = \ell_1^0(\mathbb{N})$. In particular, $d_{BM}(D, \ell_1^0(\mathbb{N})) = 1$.*

Proof. Since ℓ_1 is infinite dimensional and D is dense in ℓ_1, the algebraic dimension of D is countably infinite. Let $(d_j)_{j\in\mathbb{N}}$ be a Hamel basis of D. Choose a sequence $(\theta_m)_{m\in\mathbb{N}}$ of numbers in $(0, 1/2]$ with $\prod_{m=1}^{\infty}(1 + \theta_m)(1 - \theta_m)^{-1} \leq 1 + \varepsilon$; then $\sum_m \theta_m < +\infty$, since $(1 + \theta)/(1 - \theta) \geq 1 + 2\theta$.

We construct by induction linear bijections $F_1, F_2, \ldots$ of ℓ_1 with $\|F_m - 1_{\ell_1}\| \leq \theta_m$ and finite dimensional subspaces $Y_0 = \{0\} \subset Y_1 \subset Y_2 \subset \ldots$ of $\ell_1^0(\mathbb{N})$ such that, with $A_0 = 1_{\ell_1}$ and $A_j = F_{2j}F_{2j-1}A_{j-1}$, for every $j \in \mathbb{N}$ the following conditions hold:

(a) $A_j(d_i) \in Y_j$ for $i \leq j$;

(b) $e_i \in Y_j$ for $i \leq j$;

(c) $Y_j \subset A_j(D)$;

(d) F_{2j-1} and F_{2j} act as the identity on Y_{j-1}.

Suppose that A_{j-1} and Y_{j-1} have been constructed (for $j = 1$ there is nothing to suppose). Let N be an integer not smaller than every element of the supports of the vectors of some basis of Y_{j-1}, so that $y = y|_N$ for all $y \in Y_{j-1}$. Apply lemma 4.2 to the vector $v = A_{j-1}(d_j)$, the integer N and $\delta = \theta_{2j-1}$, and let $F_{2j-1} = 1_{\ell_1} - R$ be the resulting operator. Then $\|F_{2j-1} - 1_{\ell_1}\| \le \theta_{2j-1}$, F_{2j-1} acts as the identity on Y_{j-1}, and $y_j = F_{2j-1}A_{j-1}(d_j) \in \ell_1^0(\mathbb{N})$. Put $A' = F_{2j-1}A_{j-1}$ and $Y' = Y_{j-1} + \mathbb{C}y_j \subset \ell_1^0(\mathbb{N})$. Note that $Y' \subset A'(D)$: indeed, $Y_{j-1} \subset A_{j-1}(D)$ by the induction hypothesis (for $j = 1$ trivially), F_{2j-1} fixes Y_{j-1}, so $Y_{j-1} = F_{2j-1}(Y_{j-1}) \subset A'(D)$, and $y_j = A'(d_j) \in A'(D)$.

If $e_j \in Y'$, put $F_{2j} = 1_{\ell_1}$ and $Y_j = Y'$. Otherwise, let C be the constant given by lemma 4.3 for Y' and $n = j$, and let N' be an integer not smaller than j and than every element of the supports of the vectors of some basis of Y'. Since A' is a linear bijection of ℓ_1 and D is dense, $A'(D)$ is dense in ℓ_1, so there is $d \in D$ with

$$\|A'(d) - e_j\| \le \frac{\theta_{2j}}{16C}.$$

Apply lemma 4.2 to the vector $A'(d)$, the integer N' and $\delta = \theta_{2j}/2$, and let $G = 1_{\ell_1} - R'$ be the resulting operator. Then G fixes Y' and e_j, $\|G - 1_{\ell_1}\| \le \theta_{2j}/2$, $\|G\| \le 5/4$ because $\theta_{2j} \le 1/2$, and $z = GA'(d) \in \ell_1^0(\mathbb{N})$ satisfies

$$\|z - e_j\| = \|G(A'(d) - e_j)\| \le \frac{5\theta_{2j}}{64C} \le \frac{1}{4C}.$$

Let S be the operator given by lemma 4.3 for Y', $n = j$ and z. Then S fixes Y', $S(z) = e_j$ and $\|S - 1_{\ell_1}\| \le 2C\|z - e_j\| \le 5\theta_{2j}/32$. Put $F_{2j} = SG$ and $Y_j = Y' + \mathbb{C}e_j$. Then

$$\|F_{2j} - 1_{\ell_1}\| \le \|(S - 1_{\ell_1})G\| + \|G - 1_{\ell_1}\| \le \frac{5\theta_{2j}}{32} \cdot \frac{5}{4} + \frac{\theta_{2j}}{2} \le \theta_{2j}.$$

Let us check (a)–(d) for j. The operator F_{2j} acts as the identity on $Y' \supset Y_{j-1}$, which together with the first step gives (d). For $i < j$ we have $A_{j-1}(d_i) \in Y_{j-1}$, so $A_j(d_i) = F_{2j}F_{2j-1}A_{j-1}(d_i) = A_{j-1}(d_i) \in Y_j$, and $A_j(d_j) = F_{2j}(y_j) = y_j \in Y_j$; this is (a). Condition (b) holds by the definition of Y_j and the induction hypothesis. Finally, $Y' \subset A'(D)$ and F_{2j} fixes Y', so $Y' \subset F_{2j}A'(D) = A_j(D)$, and $e_j = SGA'(d) = A_j(d) \in A_j(D)$ (in the case $e_j \in Y'$ there is nothing to add); this is (c).

By lemma 4.4 the operators A_j converge in norm to a linear bijection T of ℓ_1 with $\|T\|\|T^{-1}\| \le \prod_m (1+\theta_m)(1-\theta_m)^{-1} \le 1+\varepsilon$. For every $i \in \mathbb{N}$ and $j \ge i$ we have $A_j(d_i) \in Y_j$ by (a), and by (d) the operators $F_{2j+1}, F_{2j+2}, \dots$ fix Y_j, so $A_k(d_i) = A_j(d_i)$ for all $k \ge j$ and $T(d_i) = A_i(d_i) \in Y_i \subset \ell_1^0(\mathbb{N})$. Hence $T(D) \subset \ell_1^0(\mathbb{N})$. Similarly, for every $i \in \mathbb{N}$ by (b) and (c) there is $d \in D$ with $A_i(d) = e_i \in Y_i$, and by (d) we get $T(d) = A_i(d) = e_i$. Hence $\ell_1^0(\mathbb{N}) \subset T(D)$. □

Now we are ready to prove the main result of this section.

Theorem 4.6. *Let P be a normed space of countable (finite or countably infinite) algebraic dimension. Then the following conditions are equivalent:*

(i) P is extremely projective;

(ii) $\overline{P}$ is isometrically isomorphic to $\ell_1(\Gamma)$ for some countable set Γ;

(iii) $d_{BM}(P, \ell_1^0(\Gamma)) = 1$ for some countable set Γ;

(iv) for every $\varepsilon > 0$ there is a set Λ with $d_{BM}(P, \ell_1^0(\Lambda)) < 1 + \varepsilon$.

Proof. $(i) \implies (ii)$ By theorem 4.1 the space $\overline{P}$ is isometrically isomorphic to $\ell_1(\Gamma)$ for some set Γ. The rational linear span of a countable Hamel basis of P is a countable dense subset of $\overline{P}$, so $\overline{P}$ is separable and Γ is countable.

$(ii) \implies (iii)$ Let $J : \overline{P} \to \ell_1(\Gamma)$ be an isometric isomorphism. If Γ is finite, then $\overline{P}$ is finite dimensional, so $P = \overline{P}$ and J maps P onto $\ell_1(\Gamma) = \ell_1^0(\Gamma)$. If Γ is infinite, we may assume that $\Gamma = \mathbb{N}$. Then $D = J(P)$ is a dense subspace of ℓ_1 of countable algebraic dimension, and by theorem 4.5 $d_{BM}(P, \ell_1^0(\mathbb{N})) = d_{BM}(D, \ell_1^0(\mathbb{N})) = 1$.

$(iii) \implies (iv)$ is obvious.

$(iv) \implies (i)$ Fix $\varepsilon > 0$ and a linear bijection $S : P \to \ell_1^0(\Lambda)$ with $\|S\|\|S^{-1}\| < 1 + \varepsilon$. Then P is a $(1+\varepsilon)$-retract of $\ell_1^0(\Lambda)$ via S^{-1} and S, so P is extremely projective by theorem 2.7. □

Corollary 4.7. *Every subspace of ℓ_1 of countable algebraic dimension which is dense in ℓ_1 is extremely projective and almost isometric to $\ell_1^0(\mathbb{N})$, that is, has Banach–Mazur distance 1 to $\ell_1^0(\mathbb{N})$.*

Corollary 4.7 may be compared with the results of Hájek and Russo on densely isomorphic normed spaces [9]: by [9, theorem 2.4] every dense subspace of $\ell_1(\Gamma)$ contains a dense subspace topologically isomorphic to $\ell_1^0(\Gamma)$. Corollary 4.7 says that for dense subspaces of ℓ_1 of countable algebraic dimension the subspace itself is almost isometric to $\ell_1^0(\mathbb{N})$.

5 Extremely projective spaces which are not free

In this section we show that extreme projectivity is not an isometric invariant of the form "P is isometrically isomorphic to $\ell_1^0(\Lambda)$". By corollary 4.7, for every subspace V of ℓ_1 of countable algebraic dimension the space $\ell_1^0(\mathbb{N}) + V$ is extremely projective. To see that such spaces are not isometrically isomorphic to free normed spaces whenever $V \not\subset \ell_1^0(\mathbb{N})$, we look at extreme points of unit balls. By $\operatorname{ext}(B)$ we denote the set of extreme points of a convex set B and by $\operatorname{conv}(M)$ the convex hull of a set M.

Lemma 5.1. *Let S be a set and E be a subspace of $\ell_1(S)$ containing $\ell_1^0(S)$, endowed with the norm of $\ell_1(S)$. Then $\operatorname{ext}(B_E) = \{\lambda e_s : s \in S,\ \lambda \in \mathbb{C},\ |\lambda| = 1\}$ and $\operatorname{conv}(\operatorname{ext}(B_E)) = B_{\ell_1^0(S)}$.*

Proof. The points λe_s with $|\lambda| = 1$ are extreme points of $B_{\ell_1(S)}$, and $B_E = B_{\ell_1(S)} \cap E$, so they are extreme points of B_E. Conversely, let $x \in B_E$ be not of this form. If $x = 0$, then $x = \frac{1}{2}e_s + \frac{1}{2}(-e_s)$ for any $s \in S$. If $0 < \|x\| < 1$, then $x = \frac{1}{2}(1+t)x + \frac{1}{2}(1-t)x$ for $t = 1 - \|x\| > 0$, and both $(1 \pm t)x$ belong to B_E. If $\|x\| = 1$, then x has at least two non-zero coordinates x_n and x_m, $n \neq m$. Put $t = \min(|x_n|, |x_m|) > 0$ and $u = t(\operatorname{sgn}(x_n)e_n - \operatorname{sgn}(x_m)e_m) \in \ell_1^0(S) \subset E$. Then $x \pm u \in E$ and

$$\|x \pm u\| = (|x_n| \pm t) + (|x_m| \mp t) + \sum_{s \neq n,m} |x_s| = \|x\| = 1,$$

so $x = \frac{1}{2}(x+u) + \frac{1}{2}(x-u)$ with $u \neq 0$. In all three cases x is not an extreme point of B_E.

Every finite convex combination of the points λe_s is a finitely supported element of $B_{\ell_1(S)}$, so $\operatorname{conv}(\operatorname{ext}(B_E)) \subset B_{\ell_1^0(S)}$. Conversely, if $x \in B_{\ell_1^0(S)}$, then

$$x = \sum_{s \in \operatorname{supp}(x)} |x_s| \operatorname{sgn}(x_s) e_s + (1 - \|x\|) \cdot 0,$$

and $0 = \frac{1}{2}e_s + \frac{1}{2}(-e_s)$, so $x \in \operatorname{conv}(\operatorname{ext}(B_E))$. □

Theorem 5.2. *Let S be a set and E be a subspace of $\ell_1(S)$ containing $\ell_1^0(S)$, endowed with the norm of $\ell_1(S)$. If E is isometrically isomorphic to $\ell_1^0(\Lambda)$ for some set Λ, then $E = \ell_1^0(S)$.*

Proof. Let $T : \ell_1^0(\Lambda) \to E$ be an isometric isomorphism. Then $T(B_{\ell_1^0(\Lambda)}) = B_E$ and T maps $\operatorname{ext}(B_{\ell_1^0(\Lambda)})$ onto $\operatorname{ext}(B_E)$. Applying lemma 5.1 to E and to $\ell_1^0(\Lambda)$ itself we get

$$B_E = T(B_{\ell_1^0(\Lambda)}) = T(\operatorname{conv}(\operatorname{ext}(B_{\ell_1^0(\Lambda)}))) = \operatorname{conv}(T(\operatorname{ext}(B_{\ell_1^0(\Lambda)}))) = \operatorname{conv}(\operatorname{ext}(B_E)) = B_{\ell_1^0(S)}.$$

Hence $E = \ell_1^0(S)$. □

Corollary 5.3. *Let V be a subspace of ℓ_1 of countable algebraic dimension with $V \not\subset \ell_1^0(\mathbb{N})$, for example $V = \mathbb{C}v$ for $v = (2^{-n})_{n\in\mathbb{N}}$. Then $\ell_1^0(\mathbb{N}) + V$ is an extremely projective normed space which is not isometrically isomorphic to $\ell_1^0(\Lambda)$ for any set Λ.*

Proof. The space $\ell_1^0(\mathbb{N}) + V$ is dense in ℓ_1 and has countable algebraic dimension, so it is extremely projective by corollary 4.7, and it is not isometrically isomorphic to any $\ell_1^0(\Lambda)$ by theorem 5.2. □

Corollary 5.3 answers the question of Helemskii from [3, section 2] and [2, remark 3.6] in the negative and disproves [8, conjecture 3.1]. Note that by theorem 5.2 every subspace E with $\ell_1^0(\mathbb{N}) \subsetneq E \subset \ell_1$ fails to be isometrically isomorphic to a free normed space, but by example 3.4 not all of these subspaces are extremely projective. Thus, between $\ell_1^0(\mathbb{N})$ and ℓ_1 there are both extremely projective and non-projective normed spaces, and the difference between them is invisible on the level of completions. The next section provides a criterion which distinguishes them.

6 Extremely projective normed spaces of uncountable dimension

Condition (ii) of theorem 4.6 shows that for normed spaces of countable dimension extreme projectivity is a property of the completion. By example 3.4 this is no longer true for spaces of uncountable dimension: the dimension restriction of theorem 3.2 is an independent necessary condition.

By theorems 3.2 and 4.1, an extremely projective normed space is, up to an isometric isomorphism, a dense subspace of some $\ell_1(\Gamma)$ all of whose separable subspaces have countable algebraic dimension; by the theorem of Grønbæk it is also topologically isomorphic to $\ell_1^0(\Gamma)$. For countable Γ the first two conditions are also sufficient by theorem 4.6. In section 7 we shall see that for uncountable Γ they are not: the chain of the subspaces $P \cap \ell_1(\delta)$, $\delta < \omega_1$, of a topologically projective dense subspace P of $\ell_1(\omega_1)$ must be continuous at almost all limit stages, and ladder systems on ω_1 produce dense subspaces for which it is not. In this section we prove the positive results: a description of the topologically projective dense subspaces of $\ell_1(\Gamma)$ by filtrations with countable blocks, and the almost isometric characterization of extremely projective normed spaces of arbitrary dimension. The main tool is a transfinite version of theorem 4.5. Its key point is that the perturbations used at different steps are controlled on pairwise disjoint blocks of coordinates, so that their "product" has norm bounded independently of the length of the construction. By a limit ordinal we always mean a non-zero ordinal which is not a successor.

Let Γ be a set, λ be an ordinal and $(\Gamma_\delta)_{\delta\le\lambda}$ be an increasing family of subsets of Γ such that $\Gamma_0 = \varnothing$, $\Gamma_\lambda = \Gamma$ and $\Gamma_\delta = \bigcup_{\delta'<\delta} \Gamma_{\delta'}$ for every limit ordinal $\delta \le \lambda$. We call such a family a filtration of Γ and put $N_\delta = \Gamma_{\delta+1} \setminus \Gamma_\delta$ for $\delta < \lambda$. Then $\Gamma_\delta = \bigsqcup_{\delta'<\delta} N_{\delta'}$ for all $\delta \le \lambda$, and $\ell_1(\Gamma_\delta)$ is the ℓ_1-sum of the spaces $\ell_1(N_{\delta'})$, $\delta' < \delta$. For $x \in \ell_1(\Gamma)$ and $A \subset \Gamma$ we denote by $x|_A$ the restriction of x to

A, extended by zero; note that $x|_A$ need not belong to a given subspace $P \ni x$. For a subspace $P \subset \ell_1(\Gamma)$ we put $P_\delta = P \cap \ell_1(\Gamma_\delta)$ and

$$D_\delta = \{x|_{N_\delta} : x \in P_{\delta+1}\} \subset \ell_1(N_\delta), \qquad \delta < \lambda.$$

Since the kernel of the restriction map $x \mapsto x|_{N_\delta}$ on $P_{\delta+1}$ is $P_{\delta+1} \cap \ell_1(\Gamma_\delta) = P_\delta$, the space D_δ is isomorphic as a vector space to $P_{\delta+1}/P_\delta$.

Theorem 6.1. *Let Γ be a set, P be a dense subspace of $\ell_1(\Gamma)$ and $(\Gamma_\delta)_{\delta\le\lambda}$ be a filtration of Γ all of whose blocks N_δ are at most countable. Suppose that*

(i) P_δ is dense in $\ell_1(\Gamma_\delta)$ for every $\delta \le \lambda$;

(ii) $P_\delta = \bigcup_{\delta'<\delta} P_{\delta'}$ for every limit ordinal $\delta \le \lambda$;

(iii) the algebraic dimension of $P_{\delta+1}/P_\delta$ is at most countable for every $\delta < \lambda$.

Then for every $\varepsilon > 0$ there exists a linear bijection $T : \ell_1(\Gamma) \to \ell_1(\Gamma)$ with $T(P) = \ell_1^0(\Gamma)$, $\|T\| \le 1+\varepsilon$ and $\|T^{-1}\| \le 1$. In particular, $d_{BM}(P, \ell_1^0(\Gamma)) = 1$.

Proof. We may assume that $\Gamma \ne \varnothing$. Fix $\theta \in (0, 1/2)$ with $(1+\theta)^2(1-\theta)^{-1} \le 1+\varepsilon$. First we construct, for every $\delta < \lambda$, a linear bijection U_δ of $\ell_1(N_\delta)$ with $U_\delta(D_\delta) = \ell_1^0(N_\delta)$, $\|U_\delta\| \le 1+\theta$ and $\|U_\delta^{-1}\| \le 1$. If $N_\delta = \varnothing$, then $\ell_1(N_\delta) = \{0\}$ and $U_\delta = 0$ works. Otherwise note that the restriction map $x \mapsto x|_{N_\delta}$ is a contractive surjection of $\ell_1(\Gamma_{\delta+1})$ onto $\ell_1(N_\delta)$ and $P_{\delta+1}$ is dense in $\ell_1(\Gamma_{\delta+1})$ by (i), so D_δ is dense in $\ell_1(N_\delta)$; moreover, D_δ has at most countable algebraic dimension by (iii). If N_δ is finite, then $D_\delta = \ell_1(N_\delta) = \ell_1^0(N_\delta)$ and $U_\delta = 1_{\ell_1(N_\delta)}$ works. If N_δ is countably infinite, we identify $\ell_1(N_\delta)$ with ℓ_1 by means of a bijection between N_δ and $\mathbb{N}$, which preserves finitely supported families, and apply theorem 4.5 to D_δ and θ: we get a linear bijection U of $\ell_1(N_\delta)$ with $U(D_\delta) = \ell_1^0(N_\delta)$ and $\|U\|\|U^{-1}\| \le 1+\theta$, and $U_\delta = \|U^{-1}\|U$ has the required properties.

Now we construct by transfinite recursion, for every $\delta < \lambda$, a bounded linear operator $V_\delta : \ell_1(N_\delta) \to \ell_1(\Gamma_\delta)$ with $\|V_\delta\| \le \theta$. For $\delta \le \lambda$ and $x \in \ell_1(\Gamma_\delta)$ put

$$W_\delta(x) = x + \sum_{\delta'<\delta} V_{\delta'}(x|_{N_{\delta'}}), \qquad U_{<\delta}(x) = \sum_{\delta'<\delta} U_{\delta'}(x|_{N_{\delta'}}), \qquad T_\delta = U_{<\delta}W_\delta.$$

Since the sets $N_{\delta'}$ are pairwise disjoint and $\|x\| = \sum_{\delta'<\delta}\|x|_{N_{\delta'}}\|$, the series defining $W_\delta(x)$ converges absolutely in $\ell_1(\Gamma_\delta)$ and $\|W_\delta - 1\| \le \theta$, so W_δ is a linear bijection of $\ell_1(\Gamma_\delta)$ with $\|W_\delta^{-1}\| \le (1-\theta)^{-1}$; the operator $U_{<\delta}$ is the ℓ_1-sum of the operators $U_{\delta'}$, so it is a linear bijection of $\ell_1(\Gamma_\delta)$ with $\|U_{<\delta}\| \le 1+\theta$ and $\|U_{<\delta}^{-1}\| \le 1$. Hence T_δ is a linear bijection of $\ell_1(\Gamma_\delta)$ with $\|T_\delta\| \le (1+\theta)^2$ and $\|T_\delta^{-1}\| \le (1-\theta)^{-1}$. Moreover, for $\delta' < \delta$ the operator T_δ coincides with $T_{\delta'}$ on $\ell_1(\Gamma_{\delta'})$, because for $x \in \ell_1(\Gamma_{\delta'})$ we have $x|_{N_{\delta''}} = 0$ for $\delta'' \ge \delta'$. The operators V_δ will be chosen so that

$$T_\delta(P_\delta) = \ell_1^0(\Gamma_\delta) \quad \text{for all } \delta \le \lambda. \tag{1}$$

Then T_λ is a linear bijection of $\ell_1(\Gamma)$ with $T_\lambda(P) = \ell_1^0(\Gamma)$, and $T = \|T_\lambda^{-1}\|T_\lambda$ satisfies $\|T^{-1}\| = 1$ and $\|T\| \le (1+\theta)^2(1-\theta)^{-1} \le 1+\varepsilon$, as required.

Let $\delta < \lambda$ and suppose that $V_{\delta'}$ has been constructed for all $\delta' < \delta$ in such a way that (1) holds for all ordinals $\le \delta$ (for $\delta = 0$ this is trivial, since $\Gamma_0 = \varnothing$). We construct V_δ so that (1) holds for $\delta+1$; note that (1) for a limit ordinal $\mu \le \lambda$ follows from (1) for all $\delta < \mu$, because by (ii) and the compatibility of the operators T_δ we have $T_\mu(P_\mu) = \bigcup_{\delta<\mu} T_\delta(P_\delta) = \bigcup_{\delta<\mu} \ell_1^0(\Gamma_\delta) = \ell_1^0(\Gamma_\mu)$.

If $N_\delta = \varnothing$, then $P_{\delta+1} = P_\delta$, $T_{\delta+1} = T_\delta$, and we put $V_\delta = 0$. So let $N_\delta \neq \varnothing$, and put $U = U_\delta$ and $y_n = U^{-1}(e_n)$ for $n \in N_\delta$. Since $U(D_\delta) = \ell_1^0(N_\delta)$, $(y_n)_{n\in N_\delta}$ is a Hamel basis of D_δ and

$$\Big\|\sum_n c_n y_n\Big\| \geq \|U\|^{-1}\Big\|\sum_n c_n e_n\Big\| \geq (1+\theta)^{-1}\sum_n |c_n|$$

for all finitely supported families $(c_n)_{n\in N_\delta}$ of scalars. For every $n \in N_\delta$ choose $x_n \in P_{\delta+1}$ with $x_n|_{N_\delta} = y_n$. Since P_δ is dense in $\ell_1(\Gamma_\delta)$ by (i), the coset $-x_n|_{\Gamma_\delta} + P_\delta$ is dense in $\ell_1(\Gamma_\delta)$, and we can choose $z_n \in \ell_1(\Gamma_\delta)$ with

$$x_n|_{\Gamma_\delta} + z_n \in P_\delta, \qquad \|z_n\| \leq \theta(1+\theta)^{-2}.$$

Define V_δ on D_δ by $V_\delta(\sum_n c_n y_n) = \sum_n c_n W_\delta(z_n)$. Then

$$\Big\|V_\delta\Big(\sum_n c_n y_n\Big)\Big\| \leq \|W_\delta\|\,\theta(1+\theta)^{-2}\sum_n |c_n| \leq \theta(1+\theta)^{-1}\sum_n |c_n| \leq \theta\Big\|\sum_n c_n y_n\Big\|,$$

so V_δ extends by continuity from the dense subspace D_δ to a bounded linear operator $V_\delta : \ell_1(N_\delta) \to \ell_1(\Gamma_\delta)$ with $\|V_\delta\| \leq \theta$ (recall that D_δ is dense in $\ell_1(N_\delta)$ by (i), since the restriction map is a contractive surjection of $\ell_1(\Gamma_{\delta+1})$ onto $\ell_1(N_\delta)$).

We claim that

$$x|_{\Gamma_\delta} + W_\delta^{-1}V_\delta(x|_{N_\delta}) \in P_\delta \quad \text{for every } x \in P_{\delta+1}. \tag{2}$$

Indeed, write $x|_{N_\delta} = \sum_n c_n y_n$ with finitely many non-zero c_n. Then $x - \sum_n c_n x_n \in P_{\delta+1}$ vanishes on N_δ, so it belongs to P_δ, and hence $x|_{\Gamma_\delta} = \sum_n c_n x_n|_{\Gamma_\delta} + p$ for some $p \in P_\delta$. Therefore

$$x|_{\Gamma_\delta} + W_\delta^{-1}V_\delta(x|_{N_\delta}) = \sum_n c_n\big(x_n|_{\Gamma_\delta} + z_n\big) + p \in P_\delta.$$

Now let $x \in P_{\delta+1}$. Since $W_{\delta+1}(x) = W_\delta(x|_{\Gamma_\delta}) + V_\delta(x|_{N_\delta}) + x|_{N_\delta}$ and $U_{<\delta+1}$ acts as $U_{<\delta}$ on $\ell_1(\Gamma_\delta)$ and as U on $\ell_1(N_\delta)$, we get

$$T_{\delta+1}(x) = U_{<\delta}W_\delta\big(x|_{\Gamma_\delta} + W_\delta^{-1}V_\delta(x|_{N_\delta})\big) + U(x|_{N_\delta}) = T_\delta\big(x|_{\Gamma_\delta} + W_\delta^{-1}V_\delta(x|_{N_\delta})\big) + U(x|_{N_\delta}).$$

By (2) and (1) for δ, the first summand belongs to $\ell_1^0(\Gamma_\delta)$, and the second one belongs to $U(D_\delta) = \ell_1^0(N_\delta)$. Hence $T_{\delta+1}(P_{\delta+1}) \subset \ell_1^0(\Gamma_{\delta+1})$. Conversely, $\ell_1^0(\Gamma_\delta) = T_\delta(P_\delta) = T_{\delta+1}(P_\delta) \subset T_{\delta+1}(P_{\delta+1})$, and for every $n \in N_\delta$ the above formula gives $T_{\delta+1}(x_n) \in \ell_1^0(\Gamma_\delta) + U(y_n) = \ell_1^0(\Gamma_\delta) + e_n$, so $e_n \in T_{\delta+1}(P_{\delta+1})$. Thus, $T_{\delta+1}(P_{\delta+1}) = \ell_1^0(\Gamma_{\delta+1})$, which completes the recursion and the proof. □

We proceed to the construction of filtrations satisfying the hypotheses of theorem 6.1. Until theorem 6.7 we fix a set Γ, a dense topologically projective subspace P of $\ell_1(\Gamma)$, a set S and bounded linear operators $\sigma : P \to \ell_1^0(S)$ and $\pi : \ell_1^0(S) \to P$ with $\pi\sigma = 1_P$ and $\|\pi\| \leq 1$; these exist by proposition 2.6. Since P is dense in $\ell_1(\Gamma)$, we can also fix for every $\gamma \in \Gamma$ a sequence $(p_n^\gamma)_{n\in\mathbb{N}}$ in P converging to e_γ. For $A \subset \Gamma$ we put $P_A = P \cap \ell_1(A)$.

Definition 6.2. *A pair (A, B) of subsets $A \subset \Gamma$ and $B \subset S$ is called closed if*

(a) $\operatorname{supp}(\pi(e_s)) \subset A$ *for all* $s \in B$;

(b) $\operatorname{supp}(\sigma(x)) \subset B$ *for all* $x \in P_A$;

(c) $\operatorname{supp}(p_n^\gamma) \subset A$ *for all* $\gamma \in A$ *and* $n \in \mathbb{N}$.

Lemma 6.3. *Let (A, B) be a closed pair. Then $P_A = \pi(\ell_1^0(B))$, the space P_A is dense in $\ell_1(A)$, and $\sigma|_{P_A} : P_A \to \ell_1^0(B)$, $\pi|_{\ell_1^0(B)} : \ell_1^0(B) \to P_A$ exhibit P_A as a retract of $\ell_1^0(B)$. In particular, P_A is topologically projective.*

Proof. By (a), $\pi(\ell_1^0(B)) \subset P \cap \ell_1(A) = P_A$. If $x \in P_A$, then $\sigma(x) \in \ell_1^0(B)$ by (b) and $x = \pi\sigma(x) \in \pi(\ell_1^0(B))$. By (c), P_A contains the sequences p_n^γ, $\gamma \in A$, so its closure contains $\ell_1^0(A)$ and P_A is dense in $\ell_1(A)$. The last claim is now obvious. □

Lemma 6.4. *Let (A, B) be a closed pair and $A' \subset \Gamma$ be a set with $A \subset A'$ such that $A' \setminus A$ is at most countable. Then the algebraic dimension of $P_{A'}/P_A$ is at most countable.*

Proof. Put $N = A' \setminus A$ and let $r : \ell_1(A') \to \ell_1(N)$ be the restriction map. The kernel of $r|_{P_{A'}}$ is $P_{A'} \cap \ell_1(A) = P_A$, so $D = r(P_{A'})$ is isomorphic as a vector space to $P_{A'}/P_A$, and D is separable as a subspace of $\ell_1(N)$. The map

$$\rho : D \to \ell_1^0(S \setminus B), \qquad \rho(y) = \sigma(x)|_{S\setminus B} \quad (x \in P_{A'},\ x|_N = y),$$

is well defined, linear and bounded. Indeed, two elements $x, x' \in P_{A'}$ with $x|_N = x'|_N$ differ by an element of P_A, whose image under σ is supported in B by condition (b) for (A, B), so ρ is well defined and linear; and given $y \in D$, $x_0 \in P_{A'}$ with $x_0|_N = y$ and $\varepsilon > 0$, the density of P_A in $\ell_1(A)$ (lemma 6.3) gives $p \in P_A$ with $\|x_0|_A + p\| \le \varepsilon$, and then $x = x_0 + p \in P_{A'}$ satisfies $x|_N = y$ and $\|x\| = \|x_0|_A + p\| + \|y\| \le \|y\| + \varepsilon$, whence $\|\rho(y)\| \le \|\sigma\|(\|y\| + \varepsilon)$. The map ρ is injective: if $\rho(y) = 0$ for $y = x|_N$, then $\sigma(x) \in \ell_1^0(B)$, so $x = \pi\sigma(x) \in \pi(\ell_1^0(B)) = P_A$ by lemma 6.3, and $y = x|_N = 0$. By lemma 3.1 the algebraic dimension of D is at most countable. □

Lemma 6.5. *Let (A, B) be a closed pair and $N \subset \Gamma$ be an at most countable set. Then there is a closed pair (A', B') with $A \cup N \subset A'$, $B \subset B'$ such that $A' \setminus A$ and $B' \setminus B$ are at most countable.*

Proof. We define increasing sequences $A \cup N = A_0 \subset A_1 \subset \dots$ and $B = B_0 \subset B_1 \subset \dots$ such that $A_k \setminus A$ and $B_k \setminus B$ are at most countable. Given A_k and B_k, by lemma 6.4 there is an at most countable set $H_k \subset P_{A_k}$ such that $P_{A_k} = P_A + \operatorname{span}(H_k)$. Put

$$B_{k+1} = B_k \cup \bigcup_{x\in H_k} \operatorname{supp}(\sigma(x)), \qquad A_{k+1} = A_k \cup \bigcup_{s\in B_{k+1}\setminus B} \operatorname{supp}(\pi(e_s)) \cup \bigcup_{\gamma\in A_k\setminus A,\, n\in\mathbb{N}} \operatorname{supp}(p_n^\gamma).$$

The sets $B_{k+1} \setminus B$ and $A_{k+1} \setminus A$ are at most countable, since all supports involved are at most countable. By condition (b) for (A, B) we have $\operatorname{supp}(\sigma(x)) \subset B_{k+1}$ for all $x \in P_{A_k}$. Put $A' = \bigcup_k A_k$ and $B' = \bigcup_k B_k$. Then $A' \setminus A$ and $B' \setminus B$ are at most countable. Condition (a) holds for (A', B'): for $s \in B$ we have $\operatorname{supp}(\pi(e_s)) \subset A$ by (a) for (A, B), and for $s \in B_{k+1} \setminus B$ we have $\operatorname{supp}(\pi(e_s)) \subset A_{k+1}$. Similarly, condition (c) holds for (A', B'), since $\operatorname{supp}(p_n^\gamma) \subset A$ for $\gamma \in A$ and $\operatorname{supp}(p_n^\gamma) \subset A_{k+1}$ for $\gamma \in A_k \setminus A$. Finally, $\bigcup_k P_{A_k}$ is dense in $\ell_1(A')$: its closure contains P_A, which is dense in $\ell_1(A)$ by lemma 6.3, and the vectors $p_n^\gamma \in P_{A_{k+1}}$, $\gamma \in A_k \setminus A$, so it contains e_γ for all $\gamma \in A'$. If $x \in P_{A'}$, choose $x_j \in \bigcup_k P_{A_k}$ with $x_j \to x$; then $\sigma(x_j) \in \ell_1^0(B_{k_j+1}) \subset \ell_1(B')$ for suitable k_j, and since σ is continuous and $\ell_1(B')$ is closed in $\ell_1(S)$, we get $\sigma(x) \in \ell_1(B') \cap \ell_1^0(S) = \ell_1^0(B')$. This is condition (b) for (A', B'). □

Note that in the last argument the support of $x \in P_{A'}$ need not be contained in a single A_k; this is why density was used instead of a direct argument. The same argument proves the following lemma.

Lemma 6.6. *Let $((A_\delta, B_\delta))_{\delta<\lambda}$ be an increasing family of closed pairs indexed by a limit ordinal λ, and let $A_\lambda = \bigcup_{\delta<\lambda} A_\delta$, $B_\lambda = \bigcup_{\delta<\lambda} B_\delta$. Then (A_λ, B_λ) is a closed pair and $P_{A_\lambda} = \bigcup_{\delta<\lambda} P_{A_\delta}$.*

Proof. Conditions (a) and (c) are inherited from the pairs (A_δ, B_δ). By lemma 6.3 each P_{A_δ} is dense in $\ell_1(A_\delta)$, so $\bigcup_{\delta<\lambda} P_{A_\delta}$ is dense in $\ell_1(A_\lambda)$. If $x \in P_{A_\lambda}$, choose $x_j \in \bigcup_\delta P_{A_\delta}$ with $x_j \to x$; then $\sigma(x_j) \in \ell_1(B_\lambda)$ by (b) for the pairs (A_δ, B_δ), hence $\sigma(x) \in \ell_1(B_\lambda) \cap \ell_1^0(S) = \ell_1^0(B_\lambda)$. This is (b) for (A_λ, B_λ). Finally, by lemma 6.3, $P_{A_\lambda} = \pi(\ell_1^0(B_\lambda)) = \bigcup_{\delta<\lambda} \pi(\ell_1^0(B_\delta)) = \bigcup_{\delta<\lambda} P_{A_\delta}$. □

We can now prove the main result of this section.

Theorem 6.7. *Let Γ be a set and P be a dense subspace of $\ell_1(\Gamma)$. Then the following conditions are equivalent:*

(i) P is extremely projective;

(ii) P is topologically projective;

(iii) there are an ordinal λ and a filtration $(\Gamma_\delta)_{\delta\le\lambda}$ of Γ, all of whose blocks $N_\delta = \Gamma_{\delta+1} \setminus \Gamma_\delta$ are at most countable, such that

(a) P_δ is dense in $\ell_1(\Gamma_\delta)$ for every $\delta \le \lambda$;

(b) $P_\delta = \bigcup_{\delta'<\delta} P_{\delta'}$ for every limit ordinal $\delta \le \lambda$;

(c) the algebraic dimension of $P_{\delta+1}/P_\delta$ is at most countable for every $\delta < \lambda$.

If these conditions hold, then for every $\varepsilon > 0$ there is a linear bijection T of $\ell_1(\Gamma)$ with $T(P) = \ell_1^0(\Gamma)$, $\|T\| \le 1+\varepsilon$ and $\|T^{-1}\| \le 1$; in particular, $d_{BM}(P, \ell_1^0(\Gamma)) = 1$.

Proof. $(i) \implies (ii)$ is trivial.

$(iii) \implies (i)$ and the last claim. By theorem 6.1, for every $\varepsilon > 0$ there is a linear bijection T of $\ell_1(\Gamma)$ with $T(P) = \ell_1^0(\Gamma)$, $\|T\| \le 1+\varepsilon$ and $\|T^{-1}\| \le 1$. Then $\zeta = T^{-1}|_{\ell_1^0(\Gamma)} : \ell_1^0(\Gamma) \to P$ and $\eta = T|_P : P \to \ell_1^0(\Gamma)$ satisfy $\zeta\eta = 1_P$ and $\|\zeta\|\|\eta\| \le 1+\varepsilon$, so P is a $(1+\varepsilon)$-retract of $\ell_1^0(\Gamma)$ for every $\varepsilon > 0$, and P is extremely projective by theorem 2.7.

$(ii) \implies (iii)$ Fix S, σ, π and the sequences p_n^γ as above, and an enumeration $(\gamma_\alpha)_{\alpha<\kappa}$ of Γ by an ordinal κ. By transfinite recursion we define an increasing family $((A_\alpha, B_\alpha))_{\alpha\le\kappa}$ of closed pairs: $A_0 = B_0 = \varnothing$, which is a closed pair since $\sigma(0) = 0$; given (A_α, B_α) with $\alpha < \kappa$, let $(A_{\alpha+1}, B_{\alpha+1})$ be a closed pair provided by lemma 6.5 for (A_α, B_α) and $N = \{\gamma_\alpha\}$, so that $A_{\alpha+1} \setminus A_\alpha$ is at most countable; for a limit ordinal $\alpha \le \kappa$ let A_α and B_α be the unions of the previous sets, which form a closed pair by lemma 6.6. Since $\gamma_\alpha \in A_{\alpha+1}$, we have $A_\kappa = \Gamma$, so $(A_\alpha)_{\alpha\le\kappa}$ is a filtration of Γ with at most countable blocks. Condition (a) holds by lemma 6.3, condition (b) by lemma 6.6, and condition (c) by lemma 6.4. □

Together with theorem 4.1, theorem 6.7 gives the main result of the paper.

Theorem 6.8. *For a normed space P the following conditions are equivalent:*

(i) P is extremely projective;

(ii) P is topologically projective and $\overline{P}$ is isometrically isomorphic to $\ell_1(\Gamma)$ for some set Γ;

(iii) $d_{BM}(P, \ell_1^0(\Gamma)) = 1$ for some set Γ;

(iv) for every $\varepsilon > 0$ there is a set Λ with $d_{BM}(P, \ell_1^0(\Lambda)) < 1 + \varepsilon$.

Moreover, in (iii) the set Γ necessarily has the cardinality of the algebraic dimension of P, and the Banach–Mazur distance in (iii) is realized by linear bijections $T : \overline{P} \to \ell_1(\Gamma)$ with $T(P) = \ell_1^0(\Gamma)$ and $\|T\|\|T^{-1}\|$ arbitrarily close to 1.

Proof. $(i) \implies (ii)$ Extremely projective spaces are topologically projective, and their completions are described by theorem 4.1.

$(ii) \implies (iii)$ Let $J : \overline{P} \to \ell_1(\Gamma)$ be an isometric isomorphism. Then $J(P)$ is a dense topologically projective subspace of $\ell_1(\Gamma)$, and theorem 6.7 gives $d_{BM}(P, \ell_1^0(\Gamma)) = d_{BM}(J(P), \ell_1^0(\Gamma)) = 1$, realized by linear bijections of the required form.

$(iii) \implies (iv)$ is obvious, and $(iv) \implies (i)$ follows from theorem 2.7, since a linear bijection $S : P \to \ell_1^0(\Lambda)$ with $\|S\|\|S^{-1}\| < 1+\varepsilon$ makes P a $(1+\varepsilon)$-retract of $\ell_1^0(\Lambda)$. Finally, if $d_{BM}(P, \ell_1^0(\Gamma)) < \infty$, then P and $\ell_1^0(\Gamma)$ are isomorphic as vector spaces, and the algebraic dimension of $\ell_1^0(\Gamma)$ is the cardinality of Γ; so the set Γ in (iii) is determined up to a bijection, and the realization of the distance obtained in $(ii) \implies (iii)$ applies to it. □

Corollary 6.9. *Every extremely projective normed space P is almost isometric to $\ell_1^0(\Gamma)$, where Γ is a set whose cardinality is the algebraic dimension of P. In particular, the conjecture of Grønbæk [8, conjecture 3.1] holds in the almost isometric form, while corollary 5.3 shows that it fails in the isometric form.*

We finish this section with some comments. Let us compare our argument with the proof of the theorem of Grønbæk, which consists of two steps: a transfinite construction which finds in every subspace Y of $\ell_1^0(\Lambda)$ a complemented subspace isomorphic to $\ell_1^0(\Lambda')$, where the cardinality of Λ' is the density character of Y [8, proposition 2.2], and the decomposition method of Pełczyński [8, proposition 2.3]. An inspection of the first step shows that it produces, for every $\varepsilon > 0$, a subspace which is $(1 + \varepsilon)$-isomorphic to $\ell_1^0(\Lambda')$ and $(1 + \varepsilon)$-complemented, so the constants are lost only in the second step. Theorems 4.5 and 6.1 replace the decomposition method by back-and-forth constructions which keep the constants, at the price of the continuity condition (ii), and lemmas 6.5 and 6.6 show that this condition can always be achieved for topologically projective spaces. Next, theorem 6.8 shows that for dense subspaces of the spaces $\ell_1(\Gamma)$ topological and extreme projectivity coincide. This is a feature of the exact norm of $\ell_1(\Gamma)$: if E is an infinite dimensional Banach space which is topologically but not almost isometrically isomorphic to $\ell_1(\Gamma)$, for instance $\ell_1(\Gamma)$ with a suitable equivalent norm, then every dense subspace of E of the form $J(\ell_1^0(\Gamma))$, J a topological isomorphism, is topologically but not extremely projective by theorem 4.1. Finally, by theorem 6.7, the question of Helemskii on dense subspaces of $\ell_1(\Lambda)$ has the following answer: a dense subspace P of $\ell_1(\Lambda)$ is extremely projective if and only if it is topologically projective, if and only if it admits a filtration with countable blocks satisfying conditions (a)–(c) of theorem 6.7; for countable Λ this means that P has countable algebraic dimension, but for uncountable Λ the continuity condition (b) is a genuine restriction, as the ladder spaces of the next section show.

7 Extremely projective normed spaces of dimension $\aleph_1$

Condition (iii) of theorem 6.7 refers only to the subspaces $P \cap \ell_1(A)$, $A \subset \Gamma$, that is, to the coordinate structure of $\ell_1(\Gamma)$. Note that this structure is intrinsic: by lemma 5.1, applied to $E = \ell_1(\Gamma)$, the extreme points of the unit ball of $\ell_1(\Gamma)$ are exactly the vectors λe_γ with $|\lambda| = 1$, so the family of the one dimensional subspaces $\mathbb{C}e_\gamma$, $\gamma \in \Gamma$, is determined by the norm of $\ell_1(\Gamma)$. For

sets Γ of cardinality $\aleph_1$ condition (iii) can be stated as a condition on a single chain of subspaces. In this section we take $\Gamma = \omega_1$; every set of cardinality $\aleph_1$ can be enumerated by ω_1, and we shall see in proposition 7.3 that the choice of the enumeration is immaterial. For $\delta \le \omega_1$ we write $P_\delta = P \cap \ell_1(\delta)$, where $\ell_1(\delta)$ is the subspace of $\ell_1(\omega_1)$ consisting of the families supported in $\delta = \{\beta : \beta < \delta\}$. Recall that a subset of ω_1 is called closed unbounded if it is unbounded in ω_1 and contains the supremum of each of its countable subsets, and stationary if it intersects every closed unbounded set; the set of limit ordinals below ω_1 is closed unbounded. We shall use the following standard facts about closed unbounded sets [13, chapter 8]: the intersection of two closed unbounded sets is closed unbounded, and for every function $h : \omega_1 \to \omega_1$ the set of ordinals $\delta < \omega_1$ such that $h(\alpha) < \delta$ for all $\alpha < \delta$ is closed unbounded; we call such ordinals closed under h. Moreover, the increasing enumeration $(c_\alpha)_{\alpha<\omega_1}$ of a closed unbounded set C is continuous, that is, $c_\alpha = \sup_{\alpha'<\alpha} c_{\alpha'}$ for every limit ordinal $\alpha < \omega_1$.

Definition 7.1. *Let P be a dense subspace of $\ell_1(\omega_1)$. We denote by $\Delta(P)$ the set of limit ordinals $\delta < \omega_1$ such that $P_\delta \ne \bigcup_{\delta'<\delta} P_{\delta'}$.*

Theorem 7.2. *Let P be a dense subspace of $\ell_1(\omega_1)$. Then P is extremely projective if and only if*

(i) P_δ has at most countable algebraic dimension for every $\delta < \omega_1$, and

(ii) the set $\Delta(P)$ is not stationary in ω_1.

Condition (i) is equivalent to the requirement that every separable subspace of P has at most countable algebraic dimension.

Proof. First we prove the last claim. Each P_δ, $\delta < \omega_1$, is separable as a subspace of $\ell_1(\delta)$. Conversely, if Z is a separable subspace of P, then the supports of the elements of a countable dense subset of Z are contained in a countable subset of ω_1, hence in some $\delta < \omega_1$, and by the continuity of the coordinate functionals $Z \subset P_\delta$.

Let P be extremely projective. Condition (i) holds by theorem 3.2. To verify (ii) we fix S, σ, π and p_n^γ as in section 6 and apply the construction of the proof of theorem 6.7 to the enumeration $\gamma_\alpha = \alpha$ of ω_1. We obtain an increasing family $((A_\alpha, B_\alpha))_{\alpha\le\omega_1}$ of closed pairs, continuous at limit ordinals, such that A_α is countable for $\alpha < \omega_1$, $\alpha \in A_{\alpha+1}$ and $A_{\omega_1} = \omega_1$. Define $f, g : \omega_1 \to \omega_1$ by $f(\alpha) = \sup A_\alpha + 1$ and $g(\beta) = \min\{\alpha : \beta \in A_\alpha\}$, and let D be the set of limit ordinals $\delta < \omega_1$ closed under f and g; it is closed unbounded. Let $\delta \in D$. Then $A_\delta = \bigcup_{\alpha<\delta} A_\alpha \subset \delta$, because $\sup A_\alpha < f(\alpha) < \delta$ for $\alpha < \delta$, and $\delta \subset A_\delta$, because $\beta < \delta$ implies $\beta \in A_{g(\beta)} \subset A_\delta$. Thus, $A_\delta = \delta$, and by lemma 6.6

$$P_\delta = P_{A_\delta} = \bigcup_{\alpha<\delta} P_{A_\alpha} \subset \bigcup_{\alpha<\delta} P_{f(\alpha)} \subset \bigcup_{\delta'<\delta} P_{\delta'} \subset P_\delta,$$

since $A_\alpha \subset f(\alpha) < \delta$. Hence $\delta \notin \Delta(P)$. So $\Delta(P) \cap D = \varnothing$, and $\Delta(P)$ is not stationary.

Conversely, assume (i) and (ii). For $\gamma < \omega_1$ fix a sequence $(p_n^\gamma)_{n\in\mathbb{N}}$ in P converging to e_γ and put $h(\gamma) = \sup \bigcup_n \operatorname{supp}(p_n^\gamma) + 1 < \omega_1$. Let C_1 be the set of ordinals $\delta < \omega_1$ closed under h; for $\delta \in C_1$ the space P_δ contains p_n^γ for all $\gamma < \delta$, so P_δ is dense in $\ell_1(\delta)$. By (ii) there is a closed unbounded set C_2 disjoint from $\Delta(P)$. Let $C = (C_1 \cap C_2) \cup \{0\}$, a closed unbounded set, and let $(c_\alpha)_{\alpha<\omega_1}$ be its increasing enumeration, so that $c_0 = 0$. Put $\Gamma_\alpha = c_\alpha$ for $\alpha < \omega_1$ and $\Gamma_{\omega_1} = \omega_1$. Then $(\Gamma_\alpha)_{\alpha\le\omega_1}$ is a filtration of ω_1: it is increasing, $\Gamma_0 = \varnothing$, continuity at limit ordinals $\alpha < \omega_1$ is the continuity of the enumeration of C, and continuity at ω_1 holds because C is unbounded. Its blocks $N_\alpha = \{\beta : c_\alpha \le \beta < c_{\alpha+1}\}$ are countable. We verify conditions (a)–(c) of theorem 6.7. Condition (a) holds for $\alpha < \omega_1$ because $c_\alpha \in C_1$ (for $\alpha = 0$ trivially), and for $\alpha = \omega_1$ because P

is dense. For a limit ordinal $\alpha < \omega_1$ the ordinal c_α is a limit ordinal in C_2, hence $c_\alpha \notin \Delta(P)$, and since the ordinals $c_{\alpha'}$, $\alpha' < \alpha$, are cofinal in c_α and the spaces P_δ increase with δ,

$$P_{c_\alpha} = \bigcup_{\delta' < c_\alpha} P_{\delta'} = \bigcup_{\alpha' < \alpha} P_{c_{\alpha'}};$$

at $\alpha = \omega_1$ condition (b) holds because every element of P has countable support and therefore belongs to $P_{c_{\alpha'}}$ for some $\alpha' < \omega_1$. Finally, (c) follows from (i), since $P_{c_{\alpha+1}}$ has at most countable algebraic dimension. By theorem 6.7 the space P is extremely projective. □

Neither of the conditions of theorem 7.2 can be dropped: the ladder spaces on stationary sets constructed in theorem 7.4 below satisfy (i) but not (ii), while the space $P = P_w + \ell_1^0(\omega_1 \setminus \omega)$, where $P_w \subset \ell_1(\omega)$ is the weighted space of example 3.4, satisfies (ii) but not (i): here $\Delta(P) = \{\omega\}$, because every element of P is the sum of an element of P_w and a finitely supported family, so the chain $(P_\delta)_\delta$ is continuous at every limit ordinal $\delta > \omega$, while $P_\omega = P_w$ has uncountable algebraic dimension.

The set $\Delta(P)$ depends on the position of P in $\ell_1(\omega_1)$, but only up to a non-stationary set.

Proposition 7.3. *Let P and P' be dense subspaces of $\ell_1(\omega_1)$ and $T : \ell_1(\omega_1) \to \ell_1(\omega_1)$ be a bounded linear bijection with $T(P) = P'$. Then $\Delta(P) \triangle \Delta(P')$ is not stationary. In particular, if P and P' are topologically isomorphic as normed spaces, then $\Delta(P) \triangle \Delta(P')$ is not stationary, and the class of $\Delta(P)$ modulo non-stationary sets does not depend on the enumeration of the index set by ω_1.*

Proof. For $\delta < \omega_1$ put

$$F(\delta) = \sup_{\beta<\delta}\big(\sup \operatorname{supp}(T(e_\beta)) + 1\big), \qquad G(\delta) = \sup_{\beta<\delta}\big(\sup \operatorname{supp}(T^{-1}(e_\beta)) + 1\big),$$

which are countable ordinals, since all supports are countable. Let D be the set of limit ordinals $\delta < \omega_1$ closed under F and G; it is closed unbounded. Let $\delta \in D$ and $\delta' < \delta$. Then $T(e_\beta) \in \ell_1(F(\delta'))$ for all $\beta < \delta'$, so by the continuity of T we have $T(\ell_1(\delta')) \subset \ell_1(F(\delta')) \subset \ell_1(\delta)$, and since $\ell_1(\delta)$ is the closure of $\bigcup_{\delta'<\delta} \ell_1(\delta')$, also $T(\ell_1(\delta)) \subset \ell_1(\delta)$. The same holds for T^{-1}, which is bounded by the open mapping theorem, with G in place of F, so $T(\ell_1(\delta)) = \ell_1(\delta)$ and $T(P_\delta) = T(P) \cap T(\ell_1(\delta)) = P'_\delta$. Moreover, $T(P_{\delta'}) \subset P' \cap \ell_1(F(\delta')) = P'_{F(\delta')}$ with $F(\delta') < \delta$, so $T\big(\bigcup_{\delta'<\delta} P_{\delta'}\big) \subset \bigcup_{\delta'<\delta} P'_{\delta'}$, and symmetrically for T^{-1}; hence T maps $\bigcup_{\delta'<\delta} P_{\delta'}$ onto $\bigcup_{\delta'<\delta} P'_{\delta'}$. Consequently, $\delta \in \Delta(P)$ if and only if $\delta \in \Delta(P')$, for every $\delta \in D$. Thus, $\Delta(P) \triangle \Delta(P')$ is disjoint from D and is not stationary.

If P and P' are topologically isomorphic, then an isomorphism $P \to P'$ extends to a bounded linear bijection of the completions, which are both $\ell_1(\omega_1)$, so the first claim applies. Finally, two enumerations of the same index set by ω_1 differ by a permutation of ω_1, which induces an isometric bijection of $\ell_1(\omega_1)$ mapping P onto the same space P regarded with respect to the other enumeration. □

We now construct the ladder spaces. Let E be a set of limit ordinals below ω_1. For every $\alpha \in E$ fix a strictly increasing sequence $(\alpha_n)_{n\in\mathbb{N}}$ with $\sup_n \alpha_n = \alpha$, a ladder at α, and put $v_\alpha = \sum_{n=1}^\infty 2^{-n} e_{\alpha_n} \in \ell_1(\omega_1)$, and let

$$P_E = \ell_1^0(\omega_1) + \operatorname{span}\{v_\alpha : \alpha \in E\} \subset \ell_1(\omega_1).$$

We call P_E the ladder space on E. Ladder systems have been used before in Banach space theory, for instance in the study of the spaces $C(K)$ of ladder system compacta [10] and in the construction of 2^κ pairwise non-isomorphic spaces $C(K)$ of weight κ [11]; here they enter through the continuity condition of theorem 7.2.

Theorem 7.4. *Let E and E' be sets of limit ordinals below ω_1. Then*

(i) *P_E is a dense subspace of $\ell_1(\omega_1)$ of algebraic dimension $\aleph_1$, and for every $\delta < \omega_1$ the space $(P_E)_\delta = \ell_1^0(\delta) + \operatorname{span}\{v_\alpha : \alpha \in E,\ \alpha \le \delta\}$ has countable algebraic dimension and is extremely projective;*

(ii) *$\Delta(P_E) = E$;*

(iii) *P_E is extremely projective if and only if E is not stationary;*

(iv) *if $E \triangle E'$ is stationary, then P_E and $P_{E'}$ are not topologically isomorphic.*

Consequently, there are $2^{\aleph_1}$ pairwise non-isomorphic dense subspaces of $\ell_1(\omega_1)$ of algebraic dimension $\aleph_1$ all of whose subspaces $P \cap \ell_1(\delta)$, $\delta < \omega_1$, are extremely projective, and exactly one of them, namely $\ell_1^0(\omega_1)$, is extremely projective itself.

Proof. *(i)* The space P_E contains $\ell_1^0(\omega_1)$, so it is dense and has algebraic dimension $\aleph_1$, being spanned by $\aleph_1$ vectors. For limit ordinals $\alpha < \beta$ the set $\operatorname{supp}(v_\alpha) \cap \operatorname{supp}(v_\beta)$ is finite: $\operatorname{supp}(v_\alpha) \subset \alpha$, while only finitely many β_n are smaller than α, because $\sup_n \beta_n = \beta > \alpha$. Let $\delta < \omega_1$ and $x = y + \sum_{\alpha\in F} c_\alpha v_\alpha \in (P_E)_\delta$, where $y \in \ell_1^0(\omega_1)$, $F \subset E$ is finite and $c_\alpha \ne 0$ for $\alpha \in F$. Suppose that $F \not\subset [0,\delta]$ and put $\alpha_0 = \max F > \delta$. The set $\operatorname{supp}(v_{\alpha_0}) \cap [\delta, \alpha_0)$ is infinite, because $\sup_n \alpha_{0,n} = \alpha_0 > \delta$, while the sets $\operatorname{supp}(y)$ and $\operatorname{supp}(v_\alpha) \cap \operatorname{supp}(v_{\alpha_0})$, $\alpha \in F \setminus \{\alpha_0\}$, are finite. Hence x has infinitely many non-zero coordinates in $[\delta, \alpha_0)$, which contradicts $x \in \ell_1(\delta)$. Therefore $F \subset [0,\delta]$, and then also $\operatorname{supp}(y) \subset \delta$. This proves the inclusion $\subset$ in the formula for $(P_E)_\delta$, and the reverse inclusion is obvious. The space $(P_E)_\delta$ has countable algebraic dimension and is dense in $\ell_1(\delta)$, because it contains $\ell_1^0(\delta)$, so it is extremely projective by theorem 4.6.

(ii) Let $\delta < \omega_1$ be a limit ordinal. By *(i)*,

$$\bigcup_{\delta'<\delta} (P_E)_{\delta'} = \ell_1^0(\delta) + \operatorname{span}\{v_\alpha : \alpha \in E,\ \alpha < \delta\},$$

because $\ell_1^0(\delta) = \bigcup_{\delta'<\delta} \ell_1^0(\delta')$ and $v_\alpha \in (P_E)_\alpha$ for $\alpha \in E$. If $\delta \notin E$, this is $(P_E)_\delta$ by *(i)*, so $\delta \notin \Delta(P_E)$. If $\delta \in E$, then $v_\delta \in (P_E)_\delta$, while $v_\delta \notin \ell_1^0(\delta) + \operatorname{span}\{v_\alpha : \alpha \in E,\ \alpha < \delta\}$: if $v_\delta = y + \sum_{\alpha\in F} c_\alpha v_\alpha$ with $y \in \ell_1^0(\delta)$ and a finite set F of limit ordinals below δ, then for all sufficiently large n the coordinate of the right hand side at δ_n vanishes, because $\operatorname{supp}(y)$ and $\operatorname{supp}(v_\alpha) \cap \operatorname{supp}(v_\delta)$, $\alpha \in F$, are finite, whereas the coordinate of v_δ at δ_n is 2^{-n}. Hence $\delta \in \Delta(P_E)$.

(iii) follows from *(i)*, *(ii)* and theorem 7.2, and *(iv)* follows from *(ii)* and proposition 7.3.

For the last claim recall that ω_1 is the union of $\aleph_1$ pairwise disjoint stationary sets $(S_\xi)_{\xi<\omega_1}$ [13, theorem 8.10]; replacing S_ξ by its intersection with the closed unbounded set of limit ordinals, we may assume that the sets S_ξ consist of limit ordinals. For $X \subset \omega_1$ put $E_X = \bigcup_{\xi\in X} S_\xi$. If $X \ne Y$, then $E_X \triangle E_Y$ contains some S_ξ and is stationary, so by *(iv)* the spaces P_{E_X}, $X \subset \omega_1$, are pairwise non-isomorphic. By *(iii)*, P_{E_X} is extremely projective only for $X = \varnothing$, and $P_{E_\varnothing} = \ell_1^0(\omega_1)$. □

Note that all the spaces P_E contain the common dense subspace $\ell_1^0(\omega_1)$. Thus, the invariant $\Delta(P)$ is not determined by the dense subspaces of P; compare with the results of [9] on densely isomorphic normed spaces.

Remark 7.5. *The results of this section are parallel to the theory of almost free abelian groups, see [14]: theorem 6.7 corresponds to the criteria of Pontryagin and Hill [15] for freeness in terms of filtrations by countable subgroups with free quotients, the class of $\Delta(P)$ modulo non-stationary*

sets corresponds to the invariant $\Gamma(G)$ of Eklof [16], and the spaces P_E correspond to the $\aleph_1$-free groups of cardinality $\aleph_1$ built from ladder systems. In that theory the situation for cardinals above $\aleph_1$ depends on set-theoretic hypotheses, and we expect the same here. An analogue of the theory of almost free groups for non-Archimedean Banach spaces, where freeness means the existence of an orthonormal basis, has recently been proposed in [17]. We do not know whether condition (iii) of theorem 6.7 can be replaced, for $|\Gamma| \geq \aleph_2$, by a condition on the subspaces $P \cap \ell_1(A)$ for A ranging over a closed unbounded family of subsets of Γ of cardinality less than $|\Gamma|$, nor whether there is a dense subspace of $\ell_1(\omega_2)$ which is not topologically projective although $P \cap \ell_1(A)$ is extremely projective for every $A \subset \omega_2$ of cardinality at most $\aleph_1$.

References

[1] *A. Ya. Helemskii* The homology of Banach and topological algebras, Kluwer, Dordrecht (1989)

[2] *A. Ya. Helemskii* Metric freeness and projectivity for classical and quantum normed modules, Sb. Math., 204:7 (2013), 1056–1083

[3] *A. Ya. Helemskii* Extreme version of projectivity for normed modules over sequence algebras, Canad. J. Math., 65:3 (2013), 559–574; arXiv:1104.2463 (the numbering of statements refers to the arXiv version, titled "Metric version of projectivity for normed modules over sequence algebras")

[4] *M. C. White* Injective modules for uniform algebras, Proc. London Math. Soc. (3), 73:1 (1996), 155–184

[5] *G. Köthe* Hebbare lokalkonvexe Räume, Math. Ann., 165 (1966), 181–195

[6] *A. Grothendieck* Une caractérisation vectorielle-métrique des espaces L_1, Canad. J. Math., 7 (1955), 552–561

[7] *H. E. Lacey* The isometric theory of classical Banach spaces, Springer, Berlin (1974)

[8] *N. Grønbæk* Lifting problems for normed spaces, Ann. Funct. Anal., 7:1 (2016), 118–126

[9] *P. Hájek, T. Russo* On densely isomorphic normed spaces, J. Funct. Anal., 279:7 (2020), 108667; arXiv:1910.01527

[10] *J. Ferrer, P. Koszmider, W. Kubiś* Almost disjoint families of countable sets and separable complementation properties, J. Math. Anal. Appl., 401:2 (2013), 939–949; arXiv:1209.0199

[11] *M. Korpalski, P. Koszmider, W. Marciszewski* Counting spaces of functions on separable compact lines, arXiv:2602.09143 (2026)

[12] *H. Löwig* Über die Dimension linearer Räume, Studia Math., 5 (1934), 18–23

[13] *T. Jech* Set theory. The third millennium edition, revised and expanded, Springer Monographs in Mathematics, Springer, Berlin (2003)

[14] *P. C. Eklof, A. H. Mekler* Almost free modules. Set-theoretic methods, revised edition, North-Holland Mathematical Library, 65, North-Holland, Amsterdam (2002)

[15] *P. Hill* On the freeness of abelian groups: a generalization of Pontryagin's theorem, Bull. Amer. Math. Soc., 76 (1970), 1118–1120

[16] *P. C. Eklof* On the existence of κ-free abelian groups, Proc. Amer. Math. Soc., 47 (1975), 65–72

[17] *T. Mihara* Almost free non-Archimedean Banach spaces and relation to large cardinals, arXiv:2604.10526 (2026)

Norbert Nemesh, Faculty of Mechanics and Mathematics, Moscow State University, Moscow 119991 Russia

E-mail address: nemeshnorbert@yandex.ru