

1 Question

$$x \in R^n$$

$$\Delta\tau = \sum_{i=1}^n \frac{\partial^2 \tau}{\partial x_i^2}$$

$$\nabla\tau = \begin{bmatrix} \frac{\partial \tau}{\partial x_1} \\ \frac{\partial \tau}{\partial x_2} \\ \dots \\ \frac{\partial \tau}{\partial x_n} \end{bmatrix}$$

We know that

$$\frac{\partial \tau}{\partial t} + \frac{r^2}{2} \Delta\tau + b \nabla\tau = 0$$

and

$$v = -\ln \tau$$

2 Solution

We get the partial derivatives based on v in regards of different variables

$$\frac{\partial v}{\partial t} = -\frac{1}{\tau} \frac{\partial \tau}{\partial t}$$

$$\frac{\partial v}{\partial x} = -\frac{1}{\tau} \frac{\partial \tau}{\partial x}$$

$$\frac{\partial^2 v}{\partial x^2} = \frac{1}{\tau^2} \left(\frac{\partial \tau}{\partial x} \right)^2 - \frac{1}{\tau} \frac{\partial^2 \tau}{\partial x^2}$$

Therefore, we get

$$\sum_{i=1}^n \frac{\partial^2 v}{\partial x_i^2} = \sum_{i=1}^n \frac{1}{\tau^2} \left(\frac{\partial \tau}{\partial x_i} \right)^2 - \sum_{i=1}^n \frac{1}{\tau} \frac{\partial^2 \tau}{\partial x_i^2}$$

By the equation

$$\frac{\partial \tau}{\partial t} + \frac{r^2}{2} \Delta\tau + b \nabla\tau = 0$$

We get

$$-\tau \frac{\partial v}{\partial t} - \frac{r^2}{2} \tau \sum_{i=1}^n \left(\frac{\partial v}{\partial x_i} \right)^2 + \frac{r^2}{2} \tau \sum_{i=1}^n \frac{\partial^2 v}{\partial x_i^2} - \tau \sum_{i=1}^n b_i \frac{\partial v}{\partial x_i} = 0$$

By eliminating τ , we get the final equation based on the partial differential of v.

$$\frac{\partial v}{\partial t} + \frac{r^2}{2} \sum_{i=1}^n \left(\frac{\partial v}{\partial x_i} \right)^2 - \frac{r^2}{2} \sum_{i=1}^n \frac{\partial^2 v}{\partial x_i^2} + \sum_{i=1}^n b_i \frac{\partial v}{\partial x_i} = 0$$