

EVERT NASEDKIN

SIMULATED INSTRUMENTAL CONSTRAINTS
ON SUB-STELLAR ATMOSPHERIC
RETRIEVALS FOR THE JAMES WEBB SPACE
TELESCOPE'S MID-INFRARED
INSTRUMENT.

Evert Nasedkin

*Simulated Instrumental Constraints on Sub-Stellar Atmospheric Retrievals for the
James Webb Space Telescope's Mid-Infrared Instrument.*

Copyright © 2020

TITLEBACK

This document was written with \LaTeX on Ubuntu using ArsClassica, designed by André Miede.

CONTACTS

✉ evertn@student.ethz.ch



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Declaration of originality

The signed declaration of originality is a component of every semester paper, Bachelor's thesis, Master's thesis and any other degree paper undertaken during the course of studies, including the respective electronic versions.

Lecturers may also require a declaration of originality for other written papers compiled for their courses.

I hereby confirm that I am the sole author of the written work here enclosed and that I have compiled it in my own words. Parts excepted are corrections of form and content by the supervisor.

Title of work (in block letters):

Authored by (in block letters):

For papers written by groups the names of all authors are required.

Name(s):

First name(s):

With my signature I confirm that

- I have committed none of the forms of plagiarism described in the '[Citation etiquette](#)' information sheet.
- I have documented all methods, data and processes truthfully.
- I have not manipulated any data.
- I have mentioned all persons who were significant facilitators of the work.

I am aware that the work may be screened electronically for plagiarism.

Place, date

Signature(s)

For papers written by groups the names of all authors are required. Their signatures collectively guarantee the entire content of the written paper.

ACKNOWLEDGEMENTS

—

ABSTRACT

Following its launch in 2021, the James Webb Telescope will provide the best infrared observations of exoplanets and brown dwarfs to date. In particular, the Mid-Infrared Instrument (MIRI), will allow for medium resolution spectroscopy across a wide wavelength band, from 4.9-28.8 μm . This will allow us to derive atmospheric properties of objects at lower temperatures than currently possible. MIRI's medium resolution spectrometer (MRS) is an integrated field unit that will perform these observations, providing both spatial and spectral information about targets. Understanding the instrumental effects is critical to analyzing data from MIRI. With that in mind, the MIRISIM instrumental simulator was developed to provide observational simulations of the various sub instruments of MIRI.

This thesis improves the implementation of a thin-film fringing model for point sources to MIRISIM, considering how the fringing effect from the detector layers varies with position. Fringing is a periodic, wavelength dependent effect, and thus has a strong impact on any spectroscopic observations. A comparison to the existing model was made, demonstrating the necessity of considering this effect when analyzing data. We will improve the fringing removal by identifying the point source location from the constructed data cube, and select the correct fringe flat for removal.

Understanding the instrumental effects is key to quantifying the ability of MIRI to derive atmospheric properties. Existing literature has considered the NIRCAM instrument and the MIRI Low-Resolution Spectrometer, but to date no retrieval studies have been performed using MIRISIM, or for the MIRI MRS, though it is critical to extend wavelength coverage to improve the results of an atmospheric retrieval. Model atmospheres will be generated using PetitRadTrans, and processed using MIRISIM and the JWST pipeline to produce a mock observation. An atmospheric retrieval will be performed, demonstrating to what extent MIRI will be able to retrieve atmospheric parameters such as temperature, pressure and composition. The posterior distributions of these parameters are compared with and without the fringing removal, again demonstrating the importance of correcting for this effect.

CONTENTS

1	INTRODUCTION	1
1.1	Exoplanets	1
1.2	Brown Dwarfs	3
1.2.1	Physics	4
1.2.2	Observational Properties	4
1.3	Motivation	4
1.3.1	Current Status of Atmospheric Characterization	4
1.3.2	JWST Studies	4
1.3.3	Biosignatures and Future Missions	4
1.4	Thesis Overview	4
2	MIRI: THE MID-INFRARED INSTRUMENT	5
2.1	The James Webb Space Telescope	5
2.2	MIRI	5
2.3	The Medium Resolution Spectrograph	6
2.3.1	Coordinates	6
2.3.2	Integrated Field Spectroscopy	7
2.3.3	Optical Systems	7
2.3.4	Detectors	7
2.4	Observations	7
2.4.1	Dithering	7
2.4.2	Exposure time calculations	7
3	FRINGING EFFECTS IN MIRI	9
3.1	Fringing	9
3.2	MIRISIM	10
3.2.1	Architecture	10
3.2.2	Data Products	11
3.2.3	Instrumental effects	11
3.2.4	Fringing	11
3.2.5	Implementation	12
3.2.6	JWST Pipeline	12
3.2.7	Cross-Correlation	12
3.2.8	Residual Statistics	13
3.3	Fringing Results	13
3.3.1	Effects of fringing on spectral extraction	13
4	ATMOSPHERIC RETRIEVALS	15
4.1	Introduction	15
4.2	Atmospheric Modeling	15
4.2.1	petitRadTrans	15
4.3	Bayesian methods	15
4.3.1	MCMC	15
4.3.2	Nested Sampling	16
4.3.3	multinest	16
4.3.4	Prior choice	16
4.3.5	Bayesian Model Selection	16

4.4	Targets	16
4.4.1	Atmospheric Parameters	16
4.4.2	petitRadTrans	16
4.5	Results	16
4.5.1	Posterior Distributions	16
5	CONCLUSIONS	17
A	APPENDICES	19
A.1	Package Requirements	19
	WORKS CITED	21

Since the first detection of a planet around a sun-like star (Mayor and Queloz) the field of exoplanets has evolved rapidly. Thousands of companions have been identified using the radial velocity and transit detection methods, and a handful have been imaged directly using both ground and space based observatories. In the last decade, many advances have been made that allow us to begin to characterize the properties of a few of these planets using spectroscopy. With the launch of the James Webb Space Telescope (JWST) in 2021, and the dawn of the era of extremely large telescopes, we will be able to peer deeper into these planets and further constrain atmospheric or geological properties, allowing us to answer questions about their formation history, climate, and even the prospects for habitability and life.

JWST will operate in near to mid infrared wavelengths, which will provide a new window into studying the atmospheres of exoplanets and brown dwarfs. The Mid Infrared Instrument (MIRI) will provide unprecedented spectral resolution in the mid infrared, allowing for the measurement of composition, pressure and temperature. Novel instrumentation does not come without challenges. Optical and instrumental effects will constrain the ability to which we can measure spectral features, which will ultimately limit the science that can be accomplished.

In this thesis, we will measure the impact of thin-film fringing in the layers of the detectors in the MIRI Medium-Resolution Spectrometer on measurements of atmospheric parameters of brown dwarfs and exoplanets. This will provide a baseline for determining the level of correction necessary to minimize the impact of fringing, as well as providing a first look into the ability of the MRS to characterize atmospheres.

1.1 EXOPLANETS

The last quarter century of observations has revealed the diversity of exoplanets and extra-solar systems. Both the architecture and individual planetary characteristics vary greatly when compared to each other, as well as to our own solar system. From the hot Jupiters initially found by Mayor and Queloz (Mayor and Queloz) to the thousands of planets discovered by the Kepler mission, the variety in exoplanets has raised questions about their formation and development, as well as their present day structure, climate, and even prospects for life. Improvements to observational techniques have allowed us to improve our understanding of these planets. Secondary eclipse and transmission spectroscopy has opened the door to the study of planets in close orbits to their host stars, while emission spectroscopy of young planet has allowed for constraints on models of planet formation. Over the next decades, new instruments will be developed that improve sensitivity,

allowing us to study smaller, colder and fainter planets: with the ultimate goal of studying atmospheric and surface features of an earth-like planet.

Of particular interest are observable features that allow us to measure physical properties of exoplanets. The radial velocity (RV) method provides a measure of the planet mass, while a transit can constrain the radius. Already these properties tell us something about the overall structure of the planet. Spectroscopy can provide insight into the composition of the planet's atmosphere, as well as its temperature and pressure. These properties are linked to its age and location of formation in the circumstellar disk. The atmosphere, combined with the distance between the planet and its star determine the climate of the planet.

Direct Imaging

While the majority of exoplanet detections have been made using the radial velocity or transit techniques, direct imaging opens up the possibility of collecting light from the planet itself. This provides a window into the planet's atmosphere and surface. Most direct imaging to date has used near-to-mid infrared wavelengths, where the contrast between the thermal emission from the planet and the star is at a minimum. This has its drawbacks: we are so far only able to image young planets that have retained some of the heat from their formation.

Direct imaging can make use of both ground and space based observatories. However, the high spatial resolution required drives the need for a large primary mirror, limiting the possibilities of space-based telescopes. On the other hand, atmospheric turbulence necessitates the use of an adaptive optics equipped facility to observe from the ground. Atmospheric absorption due to telluric lines (absorption lines of Earth's atmosphere) also restrict infrared observations to narrow bands.

In addition to the requiring high spatial resolution, it is also challenging to separate the light emitted by the planet from that of the star. Imaging techniques such as Angular Differential Imaging (ADI) (Marois2007) and Reference Differential Imaging (RDI) (Lefreniere2009; Soummer2012) provide methods for reducing the stellar point-spread-function (PSF). Coronagraphs are optical elements which suppress the stellar PSF through self-destructive interference or physical occultation, depending on the position in the optical path. The difference in spectra between the planet and the star can also be used to separate the two sources.

Presently, 10m class telescopes such as the Very Large Telescope (VLT) in Paranal, Chile or the Gemini Observatory split between Hawaii and Chile provide the best combination of resolution and instrumentation to perform direct imaging of exoplanets. The NACO instrument at the VLT provided the first image of an exoplanet in 2004 (Chauvin et al.). These observatories are among those equipped with an adaptive optics system, coronagraphic instrumentation and near to mid infrared imaging and spectroscopic capabilities to directly image exoplanets, with several exemplar systems becoming standard objects of interest. The parameters of some directly imaged exoplanets are summarized in table 1

Name
2M1207b
β Pic b
HR8799b
HR8799c
HR8799d
HR8799e
Fomalhaut b
2M J044144b
LkCa 15b
HD 95086b
Gliese 504b
51 Eridani b
PDS 70b
PDS 70c

Table 1: Summary of directly imaged planet parameters [Bowler](#) and references therein.

(Beichman et al., “Direct Imaging and Spectroscopy of Exoplanets with the James Webb Space Telescope”) ([Lagage2015](#)) ([Macintosh2006](#)) ([Macintosh2015](#))

Beta Pictoris b

Discovered in 2008 ([Lagrange et al.](#)), β Pic b is one of the most well-studied exoplanet systems ([Hoeihmakers2018](#); [Quanz et al.](#) [Chilcote et al.](#) [Chilcote et al.](#)). Due to its young age of 23 ± 3 Myr ([Mamajek and Bell](#)) and high mass of $12.9 M_J$ ([Chilcote et al.](#)) it remains hot, between 1590K and 1847K ([Nowak2019](#)). Combined with its typically wide separation on its 9.2AU orbit ([Chauvin et al.](#)), this allows for relatively easy imaging and spectroscopy in the near infrared. Recent RV observations indicate the possibility of an additional planet at 2.3 AU ([Lagrange et al.](#)).

The quantity and quality of spectral data for β Pic b has allowed for the study and characterization of its atmosphere: measurements of its metallicity and C/O ratio can trace the formation history of the planet ([Nowak2019](#); [Chilcote et al.](#)). It also provides insight into its current atmospheric structure and composition, which can in turn inform models of exoplanet atmospheres.

HR 8799 System

PDS 70 b,c

1.2 BROWN DWARFS

([Oliveira](#)) ([Helling and Casewell](#)) ([Cooper2014](#)) ([Madhusudhan2018a](#)) ([Burrows2003](#)) ([Marley2014](#)) ([Manjavacas](#)) ([B. Biller](#)) ([Faherty2018](#)) ([Morley et al.](#))

1.2.1 Physics

1.2.2 Observational Properties

T-Type

L-Type

Y-Type

1.3 MOTIVATION

1.3.1 Current Status of Atmospheric Characterization

(Kreidberg)(B. A. Biller and Bonnefoy) (Bozza, Mancini, and Sozzetti) (Danielski et al.) (Madhusudhan et al.)

Transmission Spectroscopy

(Lee, Fletcher, and P. G. Irwin) (MacDonald and Madhusudhan) (Madhusudhan, Apai, and Gandhi)

Emission Spectroscopy

1.3.2 JWST Studies

(Beichman et al., “Direct Imaging and Spectroscopy of Exoplanets with the James Webb Space Telescope”)

1.3.3 Biosignatures and Future Missions

1.4 THESIS OVERVIEW

2

MIRI: THE MID-INFRARED INSTRUMENT

MIRI is an instrument that will provide unique capabilities for studying exoplanets and other cold and distant objects. This chapter will provide a detailed overview of the technical details and capabilities of the instrument. A complete description of MIRI is provided in (10.1086/6822554; Rieke et al., “The Mid-Infrared Instrument for the James Webb Space Telescope , I: Introduction” Wright et al. P. Lagage et al. Boccaletti et al. Wells et al. Rieke et al. Ressler et al. Glasse et al.)

2.1 THE JAMES WEBB SPACE TELESCOPE

JWST is a 6.5m space based observatory built in collaboration between NASA, ESA and CSA that will be located in a halo orbit at the L2 Earth-Sun Lagrange point. As the successor to the Hubble Space Telescope and the Spitzer Space Telescope, it will provide a new perspective for infrared astronomy. It is currently scheduled to launch in March 2021. It will operate at a temperature of 6.7K to reduce instrumental backgrounds over its wavelength range of 0.6-28.8 micron. A large sun-shield will help maintain the cold operating temperature of the telescope by blocking the solar infrared radiation.

There are four primary instruments that constitute the Integrated Science Instrument Module (ISIM). Near-Infrared Camera (NIRCam), which provides imaging with coronagraphic capabilities from 0.6-5 micron. The Near-Infrared Spectrograph (NIRSpec) provides fixed slit and integrated field unit spectroscopy capable of analyzing multiple objects simultaneously, and operates in the same wavelength range as NIRCam. The Fine Guidance Sensor/ Near-Infrared Imager and Slitless Spectrograph (FGS/NIRISS) allows for low and medium resolution spectroscopy with high photometric stability, as well as aperture masking interferometry. The final instrument, MIRI, is the subject of this thesis.

2.2 MIRI

The Mid-Infrared Instrument (MIRI) provides imaging, fixed slit and integrated field spectroscopy between 4.8 and 28 micron (Rieke2015).

(Kendrew2015)



Figure 1: The James Webb Space Telescope during integration of the telescope into the Spacecraft Element (**assembled**).

Subsystem
Imaging
4QPM Coronagraphic Imaging
Lyot Coronagraphic Imaging
Low Resolution Spectroscopy
Medium Resolution Spectroscopy

Table 2: Summary of MIRI observing modes.

2.3 THE MEDIUM RESOLUTION SPECTROGRAPH

The Medium Resolution Spectrograph (MRS) consists of four integrated field spectrographs projected onto two detectors, covering 4.8-28 micron with a spectral resolution varying from $R=1700$ to $R=3500$. (Wells2015)

2.3.1 Coordinates

There are three primary coordinate systems in use with JWST/MIRI-MRS, of which two will be relevant for this thesis, with the detector and local MRS coordinates described in Fig. 2 (Argyriou et al.).

The detector coordinate grid is formed by counting x/y pixels, as well as the slice number. Each of the two MRS detectors is an array of 1032×1024 pixels, though only 1024 are photosensitive in the horizontal direction. Each image slice from the IFU appears as a curved stripe on the detector, though neighboring stripes on the detector do not correspond to neighbouring slices of the image.

The local MRS coordinate system is described in terms of α , β and λ . The continuous α coordinate is the along slice direction, while beta is perpen-

Channel	Sub-band	Band	Detector	Wavelength Range [μm]	FOV [as]	$\lambda/\Delta\lambda$
1	Short	1A	SW	4.83 - 5.82	3.46×3.72	3500
	Medium	1B		5.62 - 6.73	3.46×3.72	3500
	Long	1C		6.46 - 7.76	3.41×3.72	3300
2	Short	2A	SW	7.44 - 8.90	4.16×4.76	3000
	Medium	2B		8.61 - 10.28	4.16×4.76	3000
	Long	2C		9.94 - 11.87	4.12×4.76	3000
3	Short	3A	LW	11.47 - 13.67	6.00×6.24	2700
	Medium	3B		13.25 - 15.80	5.96×6.24	2300
	Long	3C		15.30 - 18.24	5.91×6.24	2300
4	Short	3A	LW	17.54 - 21.10	7.14×7.87	1700
	Medium	3B		20.44 - 24.72	7.06×7.06	1700
	Long	3C		23.84 - 28.82	6.99×7.87	1500

Table 3: Properties of the MIRI MRS channels (Wells et al.).

dicular and discrete, corresponding to the slice number. λ is the wavelength. Both α and λ are fit by a second order polynomial to account for along and across slice distortion (Wells et al.). Each detector sub array has its own mapping to α, β, λ space, due to the differences in FoV, slice count, distortion and spectral resolution.

The third coordinate frame is the global coordinate system of JWST itself, V_1, V_2, V_3 . The V_1 coordinate refers to the symmetry axis of the telescope, V_3 points towards the foldable secondary mirror support structure strut. V_2 completes the coordinate system, being orthogonal to V_1 and V_3 . This coordinate system will not be used in this thesis.

2.3.2 Integrated Field Spectroscopy

2.3.3 Optical Systems

Channels, bands, etc ref:Chen2019

2.3.4 Detectors

Readout Modes

2.4 OBSERVATIONS

2.4.1 Dithering

2.4.2 Exposure time calculations




Figure 2: Description of the MRS detector (x, y, s) coordinate system to the local MRS (α, β, λ) cube coordinates. **Top:** Detector coordinates. Note that the consecutive stripe numbers s_i, s_{i+1} correspond to neighbouring image slices. **Bottom:** Description of the (invertible) detector-to-cube transformation (Argyriou et al.).

Figure 3: Detector images of a spatially and spectrally flat calibration source for the SW detector (left) and LW detector (right).

Understanding optical and instrumental effects is critical for creating accurate simulated observations and for characterizing systematics. These systematics and uncertainties in turn impact the potential science results from any instrument by biasing measurements, reducing the signal to noise ratio of measurements or by injecting non-physical signals and correlations. The aim of this chapter is to examine fringing in the MIRI detectors and how this effect is modeled in the instrumental simulator (MIRISIM).

3.1 FRINGING

Thin film interference occurs when light is coherently reflected at the boundary between two layers and interferes with the incident light. This is the principle on which Fabry-Pérot interferometers function. As we wish to determine the effect of fringing on the amplitude of the signal received by the detector, we are effectively interested in the transmittance of a series of Fabry-Pérot interferometers. Assuming an ideal plane-parallel optical cavity with a reflectance R at both boundaries, thickness D , and an angle θ at which the light travels within the cavity, we can compute the transmittance as:

$$T_c = \frac{1}{1 + \frac{4R}{(1-R)^2} \sin^2\left(\frac{\delta}{2}\right)} \quad (1)$$

Where the phase δ at half a wavelength ($\phi = \pi$), with wavenumber σ is:

$$\delta = 4\pi\sigma D \cos \theta - (\phi - \pi) \quad (2)$$

Systems with a spacing on the order of millimeters produces significant interference for infrared light (Lahuis and Boogert).

The detectors of the MRS consist of several layers, as shown in Fig. 4, with a characteristic thickness of $500\mu\text{m}$, which results in significant (10%-30%) ‘fringing’ in a spectrally flat signal - visible in Fig. 3. While this is typical for infrared detectors such as those in the Spitzer Space Telescope (Lahuis and Boogert) or in the Space Telescope Imaging Spectrograph on board HST (Malumuth2003), the sensitivity and spectral resolution of the MRS increase the significance of this issue. The MIRI consortium has stated that the error budget for all detector effects must be 3.3% or less. Present fringing corrections result in a 5% deviation from a photometrically accurate signal, and can introduce correlated noise which will degrade any measured spectrum. Therefore it is critical to examine the impact of fringing on a signal, the parameters that influence the fringing strength and phase, and possible solutions for fringe correction. While a more complete treatment of proposed fringe correction can be found in (Argyriou2020), this work will examine the implementation of fringing into the MIRI instrumental simulator and




Figure 4: Layers of the MIRI MRS detectors. Note that thicknesses are not to scale (Rieke et al.).

address the current state of fringe correction in the JWST Data Calibration Pipeline.

Lahuis and Muller

3.2 MIRISIM

Consortium2018 Cossou2019 The MIRI instrument has been modeled in python as a program known as MIRISIM. This program takes in an astronomical ‘scene’ along with some configuration parameters to output a detector data product, similar to what will be produced by the actual instrument. This is relatively full-featured simulator, modelling the instrumental PSF, various noise sources and distortion maps, among other effects. While MIRISIM is functional for all of the MIRI sub-instruments, this report will only deal with the Medium-Resolution Spectrometer (MRS) sub instrument, described in section 2.3. The goal of this section is to describe the implementation and testing of an updated optical model of the ‘fringing’ effect - an optical effect caused by thin film interference from the multiple layers of the detector.

3.2.1 Architecture

SCENE - SEDs SIMULATOR PYSPECSIM

Figure 5

Table 4

3.2.2 Data Products

3.2.3 Instrumental effects

3.2.4 Fringing

Lahuis and Boogert VanderPlas A key effect on spectral data is fringing, described in **ref:Argyriou2018**. MIRI uses a total of three Si:As impurity band conduction detector arrays, two of which are used by the MRS. These detectors consist of 7 layers, listed in table 4 and illustrated in Fig. 4.

If all of the parameters were known, this would be sufficient to numerically solve for the fringing pattern within MIRI. Unfortunately, uncertainties in the thickness in the detector layers, variations in the layer deposition thickness, and the uncertainty of transmittance and reflectance of the materials used at cryogenic temperatures prevents the implementation of such a numerical model.

Instead, we turn to calibration data taken to characterize the fringing pattern.

DESCRIBE CURRENT MODEL - GENERIC FRINGING

However, due to the dependance of fringing on the incident angle of the light, a single model of fringing is insufficient to describe the full effect. Therefore, we use data taken in XXXXXXXXXX at various points across the detector and quantify how this changes the extracted spectra after processing in the JWST pipeline.

DESCRIBE HOW THE DATA WAS TAKEN HERE. - Problems with point vs extended sources - multiple collection runs

Ultimately this data collection produced a series of 'fringe-flats' of an almost point like at various position across the detector and in each channel. We implemented a new routine into the pySpecSim portion of MIRISIM to read in the location of point sources within a scene, and apply the correct position dependent fringe flat. This implementation comes with several caveats: namely that the fringing model is not yet fully developed, so it can only be considered accurate for point sources located at the same (α, β) location as the source used to produce the fringe flat. Additionally, the source used to generate the data is not a true point source, nor are there fringe flats produced for the full MRS wavelength range. We stress that the goal of this testing is to demonstrate the significance of this effect to justify the need for a more complete model along with additional calibration data to constrain the detector layer parameters.

FM Data

CV Data

3.2.5 Implementation

3.2.6 JWST Pipeline

[Bushouse, Droettboom, and Greenfield Labiano-Ortega2016](#)

Stage 1 Processing

Stage 2 Processing

Fringing correction

[Carnall](#)

Aperture Photometry

3.2.7 Cross-Correlation

[I. Snellen et al.](#)[Simkin Tonry1979](#) [Petermann Bodis](#) To quantify the similarity of the spectrum output by the JWST pipeline to the input into MIRISIM, we rely on the technique of cross correlation. For two arbitrary, complex-valued functions $f(t)$ and $g(t)$, we can compute the cross correlation as a function of the shift τ between the functions (typically in time or velocity space):

$$(f \star g)(\tau) \equiv \int_{-\infty}^{\infty} f^*(t)g(t + \tau)dt \quad (3)$$

Our signals of interest are astrophysical spectra, measured in a finite number of discrete wavelength bins. For such a signal with M bins:

$$(f \star g)[n] \equiv \sum_{m=0}^M f^*[m]g[m + n] \quad (4)$$

Care must be taken when cross-correlating signals, as differences in normalization can result in changes in the correlation coefficient. Our procedure takes in two spectra. The first is an emission spectrum produced by the petiRadTrans program [Mollière et al.](#), which provides our forward model with which we compare our data spectrum. Our data is the result of passing the template spectrum through MIRISIM, and extracting it from the resulting detector image using the JWST pipeline. We then rebin the high-resolution input spectrum to the same wavelength bins as the data spectrum, using the `spectres` package. Prior to normalization, we remove any outliers from the spectrum (due to binning errors or instrumental effects) by setting any data points separater by more than 15 standard deviations from the mean to the median value of the spectrum. For each spectrum, we subtract the minimum value to remove any offset in the spectrum, and divide by the maximum value to restrict the range to $[0,1]$. We then use apply a Savitzky-Golay filter with a window of $1/4$ the length of the spectrum and a polynomial order of 3, which we then subtract from the unfiltered spectrum. This removes the

continuum emission from the spectrum, and centers it around 0. We then renormalize the spectrum by dividing by the maximum absolute value such that the range is in $[-1,1]$. The cross correlation between the forward model and itself is computed, excluding the region of interest around 0 offset. This 'autocorrelation' is subtracted from the cross correlation between the forward model and the data spectrum in order to remove secondary peaks. Finally, we normalize the cross correlation by the standard deviation of the cross correlation (excluding the central peak), giving an output measured as a signal to noise ratio.

3.2.8 Residual Statistics

In addition to computing the cross correlation between the forward model and the data spectrum, we also examine the residuals between the two spectra. Here we can see any unexpected variations between the two (periodic signals, offsets or other features). We can also examine a histogram of the residuals, normalized by the standard deviation of the data spectrum. This provides us with a distribution which should have a mean of 0 and unit width if the data are unbiased and share a distribution with the true input spectrum.

3.3 FRINGING RESULTS

1. A stronger input signal results in a stronger correlation.
2. Fringing does NOT necessarily degrade the cross correlation SNR, but rather increases it. The scale of this increase seems to depend on the absolute magnitude of the correlation (ie, a larger increase at higher SNR)
3. The residuals from subtracting the template from the data has structure.
4. If the residuals are histogrammed (and normalized by the standard deviation of the data), the width of the distribution may correspond to the cross correlation SNR (wider distribution = lower SNR)
5. Only when strongly increasing the fringing effect does the SNR decrease.
6. Correcting for fringing using the standard JWST fringe map decreases the SNR when compared to the case of fringing with no correction, but is typically still above the no-fringing case.
7. The JWST correction performs worse in the off axis case, as the fringe pattern begins to vary more when compared to the CV fringing model.

3.3.1 Effects of fringing on spectral extraction

4

ATMOSPHERIC RETRIVALS

Schlawin et al. Madhusudhan, “Atmospheric Retrieval of Exoplanets” Fisher et al. Oreshenko2019 Barman et al. Benneke2013 Benneke and Seager Blanco-Cuaresma2018 Konopacky et al. Morley2018 Lupu et al. Gandhi and Madhusudhan Baudino et al. Line et al. Madhusudhan2018b P. G. Irwin et al. Robinson, Stapelfeldt, and Marley Waldmann et al. Waldmann et al. Line et al. Line et al. Zalesky et al.

4.1 INTRODUCTION

Batalha et al. Feng et al. Mollière et al.

4.2 ATMOSPHERIC MODELING

Behmard, Petigura, and Howard Guillot Chen et al. Mollière et al. M. Zhang et al.

4.2.1 petitRadTrans

Mollière et al.

Radiative Transfer

Line-by-line

Correlated K

Goody et al. Lacis and Oinas

Clouds

Line and Parmentier Faherty2018 Morley et al.

4.3 BAYESIAN METHODS

Lavie et al.

4.3.1 MCMC

Foreman-Mackey et al. Speagle MacKay2003

4.3.2 Nested Sampling

[Skilling](#) [Feroz2007](#) [Feroz2008](#) [Feroz2014](#) [Feroz et al.](#)

4.3.3 multineest

Hyperparameters

4.3.4 Prior choice

4.3.5 Bayesian Model Selection

4.4 TARGETS

4.4.1 Atmospheric Parameters

Madhusudhan, “C/O ratio as a dimension for characterizing exoplanetary atmospheres”[Moses2012](#)[Garland and P. G. J. Irwin](#) [Bowler](#) [Fegley](#) and [Lodders](#) [Tokunaga](#)

4.4.2 petitRadTrans

4.5 RESULTS

4.5.1 Posterior Distributions

5 | CONCLUSIONS

A | APPENDICES

A.1 PACKAGE REQUIREMENTS

```
numpy=1.4.0  
scipy=1.5
```


WORKS CITED

- Argyriou, Ioannis, Ruymán Azzollini, and Bart Vandenbussche. "Spectrum extraction from detector plane images for the medium-resolution spectrometer of the mid-Infrared Instrument on-board the James Webb Space Telescope." July 2018 (2018): 124. Print.
- Argyriou, Ioannis, Bart Vandenbussche, and Martyn Wells. "Fringing solution for the mid-infrared instrument on-board the James Webb Space Telescope". July 2018 (2018): 127. Print.
- Argyriou, Ioannis, et al. "Nature of point source fringes in mid-infrared spectra acquired with the James Webb Space Telescope". *Astronomy & Astrophysics* (): 1–13. Print.
- Barman, Travis S., et al. "Simultaneous detection of water, methane, and carbon monoxide in the atmosphere of Exoplanet hr 8799 b". *Astrophysical Journal* 804.1 (2015): 1–10. arXiv: [1503.03539](#). Web.
- Batalha, Natasha E., et al. "Strategies for Constraining the Atmospheres of Temperate Terrestrial Planets with JWST ". *The Astrophysical Journal* 856.2 (2018): L34. arXiv: [1803.07983](#). Web.
- Baudino, Jean-loup, et al. "Toward the Analysis of JWST Exoplanet Spectra : Identifying Troublesome Model Parameters". *The Astrophysical Journal* 850.2 (2017): 150. Web.
- Behmard, Aida, Erik A. Petigura, and Andrew W. Howard. "Data-driven Spectroscopy of Cool Stars at High Spectral Resolution". *The Astrophysical Journal* 876.1 (2019): 68. arXiv: [arXiv:1904.00094v1](#).
- Beichman, Charles A., Pierre-olivier Lagage, and Marie Ygouf. "1188 - Spectroscopy of Young , Widely Separated Exoplanets". 4. Print.
- Beichman, Charles A., et al. "1194 - Characterization of the HR 8799 planetary system and planet search". Print.
- Beichman, Charles A and Thomas P Greene. "Observing Exoplanets with the James Webb Space Telescope". *Handbook of Exoplanets* 2009 (2018): 1283–1308. Print.
- Beichman, Charles A, et al. "Astro2020 Science White Paper: Direct Imaging and Spectroscopy of Exoplanets with the James Webb Space Telescope" (2019). Web.
- . "Direct Imaging and Spectroscopy of Exoplanets with the James Webb Space Telescope". *Astro2020 Science White Paper* (2019). Web.
- Benneke, Bjoern and Sara Seager. "Atmospheric retrieval for super-earths: Uniquely constraining the atmospheric composition with transmission spectroscopy". *Astrophysical Journal* 753.2 (2012). arXiv: [1203.4018](#).
- Biller, Beth. "The time domain for brown dwarfs and directly imaged giant exoplanets: the power of variability monitoring". *Astronomical Review* 13.1 (2017): 1–27. Web.
- Biller, Beth A. and Mickaël Bonnefoy. "Exoplanet Atmosphere Measurements from Direct Imaging". *Handbook of Exoplanets* (2018): 2107–2135. arXiv: [arXiv:1807.05136v1](#).

- Blanco-cuaresma, Sergi. "Modern stellar spectroscopy caveats". *Monthly Notices of the Royal Astronomical Society* 486.2 (2019): 2075–2101. arXiv: [arXiv:1902.09558v2](#).
- Boccaletti, A., et al. "The Mid-Infrared Instrument for the James Webb Space Telescope , V: Predicted Performance of the MIRI Coronagraphs". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 633–645. arXiv: [1508.02352](#).
- Bodis, Lorant. "Quantification of spectral similarity towards automatic spectra verification". Diss. ETH Zurich, 2007. [Web](#).
- Bouchet, Patrice, et al. "The Mid-Infrared Instrument for the James Webb Space Telescope, III: MIRIM, The MIRI Imager". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 612–622. [Web](#).
- Bowler, Brendan P. "Imaging Extrasolar Giant Planets". *Publications of the Astronomical Society of the Pacific* 128.968 (2016): 1–38. [Web](#).
- Bozza, Valerio, Luigi Mancini, and Alessandro Sozzetti. *Astrophysics of Exoplanetary Atmospheres*. Print.
- Bushouse, Howard, Michael Droettboom, and Perry Greenfield. "The James Webb Space Telescope Data Calibration Pipeline". *14th Python in Science Conf*. 2015. 289. arXiv: [0606175 \[astro-ph\]](#).
- Carnall, A. C. "SpectRes: A Fast Spectral Resampling Tool in Python". 2 (2017): 1–4. arXiv: [1705.05165](#). [Web](#).
- Chauvin, G., et al. "A giant planet candidate near a young brown dwarf Direct VLT/NACO observations using IR wavefront sensing". *Astronomy and Astrophysics* 425.2 (2004): 29–32. Print.
- Chauvin, G., et al. "Orbital characterization of the β Pictoris b giant planet". *Astronomy and Astrophysics* 542 (2012): 1–9. arXiv: [1202.2655](#).
- Chen, Howard, et al. "Habitability and Spectroscopic Observability of Warm M-dwarf Exoplanets Evaluated with a 3D Chemistry-Climate Model" (2019). arXiv: [1907.10048](#). [Web](#).
- Chilcote, Jeffrey, et al. "1–2.4 μ m Near-IR Spectrum of the Giant Planet β Pictoris b Obtained with the Gemini Planet Imager ". *The Astronomical Journal* 153.4 (2017): 182. arXiv: [1703.00011](#). [Web](#).
- Chilcote, Jeffrey, et al. "The first H-band spectrum of the giant planet β Pictoris b". *Astrophysical Journal Letters* 798.1 (2015): 1–5. Print.
- Christiaens, Valentin, et al. "Evidence for a Circumplanetary Disk around Protoplanet PDS 70 b". *The Astrophysical Journal* 877.2 (2019): L33. arXiv: [1905.06370](#). [Web](#).
- Citation, Suggested. *Exoplanet science strategy*. 2019. 1–172. Print.
- Consortium, Miri European. "MIRISim Documentation" (2018). Print.
- Cooper, Curtis S, et al. "Modeling the Formation of Clouds in Brown Dwarf Atmospheres". *The Astrophysical Journal* 156.2 (2003): 1320–1337. arXiv: [0205192v3 \[arXiv:astro-ph\]](#).
- Cossou, Christophe. "MIRISim Spreadsheet and Quick examples" (2018): 1–12. Print.
- Danielski, Camilla, et al. " Atmospheric Characterization of Directly Imaged Exoplanets with JWST /MIRI ". *The Astronomical Journal* 156.6 (2018): 276. arXiv: [1810.00894](#).

- Faherty, Jacqueline K, et al. "Indications of water clouds in the coldest known brown dwarf". *The Astrophysical Journal Letters* 793.1 (2014). arXiv: [arXiv:1408.4671v2](#).
- Fegley, Bruce and Katharina Lodders. "Chemical Models of the Deep Atmospheres of Jupiter and Saturn". *Icarus* 110 (1994): 117–154. Print.
- Feng, Y. Katherina, et al. "Characterizing Earth Analogs in Reflected Light: Atmospheric Retrieval Studies for Future Space Telescopes". *The Astronomical Journal* 155.5 (2018): 200. [Web](#).
- Feroz, F., M. P. Hobson, and M. Bridges. "MultiNest: An efficient and robust Bayesian inference tool for cosmology and particle physics". *Monthly Notices of the Royal Astronomical Society* 398.4 (2009): 1601–1614. arXiv: [0809.3437](#).
- Feroz, Farhan, et al. "Importance Nested Sampling and the MultiNest Algorithm". *The Open Journal of Astrophysics* 2.1 (2019): 1–17. arXiv: [1306.2144](#).
- Fisher, Chloe, et al. "Interpreting High-Resolution Spectroscopy of Exoplanets Using Cross-Correlations and Supervised Machine Learning" (2019): 1–15. arXiv: [1910.11627](#). [Web](#).
- Foreman-Mackey, Daniel, et al. "emcee : The MCMC Hammer ". *Publications of the Astronomical Society of the Pacific* 125.925 (2013): 306–312. arXiv: [1202.3665](#).
- Gandhi, Siddharth and Nikku Madhusudhan. "Retrieval of exoplanet emission spectra with HyDRA". *Monthly Notices of the Royal Astronomical Society* 474.1 (2018): 271–278. arXiv: [1710.06433](#).
- Garland, R. and P. G. J. Irwin. "Effectively Calculating Gaseous Absorption in Radiative Transfer Models of Exoplanetary and Brown Dwarf Atmospheres". March (2019). arXiv: [1903.03997](#). [Web](#).
- Glasse, Alistair, et al. "The Mid-Infrared Instrument for the James Webb Space Telescope , IX: Predicted Sensitivity". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 686–695. arXiv: [1508.02427](#).
- Goody, Richard, et al. "The correlated-k method for radiation calculations in nonhomogeneous atmospheres". *Journal of Quantitative Spectroscopy and Radiative Transfer* 42.6 (1989): 539–550. Print.
- GRAVITY Collaboration, et al. "Peering into the formation history of beta Pictoris b with VLTI/GRAVITY long baseline interferometry". 2011 (2019): 1–21. arXiv: [1912.04651](#). [Web](#).
- Greenbaum, Alexandra Z., et al. " GPI Spectra of HR 8799 c, d, and e from 1.5 to 2.4 μ m with KLIP Forward Modeling ". *The Astronomical Journal* 155.6 (2018): 226. arXiv: [1804.07774](#).
- Guillot, T. "On the radiative equilibrium of irradiated planetary atmospheres". *Astronomy and Astrophysics* 520.18 (2010): 1–13. arXiv: [1006.4702](#).
- Haffert, S. Y., et al. "Two accreting protoplanets around the young star PDS 70". *Nature Astronomy* 3.8 (2019): 749–754. [Web](#).
- Helling, Christiane and Sarah Casewell. "Atmospheres of Brown Dwarfs". *The Astronomy and Astrophysics Review* 22 (2014): 1–53. arXiv: [arXiv:1410.6029v2](#).
- Hoeijmakers, H. J., et al. "Medium-resolution integral-field spectroscopy for high-contrast exoplanet imaging: Molecule maps of the β Pictoris system

- with SINFONI". *Astronomy and Astrophysics* 617 (2018): 1–11. arXiv: [1802.09721](#).
- Irwin, P. G.J., et al. "The NEMESIS planetary atmosphere radiative transfer and retrieval tool". *Journal of Quantitative Spectroscopy and Radiative Transfer* 109.6 (2008): 1136–1150. Print.
- Keppler, M., et al. "Discovery of a planetary-mass companion within the gap of the transition disk around PDS 70". *Astronomy and Astrophysics* 617 (2018): 1–21. arXiv: [1806.11568](#).
- Konopacky, Quinn M., et al. "Detection of carbon monoxide and water absorption lines in an exoplanet atmosphere". *Science* 339.6126 (2013): 1398–1401. Print.
- Kreidberg, Laura. "Exoplanet Atmosphere Measurements from Transmission Spectroscopy and Other Planet Star Combined Light Observations". *Handbook of Exoplanets* (2018): 2083–2105. arXiv: [arXiv:1709.05941v2](#).
- Labiano-Ortega, Alvaro, et al. "The MIRI Medium Resolution Spectrometer calibration pipeline". *exposure 1* (2016): 117. Print.
- Lacis, A. A. and V. Oinas. "A description of the correlated k distribution method for modeling nongray gaseous absorption, thermal emission, and multiple scattering in vertically inhomogeneous atmospheres". *Journal of Geophysical Research* 96.D5 (1991): 9027–9063. Print.
- Lacour, S., et al. "First direct detection of an exoplanet by optical interferometry". *Astronomy & Astrophysics* 623 (2019): L11. Print.
- Lafrenière, David, et al. "HST/nicmos detection of HR 8799 B in 1998". *Astrophysical Journal* 694.2 (2009): 148–152. Print.
- Lagage, Pierre-olivier and David Barrado. "1278 - MIRI Spectroscopic Observations of Brown Dwarfs". Print.
- Lagage, Pierre-olivier, J. Bouwman, and Fred Lahuis. "1275 - Spectroscopic characterization of PSO J318". Print.
- Lagage, Pierre-olivier and Thomas L Roellig. "1276 - Spectroscopic Observations of WD 0806-661B". Print.
- Lagage, P, et al. "The Mid-Infrared Instrument for the James Webb Space Telescope , IV : The Low Resolution Spectrometer". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 1–23. arXiv: [1512.03000](#).
- Lagrange, A. M., et al. "A probable giant planet imaged in the β Pictoris disk* VLT/NaCo deep L'-band imaging". *Astronomy and Astrophysics* 493.2 (2009): 21–25. Print.
- Lagrange, A. M., et al. "Evidence for an additional planet in the β Pictoris system". *Nature Astronomy* 3.12 (2019): 1135–1142. [Web](#).
- Lahuis, F and A Boogert. "How to Get Rid of Fringes in SIRTf/{IR}S Data". *SFChem 2002: Chemistry as a Diagnostic of Star Formation* January 2003 (2003): 335. Print.
- Lavie, Baptiste, et al. " HELIOS-RETRIEVAL: An Open-source, Nested Sampling Atmospheric Retrieval Code; Application to the HR 8799 Exoplanets and Inferred Constraints for Planet Formation ". *The Astronomical Journal* 154.3 (2017): 91. arXiv: [1610.03216](#).
- Lee, J. M., L. N. Fletcher, and P. G.J. Irwin. "Optimal estimation retrievals of the atmospheric structure and composition of HD189733b from sec-

- ondary eclipse spectroscopy". *Monthly Notices of the Royal Astronomical Society* 420.1 (2012): 170–182. arXiv: [1110.2934](#).
- Li, Jianfeng, et al. "A comparative study of point-to-point algorithms for matching spectra". *Chemometrics and Intelligent Laboratory Systems* 82.1-2 SPEC. ISS (2006): 50–58. Print.
- Line, Michael R. and Vivien Parmentier. "the Influence of Nonuniform Cloud Cover on Transit Transmission Spectra". *The Astrophysical Journal* 820.1 (2016): 78. arXiv: [1511.09443](#). [Web](#).
- Line, Michael R., et al. "A systematic retrieval analysis of secondary eclipse spectra. I. A comparison of atmospheric retrieval techniques". *Astrophysical Journal* 775.2 (2013). arXiv: [1304.5561](#).
- Line, Michael R., et al. "UNIFORM ATMOSPHERIC RETRIEVAL ANALYSIS of ULTRACOOOL DWARFS. I. CHARACTERIZING BENCHMARKS, GJ 570D and HD 3651B". *Astrophysical Journal* 807.2 (2015): 183. arXiv: [1504.06670](#). [Web](#).
- Line, Michael R., et al. "Uniform Atmospheric Retrieval Analysis of Ultracool Dwarfs. II. Properties of 11 T dwarfs". *The Astrophysical Journal* 848.2 (2017): 83. arXiv: [1612.02809](#).
- Lupu, Roxana E, et al. "DEVELOPING ATMOSPHERIC RETRIEVAL METHODS FOR DIRECT IMAGING SPECTROSCOPY OF GAS GIANTS IN REFLECTED LIGHT . I . METHANE ABUNDANCES AND BASIC CLOUD PROPERTIES". *The Astronomical Journal* 10.October (2018): 1–29. arXiv: [arXiv:1306.2144v2](#). [Web](#).
- MacDonald, Ryan J. and Nikku Madhusudhan. "HD 209458b in new light: evidence of nitrogen chemistry, patchy clouds and sub-solar water". *Monthly Notices of the Royal Astronomical Society* 469.2 (2017): 1979–1996. arXiv: [1701.01113](#).
- Madhusudhan, N and S Seager. "A TEMPERATURE AND ABUNDANCE RETRIEVAL METHOD FOR EXOPLANET ATMOSPHERES". *The Astrophysical Journal* 707.1 (2009): 24–39. arXiv: [arXiv:0910.1347v2](#).
- Madhusudhan, Nikku. "Atmospheric Retrieval of Exoplanets". *Handbook of Exoplanets* (2018): 2153–2182. arXiv: [1808.04824](#).
- . "C/O ratio as a dimension for characterizing exoplanetary atmospheres". *Astrophysical Journal* 758.1 (2012). arXiv: [1209.2412](#).
- Madhusudhan, Nikku, et al. "Exoplanetary Atmospheres—Chemistry, Formation Conditions, and Habitability". *Space Science Reviews* 205.1-4 (2016): 285–348. arXiv: [arXiv:1604.06092v1](#).
- Madhusudhan, Nikku, et al. "High C / O Ratio and Weak Thermal Inversion in the Very Hot Atmosphere of Exoplanet WASP-12b". *Nature* 469.7328 (2011): 64–67. arXiv: [arXiv:1012.1603](#).
- Mamajek, Eric E. and Cameron P.M. Bell. "On the age of the β pictoris moving group". *Monthly Notices of the Royal Astronomical Society* 445.3 (2014): 2169–2180. Print.
- Manjavacas, Elena. "Physical characterization of brown dwarfs". Diss. Ruperto-Carola-University of Heidelberg, 2014. Print.
- Marley, M S and T D Robinson. "On the Cool Side : Modeling the Atmospheres of Brown Dwarfs and Giant Planets arXiv : 1410 . 6512v1 [astro-ph

- . EP] 23 Oct 2014". *Annual Review of Astronomy and Astrophysics* 53 (2015): 279–323. arXiv: [arXiv:1410.6512v1](#).
- Marois, Christian, et al. "ANGULAR DIFFERENTIAL IMAGING : A POWERFUL HIGH-CONTRAST IMAGING TECHNIQUE 1". *The Astrophysical Journal* 641 (2006): 556–564. Print.
- Marois, Christian, et al. "Direct imaging of multiple planets orbiting the star HR 8799". *Science* 322.5906 (2008): 1348–1352. arXiv: [0811.2606](#).
- Mayor, M and D Queloz. "A Jupiter-mass companion to a solar-type star". *Nature* 378 (1995): 667–668. Print.
- Mesa, D., et al. "VLT/SPHERE exploration of the young multiplanetary system PDS70". *Astronomy and Astrophysics* 632 (2019): 1–12. arXiv: [1910.11169](#).
- Molliere, P. and I. A.G. Snellen. "Detecting isotopologues in exoplanet atmospheres using ground-based high-dispersion spectroscopy". *Astronomy and Astrophysics* 622 (2019). arXiv: [1809.01156](#).
- Mollière, P., et al. "petitRADTRANS: a Python radiative transfer package for exoplanet characterization and retrieval". 67 (2019): 1–17. arXiv: [1904.11504](#). Web.
- Mollière, P, et al. "MODEL ATMOSPHERES OF IRRADIATED EXOPLANETS : THE INFLUENCE OF STELLAR PARAMETERS , METALLICITY , AND THE C / O RATIO". *The Astrophysical Journal* 813.1 (2015): 47. Web.
- Morley, Caroline V., et al. "Water clouds in y dwarfs and exoplanets". *Astrophysical Journal* 787.1 (2014). arXiv: [1404.0005](#).
- Moroz, V.I. "On the infrared spectra of Jupiter and Saturn (0.9-2.5um)". *Soviet Astronomy* 5.6 (1962). Print.
- Moses, J. I., et al. "Chemical consequences of the C/O ratio on hot jupiters: Examples from WASP-12b, CoRoT-2b, XO-1b, and HD 189733b". *Astrophysical Journal* 763.1 (2013). arXiv: [1211.2996](#).
- Oreshenko, Maria, et al. "Supervised Machine Learning for Inter-comparison of Model Grids of Brown Dwarfs: Application to GJ 570D and the Epsilon Indi B Binary System". *The Astronomical Journal* 159.1 (2020): 15. arXiv: [arXiv:1910.11795v1](#).
- Petermann, R. "Atmospheric characterization of terrestrial planets using high-resolution spectroscopy with ELT / METIS A study of Proxima Cen b". Diss. ETH Zurich, 2019. Print.
- Quanz, Sascha P., et al. "First results from very large telescope NACO apodizing phase plate: 4 μ m images of the exoplanet β Pictoris b". *Astrophysical Journal Letters* 722.1 PART 2 (2010): 49–53. Print.
- Ressler, M. E., et al. "The Mid-Infrared Instrument for the James Webb Space Telescope , VIII: The MIRI Focal Plane System". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 675–685. Print.
- Rieke, G. H., et al. "The Mid-Infrared Instrument for the James Webb Space Telescope , I: Introduction". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 584–594. Print.
- . "The Mid-Infrared Instrument for the James Webb Space Telescope , I: Introduction ". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 584–594. Print.

- Rieke, G. H., et al. "The Mid-Infrared Instrument for the James Webb Space Telescope , VII: The MIRI Detectors". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 665–674. Print.
- Robinson, Tyler D., Karl R. Stapelfeldt, and Mark S. Marley. "Characterizing rocky and gaseous exoplanets with 2m class space-based coronagraphs". *Publications of the Astronomical Society of the Pacific* 128.960 (2016): 1–22. [Web](#).
- Schlawin, Everett, et al. "Clear and Cloudy Exoplanet Forecasts for JWST : Maps, Retrieved Composition, and Constraints on Formation with MIRI and NIRC2". *The Astronomical Journal* 156.1 (2018): 40. arXiv: [1803.08173](#).
- Seager, Sara. "Bayesian Atmospheric Retrieval for Exoplanets" (2013). Print.
- Simkin, Susan M. "Measurements of Velocity Dispersions and Doppler Shifts from Digitized Optical Spectra". *Astronomy & Astrophysics* 32 (1974): 129–136. Print.
- Skilling, John. "Nested Sampling". 395.November 2004 (2004): 395–405. Print.
- Snellen, I., et al. "Combining high-dispersion spectroscopy with high contrast imaging: Probing rocky planets around our nearest neighbors". *Astronomy and Astrophysics* 576.3 (2015). arXiv: [1503.01136](#).
- Snellen, Ignas, et al. "The fast spin-rotation of a young extrasolar planet". *Nature* 509.7498 (2014): 63–65. arXiv: [1404.7506 \[astro-ph.EP\]](#). [Web](#).
- Soummer, Rémi, et al. "Orbital motion of HR 8799 b, c, d using hubble space telescope data from 1998: Constraints on inclination, eccentricity, and stability". *Astrophysical Journal* 741.1 (2011). arXiv: [1110.1382](#).
- Speagle, Joshua S. "A Conceptual Introduction to Markov Chain Monte Carlo Methods" (2019). arXiv: [1909.12313](#). [Web](#).
- Tokunaga, Alan T. "HIGH-RESOLUTION INFRARED SPECTROSCOPY OF PLANETARY ATMOSPHERES". *Publications of the Astronomical Society of the Pacific* 95.October (1983): 691–699. Print.
- VanderPlas, Jacob T. "Understanding the Lomb–Scargle Periodogram". *The Astrophysical Journal Supplement Series* 236.1 (2018): 16. arXiv: [1703.09824](#). [Web](#).
- Waldmann, I. P., et al. "T-REx. II. RETRIEVAL of EMISSION SPECTRA". *Astrophysical Journal* 813.1 (2015): 13. arXiv: [1508.07591](#). [Web](#).
- Waldmann, I. P., et al. "Tau-REx I: A next generation retrieval code for exoplanetary atmospheres". *Astrophysical Journal* 802.2 (2015): 107. arXiv: [1409.2312](#). [Web](#).
- Wells, Martyn, et al. "The Mid-Infrared Instrument for the James Webb Space Telescope , VI: The Medium Resolution Spectrometer". *Publications of the Astronomical Society of the Pacific* 127.953 (2015): 646–664. arXiv: [1508.03070](#).
- Wright, G S, et al. "The Mid-Infrared Instrument for JWST , II : Design and Build" (). 1–44. Print.
- Zalesky, Joseph A., et al. "A Uniform Retrieval Analysis of Ultra-cool Dwarfs. III. Properties of Y Dwarfs". *The Astrophysical Journal* 877.1 (2019): 24. arXiv: [1903.11658](#).
- Zhang, Le and Raja R.A. Issa. *Development of IFC-based construction industry ontology for information retrieval from IFC models*. 2014. Print.

- Zhang, Michael, et al. "Forward modeling and retrievals with PLATON, a fast open-source tool". *Publications of the Astronomical Society of the Pacific* 131.997 (2019). arXiv: [1811.11761](#).
- Zucker, S. "Cross-correlation and maximum-likelihood analysis : a new approach to combining cross-correlation functions 2 F R O M M A X I M U M L I K E L I H O O D T O C R O S S - C O R R E L A T I O N 1292". *Monthly Notices of the Royal Astronomical Society* 342 (2003): 1291–1298. Print.
- Zurlo, A., et al. "First light of the VLT planet finder SPHERE: III. New spectrophotometry and astrometry of the HR 8799 exoplanetary system". *Astronomy and Astrophysics* 587 (2016): 1–13. arXiv: [1511.04083](#).