

EVERT NASEDKIN

SIMULATED INSTRUMENTAL CONSTRAINTS
ON SUB-STELLAR ATMOSPHERIC
RETRIEVALS FOR THE JAMES WEBB SPACE
TELESCOPE'S MID-INFRARED
INSTRUMENT.

Evert Nasedkin

*Simulated Instrumental Constraints on Sub-Stellar Atmospheric Retrievals for the
James Webb Space Telescope's Mid-Infrared Instrument.*

Copyright © 2020

TITLEBACK

This document was written with \LaTeX on Ubuntu using ArsClassica, designed by André Miede.

CONTACTS

✉ evertn@student.ethz.ch



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Declaration of originality

The signed declaration of originality is a component of every semester paper, Bachelor's thesis, Master's thesis and any other degree paper undertaken during the course of studies, including the respective electronic versions.

Lecturers may also require a declaration of originality for other written papers compiled for their courses.

I hereby confirm that I am the sole author of the written work here enclosed and that I have compiled it in my own words. Parts excepted are corrections of form and content by the supervisor.

Title of work (in block letters):

Authored by (in block letters):

For papers written by groups the names of all authors are required.

Name(s):

First name(s):

With my signature I confirm that

- I have committed none of the forms of plagiarism described in the '[Citation etiquette](#)' information sheet.
- I have documented all methods, data and processes truthfully.
- I have not manipulated any data.
- I have mentioned all persons who were significant facilitators of the work.

I am aware that the work may be screened electronically for plagiarism.

Place, date

Signature(s)

For papers written by groups the names of all authors are required. Their signatures collectively guarantee the entire content of the written paper.

ACKNOWLEDGEMENTS

—

ABSTRACT

Following its launch in 2021, the James Webb Telescope will provide the best infrared observations of exoplanets and brown dwarfs to date. In particular, the Mid-Infrared Instrument (MIRI), will allow for medium resolution spectroscopy across a wide wavelength band, from 4.9-28.8 μm . This will allow us to derive atmospheric properties of objects at lower temperatures than currently possible. MIRI's medium resolution spectrometer (MRS) is an integrated field unit that will perform these observations, providing both spatial and spectral information about targets. Understanding the instrumental effects is critical to analyzing data from MIRI. With that in mind, the MIRISIM instrumental simulator was developed to provide observational simulations of the various sub instruments of MIRI.

This thesis improves the implementation of a thin-film fringing model for point sources to MIRISIM, considering how the fringing effect from the detector layers varies with position. Fringing is a periodic, wavelength dependent effect, and thus has a strong impact on any spectroscopic observations. A comparison to the existing model was made, demonstrating the necessity of considering this effect when analyzing data. We will improve the fringing removal by identifying the point source location from the constructed data cube, and select the correct fringe flat for removal.

Understanding the instrumental effects is key to quantifying the ability of MIRI to derive atmospheric properties. Existing literature has considered the NIRCAM instrument and the MIRI Low-Resolution Spectrometer, but to date no retrieval studies have been performed using MIRISIM, or for the MIRI MRS, though it is critical to extend wavelength coverage to improve the results of an atmospheric retrieval. Model atmospheres will be generated using PetitRadTrans, and processed using MIRISIM and the JWST pipeline to produce a mock observation. An atmospheric retrieval will be performed, demonstrating to what extent MIRI will be able to retrieve atmospheric parameters such as temperature, pressure and composition. The posterior distributions of these parameters are compared with and without the fringing removal, again demonstrating the importance of correcting for this effect.

CONTENTS

1	INTRODUCTION	1
1.1	Brown Dwarfs	1
1.1.1	Physics	1
1.1.2	Observational Properties	1
1.2	Exoplanets	1
1.2.1	Direct imaging	1
1.2.2	Characteristic Examples	1
1.3	Current Status of Atmospheric Characterization	1
1.3.1	Transmission Spectroscopy	1
1.3.2	Emission Spectroscopy	1
1.3.3	JWST Studies	1
1.3.4	Biosignatures and Future Missions	1
2	MIRI: THE MID-INFRARED INSTRUMENT	3
2.1	JWST	3
2.2	MRS	3
2.2.1	Integrated Field Spectroscopy	3
2.2.2	Optical Systems	3
2.2.3	Detectors	3
2.3	Coordinates	3
2.4	Observations	3
2.4.1	Dithering	3
2.4.2	Readout Modes	3
2.4.3	Exposure time calculations	3
3	FRINGING EFFECTS IN MIRISIM	5
3.1	MIRISIM	5
3.1.1	Architecture	5
3.1.2	Data Products	5
3.1.3	Instrumental effects	5
3.1.4	Fringing	5
3.1.5	Implementation	7
3.1.6	JWST Pipeline	7
3.1.7	Cross-Correlation	7
3.2	Fringing Results	7
3.2.1	Effects of fringing on spectral extraction	7
4	ATMOSPHERIC RETRIEVALS	9
4.1	Introduction	9
4.2	Atmospheric Modeling	9
4.2.1	petitRadTrans	9
4.3	Bayesian methods	9
4.3.1	MCMC	9
4.3.2	Nested Sampling	9
4.3.3	multinest	9
4.4	Targets	9
4.4.1	Atmospheric Parameters	9

	4.4.2	petitRadTrans	9
4.5		Results	9
	4.5.1	Posterior Distributions	9
5		CONCLUSIONS	11
		BIBLIOGRAPHY	13

1

INTRODUCTION

1.1 BROWN DWARFS

1.1.1 Physics

1.1.2 Observational Properties

T-Type

L-Type

Y-Type

1.2 EXOPLANETS

1.2.1 Direct imaging

1.2.2 Characteristic Examples

β *Pictoris b*

HR 8799 System

PDS 70 b,c

1.3 CURRENT STATUS OF ATMOSPHERIC CHARACTERIZATION

1.3.1 Transmission Spectroscopy

1.3.2 Emission Spectroscopy

1.3.3 JWST Studies

1.3.4 Biosignatures and Future Missions

2 | MIRI: THE MID-INFRARED INSTRUMENT

2.1 JWST

2.2 MRS

2.2.1 Integrated Field Spectroscopy

2.2.2 Optical Systems

Channels, bands, etc **ref:Chen2019**

2.2.3 Detectors

2.3 COORDINATES

2.4 OBSERVATIONS

2.4.1 Dithering

2.4.2 Readout Modes

2.4.3 Exposure time calculations

3

FRINGING EFFECTS IN MIRISIM

3.1 MIRISIM

The MIRI instrument has been modeled in python as a program known as MIRISIM. This program takes in an astronomical ‘scene’ along with some configuration parameters to output a detector data product, similar to what will be produced by the actual instrument. This is relatively full-featured simulator, modelling the instrumental PSF, various noise sources and distortion maps, among other effects. While MIRISIM is functional for all of the MIRI sub-instruments, this report will only deal with the Medium-Resolution Spectrometer (MRS) sub instrument, described in section 2.2. The goal of this section is to describe the implementation and testing of an updated optical model of the ‘fringing’ effect - an optical effect caused by thin film interference from the multiple layers of the detector.

3.1.1 Architecture

SCENE - SEDs SIMULATOR PYSPECSIM

3.1.2 Data Products

3.1.3 Instrumental effects

3.1.4 Fringing

A key effect on spectral data is fringing, described in **ref:Argyriou2018**. MIRI uses a total of three Si:As impurity band conduction detector arrays, two of which are used by the MRS. These detectors consist of 7 layers, listed in table 1 and illustrated in Fig. 1.

Thin film interference occurs when light is coherently reflected at the boundary between two layers and interferes with the incident light. This is the principle on which Fabry-Pérot interferometers function. As we wish to determine the effect of fringing on the amplitude of the signal received by the detector, we are effectively interested in the transmittance of a series of Fabry-Pérot interferometers. Assuming an ideal plane-parallel optical cavity with a reflectance R at both boundaries, thickness D , and an angle θ at which the light travels within the cavity, we can compute the transmittance as:

$$T_c = \frac{1}{1 + \frac{4R}{(1-R)^2} \sin^2\left(\frac{\delta}{2}\right)} \quad (1)$$

Figure 1

Table 1

Figure 2

Where the phase δ at half λ ($\psi = \pi$) is:

$$\delta = 4\pi\sigma D \cos \theta - (\phi - \pi) \quad (2)$$

If all of the parameters were known, this would be sufficient to numerically solve for the fringing pattern within MIRI. Unfortunately, uncertainties in the thickness in the detector layers, variations in the layer deposition thickness, and the uncertainty of transmittance and reflectance of the materials used at cryogenic temperatures prevents the implementation of such a numerical model.

Instead, we turn to calibration data taken to characterize the fringing pattern.

DESCRIBE CURRENT MODEL - GENERIC FRINGING

However, due to the dependance of fringing on the incident angle of the light, a single model of fringing is insufficient to describe the full effect. Therefore, we use data taken in XXXXXXXXXX at various points across the detector and quantify how this changes the extracted spectra after processing in the JWST pipeline.

DESCRIBE HOW THE DATA WAS TAKEN HERE. - Problems with point vs extended sources - multiple collection runs

Ultimately this data collection produced a series of ‘fringe-flats’ of an almost point like at various position across the detector and in each channel. We implemented a new routine into the pySpecSim portion of MIRISIM to read in the location of point sources within a scene, and apply the correct position dependent fringe flat. This implementation comes with several caveats: namely that the fringing model is not yet fully developed, so it can only be considered accurate for point sources located at the same (α, β) location as the source used to produce the fringe flat. Additionally, the source used to generate the data is not a true point source, nor are there fringe flats produced for the full MRS wavelength range. We stress that the goal of this testing is to demonstrate the significance of this effect to justify the need for a more complete model along with additional calibration data to constrain the detector layer parameters.

FM Data

CV Data

3.1.5 Implementation

3.1.6 JWST Pipeline

Stage 1 Processing

Stage 2 Processing

Fringing correction

Aperture Photometry

3.1.7 Cross-Correlation

To quantify the similarity of the spectrum output by the JWST pipeline to the input into MIRISIM, we rely on the technique of cross correlation.

3.2 FRINGING RESULTS

3.2.1 Effects of fringing on spectral extraction

4 | ATMOSPHERIC RETRIVALS

4.1 INTRODUCTION

4.2 ATMOSPHERIC MODELING

4.2.1 petitRadTrans

Radiative Transfer

Line-by-line

Correlated K

Clouds

4.3 BAYESIAN METHODS

4.3.1 MCMC

4.3.2 Nested Sampling

4.3.3 multinest

Hyperparameters

4.4 TARGETS

4.4.1 Atmospheric Parameters

4.4.2 petitRadTrans

4.5 RESULTS

4.5.1 Posterior Distributions

5 | CONCLUSIONS

[1]

BIBLIOGRAPHY

- [1] I. Argyriou, R. Azzollini, and B. Vandenbussche, "Spectrum extraction from detector plane images for the medium-resolution spectrometer of the mid-Infrared Instrument on-board the James Webb Space Telescope.", no. July 2018, p. 124, 2018. DOI: [10.1117/12.2312697](https://doi.org/10.1117/12.2312697).
- [2] L. Zhang and R. R. Issa, *Development of IFC-based construction industry ontology for information retrieval from IFC models*. 2014, ISBN: 9789036532167.
- [3] D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman, "emcee : The MCMC Hammer ", *Publications of the Astronomical Society of the Pacific*, vol. 125, no. 925, pp. 306–312, 2013, ISSN: 00046280. DOI: [10.1086/670067](https://doi.org/10.1086/670067). arXiv: [1202.3665](https://arxiv.org/abs/1202.3665).
- [4] C. V. Morley, M. S. Marley, J. J. Fortney, R. Lupu, D. Saumon, T. Greene, and K. Lodders, "Water clouds in y dwarfs and exoplanets", *Astrophysical Journal*, vol. 787, no. 1, 2014, ISSN: 15384357. DOI: [10.1088/0004-637X/787/1/78](https://doi.org/10.1088/0004-637X/787/1/78). arXiv: [1404.0005](https://arxiv.org/abs/1404.0005).
- [5] M. R. Line, M. S. Marley, M. C. Liu, B. Burningham, C. V. Morley, N. R. Hinkel, J. Teske, J. J. Fortney, R. Freedman, and R. Lupu, "Uniform Atmospheric Retrieval Analysis of Ultracool Dwarfs. II. Properties of 11 T dwarfs", *The Astrophysical Journal*, vol. 848, no. 2, p. 83, 2017, ISSN: 1538-4357. DOI: [10.3847/1538-4357/aa7ff0](https://doi.org/10.3847/1538-4357/aa7ff0). arXiv: [1612.02809](https://arxiv.org/abs/1612.02809).
- [6] J. A. Zalesky, M. R. Line, A. C. Schneider, and J. Patience, "A Uniform Retrieval Analysis of Ultra-cool Dwarfs. III. Properties of Y Dwarfs", *The Astrophysical Journal*, vol. 877, no. 1, p. 24, 2019, ISSN: 1538-4357. DOI: [10.3847/1538-4357/ab16db](https://doi.org/10.3847/1538-4357/ab16db). arXiv: [1903.11658](https://arxiv.org/abs/1903.11658).
- [7] M. R. Line, M. S. Marley, M. C. Liu, B. Burningham, C. V. Morley, N. R. Hinkel, J. Teske, J. J. Fortney, R. Freedman, and R. Lupu, "Uniform Atmospheric Retrieval Analysis of Ultracool Dwarfs. II. Properties of 11 T dwarfs", *The Astrophysical Journal*, vol. 848, no. 2, p. 83, 2017, ISSN: 1538-4357. DOI: [10.3847/1538-4357/aa7ff0](https://doi.org/10.3847/1538-4357/aa7ff0). arXiv: [1612.02809](https://arxiv.org/abs/1612.02809). [Online]. Available: <http://dx.doi.org/10.3847/1538-4357/aa7ff0>.
- [8] M. R. Line, J. Teske, B. Burningham, J. J. Fortney, and M. S. Marley, "UNIFORM ATMOSPHERIC RETRIEVAL ANALYSIS of ULTRA-COOL DWARFS. I. CHARACTERIZING BENCHMARKS, Gl 570D and HD 3651B", *Astrophysical Journal*, vol. 807, no. 2, p. 183, 2015, ISSN: 15384357. DOI: [10.1088/0004-637X/807/2/183](https://doi.org/10.1088/0004-637X/807/2/183). arXiv: [1504.06670](https://arxiv.org/abs/1504.06670). [Online]. Available: <http://dx.doi.org/10.1088/0004-637X/807/2/183>.
- [9] B. Biller, "The time domain for brown dwarfs and directly imaged giant exoplanets: the power of variability monitoring", *Astronomical Review*, vol. 13, no. 1, pp. 1–27, 2017, ISSN: 2167-2857. DOI: [10.1080/21672857.2017.1303105](https://doi.org/10.1080/21672857.2017.1303105). [Online]. Available: <http://dx.doi.org/10.1080/21672857.2017.1303105>.

- [10] I. P. Waldmann, G. Tinetti, M. Rocchetto, E. J. Barton, S. N. Yurchenko, and J. Tennyson, “Tau-REx I: A next generation retrieval code for exoplanetary atmospheres”, *Astrophysical Journal*, vol. 802, no. 2, p. 107, 2015, ISSN: 15384357. DOI: [10.1088/0004-637X/802/2/107](https://doi.org/10.1088/0004-637X/802/2/107). arXiv: [1409.2312](https://arxiv.org/abs/1409.2312). [Online]. Available: <http://dx.doi.org/10.1088/0004-637X/802/2/107>.
- [11] I. P. Waldmann, M. Rocchetto, G. Tinetti, E. J. Barton, S. N. Yurchenko, and J. Tennyson, “T-REx. II. RETRIEVAL of EMISSION SPECTRA”, *Astrophysical Journal*, vol. 813, no. 1, p. 13, 2015, ISSN: 15384357. DOI: [10.1088/0004-637X/813/1/13](https://doi.org/10.1088/0004-637X/813/1/13). arXiv: [1508.07591](https://arxiv.org/abs/1508.07591). [Online]. Available: <http://dx.doi.org/10.1088/0004-637X/813/1/13>.
- [12] B. Benneke and S. Seager, “Atmospheric retrieval for super-earths: Uniquely constraining the atmospheric composition with transmission spectroscopy”, *Astrophysical Journal*, vol. 753, no. 2, 2012, ISSN: 15384357. DOI: [10.1088/0004-637X/753/2/100](https://doi.org/10.1088/0004-637X/753/2/100). arXiv: [1203.4018](https://arxiv.org/abs/1203.4018).
- [13] L. Vicum, “Research Collection: Theses”, *PhD thesis*, 2005. DOI: [10.3929/ethz-a-010782581](https://doi.org/10.3929/ethz-a-010782581).
- [14] T. D. Robinson, K. R. Stapelfeldt, and M. S. Marley, “Characterizing rocky and gaseous exoplanets with 2m class space-based coronagraphs”, *Publications of the Astronomical Society of the Pacific*, vol. 128, no. 960, pp. 1–22, 2016, ISSN: 00046280. DOI: [10.1088/1538-3873/128/960/025003](https://doi.org/10.1088/1538-3873/128/960/025003). [Online]. Available: <http://dx.doi.org/10.1088/1538-3873/128/960/025003>.
- [15] M. Zhang, Y. Chachan, E. M. Kempton, and H. A. Knutson, “Forward modeling and retrievals with PLATON, a fast open-source tool”, *Publications of the Astronomical Society of the Pacific*, vol. 131, no. 997, 2019, ISSN: 00046280. DOI: [10.1088/1538-3873/aaf5ad](https://doi.org/10.1088/1538-3873/aaf5ad). arXiv: [1811.11761](https://arxiv.org/abs/1811.11761).
- [16] M. R. Line and V. Parmentier, “the Influence of Nonuniform Cloud Cover on Transit Transmission Spectra”, *The Astrophysical Journal*, vol. 820, no. 1, p. 78, 2016, ISSN: 1538-4357. DOI: [10.3847/0004-637x/820/1/78](https://doi.org/10.3847/0004-637x/820/1/78). arXiv: [1511.09443](https://arxiv.org/abs/1511.09443). [Online]. Available: <http://dx.doi.org/10.3847/0004-637X/820/1/78>.
- [17] J. Skilling, “Nested Sampling”, vol. 395, no. November 2004, pp. 395–405, 2004. DOI: [10.1063/1.1835238](https://doi.org/10.1063/1.1835238).
- [18] P. G. Irwin, N. A. Teanby, R. de Kok, L. N. Fletcher, C. J. Howett, C. C. Tsang, C. F. Wilson, S. B. Calcutt, C. A. Nixon, and P. D. Parrish, “The NEMESIS planetary atmosphere radiative transfer and retrieval tool”, *Journal of Quantitative Spectroscopy and Radiative Transfer*, vol. 109, no. 6, pp. 1136–1150, 2008, ISSN: 00224073. DOI: [10.1016/j.jqsrt.2007.11.006](https://doi.org/10.1016/j.jqsrt.2007.11.006).
- [19] J. S. Speagle, “A Conceptual Introduction to Markov Chain Monte Carlo Methods”, 2019. arXiv: [1909.12313](https://arxiv.org/abs/1909.12313). [Online]. Available: <http://arxiv.org/abs/1909.12313>.

- [20] M. R. Line, A. S. Wolf, X. Zhang, H. Knutson, J. A. Kammer, E. Ellison, P. Deroo, D. Crisp, and Y. L. Yung, “A systematic retrieval analysis of secondary eclipse spectra. I. A comparison of atmospheric retrieval techniques”, *Astrophysical Journal*, vol. 775, no. 2, 2013, ISSN: 15384357. DOI: [10.1088/0004-637X/775/2/137](https://doi.org/10.1088/0004-637X/775/2/137). arXiv: [1304.5561](https://arxiv.org/abs/1304.5561).
- [21] S. Gandhi and N. Madhusudhan, “Retrieval of exoplanet emission spectra with HyDRA”, *Monthly Notices of the Royal Astronomical Society*, vol. 474, no. 1, pp. 271–278, 2018, ISSN: 13652966. DOI: [10.1093/mnras/stx2748](https://doi.org/10.1093/mnras/stx2748). arXiv: [1710.06433](https://arxiv.org/abs/1710.06433).
- [22] R. J. MacDonald and N. Madhusudhan, “HD 209458b in new light: evidence of nitrogen chemistry, patchy clouds and sub-solar water”, *Monthly Notices of the Royal Astronomical Society*, vol. 469, no. 2, pp. 1979–1996, 2017, ISSN: 0035-8711. DOI: [10.1093/mnras/stx804](https://doi.org/10.1093/mnras/stx804). arXiv: [1701.01113](https://arxiv.org/abs/1701.01113).
- [23] J. M. Lee, L. N. Fletcher, and P. G. Irwin, “Optimal estimation retrievals of the atmospheric structure and composition of HD189733b from secondary eclipse spectroscopy”, *Monthly Notices of the Royal Astronomical Society*, vol. 420, no. 1, pp. 170–182, 2012, ISSN: 00358711. DOI: [10.1111/j.1365-2966.2011.20013.x](https://doi.org/10.1111/j.1365-2966.2011.20013.x). arXiv: [1110.2934](https://arxiv.org/abs/1110.2934).
- [24] R. E. Lupu, M. S. Marley, N. Lewis, M. Line, W. A. Traub, K. Zahnle, I. M. Jan, F. Feroz, M. P. Hobson, M. Bridges, B. Macintosh, J. R. Graham, T. Barman, R. J. D. Rosa, Q. Konopacky, M. S. Marley, C. Marois, E. L. Nielsen, L. Pueyo, A. Rajan, J. Rameau, D. Saumon, J. J. Wang, J. Patience, M. Ammons, P. Arriaga, E. Artigau, S. Beckwith, J. Brewster, S. Bruzzone, J. Bulger, B. Burningham, A. S. Burrows, C. Chen, E. Chiang, J. K. Chilcote, R. I. Dawson, R. Dong, R. Doyon, Z. H. Draper, T. M. Esposito, D. Fabrycky, M. P. Fitzgerald, K. B. Follette, J. J. Fortney, B. Gerard, S. Goodsell, A. Z. Greenbaum, P. Hibon, S. Hinkley, T. H. Cotten, P. Ingraham, P. Kalas, D. Lafreniere, J. E. Larkin, J. Lee, M. Line, D. Long, J. Maire, F. Marchis, B. C. Matthews, C. E. Max, S. Metchev, T. Mittal, C. V. Morley, K. M. Morzinski, R. Oppenheimer, D. W. Palmer, R. Patel, M. D. Perrin, L. A. Poyneer, R. R. Rafikov, E. L. Rice, P. Rojo, A. R. Rudy, M. T. Ruiz, N. Sadakuni, L. Saddlemyer, M. Salama, D. Savransky, A. C. Schneider, A. Sivaramakrishnan, I. Song, R. Soummer, S. Thomas, G. Vasisht, J. K. Wallace, S. J. Wiktorowicz, S. G. Wolff, B. Zuckerman, E. Planetary, S. Congress, S. Seager, S. Cruz, P. Torinese, G. H. Road, M. View, M. View, A. S. Burrows, D. Sudarsky, and J. I. Lunine, “DEVELOPING ATMOSPHERIC RETRIEVAL METHODS FOR DIRECT IMAGING SPECTROSCOPY OF GAS GIANTS IN REFLECTED LIGHT . I . METHANE ABUNDANCES AND BASIC CLOUD PROPERTIES”, *The Astronomical Journal*, vol. 10, no. October, pp. 1–29, 2018, ISSN: 1538-3881. DOI: [10.3847/0004-6256/152/6/217](https://doi.org/10.3847/0004-6256/152/6/217). arXiv: [arXiv:1306.2144v2](https://arxiv.org/abs/1306.2144v2). [Online]. Available: <http://dx.doi.org/10.3847/0004-6256/152/6/217>.
- [25] E. Schlawin, T. P. Greene, M. Line, J. J. Fortney, and M. Rieke, “ Clear and Cloudy Exoplanet Forecasts for JWST : Maps, Retrieved Composition, and Constraints on Formation with MIRI and NIRCam ”, *The*

- Astronomical Journal*, vol. 156, no. 1, p. 40, 2018, ISSN: 0004-6256. DOI: [10.3847/1538-3881/aac774](https://doi.org/10.3847/1538-3881/aac774). arXiv: [1803.08173](https://arxiv.org/abs/1803.08173).
- [26] Q. M. Konopacky, T. S. Barman, B. A. Macintosh, and C. Marois, "Detection of carbon monoxide and water absorption lines in an exoplanet atmosphere", *Science*, vol. 339, no. 6126, pp. 1398–1401, 2013, ISSN: 10959203. DOI: [10.1126/science.1232003](https://doi.org/10.1126/science.1232003).
 - [27] S. Seager, "Bayesian Atmospheric Retrieval for Exoplanets", 2013.
 - [28] C. A. D. Oliveira, "Peering into the physics of brown dwarfs : spectroscopy with JWST / NIRSpec",
 - [29] T. S. Barman, Q. M. Konopacky, B. Macintosh, and C. Marois, "Simultaneous detection of water, methane, and carbon monoxide in the atmosphere of Exoplanet hr 8799 b", *Astrophysical Journal*, vol. 804, no. 1, pp. 1–10, 2015, ISSN: 15384357. DOI: [10.1088/0004-637X/804/1/61](https://doi.org/10.1088/0004-637X/804/1/61). arXiv: [1503.03539](https://arxiv.org/abs/1503.03539). [Online]. Available: <http://dx.doi.org/10.1088/0004-637X/804/1/61>.
 - [30] T. L. Roellig, N. Ames, T. P. Greene, N. Ames, C. Beichman, M. Meyer, E. T. H. Zurich, M. Rieke, and U. Arizona, "Studies of the Coldest Brown Dwarfs with the James Webb Space Telescope", p. 2014, 2014.
 - [31] J. Li, D. B. Hibbert, S. Fuller, and G. Vaughn, "A comparative study of point-to-point algorithms for matching spectra", *Chemometrics and Intelligent Laboratory Systems*, vol. 82, no. 1-2 SPEC. ISS, pp. 50–58, 2006, ISSN: 01697439. DOI: [10.1016/j.chemolab.2005.05.015](https://doi.org/10.1016/j.chemolab.2005.05.015).
 - [32] I. Argyriou, R. Azzollini, and B. Vandenbussche, "Spectrum extraction from detector plane images for the medium-resolution spectrometer of the mid-Infrared Instrument on-board the James Webb Space Telescope.", no. July 2018, p. 124, 2018, ISSN: 1996756X. DOI: [10.1117/12.2312697](https://doi.org/10.1117/12.2312697).
 - [33] T. Guillot, "On the radiative equilibrium of irradiated planetary atmospheres", *Astronomy and Astrophysics*, vol. 520, no. 18, pp. 1–13, 2010, ISSN: 00046361. DOI: [10.1051/0004-6361/200913396](https://doi.org/10.1051/0004-6361/200913396). arXiv: [1006.4702](https://arxiv.org/abs/1006.4702).
 - [34] R. Garland and P. G. J. Irwin, "Effectively Calculating Gaseous Absorption in Radiative Transfer Models of Exoplanetary and Brown Dwarf Atmospheres", no. March, 2019. arXiv: [1903.03997](https://arxiv.org/abs/1903.03997). [Online]. Available: <http://arxiv.org/abs/1903.03997>.
 - [35] P. Mollière, J. P. Wardenier, R. van Boekel, T. Henning, K. Molaverdikhani, and I. A. G. Snellen, "petitRADTRANS: a Python radiative transfer package for exoplanet characterization and retrieval", vol. 67, pp. 1–17, 2019, ISSN: 0004-6361. DOI: [10.1051/0004-6361/201935470](https://doi.org/10.1051/0004-6361/201935470). arXiv: [1904.11504](https://arxiv.org/abs/1904.11504). [Online]. Available: <http://arxiv.org/abs/1904.11504> <http://dx.doi.org/10.1051/0004-6361/201935470>.
 - [36] P. Lagage, F. Lahuis, S. Ronayette, D. Wright, and G. S. Wright, "The Mid-Infrared Instrument for the James Webb Space Telescope , IV : The Low Resolution Spectrometer", pp. 1–23,

- [37] G. H. Rieke, G. S. Wright, T. Böker, J. Bouwman, L. Colina, A. Glasse, K. D. Gordon, T. P. Greene, M. Güdel, T. Henning, K. Justtanont, P.-O. Lagage, M. E. Meixner, H.-U. Nørgaard-Nielsen, T. P. Ray, M. E. Ressler, E. F. van Dishoeck, and C. Waelkens, “The Mid-Infrared Instrument for the James Webb Space Telescope , I: Introduction ”, *Publications of the Astronomical Society of the Pacific*, vol. 127, no. 953, pp. 584–594, 2015, ISSN: 00046280. DOI: [10.1086/682252](https://doi.org/10.1086/682252).
- [38] J. T. VanderPlas, “Understanding the Lomb–Scargle Periodogram”, *The Astrophysical Journal Supplement Series*, vol. 236, no. 1, p. 16, 2018, ISSN: 0067-0049. DOI: [10.3847/1538-4365/aab766](https://doi.org/10.3847/1538-4365/aab766). arXiv: [1703.09824](https://arxiv.org/abs/1703.09824). [Online]. Available: <http://dx.doi.org/10.3847/1538-4365/aab766>.
- [39] A. A. Lacis and V. Oinas, “A description of the correlated k distribution method for modeling nongray gaseous absorption, thermal emission, and multiple scattering in vertically inhomogeneous atmospheres”, *Journal of Geophysical Research*, vol. 96, no. D5, pp. 9027–9063, 1991, ISSN: 01480227. DOI: [10.1029/90JD01945](https://doi.org/10.1029/90JD01945).
- [40] J. Mather, “The James Webb Space Telescope”, *Adaptive Optics: Analysis, Methods and Systems, AO 2015*, no. Scipy, p. 289, 2015, ISSN: 00386308. DOI: [10.1364/AOMS.2015.JT1A.1](https://doi.org/10.1364/AOMS.2015.JT1A.1). arXiv: [0606175](https://arxiv.org/abs/0606175) [astro-ph].
- [41] Qiang Fu and K. N. Liou, *On the correlated k-distribution method for radiative transfer in nonhomogeneous atmospheres*, 1992. DOI: [10.1175/1520-0469\(1992\)049<2139:otcdf>2.0.co;2](https://doi.org/10.1175/1520-0469(1992)049<2139:otcdf>2.0.co;2).
- [42] F. Lahuis and A. Boogert, “How to Get Rid of Fringes in SIRTf/{IR}S Data”, *SFChem 2002: Chemistry as a Diagnostic of Star Formation*, no. January 2003, p. 335, 2003.
- [43] R. Goody, R. West, L. Chen, and D. Crisp, “The correlated-k method for radiation calculations in nonhomogeneous atmospheres”, *Journal of Quantitative Spectroscopy and Radiative Transfer*, vol. 42, no. 6, pp. 539–550, 1989, ISSN: 00224073. DOI: [10.1016/0022-4073\(89\)90044-7](https://doi.org/10.1016/0022-4073(89)90044-7).
- [44] N. Madhusudhan, “Atmospheric Retrieval of Exoplanets”, *Handbook of Exoplanets*, pp. 2153–2182, 2018. DOI: [10.1007/978-3-319-55333-7_104](https://doi.org/10.1007/978-3-319-55333-7_104). arXiv: [1808.04824](https://arxiv.org/abs/1808.04824).
- [45] C. Danielski, J.-L. Baudino, P.-O. Lagage, A. Boccaletti, R. Gastaud, A. Coulais, and B. Bézard, “Atmospheric Characterization of Directly Imaged Exoplanets with JWST /MIRI ”, *The Astronomical Journal*, vol. 156, no. 6, p. 276, 2018, ISSN: 1538-3881. DOI: [10.3847/1538-3881/aae651](https://doi.org/10.3847/1538-3881/aae651). arXiv: [1810.00894](https://arxiv.org/abs/1810.00894).
- [46] V. Bozza, L. Mancini, and A. Sozzetti, *Astrophysics of Exoplanetary Atmospheres*, ISBN: 9783319897004.
- [47] C. A. Beichman, D. Barrado, R. Belikov, B. Biller, A. Boccaletti, A. Burrows, C. Danielski, R. Doyon, J. Fortney, A. Gaspar, A. Glasse, T. P. Greene, S. Hinkley, R. Hu, T. Kataria, J. Krist, D. Lafreni, P.-o. Lagage, J. Lunine, M. Marley, D. Mawet, T. Meshkat, M. Meyer, R. Oppenheimer, M. Perrin, L. Pueyo, M. Ressler, G. Rieke, M. Rieke, T. Roellig, E. Serabyn, J. E. Schlieder, A. Skemer, R. Soummer, K. Su, P. Tremblin, O.

- Venot, and M. Ygouf, "Direct Imaging and Spectroscopy of Exoplanets with the James Webb Space Telescope", *Astro2020 Science White Paper*, 2019. [Online]. Available: <https://jwst.stsci.edu/observing-programs/approved-gto-programs..>
- [48] T. Guillot, "On the radiative equilibrium of irradiated planetary atmospheres", *Astronomy and Astrophysics*, vol. 520, no. 18, pp. 1–13, 2010, ISSN: 00046361. DOI: [10.1051/0004-6361/200913396](https://doi.org/10.1051/0004-6361/200913396). arXiv: [1006.4702](https://arxiv.org/abs/1006.4702).
- [49] Qiang Fu and K. N. Liou, *On the correlated k-distribution method for radiative transfer in nonhomogeneous atmospheres*, 1992. DOI: [10.1175/1520-0469\(1992\)049<2139:otcdmf>2.0.co;2](https://doi.org/10.1175/1520-0469(1992)049<2139:otcdmf>2.0.co;2).
- [50] I. Argyriou, B. Vandenbussche, and M. Wells, "Fringing solution for the mid-infrared instrument on-board the James Webb Space Telescope", no. July 2018, p. 127, 2018. DOI: [10.1117/12.2310268](https://doi.org/10.1117/12.2310268).
- [51] C. A. Beichman and T. P. Greene, "Observing Exoplanets with the James Webb Space Telescope", *Handbook of Exoplanets*, no. 2009, pp. 1283–1308, 2018. DOI: [10.1007/978-3-319-55333-7_85](https://doi.org/10.1007/978-3-319-55333-7_85).
- [52] G. H. Rieke, G. S. Wright, T. Böker, J. Bouwman, L. Colina, A. Glasse, K. D. Gordon, T. P. Greene, M. Güdel, T. Henning, K. Justtanont, P.-O. Lagage, M. E. Meixner, H.-U. Nørgaard-Nielsen, T. P. Ray, M. E. Ressler, E. F. van Dishoeck, and C. Waelkens, "The Mid-Infrared Instrument for the James Webb Space Telescope , I: Introduction ", *Publications of the Astronomical Society of the Pacific*, vol. 127, no. 953, pp. 584–594, 2015, ISSN: 00046280. DOI: [10.1086/682252](https://doi.org/10.1086/682252).
- [53] I. Argyriou, B. Vandenbussche, and M. Wells, "Fringing solution for the mid-infrared instrument on-board the James Webb Space Telescope", no. July 2018, p. 127, 2018. DOI: [10.1117/12.2310268](https://doi.org/10.1117/12.2310268).
- [54] Y. K. Feng, T. D. Robinson, J. J. Fortney, R. E. Lupu, M. S. Marley, N. K. Lewis, B. Macintosh, and M. R. Line, "Characterizing Earth Analogs in Reflected Light: Atmospheric Retrieval Studies for Future Space Telescopes", *The Astronomical Journal*, vol. 155, no. 5, p. 200, 2018, ISSN: 1538-3881. DOI: [10.3847/1538-3881/aab95c](https://doi.org/10.3847/1538-3881/aab95c). [Online]. Available: <http://dx.doi.org/10.3847/1538-3881/aab95c>.
- [55] P. Lagage, F. Lahuis, S. Ronayette, D. Wright, and G. S. Wright, "The Mid-Infrared Instrument for the James Webb Space Telescope , IV : The Low Resolution Spectrometer", pp. 1–23,
- [56] M. E. Consortium, "MIRISim Documentation", 2018.
- [57] H. Chen, E. T. Wolf, Z. Zhan, and D. E. Horton, "Habitability and Spectroscopic Observability of Warm M-dwarf Exoplanets Evaluated with a 3D Chemistry-Climate Model", 2019. arXiv: [1907.10048](https://arxiv.org/abs/1907.10048). [Online]. Available: <http://arxiv.org/abs/1907.10048>.
- [58] A. Labiano-Ortega, D. Dicken, B. Vandenbussche, F. Lahuis, M. Muller, S. M. Beard, K. Justtanont, R. Azzollini, D. R. Law, K. D. Gordon, A. C. H. Glasse, G. S. Wright, G. H. Rieke, P. Klaassen, A. M. Glauser, J. Morrison, V. Geers, J. I. Bailey, and M. Garcia-Marin, "The MIRI Medium Resolution Spectrometer calibration pipeline", no. exposure 1, p. 117, 2016. DOI: [10.1117/12.2232554](https://doi.org/10.1117/12.2232554).

- [59] N. Madhusudhan, M. Agúndez, J. I. Moses, and Y. Hu, “Exoplanetary Atmospheres—Chemistry, Formation Conditions, and Habitability”, *Space Science Reviews*, vol. 205, no. 1-4, pp. 285–348, 2016, ISSN: 15729672. DOI: [10.1007/s11214-016-0254-3](https://doi.org/10.1007/s11214-016-0254-3). arXiv: [arXiv:1604.06092v1](https://arxiv.org/abs/1604.06092v1).
- [60] A. Behmard, E. A. Petigura, and A. W. Howard, “Data-driven Spectroscopy of Cool Stars at High Spectral Resolution”, *The Astrophysical Journal*, vol. 876, no. 1, p. 68, 2019. DOI: [10.3847/1538-4357/ab14e0](https://doi.org/10.3847/1538-4357/ab14e0). arXiv: [arXiv:1904.00094v1](https://arxiv.org/abs/1904.00094v1).
- [61] B. A. Biller and M. Bonnefoy, “Exoplanet Atmosphere Measurements from Direct Imaging”, *Handbook of Exoplanets*, pp. 2107–2135, 2018. DOI: [10.1007/978-3-319-55333-7_101](https://doi.org/10.1007/978-3-319-55333-7_101). arXiv: [arXiv:1807.05136v1](https://arxiv.org/abs/1807.05136v1).
- [62] L. Kreidberg, “Exoplanet Atmosphere Measurements from Transmission Spectroscopy and Other Planet Star Combined Light Observations”, *Handbook of Exoplanets*, pp. 2083–2105, 2018. DOI: [10.1007/978-3-319-55333-7_100](https://doi.org/10.1007/978-3-319-55333-7_100). arXiv: [arXiv:1709.05941v2](https://arxiv.org/abs/1709.05941v2).