

Atmospheric Retrievals with petitRADTRANS

Evert Nasedkin^{1*}, Paul Mollière^{1*}, Doriann Blain^{1*}, Eleonara Alei², Tomas Stolker³, Nick Wogan⁴, Karan Molaverdikhani⁵, Mantas Zilinskas⁶, and Francois Rozet⁷

¹ Max Planck Institut für Astronomie, Germany ¶ Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ¶
- [Repository](#) ¶
- [Archive](#) ¶

Editor: [Open Journals](#) ¶

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

The petitRADTRANS (pRT) codebase has undergone significant updates since its initial publication in @molliere2019. A retrieval module combining the pRT spectrum calculations with the MultiNest [@feroz2008; @feroz2009; @feroz2013] and UltraneSt [@buchner2014] samplers has been included to streamline retrievals of exoplanet atmospheres in emission and transmission.

SUMMARY HERE # Statement of need

Retrievals are important. Faster sampling using nested sampling. Unique in ability to do c-k, line by line, transmission and emission, or any combination thereof. Widely used in the community (e.g. ...) # petitRADTRANS Retrieval Module

Brief Intro

Multiple datasets can be included into a single retrieval, with each dataset receiving its own RadTrans object used for the radiative transfer calculation, allowing for highly flexible retrievals where multiple spectral resolutions, wavelength ranges and even atmospheric models can be combined in a single retrieval. Each dataset can also receive scaling factors (for the flux, uncertainties or both), error inflation factors and offsets. Several atmospheric models are built into the models module, allowing for a wide range of P-T, cloud and chemistry parameterizations. These models are used to compute a spectrum \vec{S} , which is convolved to the instrumental resolution and binned to the wavelength bins of the data using a custom binning function to account for non-uniform bin sizes. The resulting spectrum compared to the data with flux \vec{F} and covariance \mathbf{C} in the likelihood function:

$$-2 \log \mathcal{L} = (\vec{S} - \vec{F})^T \mathbf{C}^{-1} (\vec{S} - \vec{F}) + \log (2\pi \det (\mathbf{C})). \quad (1)$$

The second term is included in the likelihood to allow for uncertainties to vary as a free parameter during the retrieval, and penalizes overly large uncertainties.

pRT can compute spectra either using line-by-line calculations, or using correlated-k tables for defining the opacities of molecular species. We include up-to-date correlated-k line lists from Exomol [@tennyson2012; mckemmish2016; polyansky2018; chubb2020] and HITEMP [@rothman2010], with the full set of available opacities listed in the online documentation. The exo-k package is used to resample the the correlated-k opacity tables to a lower spectral resolution in order to reduce the computation time [@leconte2021].

Included in pRT is an option to use an adaptive pressure grid with a higher resolution around the location of the cloud base, and a lower resolution elsewhere. The higher resolution grid is 10 times as fine as the remaining grid, and replaces one grid cell above and below the cloud base layer, as well as the cloud base layer cell itself. This allows for more precise positioning of the cloud layers within the atmosphere. Including this adaptive mesh, our pressure grid

41 contains a total of 154 layers when two cloud species are used, which is the standard grid used
42 in this work.

43 Finally, photometric data are fully incorporated into the retrieval process. As with spectroscopic
44 data, a model is computed using a user-defined function. This model spectrum is then
45 multiplied by a filter transmission profile from the SVO database using the species package
46 [stolker2020]. This results in accurate synthetic photometry, which can be compared to
47 the values specified by the user with the `add_photometry` function.

48 Combining the c-k opacities of multiple species requires mixing the distributions in g space.
49 Previously, this was accomplished by taking 1000 samples of each distribution. This sampling
50 process resulted in non-deterministic spectral calculations, resulting in unexpected behaviour
51 from the nested sampling process, as the same set of parameters could result in varying
52 log-likelihood. This has been updated to fully mix the c-k distributions. Considering the first
53 species, the second species is added in, and the resulting grid is sorted. The cumulative opacity
54 grid is then mixed with the next species, a process which iterates until every species with
55 significant opacity contributions ($>0.1\%$ of the current opacity in any bin) is mixed in to
56 the opacity grid. Once complete, the resulting grid is linearly interpolated back to the 16 g
57 points at each pressure and frequency bin as required by pRT. This fully deterministic process
58 stabilized the log-likelihood calculations in the retrievals, and resulted in a $5\times$ improvement in
59 the speed of the c-k mixing function.

60 Acknowledgements

61 We acknowledge contributions

62 References