AGRISOURCE

WARNING

KEEP OUT OF REACH OF CHILDREN

THIRAM 80 WDG

FUNGICIDE

Fungicide for control of diseases on fruit, vegetable, ornamental crops, and turf

Active ingredient
Contains 800 g/kg THIRAM in the form of a water dispersible granule.

GROUP	M3	FUNGICIDE

Net contents: 20 kg

Bar code to be inserted

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

19/02/2024

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of registration.
Anyone who mixes, loads, or applies this product, either by air or ground based methods, is required to be suitably qualified.
Refer to Hazardous Substances (Hazardous Property Controls) notice 2017 for a list of suitable qualifications.

WARNING
Harmful if swallowed.
Harmful if inhaled.
Causes serious eye irritation.
May cause an allergic skin reaction.
Suspected of damaging fertility or the unborn child.
May cause damage to organs through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.
Hazardous to terrestrial vertebrates.

PRECAUTIONARY STATEMENTS
PREVENTION
Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
Use personal protective equipment as required.
Wash hands thoroughly after handling.
Wear eye protection.
Wear protective gloves.
Avoid breathing spray.
Do not eat, drink or smoke when using this product.
Use only outdoors or in a well-ventilated area.
Contaminated work clothing should not be allowed out of the workplace.
Avoid release to the environment.
Do not apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.
Ensure any seeds treated by this substance are not accessible to birds.
Not to be used for human or animal consumption.
Ensure any granules are not accessible to birds.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/02/2024

RESPONSE
IF SWALLOWED: Call a POISON CENTER or doctor if you feel unwell.
Rinse mouth.
IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.
Call a POISON CENTER or doctor if you feel unwell.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
If eye irritation persists: Get medical advice.
IF ON SKIN: Wash with plenty of soap and water.
If skin irritation or rash occurs: Get medical advice.
Wash contaminated clothing before reuse.
IF exposed or concerned: Get medical advice.
Get medical advice if you feel unwell.
Collect spillage.

STORAGE
Store locked up, in original container tightly closed and in a dry, cool, well-ventilated area away from fertilisers, feed, seeds and foodstuffs. Store in accordance with NZS 8409 Management of Agrichemicals.

DISPOSAL
Dispose of this product in accordance with local regulations.
Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.
Container disposal: Ensure container is completely empty by adding residue to spray tank. Crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
Treated Crops and seeds: Unwanted pre-plant dip crops and seeds are to be treated as hazardous substances and disposed of accordingly.

FIRST AID
If swallowed **do not** induce vomiting. For advice, contact the National Poisons Centre 0800 Poison (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. **Do not scrub the skin**. If concentrate is splashed in the eyes, flush with running water immediately.

SPILLAGE
Move non-essential people from the area. Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

SHELF LIFE
When stored appropriately, this product should show no significant degradation for two years from the date of manufacture. When using this product beyond this shelf life, contact the registrant for further information.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at **www.upl-ltd.com/nz**.

EXCLUSION OF LIABILITY – PLEASE READ
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to:
Agri Solutions Ltd
45 Kitchener Road, Pukekohe 2120, New Zealand

Distributed by:
UPL New Zealand Ltd
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
www.upl-ltd.com/nz

Registered pursuant to the ACVM Act 1997, No P000677
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996. Approval Code HSR101528
See www.epa.govt.nz for approval controls

© Copyright UPL New Zealand Limited 2020, all rights reserved.
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited.

® Registered trademark

Batch number:	
Date of Manufacture:	

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622)
UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (THIRAM)
MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2XE

19/02/2024

GENERAL INFORMATION

THIRAM 80 WDG is a protectant fungicide that prevents disease establishment on treated plant surfaces. Good coverage is essential for satisfactory results.

RESISTANCE MANAGEMENT

THIRAM 80 WDG contains the group M3 fungicide, thiram. Resistance to pesticides can develop from excessive use. To minimise this risk, avoid using this fungicide exclusively all season, avoid unnecessary spraying and endeavour to alternate with fungicides from other groups within the crop and in subsequent seasons for a sprayed paddock. Use strictly in accordance with label instructions. Maintain good cultural practices.

DIRECTIONS FOR USE

Apply as a high-volume spray to the point of runoff. Complete and uniform coverage is essential. Rates suggested are for dilute spraying and require adjusting for concentrate spraying.

Record Keeping

If 3 kg or more of the substance is applied within 24 hours, in a place where the substance is likely to enter air or water and leave the application area, a PCBU with management or control of the substance must ensure that a written record is kept of each application of the substance.

Use restrictions:

1. This substance must be applied using ground-based methods only.
2. Seed treatment - information requirements: Information must be provided with seeds treated with this substance that indicates the substance that the seeds have been treated with, its hazardous properties and precautions to be taken in handling the seeds.

CROP	DISEASE	RATE PER 100L OF WATER	COMMENTS
FRUIT			
Apples & pears	Blackspot	75 – 150 g	Use the higher rate if applying early in the season. WARNING: Fruit thinning may occur if **THIRAM 80 WDG** is applied to Granny Smith apples earlier than December.
Stonefruit	Brown rot, rust, peach scab	150 g	May be applied at any time after petal fall.
	Leaf curl	160 g	Apply at start of leaf bud movement.
Grapes	Blackspot, botrytis	150 – 200 g	Apply at start of leaf bud movement.
Strawberries	Botrytis, Leafspot (*Mycosphaerella*)	150 g	Apply at 7-day intervals from the start of flowering.
Other berryfruit	Botrytis, Canefruit leafspot (*Septoria*)	150 – 200 g	Apply at green tip and continue at 10–14 day intervals. Use the higher rate over flowering or in wet weather.
VEGETABLES, ORNAMENTALS & TURF			
Tomatoes	Botrytis, leaf mould	150 g	Apply when symptoms first appear and repeat at 7-14 day intervals.
Bulbs	Botrytis, corms and tubers	1.2 kg	Dip at lifting and again pre-planting. Leave immersed in dip for 4 minutes.
Roses	Blackspot	150 g	Apply at 10 -14 day intervals during humid weather.
Turf	Brown patch, damping off, rust, leaf spot (*Helminthosporium*)		Use 15 g per 10 L of water per 20 sq. m. Apply at 7-10 day intervals until spread of the disease is halted.
Seed Beds	Damping-off		Use 40 g per 10 sq. m as a soil drench before planting or mix 1 part **THIRAM 80 WDG** with 4 parts sand and incorporate into the soil mix.
Seed treatment	Pythium (onions), soil borne seed and damping-off diseases (maize, turf)		Add 50 g of **THIRAM 80 WDG** per 1 kg of seed in a slurry form. **NB:** Use specialist seed treatment equipment to ensure complete coverage.
RABBIT / HARE REPELLENT			
Pipfruit and stonefruit.			Mix 125 g in water sufficient to make a cream suitable for painting on to the trunks/stems/buds accessible to these animals.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

It is an offence to use this product on animals.

The following maximum application rates have been set by EPA and must be followed

Crop	Max Rate	Max No. of Applications
Fruiting trees (e.g. apples, pears, stonefruit)	3.2 kg /ha	1
Grapes	0.6 kg /ha	1
Strawberries	2.25 kg /ha	5 (min 7 day interval)
Other berryfruit	1 kg /ha	5 (min 10 day interval)
Tomatoes	0.75 kg /ha	5 (min 7 day interval)
Bulbs, corms, and tubers	1.2 kg /ha	-
Roses	0.75 kg /ha	5 (min 10 day interval)
Seeds treated with this substance must not be sowed at a density that would result in concentrations exceeding 1.73 kg ai/ha.		

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

19/02/2024

MIXING

Add the required quantity of **THIRAM 80 WDG** to a small volume of water to form a slurry. Then add to the rest of the water while agitating.

EQUIPMENT

Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409). Flush spray equipment thoroughly with water after use.

When used as a seed treatment, this substance must only be applied in a purpose-built seed coating machine in a treatment plant.

COMPATIBILITY
THIRAM 80 WDG is compatible with most commonly used pesticides. Do not combine with hydrated lime, bordeaux mixture or lime sulphur.

WITHHOLDING PERIODS

Tomatoes	3 days
Strawberries	2 days
All other crops	7 days

® Registered trademarks

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

19/02/2024