DANGER

KEEP OUT OF REACH OF CHILDREN

Karate Zeon®

Insecticide

syngenta®

Active Ingredient:
KARATE ZEON® contains 250 g/litre LAMBDA-CYHALOTHRIN in the form of a capsule suspension.
Also contains: 13.2 g/litre Oxirane, methyl, polymer with oxirane, monobutyl ether.

GROUP	3	INSECTICIDE

For insect control in Beans, Cereals, Citrus, Grapes, Forage and Vegetable Brassicas, Fodder Beet (seedlings), Tomatoes, Onions, Potatoes, Pumpkins and Winter Squash (seedlings,) Maize, Sweetcorn (seedlings), and White Clover seed crops.

Net Contents: 250 mL, 1 litre, 5 litre

Registered pursuant to the ACVM Act 1997, No. P3495.
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996, Approval Code HSR000336.

Syngenta Crop Protection Limited
Level 4, 60 Parnell Road, Parnell, Auckland 1052

®

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/04/2024

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.
Any person mixing, loading or handling this product must be a qualified loader or under the guidance of a qualified loader.
Any person applying this product must be a qualified contractor/person or under the guidance of a qualified contractor/person.
This substance is restricted to use in the workplace only.

GHS Classifications
Acute oral toxicity, Category 3
Acute inhalation toxicity, Category 4
Eye irritation, Category 2
Specific target organ toxicity (repeated exposure), Category 1
Hazardous to the aquatic environment acute, Category 1
Hazardous to the aquatic environment chronic Category 1
Hazardous to terrestrial vertebrates
Hazardous to terrestrial invertebrates

DANGER
Toxic if swallowed.
Harmful if inhaled.
May cause skin irritation. Skin contact may cause a temporary sensation of the skin, such as numbness, tingling, pricking, burning or creeping on the skin.
Causes serious eye irritation.
May cause neurotoxicity and lung damage from repeated oral and inhalation exposure at high doses.
Very toxic to aquatic life with long lasting effects.
Toxic to terrestrial vertebrates.
Very toxic to terrestrial invertebrates.

PRECAUTIONS
Store in original container, tightly closed, away from foodstuffs and under lock and key.
Use only outdoors or in a well ventilated area.
When mixing or applying wear protective clothing including overalls, chemical resistant gloves, face shield and footwear.
Avoid contact with skin and eyes and inhalation of the spray mist.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking and after work.
Wash protective clothing after use.
Apply with well maintained and calibrated spray equipment.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
Records of use as described in HSW Hazardous Substances Regulations and EPA HS Notices must be kept if 3kg or more of this substance is applied within 24 hours in a place where the substance is likely to enter air or water and leave the application area.
DO NOT carry on a passenger service vehicle.
Avoid contamination of any water supply with chemical or empty container. DO NOT apply onto or into water.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/04/2024

Container disposal: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility. For additional information refer to EPA Notice: Hazardous Substances (Disposal) Notice 2017.

FIRST AID
If SWALLOWED, DO NOT induce vomiting. Rinse Mouth. Immediately call the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.
If SKIN or HAIR contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If skin irritation occurs seek medical advice.
If in EYES, wash out immediately with water. Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.
If INHALED, move the victim to fresh air immediately and keep at rest in a position comfortable for breathing. Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth. Obtain medical attention. Take to hospital without delay.
For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

SPILLAGE
Collect spillage. Wear suitable protective clothing including face shield, chemical resistant gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

GENERAL INFORMATION
KARATE ZEON® is a fast acting stomach and contact insecticide. KARATE ZEON® is relatively stable to degradation in sunlight and hence gives extended control of insects on plant surfaces. It has short persistence in the soil. As KARATE ZEON® is primarily a contact insecticide, thorough coverage of plants is important for optimum results.

RESISTANCE MANAGEMENT

GROUP	3	INSECTICIDE

KARATE ZEON® Insecticide contains a member of the pyrethroid group of insecticides. KARATE ZEON® is a Group 3 insecticide. Resistance to this insecticide could develop from repeated use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies. Since the occurrence of resistant insects is difficult to detect prior to use, Syngenta Crop Protection Ltd accepts no liability for any losses that may result from the failure of KARATE ZEON® to control resistant insects. Consult your local supplier, consultant, a Syngenta representative or New Zealand Plant Protection Society (Inc), www.resistance.nzpps.org for alternate modes of action and the details of resistance management strategies for the crops listed on the label.

Avoid using this insecticide exclusively all season, and avoid unnecessary spraying. Maintain good cultural control practices.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/04/2024

DIRECTIONS FOR USE
It is an offence to use this product on animals.
The rates suggested are for high volume spraying to run off. For concentrate spraying, adjust dilution rate accordingly.

Crop	Pest	Rate	Critical Comments
Beans	Tomato Fruitworm	40 mL/ha in a minimum of 500 litres of water/ha	Apply when insects first appear and if necessary 2 to 3 weeks later.
Beans, Forage Brassicas, Fodder Beet (seedlings), Maize, Onions, Pumpkins and Winter squash (seedlings), Sweetcorn (seedlings), Tomatoes and Vegetable Brassicas	Cutworm	40 mL/ha in a minimum of 300 litres of water/ha ***Aerial***: Use 100 to 150 litres of water/ha	Apply immediately damage occurs, preferably in the evening as a directed spray to thoroughly cover the base of the plants and the surrounding soil.
Cereals	Cereal Aphid, Rose Grain Aphid	20 to 40 mL/ha in a minimum of 200 litres of water/ha	Apply at first sign of aphids. Apply the lower rate when aphid pressure is low and shorter residual activity is required. A second application may be needed if aphids continue to migrate into the crop. Ensure sufficient water volume is used to give thorough crop penetration and coverage.
Citrus	Fullers Rose Weevil	300 mL/100 litres as a directed trunk spray	Apply sufficient spray to form a complete band around the trunk from the base to 30 cm high, using up to 300 mL/trunk. Apply at monthly intervals, using directed spray equipment. Commence prior to adult weevil emergence from ground and continue over duration of pest activity, usually December to June unless otherwise ascertained by monitoring. Ensure trees are well skirted up and tree rows are free of weeds to assist in control.
Field Tomatoes	Tomato Fruitworm	40 mL/ha in a minimum of 1000 litres of water/ha	Apply at 3 week intervals from transplanting or thinning.
Forage and Vegetable Brassicas	Diamondback Moth, White Butterfly Caterpillars	40 mL/ha in a minimum of 700 litres of water/ha	Apply to seedbeds and following transplanting at 2 week intervals or as indicated by pest activity.
Grapes	Thrips	4 mL/100 litres as a dilute spray	Apply at early flowering. Apply in sufficient water to achieve good coverage of flower clusters.
	Bronze Beetle, Grass Grub Beetle	4 mL/100 litres as a dilute spray	Apply at dusk, when beetle flights begin. Apply in sufficient water to obtain coverage. Repeat after 14 days if flights are still continuing.
Onions	Onion Thrip	40 mL/ha in a minimum of 500 litres of water/ha	Apply at first sighting of Onion Thrip and repeat as necessary.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/04/2024

Potatoes	Potato Tuber Moth	40 mL/ha in a minimum of 500 litres of water/ha	Apply at first sighting of moths then at 2 week intervals. Maintain adequate soil moisture and soil cover over developing tubers to assist prevention of tuber mining.
	Tomato/Potato Psyllid	100 mL/ha in a minimum of 500 litres of water/7	Apply at first sign of psyllid then at 7 to 14 day intervals. Use the closer interval under high pressure. Ensure sufficient water volume is used to give thorough crop penetration and coverage.
White Clover seed crops	Blue-Green Aphid, Potato Mirid, Thrips	40 mL/ha in a minimum of 200 litres of water/ha	Apply at first flower initiation, and repeat again 10 to 14 days later if required.

NOTE: Add a non-ionic low foam wetting agent at label rates.

SPRAY DRIFT
Take all reasonable steps to ensure that KARATE ZEON® does NOT cause any adverse effects to the environment beyond the application area.

POLLINATOR SAFETY
DO NOT apply KARATE ZEON® to plants if:
(a) Bees are foraging; or
(b) The plants are in flower or part flower, and are likely to be visited by non-target invertebrate pollinators (including bees).

When applied during non-foraging periods such as early morning or evening, application will not interfere with the activity of honey bees once the spray has dried, other than causing a short 2 to 3 day reduced foraging effect.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/04/2024

MIXING
Pour the measured quantity of KARATE ZEON® into a partly filled tank and agitate the spray mixture while adding the remaining water. Maintain agitation during spraying. If applying KARATE ZEON® in mixture, always add the KARATE ZEON® last.

APPLICATION
KARATE ZEON® may be applied through all conventional ground and aerial spray equipment, calibrated to ensure thorough coverage of the crop.

Application in grapes
The application rate for grapes is for high volume spraying to the point of runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved.
Set up and operate the sprayer to achieve even coverage throughout the crop canopy, applying sufficient water to cover the crop to just prior to the point of runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

COMPATIBILITY
KARATE ZEON® mixes readily with water and is compatible with commonly used fungicides except strong alkaline materials such as Bordeaux mixture and lime sulphur.

WITHHOLDING PERIODS
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

Beans	3 days
Citrus	Avoid contamination of fruit
Cereals (grain)	70 days
Cereals (Barley and wheat forage)	28 days
Field Tomatoes	3 days
Forage Brassicas	DO NOT allow stock to graze within 14 days of last application.
Fodder Beet	Seedlings only
Grapes	DO NOT use after capfall
Maize, Sweetcorn	Seedlings only
Onions	14 days
Potatoes	14 days
Pumpkins and Winter squash	Seedlings only
Vegetable Brassicas	3 days
White Clover seed crops	DO NOT allow stock to graze within 70 days of last application

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/04/2024

STORAGE
Store in a cool dry place avoiding temperatures above 35°C. Store locked up. See Safety Data Sheet for additional storage information. Stores containing quantities of 100 litres or more require signage and secondary containment.

SAFETY DATA SHEET
If additional hazard information is required, refer to the Safety Data Sheet. For a copy, visit our website at www.syngenta.co.nz

DISCLAIMER
This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland 1052. Phone 0800 333 336.

Bar Code:

Batch Number:	
Date of Manufacture:	

UN 3352
PYRETHROID PESTICIDE, LIQUID, TOXIC
(lambda-cyhalothrin) MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2X
CONTAIN SPILLAGE

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours).

AGRECOVERY LOGO

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/04/2024