MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/02/2026

DANGER

KEEP OUT OF REACH OF CHILDREN

DITHANE™ RAINSHIELD™

Neo Tec™

Fungicide

For the control of fungal diseases in agricultural, horticultural and ornamental crops

Active ingredient
Contains 750 g/kg MANCOZEB in the form of a water dispersible granule

GROUP	M3	FUNGICIDE

Net contents: 10 kg

Bar code to be inserted

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of registration.
Any person purchasing, handling or disposing of this product must be suitably qualified.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/02/2026

DANGER
Causes serious eye irritation. May cause allergy or asthma symptoms or breathing difficulties if inhaled. May cause an allergic skin reaction. May cause damage to organs (thyroid, liver) through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS

PREVENTION
DO NOT breathe dust, vapours or spray mist. In case of inadequate ventilation wear respiratory protection. Contaminated work clothing should not be allowed out of the workplace.
Wash hands and face thoroughly after handling.
When mixing or applying wear eye protection, chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, chemical resistant boots and cotton overalls buttoned to the neck and wrist. In case of inadequate ventilation wear respiratory protection. Contaminated work clothing should not be allowed out of the workplace.
DO NOT apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

RESPONSE
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
If eye irritation persists: Get medical advice.
IF INHALED: remove person to fresh air and keep comfortable for breathing.
If experiencing respiratory symptoms: Call a POISON CENTRE.
IF ON SKIN: Wash with plenty of soap and water.
If skin irritation or rash occurs: Get medical advice.
Take off contaminated clothing and wash it before re-use.
Get medical advice/attention if you feel unwell.
Records of use must be kept.
Collect spillage.

STORAGE
Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.
DO NOT allow package to become wet or overheated in storage; decomposition, impaired activity or fire may result. DO NOT store near food, feedstuffs, fertilisers or seed.

DISPOSAL
Dispose of contents/container in accordance with local regulations.
Container disposal: Puncture or shred and bury empty bags in a local authority landfill. If no landfill is available, bury the bags below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.
Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID
If medical advice is needed, have the product container or label at hand. If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre (0800 POISON) 0800 764 766 or a doctor immediately. If INHALED and breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing. Call a doctor. If experiencing respiratory symptoms: Call a poison centre.

SPILLAGE
Wear appropriate protective clothing and prevent material from entering waterways. Sweep up solids without generating dust. Contain liquid spills and absorb with inert material. Dispose of waste safely in an appropriate landfill.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at **www.uplcorp.com/nz**.

EXCLUSION OF LIABILITY – PLEASE READ
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to:
UPL New Zealand Ltd
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
www.uplcorp.com/nz

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/02/2026

Registered pursuant to the ACVM Act 1997, No. P004053.
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996. Approval Code HSR000737.
See www.epa.govt.nz for approval controls.

© Copyright UPL New Zealand Limited 2023, all rights reserved.
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited.
® Registered trademark

Batch number:	
Date of Manufacture:	

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622)

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Mancozeb 75%)
MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2X

Label – Back



MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/02/2026

GENERAL INFORMATION

RESISTANCE MANAGEMENT

DITHANE RAINSHIELD Neo Tec Fungicide is a dithio carbamate fungicide (Group M3). Some naturally occurring individual fungi may become resistant when excessive use of the same fungicide occurs. To maximize the effectiveness of fungicides, products containing alternate modes of action should be used in the spray program. Consult http://resistance.nzpps.org/ for alternate modes of action and the details of resistance management strategies (if any) for the crops listed on the label.

DIRECTIONS FOR USE

FRUIT AND BERRIES

CROP	DISEASE	RATE	COMMENTS
Apples, pears	Black spot, *Glomerella*, ripe spot, sooty blotch	160 g/100 L	Apply every 10-14 days beginning at green tip. In regions where *Glomerella* is not a problem, reduce late cover spray rates to 110 g/100 L.
Boysenberries, loganberries, raspberries	Leaf spot, rust, downy mildew (dry berry).	200 g/100 L	Apply every 10-14 days from green tip.
Citrus	Brown rot		Apply in May. If infection is serious begin applications in April and continue at 4-6 week intervals until late winter.
	Alternaria		Apply from October to April every 14-28 days.
	Verrucosis (Citrus Scab)		Apply at petal fall.
	Citrus Rust Mite		Apply at the first sign of mite activity around December- January. Reapply after 21-28 days if necessary.
Currants (red and black), gooseberries	Leaf spot		Apply at green tip, pre- flowering and thereafter at 2- 3 week intervals.
Grapes	Anthracnose (black spot), downy mildew, dead arm disease		Apply every 10-14 days commencing at bud burst, or in accordance with local recommendations.
Peaches, nectarines	Rust		Apply in early November. reapply if disease is severe. do not apply **DITHANE RAINSHIELD NEO TEC** to plums as defoliation may occur.
Strawberries	*Mycosphaerella* leaf spot		Apply every 10-14 days beginning in mid-June. In wet conditions, reduce spraying interval to 7 days.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/02/2026

ORNAMENTALS

CROP	DISEASE	RATE	COMMENTS
Ornamentals	Black spot of roses, flower blight and rust of chrysanthemums, septoria leaf blight of gladioli and gerberas, rust of	210 g/100 L	Apply at 7-10 day intervals

CROP	DISEASE	RATE	COMMENTS
	antirrbinum, pelargonium and hollyhock		

VEGETABLES

CROP	DISEASE	RATE	COMMENTS
Beans	Rust, Anthracnose	160 – 210 g/100 L	Apply at 7-10 day intervals following the first signs of disease.
Carrots	Cercospora leaf spot		Begin applications on appearance of disease and repeat at 7-10 day intervals.
Cabbage, cauliflower, broccoli, brussels sprouts	Downy mildew		Apply at 7-10 day intervals in the seedbed and field. In wet weather, reduce the spray interval to 3-7 days.
Celery	Septoria leaf spot		Apply at 3-5 day intervals in the seedbed and 7-10 day intervals in the field.
Cucurbits	Downy mildew, cercospora leaf spot, anthracnose, alternaria		Apply at 5-7 day intervals beginning when runners form or at the first sign of disease.
Lettuce	Downy mildew, pleaspora spot, ring spot		Apply at the first sign of disease and repeat at 7-10 day intervals.
Onions	Downy mildew		Apply at 7-10 day intervals.
Potatoes	Early and late blight	1.6 - 2.1 kg/ha	Apply 1.6 kg/ha at crop emergence and increase rate to 2.1 kg /ha as vines increase in size. Spray every 7-10 days throughout the season, reducing the spray interval to 3-4 days in wet conditions conducive to Blight infection.
Sweetcorn, maize	Northern leaf blight	2.1 kg/ha	Apply by aerial application at the first sign of disease. Repeat sprays as necessary.
Tomatoes	Early and late blight, anthracnose, leaf mould, septoria leaf spot	160 – 210 g/100 L	Apply every 7-10 days. Increase the rate to 210 g/100 L or reduce spray intervals under severe infection conditions.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/02/2026

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residues Levels for Agricultural Compounds.

WITHHOLDING PERIODS

Beans, Boysenberries, Broccoli, Brussel Sprouts, Cabbage, Carrots, Cauliflower, Celery, Citrus, Currants (Red and Black), Gooseberries, Grapes, Lettuce, Loganberries, Maize, Nectarines, Onions, Peaches, Potatoes, Raspberries, Strawberries, Sweetcorn: DO NOT apply within 14 days of harvest.
Cucurbits and Tomatoes: DO NOT apply within 1 day of harvest.

Grazing Restriction

The following restrictions on the use of this product must be observed. DO NOT feed treated crops or pasture to stock other than sheep used for leaf plucking in vineyards. Take care not to allow this product to drift onto pasture or crops that may be grazed.
DO NOT graze stock in orchards which have been treated with **DITHANE RAINSHIELD Neo Tec**.

Vineyards: DO NOT graze stock other than sheep in vineyards which have been treated with **DITHANE RAINSHIELD Neo Tec**. For the purpose of leaf plucking, sheep may graze vineyards providing they are not slaughtered for 2 months after leaving the vineyard.

RESTRICTED ENTRY INTERVALS

The following restricted entry intervals (REI) have been set for this product. Anyone entering the application area during the REI must use appropriate PPE.

– Apples, pears 5 days

– Stonefruit 6 days

– Grapes 8 days

– Onions 24 hrs

– Broccoli, brussels sprouts, cabbage, cauliflower 10 days

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/02/2026

MIXING
DO NOT premix **DITHANE RAINSHIELD Neo Tec** into a smaller tank. Pour **DITHANE RAINSHIELD Neo Tec** into a ¼ filled spray tank while the agitator is running then add remaining water. DO NOT let **DITHANE RAINSHIELD Neo Tec** settle out in an unagitated spray tank. When preparing spray solutions for use in a hand sprayer, premix as a slurry in a small container and add to sprayer containing ¼ - ½ final water volume and then add remaining water.

APPLICATION
Apply **DITHANE RAINSHIELD Neo Tec** at 160 – 210 g/100 L of water as a full coverage, dilute spray unless otherwise specified.
For low volume sprayers, adjust the mixing rate to give equivalent rates per hectare. Thorough, uniform coverage is essential for good disease control.
Using the Spray:
IMPORTANT: In order to select the appropriate nozzles for the required spray quality and to operate the equipment to minimize spray drift, this product must be applied in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409).

EQUIPMENT
Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).
Flush equipment thoroughly with clean water after use according to Cleaning Spray Equipment instructions below. Dispose of cleaning water according to the Disposal instructions below.
Cleaning of Spray Equipment
Rinsing: Empty the spray tank completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain spray unit and clean any filters in the tank, pump, lines, hoses and nozzles.
After cleaning the spray units as above, quarter fill with clean water and circulate the water through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.
Discard rinse water on land already sprayed or on wasteland away from desirable plants and water sources.
Decontamination: Following an application of **DITHANE RAINSHIELD Neo Tec** all spraying equipment must be thoroughly cleaned both inside and out with a suitable spray tank cleaner. Follow the instructions on the tank cleaner label for the decontamination process. DO NOT use chlorine-based tank cleaners.
Once decontamination has been completed, cleaning water should be discharged onto a designated disposal area, or onto unused land away from desirable plants and water sources.

COMPATIBILITY

DITHANE RAINSHIELD Neo Tec is compatible with most commonly used insecticides and miticides, however compatibility should be checked before using EC formulations. **DITHANE RAINSHIELD Neo Tec** is not compatible with Bordeaux mixture or lime sulphur.