DANGER
KEEP OUT OF REACH OF CHILDREN

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/08/2025

PENNCOZEB® DF
Fungicide

A protective fungicide for the control of fungous diseases on a wide range of crops, ornamentals and turf

Active ingredient
Contains 750g/kg MANCOZEB in the form of a water dispersible granule, also contains 25g/kg HEXAMINE

GROUP	M3	FUNGICIDE

Net contents: 10 kg

Bar code to be inserted

Public

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.
Any person purchasing, handling or disposing of this product must be suitably qualified.



MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/08/2025

DANGER
Causes serious eye irritation.
May cause allergy or asthma symptoms or breathing difficulties if inhaled.
May cause an allergic skin reaction.
May cause damage to organs (Thyroid, Liver) through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS

PREVENTION
DO NOT breathe dust, vapours or spray mist.
In case of inadequate ventilation wear respiratory protection.
Contaminated work clothing should not be allowed out of the workplace.
Wash hands and face thoroughly after handling.
When mixing or applying wear eye protection, chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, chemical resistant boots and cotton overalls buttoned to the neck and wrist. In case of inadequate ventilation wear respiratory protection.
Contaminated work clothing should not be allowed out of the workplace.
DO NOT apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

RESPONSE
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
If eye irritation persists: Get medical advice.
IF INHALED: remove person to fresh air and keep comfortable for breathing.
If experiencing respiratory symptoms: Call a POISON CENTER.
IF ON SKIN: Wash with plenty of soap and water.
If skin irritation or rash occurs: Get medical advice.
Take off contaminated clothing and wash it before re-use.
Get medical advice if you feel unwell.
Records of use must be kept.
Collect spillage.

STORAGE
Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. DO NOT allow package to become wet or overheated in storage; decomposition, impaired activity or fire may result. DO NOT store near food, feedstuffs, fertilisers or seed.

DISPOSAL
Dispose of contents/container in accordance with local regulations.

Public

Container disposal: Puncture or shred and bury empty bags in a local authority landfill. If no landfill is available, bury the bags below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID
If medical advice is needed, have the product container or label at hand. If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre (0800 POISON) 0800 764 766 or a doctor immediately. If INHALED and breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing. Call a doctor. If experiencing respiratory symptoms: Call a poison centre.

SPILLAGE
Wear appropriate protective clothing and prevent material from entering waterways. Sweep up solids without generating dust. Contain liquid spills and absorb with inert material. Dispose of waste safely in an appropriate landfill.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at **www.uplcorp.com/nz**.

EXCLUSION OF LIABILITY - PLEASE READ
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as registrant, nor any distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to:
UPL New Zealand Ltd
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
www.uplcorp.com/nz/

Registered pursuant to the ACVM Act 1997, No. P004470
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval Code HSR000737
See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/08/2025

© Copyright UPL New Zealand Limited 2020, all rights reserved.
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited.

® Penncozeb is a registered trademark of Cerexagri BV

Batch number:	
Date of Manufacture:	
Expiry date:	

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622)

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (Mancozeb 75%)
MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2X



Public

GENERAL INFORMATION

DIRECTIONS FOR USE:

The rates suggested are for high volume spraying to just before run-off. For concentrate spraying adjust dilution rate accordingly. Use the higher rates and/or shorter spray intervals when disease pressure is high.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/08/2025

Knapsacks:

(1) Hydraulic and Knapsack sprayers. Use 15-20g per 10 L of water.

(2) Motorised mistblower. Use 45-60g per 10 L of water. Use middle settings on the restrictor and operate at sufficient throttle to give small droplets. Move all foliage in the airstream.

FRUIT CROPS

CROP	DISEASE	RATE	COMMENTS
Apples, pears	Blackspot, Glomerella, ripe spot, sooty blotch	150-200 g/100 L water	Begin at green tip and repeat every 10-14 days until harvest. Maintain higher rate where Glomerella is a problem. N.B. Early season use under frosty conditions may cause russet.
Bushfruit, canefruit	Leaf spot, rust, downy mildew (dryberry)	200 g/100 L water	Begin at green tip and repeat at 10-14 day intervals until the start of flowering.
Citrus	Brown rot, Alternaria, citrus rust mite	200 g/100 L water	Apply every 4-6 weeks from April until late winter. Apply from October to April at 14-28 days intervals. Apply at first signs of mite activity, normally December/January. Repeat 21-28 days later if necessary.
Grapes	Downy mildew, Anthracnose	200 g/100 L water	Apply pre-bloom, post bloom and repeat at 10-14 day intervals.
Peaches, nectarines	Rust	200 g/100 L water	Apply early November. A second or third application may be necessary. N.B. Do not allow drift onto plums as foliage damage and defoliation may occur.
Strawberries	Mycosphaerella leaf spot (frog's eye)	200 g/100 L water or 2 kg/ha	Apply every 10-14 days from mid-June until the start of flowering. During wet weather spray every 7 days.

ORNAMENTALS

CROP	DISEASE	RATE	COMMENTS
Roses	Black spot, rust	150-200 g/100 L water	Begin spraying at bud burst and continue at 7-14 day intervals throughout the growing season.
Other ornamentals	Rust, Septoria leaf spot	200 g/100 L water	Apply on appearance of disease and as required until disease is under control.

Public

VEGETABLES

CROP	DISEASE	RATE	COMMENTS
Beans	Anthracnose, rust	150-200 g/100 L water	Apply at first signs of disease and repeat every 7-10 days until the start of flowering.
Cabbage **Cauliflower** **Broccoli** **Brussels sprouts**	Downy mildew	150-200 g/100 L water or 2-3 kg/ha	Plant beds. Apply every 3-4 days in wet weather, otherwise every 7-10 days.
Celery	Septoria leaf spot	(a) 150-200 g/100 L water (b) 2-3 kg/ha	(a) Plant beds. Apply every 3-4 days in wet weather, otherwise every 7-10 days. (b) Field. Apply every 7-10 days from first appearance of disease.
Carrots	Cercospora leaf spot	2-3 kg/ha	Apply every 7-10 days or as required.
Pumpkin **Cucumber** **Melons** **Squash**	Downy mildew, Cercospora leaf spot, Anthracnose	2-3 kg/ha or 200 g/100 L water (high volume)	Begin spraying when runners begin to form and repeat as required.
Lettuce	Downy mildew, ring spot, leaf spot	2 kg/ha	Spray on appearance of disease and repeat at 7-10 day intervals.
Onions	Downy mildew, Botrytis	2-3 kg/ha	Begin spraying at the 2 leaf stage and repeat at 7-10 day intervals.
Maize **Sweetcorn**	Northern leaf blight	2 kg/ha	As an aerial treatment in sufficient water to cover. Repeat as necessary.
Potatoes	Late blight, early blight (target spot)	1.5 kg/ha (crop emergence) 2.0 kg/ha (as vines size)	In wet weather apply every 3-4 days, otherwise every 7-10 days.
Tomatoes	Late blight, early blight, Septoria leaf spot	150-200 g/100 L water (high volume) or 1 kg/ha (planting) to 3 kg/ha (as vines size)	Apply at 7-10 day intervals or more frequently in wet weather. Do not use on young tomato seedlings in glasshouses or tunnel houses as injury may occur.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/08/2025

Public

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residues Levels for Agricultural Compounds.

It is an offence to use this product on animals.

MIXING
When filling the spray tank wash the correct amount of **PENNCOZEB DF** through the "filler screen" of the spray tank or thoroughly premix in a bucket and pour into the partly filled tank through a filter. Keep agitated during use. Agitate thoroughly after a stoppage before recommencing spraying. For hard-to-wet vegetables (e.g. cabbage, cauliflower, carrots and onions), the addition of a non-ionic wetting agent will improve coverage.

RESISTANCE MANAGEMENT
PENNCOZEB DF is a dithio carbamate fungicide (Group M3). Some naturally occurring individual fungi may become resistant when excessive use of the same fungicide occurs. To maximize the effectiveness of fungicides, products containing alternate modes of action should be used in the spray program. Consult http://resistance.nzpps.org/ for alternate modes of action and the details of resistance management strategies (if any) for the crops listed on the label.

EQUIPMENT
Apply using well maintained spray equipment. Clean equipment thoroughly after use. **PENNCOZEB DF** is suitable for use in most agricultural sprayers. Do not use strainers finer than 300 micrometres. Calibrate the equipment to give the desired application rate and full coverage of the foliage. Fine droplets are more effective than coarse (large) droplets. Ensure the undersides of the plant are treated.

COMPATIBILITY: PENNCOZEB DF is compatible with most commonly used insecticides and fungicides. Do NOT use with lime sulphur or Bordeaux mixture.

Leaf plucking in vineyards: Sheep must not be sent for slaughter or milked for 6 months after being removed from the vineyard (and placed on clean feed).

WITHHOLDING PERIODS:
Potatoes: 7 days
Tomatoes: 7 days (field), 2 days (indoor)
Cucurbits: 14 days (field), 2 days (indoor)
Pipfruit, citrus, onions, celery, lettuce: 14 days
Grapes: 14 days
Strawberries, bushfruit, canefruit: Use only up until the start of flowering
Vegetable brassicas: Seedlings only
Stonefruit: To avoid spray deposits on fruit, do not spray within 6 weeks of harvest.

RESTRICTED ENTRY INTERVALS
The following restricted entry intervals (REI) have been set for this product. Anyone entering the application area during the REI must use appropriate PPE.
– Apples, pears 5 days
– Stonefruit 6 days
– Grapes 8 days
– Onions 24 hrs
– Broccoli, brussels sprouts, cabbage, cauliflower 10 days

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/08/2025

Public

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/08/2025

Public