18.5”

0.625” | 3.8125” RIGHT Back | 8.625” Front | 4.4375” LEFT Back | 0.375” | 0.625”

Eyespot: 0.25” x 0.625”

Delfin®

BIOLOGICAL INSECTICIDE

For the control of caterpillars on APPLES, KIWIFRUIT, AVOCADOS, GRAPES, CITRUS, BERRY FRUITS, VEGETABLE BRASSICAS, and TOMATOES

APPLICATIONS METHOD
For dilute application, spraying to point of run-off is recommended to ensure all plant surfaces are adequately covered. Where concentrate airblast application is made, it is important that uniform distribution of spray droplets is achieved.

Crops	Pest	Rate/100 Litres	Comments
Apples	Leafrollers	50g/100 litres water	Make the first application in late October to coincide with the first generation egg hatch. A programme of three applications at 7 - 10 day intervals is recommended. Further applications should be made whenever leafroller monitoring thresholds are exceeded.
Avocados	Leafrollers	50g/100 litres water	Ensure complete spray coverage. Apply at early flowering mid and post blossom. From late November till harvest monitor for leafrollers and repeat applications at 10 - 14 day intervals if thresholds are exceeded.
Berry Fruits	Leafrollers	60g/100 litres water	Apply at 10 - 14 day intervals over flowering. Then monitor for leafroller and repeat applications if thresholds are exceeded.
Citrus	Leafrollers	50g/100 litres water	Apply at 10 - 14 day intervals from flowering until late January. Then monitor for leafroller and repeat applications if thresholds are exceeded.
Grapes	Leafrollers	50g/100 litres water	Apply at 80% capfall repeat at 10 - 14 day intervals. From late December monitor for leafrollers and repeat applications if thresholds are exceeded.
Kiwifruit	Leafrollers	50g/100 litres water	Apply the first application at flowering followed by two further sprays at 14 day intervals. From January to harvest, monitor for leafrollers and repeat applications if thresholds are exceeded.
Tomatoes	Tomato fruitworm	50g/100 litres water (minimum 500g/ha)	Apply at first sign of insect attack. Repeat at 10 - 14 day intervals while caterpillars persist.
Vegetable brassicas	White butterfly, Diamondback moth	50g/100 litres water (minimum 500g/ha)	Apply at 7 - 10 day intervals from transplanting. Use the longer interval if insect pressure is low. Add a wetting agent to aid coverage. Resistance management for diamondback moth is to use Bt products sush as Delfin against young larvae on small plants.

WITHHOLDING PERIOD..... 1 day all crops.

Registered to: Hilado Pty Ltd, 7 Marriott Grove, Castle Hill, NSW 2154, Australia.
NZ Agent: Mitsul & Co. (NZ) Ltd, Level 14, 23 Customs Street East , Auckland 1010.

Net Contents: 1.0 kg

GENERAL INFORMATION
DELFIN must be eaten by the caterpillar to be effective and shortly after ingestion the larvae stop feeding. The treated larvae will die after 1 to 3 days. DELFIN is not harmful to the following beneficial insects: ladybugs, lacewing, syrphid flies, bees, and wasps.

Integrated Mite Control and Integrated Pest Management:
DELFIN is compatible with IMC and IPM programmes involving *T.pyri, T.occidentalis, A.fallacis, P.persimilis,* and *Encarsia formosa.*

DIRECTIONS FOR USE

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS

TIME OF APPLICATION
DELFIN should be applied at the first sign of insect infestation or damage, thereby treating hatching eggs and the first larval stages. Apply DELFIN during active feeding periods to maintain control and protect fruit and new foliage. Repeat at 10-14 day intervals. Use the shorter interval during wet conditions.

APPLICATION METHOD
For dilute application, spraying to point of run-off is recommended to ensure all plant surfaces are adequately covered. Where concentrate airblast application is made, it is important that uniform distribution of spray droplets is achieved.

MIXING
Half fill the spray tank with water and start agitation. Pour the amount of product onto the surface of the water whilst maintaining continuous agitation. Add other spray materials and balance of water until the tank is full. Agitation should be maintained during spraying. Do not allow diluted spray to stand in the sprayer tank more than 12 hours prior to usage.

COMPATIBILITY
DELFIN is compatible with commonly used oils, fungicides and insecticides but not with products that are either alkaline in reaction (e.g. Bordeaux mixture, lime sulphur) or based on heavy metals (e.g. copper based products). Mixing with foliar nutrients is not recommended.

WARRANTY
This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Item: 110163-NZ
Package: 516801-NZ
REV-A

Date of Manufacture:

Batch Number:

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
18/09/2025

Active Ingredient: Contains a minimum of 37,000 International Units/mg *Bacillus thuringiensis,* subspecies *kurstaki* (serotype 3a, 3b, strain SA 11) in the form of a water dispersible granule.

EMERGENCY RESPONSE (ALL HOURS)
0800 CHEMCALL or 0800 243 622.
Contact PGG Wrightson Freephone 0800 10 22 76.

PLEASE READ THE LABEL COMPLETELY BEFORE USE

HSNO CLASSIFICATIONS:
Designed for biocidal action

ECOTOXIC
Harmful to Lepidoptera: Do not apply this substance in a way that adversely affects native Lepidoptera of a species found only in the region of application. Do not apply directly into or onto water.

HANDLING PRECAUTIONS
STORAGE: Keep out of reach of children. Store in original container, tightly closed in a locked, cool, dry, well ventilated place. Prolonged exposure above 30°C may damage this product. Storage must be in accordance with current version of NZS 8409 Management of Agrichemicals. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.
PERSONAL PROTECTION: May cause mild eye irritation and mild skin irritation. Avoid contact with eyes, skin and open wounds. When mixing or applying, wear protective clothing including water proof gloves and goggles. Wash hands and exposed skin before meals and after work.
EQUIPMENT: Apply using accurately calibrated equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS8409).
PRODUCT DISPOSAL: Dispose of this product only by using according to the label or at an approved landfill.
CONTAINER DISPOSAL: Triple rinse container and add rinsate to spray tank. Burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise bury in a suitable landfill. Avoid contamination of any water supply with chemical or empty container.

FIRST AID
If swallowed do not induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. Do not scrub the skin. If splashed in the eyes, wash out immediately with water.

SPILLAGE: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely in approved landfill.

Registered pursuant to the ACVM Act 1997, No. P4529.
Refer to www.foodsafety.govt.nz for registration conditions.
HSNO Approval Code HSR000699.
See www.epa.govt.nz for approval controls.

PGG Wrightson

PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport 8053
PO Box 292, Christchurch 8140
www.pggwrightson.co.nz

1.875

14.0000”
Repeat

0.875

0.1875”