KEEP OUT OF REACH OF CHILDREN

MICROTHIOL® DISPERSS

Fungicide

For the control of certain fungal diseases in apples, grapes, stonefruit, tamarillos, vegetables, and ornamentals.

ACTIVE INGREDIENT
800 g/kg SULPHUR in the form of a water dispersible granule

GROUP	M2	FUNGICIDE

Net contents: 15 kg

Bar code to be inserted

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
06/12/2024

PLEASE READ THE LABEL COMPLETELY BEFORE USE

Designed for biocidal action.

PRECAUTIONARY STATEMENTS
PREVENTION
When handling wear eye protection and gloves.
Wash hands thoroughly after handling.
Do not eat drink or smoke when using this product.
RESPONSE
Collect spillage.
STORAGE
Keep out of reach of children. Store in accordance with NZS8409 Management of Agrichemicals.
When stored appropriately this product should show no significant degradation for 3 years. Contact your supplier for further information about the use of any product that is older than this.
DISPOSAL
Ensure container is completely empty. Bury in a suitable landfill. Dispose of product only by using according to the label, or at an appropriate landfill.

FIRST AID
If swallowed, do NOT induce vomiting. For advice contact the National Poisons Centre (0800 POISON) 0800 764 766 or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If in the eyes, wash out immediately with water.

SPILLAGE
Wear appropriate protective clothing and prevent material from entering waterways. Sweep up solids without generating dust. Contain liquid spills and absorb with inert materials. Dispose of waste safely in an appropriate landfill.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at **www.upl-ltd.com/nz**

EXCLUSION OF LIABILITY
Please read these conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened.
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to:
UPL New Zealand Ltd
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
www.upl-ltd.com/nz

Registered pursuant to the ACVM Act 1997, No. P004795.
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000816.
See www.epa.govt.nz for approval controls.

© Copyright UPL New Zealand Limited 2020, all rights reserved.
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited
® Microthiol is a registered trademark of Cerexagri.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
06/12/2024

Batch number:	
Date of Manufacture:	

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622).
UN-Free

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
06/12/2024

DIRECTIONS FOR USE

CROP	DISEASE	RATE	COMMENTS
Apples (Do NOT apply to Pears)	Powdery mildew	100-200g/100 litres water	Apply from bud movement until harvest at 10-14 day intervals. Do NOT apply in combination with captan on sulphur sensitive varieties such as Kidd's Orange Red, Cox's Orange Pippin or Red Delicious. Fruit finish my be affected on some apple varieties. Do NOT use on apples post-blossom unless one sulphur containing spray has been used pre-blossom. Do NOT apply in temperatures above 25°C.
Grapes	Powdery mildew, Erineum mite	150-300g/100 litres water	Apply at 10-14 day intervals from bud-swell until 6 weeks before harvest. Use the higher rate when conditions favour disease.
Stonefruit (Do NOT apply to apricots)	Rust	100-200g/100 litres water	Apply at 21 day intervals from fruit set until harvest. Use the higher rate when conditions favour the disease.
Tamarillos	Powdery mildew	100-200g/100 litres water	Apply at 21 day intervals throughout the growing season.
Vegetables (Do NOT apply to Cucumbers, Rockmelons & Tomatoes)	Powdery mildew, rust	100-200g/100 litres water	Apply every 10-14 day intervals when disease infection is likely.
Ornamentals	Powdery mildew, Rust	100-200g/100 litres water	Apply every 10-14 day intervals when disease infection is likely.

It is an offence to use this product on animals.

MIXING
The rates suggested are for high volume spraying to just before run-off. For concentrate spraying adjust dilution rate accordingly. Use the higher rates and/or shorter spray intervals when disease pressure is high. The recommended quantity of **MICROTHIOL DISPERSS** may be added directly to the water in the spray tank with the agitation running. If the leaves are hairy and/ or waxy, or where the water is hard, add a suitable wetting agent at recommended rates.

RESISTANCE MANAGEMENT
MICROTHIOL DISPERSS contains the group M2 fungicide, sulphur. To minimise risk maintain the label dose rate. Use as part of a spray program in combination with other fungicides with different modes of action where relevant.

EQUIPMENT
MICROTHIOL DISPERSS is suitable for use in most agricultural sprayers. Do not use strainers finer than 300 micrometres. Calibrate equipment to give the desired application rate and full coverage of foliage. Fine droplets are more effective than coarse (large) droplets.

COMPATIBILITY
MICROTHIOL DISPERSS should not be used in combination oils or formulations containing oils, or used within 21 days of an oil application. The product is compatible with most other formulations of fungicides and insecticides.

WITHHOLDING PERIOD/S
All crops: 1 day.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
06/12/2024

Leaf plucking in vineyards: Sheep must not be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
06/12/2024