WARNING

KEEP OUT OF REACH OF CHILDREN

Switch®
Fungicide

syngenta®

Active Ingredients:
SWITCH® contains 375 g/kg CYPRODINIL and 250 g/kg FLUDIOXONIL in the form of a water dispersible granule.

GROUP	9	12	Fungicide

For the control of Botrytis in Blackcurrants, Bulb Onions and Grapes; Botrytis and Anthracnose in Blueberries; and Botrytis, Anthracnose and Rhizopus in Strawberries

Net Contents: **1 kg and 5 kg**

Registered pursuant to the ACVM Act 1997, No. P5113.
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code HSR000645.
See www.epa.govt.nz for approval controls.

Syngenta Crop Protection Limited
Level 4, 60 Parnell Road, Parnell, Auckland 1052

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
17/12/2024

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.
ny person mixing, loading or handling this product MUST be a qualified loader or under the guidance of a qualified loader.
Any person applying this product MUST be a qualified contractor/person or under the guidance of a qualified contractor/person.

GHS Classifications:
Specific target organ toxicity (repeated exposure) Category 2
Hazardous to the aquatic environment acute Category 1
Hazardous to the aquatic environment chronic Category 1
WARNING
May cause organ damage through prolonged or repeated oral exposure at high doses.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONS
Store in original container, tightly closed, away from foodstuffs.
When mixing or applying wear protective clothing, including face shield, chemical resistant gloves and footwear.
Avoid contact with skin and eyes and inhalation of concentrate or spray mist.
If clothing becomes contaminated with product, remove clothing immediately.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking.
Wash protective clothing daily after work.
Apply with well-maintained and calibrated spray equipment.
DO NOT re-enter site until spray has dried.
Records of use MUST be kept as per Clause 48 of the Hazardous Property Controls Notice 2017.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
DO NOT carry on a passenger service vehicle.
Avoid contamination of any water supply with chemical or empty container. DO NOT apply into or onto water.

Container disposal: ***For hermetic boxes:*** Ensure the container is empty. Rinse empty container and add the residue to the spray tank. If allowed under local Territorial Authority District Plans, burn if circumstances, especially wind direction permit, otherwise crush and bury in an approved local authority facility. DO NOT reuse this container for any other purpose.
For HDPE Bottles: Ensure the container is empty. Triple rinse the empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in an approved local authority facility. DO NOT reuse this container for any other purpose.
Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility. For additional information refer to EPA Notice: Hazardous Substances (Disposal) Notice 2017.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
17/12/2024

FIRST AID
If SWALLOWED, DO NOT induce vomiting. Rinse Mouth.
If SKIN or HAIR contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin.
If splashed in the EYES, wash out immediately with water. Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.
If INHALED, move the victim to fresh air immediately.
Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth. Obtain medical attention. Take to hospital without delay.
For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

SPILLAGE
Collect spillage. Wear suitable protective clothing including face shield, chemical resistant gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

GENERAL INFORMATION
SWITCH® is a mixture of aniline-pyrimidine and phenylpyrrole fungicides, which in combination have protectant and systemic activity against *Botrytis* (grey mould) infections in blackcurrants, bulb onions and grapes and *Botrytis* (grey mould) and *Colletotrichum* (anthracnose) in blueberries and *Colletotrichum* (anthracnose) and *Rhizopus* (leak) infections in strawberries. SWITCH® has multi-site action, inhibiting spore germination, penetration of the germ tube and mycelial growth. It is rainfast within 2 hours following application.

RESISTANCE MANAGEMENT | **GROUP 9 12 FUNGICIDE**
SWITCH® contains the GROUP 9 fungicide, cyprodinil, belonging to the aniline-pyrimidine class of chemistry) and the GROUP 12 fungicide, fludioxonil, belonging to the phenylpyrrole class of chemistry). Resistance to this fungicide and related anilinopyrimidine products could develop in some situations from repeated use. To minimise this risk use strictly in accordance with the label instructions.

Blackcurrants, Blueberries, Grapes and Bulb Onions: DO NOT use more than 2 applications of SWITCH® or any other anilinopyrimidine fungicide on the same crop during the season.

Strawberries: A maximum of 3 applications of SWITCH® or any other anilinopyrimidine fungicide per season can be made if more than 6 sprays in total are made for *botrytis* control. If 6 or less *botrytis* control sprays are made then DO NOT apply more than 2 applications of SWITCH® or any other anilinopyrimidine fungicide per season.

Refer to the NZPPS web site (http://resistance.nzpps.org) for more detailed information.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
17/12/2024

DIRECTIONS FOR USE
It is an offence to use this product on animals.

Crop	Disease	Rate	Critical comments
Blackcurrants	Grey Mould (*Botrytis)*,	160 g/100 litres of water applied as a dilute spray to the point of runoff	A maximum of 2 applications of SWITCH® can be made per season. Applications should be made using a water rate sufficient to obtain good coverage without runoff (approximately 500 L per hectare). Make 1 to 2 applications over flowering according to disease pressure.
Blueberries	Grey Mould (*Botrytis)*, Anthracnose (*Colletotrichum*)	80 g/100 litres of water applied as a dilute spray to the point of runoff	A maximum of 2 applications of SWITCH® can be made per season. Applications should be made using a water rate sufficient to obtain good coverage without runoff (approximately 1000 L per hectare). The first application should be made at full flower with the second application no later than 14 days after the completion of flowering.
Bulb Onions	Botrytis	800 g/ha	A maximum of 2 applications of SWITCH® can be made per season at key timings for botrytis control i.e. flag leaf and pre harvest. Apply in sufficient water to ensure thorough and uniform coverage. Applications MUST be made with the addition of a non-ionic wetting agent at manufacturer's label rates.
Grapes	Botrytis	80 g/100 litres of water applied as a dilute spray to the point of runoff	The rate suggested is for high volume spraying to run off. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage of the bunch zone is achieved. The addition of a non-ionic wetting agent at manufacturer's label rates will assist coverage. A maximum of 2 applications per season can be made. SWITCH® can be applied at either early flowering (5% capfall), late flowering (80% capfall), pre bunch closure or in the period from the beginning of berry ripening up until 28 days before harvest. Apply an approved fungicide from a different chemical group for the remainder of the season.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
17/12/2024

Crop	Disease	Rate	Critical comments
Strawberries	Grey Mould (*Botrytis)*, Anthracnose (*Colletotrichum*) and leak (*Rhizopus*)	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	The rate suggested is for high volume spraying to run off. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved. The addition of a non-ionic wetting agent at manufacturer's label rates will assist coverage. SWITCH® can be applied at 7 to 14 day intervals from flowering onwards. The closer interval is recommended when conditions are conducive to disease infection. SWITCH® applications may be applied in a block or in alternation with other non anilinopyrimidine fungicides registered for *botrytis* control. See resistance management for the maximum number of applications.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
17/12/2024

SPRAY DRIFT
Take all reasonable steps to ensure that SWITCH® does NOT cause any adverse effects to the environment beyond the application area.

MIXING
Add the required amount of SWITCH® into the mixing chamber of the partly filled spray tank and add the remainder of the water ensuring complete mixing before spraying. Continue agitation during spraying.

COMPATIBILITY
SWITCH® is compatible with other commonly used fungicides and insecticides.

WITHHOLDING PERIOD
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds
Blackcurrants ..... DO NOT apply after the completion of flowering / petal fall.
Blueberries ..... Last application no later than 14 days after flowering and at least 28 days before harvest.
Bulb Onions ..... 7 days
Grapes ..... 28 days
Strawberries ..... 3 days
Leaf plucking in vineyards
Sheep MUST NOT be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

STORAGE
Store in a cool, dry place, avoiding temperatures below minus 10°C or above 35°C. See Safety Data Sheet for additional storage information. Stores containing quantities of 100 kg or more require signage and secondary containment.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at www.syngenta.co.nz.

DISCLAIMER
This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage

and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland 1052. Phone 0800 333 336.

Bar code:

Batch Number:	
Date of Manufacture	

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
17/12/2024

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S.
(cyprodinil and fludioxonil) MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2Z

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours)

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
17/12/2024