MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/05/2025

DANGER
KEEP OUT OF REACH OF CHILDREN

Topas® 200 EW
Fungicide

syngenta®

Active Ingredient:
TOPAS® 200 EW contains 200 g/litre PENCONAZOLE in the form of an oil in water emulsion. Also contains: 139 g/litre hydrocarbon liquids.

GROUP	3	FUNGICIDE

For the control of Powdery Mildew in Grapes, Cucurbits, Peas and Apples and Blackspot in Apples and Pears.

Net Contents: 1 litre

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT.

Registered pursuant to the ACVM Act 1997, No. P5374.
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code HSR000592.
See www.epa.govt.nz for approval controls.

Syngenta Crop Protection Limited
Level 4, 60 Parnell Road, Parnell, Auckland 1052

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/05/2025

GHS Classifications
Eye irritation Category 2
Specific target organ toxicity (repeated exposure) Category 1
Hazardous to the aquatic environment chronic Category 2
DANGER
Causes serious eye irritation.
Causes liver damage through prolonged or repeated oral exposure at high doses.
Toxic to aquatic life with long lasting effects.

PRECAUTIONS
Store in original container, tightly closed, away from foodstuffs.
When mixing or applying wear protective clothing including, overalls, chemical resistant gloves, face shield and footwear.
Avoid contact with skin and eyes and inhalation of the spray mist.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking.
Wash protective clothing after use.
Apply with well maintained and calibrated spray equipment.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
DO NOT carry on a passenger service vehicle.
Avoid contamination of any water supply with chemical or empty container.
DO NOT apply this product into or onto water.

Container disposal: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle punctured container without caps through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility. For additional information refer to EPA Notice: Hazardous Substances (Disposal) Notice 2017.

FIRST AID
If SWALLOWED, DO NOT induce vomiting. Rinse Mouth.
If SKIN or HAIR contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If skin irritation occurs seek medical advice.
If in EYES, wash out immediately with water. Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.
If INHALED, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth. Obtain medical attention. Take to hospital without delay.
Get medical advice/attention if you feel unwell. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

SPILLAGE
Collect spillage. Wear suitable protective clothing including face shield, chemical resistant gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

GENERAL INFORMATION

TOPAS® 200 EW is a systemic triazole (DMI) fungicide with preventative and curative properties for the control of Powdery Mildew in a range of crops and Blackspot in pipfruit. TOPAS® 200 EW penetrates rapidly into the leaf tissue, where it acts curatively on the developing fungal infections by inhibiting the creation of sterols in fungal cell walls. In apples, this action provides control of primary mildew infections and reduces both the infection of the fruit, which can occur soon after petal fall, and infected buds, which are a source of infection for the following season. In addition, TOPAS® 200 EW in combination with a protectant fungicide, controls Blackspot by both a contact action on fungal spore germination and curatively within the leaf during the incubation period before symptoms appear.

RESISTANCE MANAGEMENT

GROUP 3 FUNGICIDE

TOPAS® 200 EW is a member of the demethylation inhibitor (DMI) group of fungicides. For fungicide resistance management TOPAS® 200 EW is a Group 3 fungicide. Some naturally occurring individual fungi resistant to TOPAS® 200 EW and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by TOPAS® 200 EW and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Crop Protection Limited accepts no liability for any losses that may result from the failure of TOPAS® 200 EW to control resistant fungi. Consult your local supplier, consultant, a Syngenta representative or New Zealand Plant Protection Society (Inc) https://resistance.nzpps.org/ for alternate modes of action and the details of resistance management strategies for the crops listed on the label.

PIPFRUIT

Apply a maximum of two DMI-containing products per season and only in mixture with an effective dose of a broad spectrum protectant fungicide that is not at risk from resistance.
A maximum of four DMI-containing applications can be made in extremely wet seasons, but only where there is no history of black spot control failure following DMI applications or where the orchard has a history of low DMI use.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/05/2025

DIRECTIONS FOR USE

It is an offence to use this product on animals

Crop	Disease	Rate	Critical Comments
Apples, Pears	Blackspot	12.5 mL/100 litres of water as dilute spray	For high volume spraying, apply as a dilute spray, ensuring complete coverage, without excessive run off. Apply in mixture with an approved protectant fungicide. For concentrate spraying, adjust dilution rate accordingly. **Preventative:** Apply in mixture with a protectant fungicide at no greater than 10 day intervals preferably from tight cluster to second cover. **Curative:** Apply TOPAS® 200 EW plus a protectant fungicide as soon as possible but within 4 days of each Blackspot infection period.
Apples	Powdery Mildew		**Preventative:** Apply in mixture with an approved protectant fungicide at 10 to 14 day intervals during the period from tight cluster to the end of extension growth.
Cucurbits	Powdery Mildew	125 mL/ha. For high volume spraying, mix 12.5 mL/100 litres of water ensuring a minimum of 1000 litres of water/ha	As a low volume broadcast spray, apply in sufficient water to ensure complete foliage cover. Make the first application as a preventative spray and repeat 14 days later. To minimise the risk of developing resistant disease strains, apply a maximum of 2 applications of TOPAS® 200 EW, preferably in mixture with an approved protectant fungicide. Use non DMI fungicides at other times. DO NOT use as an eradicant when Powdery Mildew is widespread.

Crop	Disease	Rate	Critical Comments
Grapes	Powdery Mildew	7.5 to 12.5 mL/100 litres of water as a dilute spray.	Early in the season when disease pressure is low, use the lower rate at 14 to 21 day intervals. Under high disease pressure, use the higher rate at 14 day intervals. To minimise the risk of developing resistant disease strains, fungicides of other chemical groups, such as sulphur, should be used from bud burst as early season sprays for Powdery Mildew control. The first TOPAS® 200 EW application should be made during early flowering, with repeat applications at 14 to 21 day intervals up to a maximum of 4 sprays, preferably in mixture with an approved protectant fungicide. No more than two DMI fungicides should be applied alone per season. DO NOT use as an eradicant.
Peas	Powdery Mildew	125 to 175 mL/ha	As a low volume broadcast spray, apply in sufficient water to give complete foliage cover without run off. Apply the higher rate if the first signs of the disease are evident. The addition of wetting agent at manufacturer's label rates is recommended. Make a single application, preferably no later than flowering, either as a preventative spray before the disease is present, or at the first sign of disease symptoms. DO NOT use as an eradicant when Powdery Mildew is widespread.

SPRAY DRIFT
Take all reasonable steps to ensure that TOPAS® 200 EW does NOT cause any adverse effects to the environment beyond the application area.

MIXING
TOPAS® 200 EW mixes readily with water. Add the required amount of TOPAS® 200 EW to the partly filled spray tank and then add the remainder of the water with agitation.

APPLICATION
Aerial Application:
Apply in a minimum of 65 litres of water/ha to ensure more than 20 droplets per square centimetre on all leaves. As there are many factors involved in aerial application, for optimum results, contact your local Syngenta Crop Protection representative for detailed information.

Application in vine crops
The application rate is for high volume spraying to the point of runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved.
Set up and operate the sprayer to achieve even coverage throughout the crop canopy, applying sufficient water to cover the crop to just prior to the point of runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

COMPATIBILITY
TOPAS® 200 EW is compatible with other fungicides and insecticides likely to be used, except oil and strongly alkaline materials such as Bordeaux mixture and lime sulphur.
Note: As formulations of other manufacturers products are beyond the control of Syngenta Crop Protection Limited, all mixtures should be tested for crop safety and physical compatibility prior to mixing commercial quantities.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/05/2025

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

Cucurbits ..... 3 days
Peas ..... 14 days
Grapes ..... 28 days
Pipfruit ..... 35 days

Leaf plucking in vineyards:

Sheep MUST NOT be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

STORAGE

Store in cool, dry place, out of direct sunlight, avoiding temperatures below minus 5°C and above 35°C. See Safety Data Sheet for additional storage information. Stores containing quantities of 1000 litres or more require signage and secondary containment.

SAFETY DATA SHEET

If additional hazard information is required, refer to the Safety Data Sheet. For a copy, visit our website at www.syngenta.co.nz

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland 1052. Phone 0800 333 336.

Bar code:

Batch Number:	
Date of Manufacture:	

Agrecovery Logo

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
15/05/2025

UN-3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.
(naphtha solvent) MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 3Z
CONTAIN SPILLAGE

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours).