Thiovit® Jet

Fungicide

syngenta®

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/05/2025

Active Ingredient:
THIOVIT® Jet contains 800 g/kg SULPHUR in the form of a water dispersible granule.

For the control of certain diseases of
Apples, Grapes, Stonefruit, Tamarillos, Vegetables
and Ornamentals.

GROUP	M2	FUNGICIDE

Net Contents: 10, 15 and 25 kg

Registered pursuant to the ACVM Act 1997, No. P5819.
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code HSR000816.
See www.epa.govt.nz for approval controls.

Syngenta Crop Protection Limited
Level 4, 60 Parnell Road, Parnell, Auckland 1052

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours).

CERTIFIED INPUT FOR ORGANICS bio gro NZ CERT
5962

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/05/2025

GHS Classification
Designed for biocidal action.

PRECAUTIONS
Keep out of reach of children.
Store in original container, tightly closed, away from foodstuffs.
When mixing or applying, wear protective clothing including overalls, chemical resistant gloves and footwear.
Avoid contact with skin and eyes and inhalation of spray mist.
Wash hands and face thoroughly with soap and water before meals and after work.
DO NOT eat, drink or smoke while using.
Wash protective clothing daily after work.
Apply with well maintained and calibrated spray equipment.
Application and storage should be as described in the Management of Agrichemicals NZS8409.
DO NOT carry on a passenger vehicle.
Avoid contamination of any water supply with chemical or empty container.
DO NOT apply this product into or onto water.

Container disposal: Ensure the container is empty. If allowed under local Territorial Authority District Plans, burn if circumstances, especially wind direction permit, otherwise crush and bury in an approved local authority facility. DO NOT reuse this container for any other purpose.
Product disposal: Dispose of this product only by using according to the label or at an approved landfill or other approved facility. For additional information refer to EPA Notices: Hazardous Substance (Disposal) Notice 2017.

FIRST AID
If SWALLOWED, DO NOT induce vomiting. Rinse Mouth.
If SKIN or HAIR contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin.
If splashed in the EYES, wash out immediately with water. Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.
If INHALED, move the victim to fresh air immediately.
Begin artificial respiration if breathing has stopped. Use mouth to nose rather than mouth to mouth. Obtain medical attention. Take to hospital without delay.
For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

SPILLAGE
Wear suitable protective clothing including face shield, chemical resistant gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert materials such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

RESISTANCE MANAGEMENT **GROUP M2 FUNGICIDE**
THIOVIT® Jet is a member of the inorganic (M2) group of fungicides. Resistance to this fungicide could develop from repeated use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies. Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Crop Protection Ltd accepts no liability for any losses that may result from the failure of THIOVIT® Jet to control resistant fungi. Consult your local supplier, consultant, a Syngenta representative or New Zealand Plant Protection Society (Inc), www.resistance.nzpps.org for alternate modes of action and the details of resistance management strategies for the crops listed on the label.

DIRECTIONS FOR USE

The rates suggested are for high volume spraying to run off. Apply the higher rate when conditions favour the specific disease.

It is an offence to use this product on animals.

Crop	Disease	Rates	Critical Comments
Apples	Powdery Mildew	100 to 200 g /100 litres of water	Apply from bud movement until harvest. DO NOT apply in temperatures above 25°C. DO NOT apply in combination with captan. DO NOT apply to sulphur sensitive varieties such as Kidds Orange Red, Cox's Orange Pippin or Red Delicious. Fruit finish may be affected on some apple varieties. To avoid "sulphur shock", DO NOT use THIOVIT® Jet on apples post-blossom unless one sulphur containing spray has been used pre-blossom. DO NOT apply on pears.
Stonefruit (except apricots)	Rust	100 to 200 g /100 litres of water	Apply every 3 weeks from fruit set until harvest.
Grapes	Powdery Mildew, Erineum Mites	150 to 300 g /100 litres of water	Apply at 10 to 14 day intervals from bud swell until 6 weeks before harvest.
Tamarillos	Powdery Mildew	100 to 200 g /100 litres of water	Apply at 10 to 14 day intervals throughout the growing season.
Ornamentals, **Vegetables**	Powdery Mildew, Rust	100 to 200 g /100 litres of water	Apply at 10 to 14 day intervals when weather conditions are conducive for infection periods to occur. DO NOT apply to cucumbers or rockmelons.

MIXING

This product MUST be mixed with water and applied by suitable equipment.

1. Fill spray tank no more than 25% full with clean water BEFORE adding THIOVIT® Jet. Begin agitating tank contents vigorously and continue agitation during entire mixing and spraying operations.
2. Pour required amount of THIOVIT® Jet STEADILY into tank. Allow vigorous bypass agitation to completely disperse product. DO NOT dump product into tank all at once.
3. After adding required quantity of THIOVIT® Jet and obtaining COMPLETE DISPERSION, continue to fill tank to desired level for spraying. Thorough agitation of the spray liquid should continue during the entire spraying operation.

NB: Spray solution should NOT be left standing in the spray tank overnight

COMPATIBILITY

DO NOT combine with oils or use within 3 weeks of applying oil sprays. If physical compatibility is unknown, a small test mix in water first is recommended.

THIOVIT® Jet is a microgranule formulation of sulphur. When mixing THIOVIT® Jet and other microgranule (MG), water dispersible granule (WG) or wettable powder (WP) formulations, ensure they are added and mixed well prior to adding emulsifiable concentrate (EC) or suspension concentrate (SC) products. Wettable powder (WP) formulations should be pre-mixed separately and then added to the spray tank.

WITHHOLDING PERIOD

All crops.....1 day

Leaf plucking in vineyards

Sheep MUST NOT be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/05/2025

STORAGE
Store in a dry place. See Safety Data Sheet for additional storage information. Stores containing quantities of 10,000 kg or more require signage and secondary containment.

SAFETY DATA SHEET
If additional hazard information is required, refer to the Safety Data Sheet. For a copy, visit our website at www.syngenta.co.nz

DISCLAIMER
This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland 1052. Phone 0800 333 336.

Batch No	
Date of Manufacture	

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/05/2025