WARNING

Keep out of reach of children

Organic JMS Stylet-Oil®

For use on Apples, Avocados, Citrus, Kiwifruit, Pears, Stonefruit to control a range of insect pests and on Grapes to control Powdery Mildew, Botrytis, Mealy bug and Mites

READ THE LABEL CAREFULLY BEFORE USE

Contains: 971 ml/litre mineral oil in the form of an emulsifiable concentrate. (Minimum unsulfonated residue 99.1%)

AGRECOVERY FOUNDING BRAND

JMS Flower Farms, Inc.

CERTIFIED INPUT FOR ORGANICS bio gro NZ CERT 4429

NET CONTENTS:
20 LITRES

Registered to:
UPL New Zealand Limited
P O Box 51584, Pakuranga 2140
Auckland, New Zealand
Telephone: 0800 100 325
www.upl-ltd.com/nz

Registered pursuant to the ACVM Act 1997, No P007067
HSNO Approval Code HSR000710

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND 31/08/2022

HSNO Class: 6.1E, 6.3B, 9.1B
ENVIRONMENTAL WARNING: TOXIC TO TERRESTRIAL INVERTEBRATES
HEALTH WARNING: This material may be harmful if swallowed. May cause skin irritation.

HANDLING PRECAUTIONS: When mixing or applying work up wind and avoid breathing vapour.
Avoid contact with skin and eyes. Keep out of the reach of children.
Wear impervious clothing including gloves and goggles or face shield when mixing or applying.
Do not eat, smoke or drink while using. Wash face and hands after application.
Wash clothing before re-use. Store in original container, tightly closed away from heat.
PRODUCT DISPOSAL: Ensure unwanted product is used by another operator strictly in accordance with label directions.
CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Submit clean empty container for recycling through Agrecovery. If this is not possible, burn if permitted and circumstances, especially wind direction allow, otherwise bury in landfill
STORAGE: Store in an original container tightly closed and in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. Storage might be in accordance with NZS8409 Management of Agrichemicals. See the HAZNOTE for further information. When stored appropriately this product should show no significant degradation for 2 years from date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

FIRST AID
Flush eyes with plenty or water for at least 15 minutes. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. Call a doctor if irritation persists. Do not give liquids. Call a doctor immediately. If inhaled, remove from exposure and call a doctor. If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

PRODUCT INFORMATION

Organic JMS Stylet-Oil is a food grade mineral oil. It is a colourless, high purity, medium viscosity oil which falls into the narrow to intermediate range of oil specifications. It contains an emulsification system to facilitate mixing with water.
Organic JMS Stylet-Oil is a protectant and curative contact product. It is not systemic so requires thorough spray coverage. The mode of action on fungi and insects is through contact with the pest, interfering with respiration and for fungi, by modifying plant surfaces making them unsuitable for the establishment of a disease pathogen.

NOTES:
1. Organic JMS Stylet-Oil may reduce the visible bloom on grapes, particularly red varieties when used close to harvest, however there is no evidence of any reduction in vintning qualities.
   DO NOT USE ON TABLE GRAPES.
2. Use of Organic JMS Stylet-Oil may delay maturity by up to one week.

DIRECTIONS FOR USE
The dilution rates expressed are for high volume spraying to the point of run-off. For low volume application, follow the Du-Wett per hectare recommendations where given.
Mixing: Half fill the spray tank with water. Start agitation and add the required quantity of **Organic JMS Stylet-Oil**. Maintain agitation while filling the tank and until application is complete. If other chemicals are being added to the tank (note compatibility below), add them after **Organic JMS Stylet-Oil** has been thoroughly mixed. No additional adjuvants need to be added at high water volumes. The addition of **Oil-Mate®** adjuvant to the **Organic JMS Stylet-Oil** PRIOR to the addition of the oil to the spray tank will assist coverage of pests at lower water volumes.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
31/08/2022

RECOMMENDATIONS:

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS
To control a range of insect pests on Apples, Avocados, Citrus, Kiwifruit, Pears, Stonefruit and to control Powdery Mildew, Botrytis, Mealy bug and Mites on Grapes.

To control a range of insect pests on Apples, Avocados, Citrus, Kiwifruit, Pears, Stonefruit and to control Powdery Mildew, Botrytis, Mealy bug and Mites on Grapes.

Crop	Pest/Disease	Rate	Remarks
Grapes	*Early Season* Powdery mildew & Erinose Mite	2 Litres/ha. Apply dilute in up to 400 litres/ha or concentrate with Du-Wett in 100 – 200 litres/ha	Early Season: Apply from first leaves present to pre – flowering. To prevent powdery mildew, apply on a 10 – 14 day schedule as required to maintain adequate control. Apply dilute in up to 400 litres of water per hectare and in 100 – 200 (max) litres of water per hectare with Du-Wett Superspreader to enhance coverage.
	Powdery mildew & Erinose Mite	2 Litres/ 100 litres	Late Season: Flowering to harvest. To prevent powdery mildew apply on a 7 – 10 day schedule as required to maintain adequate control of powdery mildew. For eradication of powdery mildew infection, apply as a high volume spray to run-off. Repeat on a 7 – 10 day schedule as required to maintain adequate control of powdery mildew. If concentrate spraying, use a minimum of 5 litres per hectare.
	Botrytis		Where disease conditions are high, or during periods of high risk (wet, humid weather) use a specific botrytis fungicide. Under low botrytis conditions through the season use Organic JMS Stylet-Oil at 2 litres to 100 litres of water. Do not mix with captan, chlorothalonil, or Shirlan® (fluazinam).
	Mealy Bug & Mites		Pests present at the time of treatment with **Organic JMS Stylet-Oil** will be affected, particularly when they are at the juvenile stage of development. Specific mealy bug chemicals may be added to **Organic JMS Stylet-Oil** when mealy bug is a problem. The addition of OilMate II will help with the control of Mealy bug hiding in bark and cavities. Add OilMate directly to the Oil before mixing in the spray tank.
Apples, Pears and Stonefruit	San Jose Scale, European Red Mites and Aphids	*Green tip to tight cluster*: 1.5-2 litres/100 litres of water *Summer* Application: 1 litre /100 litres of water	To control high pest populations, apply with oil compatible insecticides. Do not apply to russet sensitive varieties later than early green tip. Do not apply to crops under stress from drought or frost. Monitor the advancement of green tip, especially on early varieties. Complete coverage is essential.
Avocados	Armoured Scale, Greenhouse Thrips, and Six-Spotted Mite.	1 litre/100 litres of water.	Make two applications from late January to February, one month apart. If pest pressure is high, apply three sprays from January to April. May be applied pre-flowering: one or two applications in September to October. Do not apply to plants under stress.
Citrus	Scale insects, Aphids, Citrus Red Mite and Thrips	1 litre/100 litres of water. Apply during spring and summer	Do not apply in temperatures over 32°C or to trees in hot dry weather when trees are suffering moisture stress. May be mixed with oil compatible insecticides to control high pest populations and mealy bug. Complete coverage is essential.
Kiwifruit	Greedy, Latania	1 litre/100 litres of water.	HAYWARD – Use 1 litre per 100 litres of water (1%).

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
31/08/2022

	and Oleander scales.		DO NOT apply from 14 days after fruit set to 35 days after fruit set. The application of oil sprays after March increases the risk of fruit drop, and care is essential. Apply when conditions favour rapid drying on the plant. ZESPRI™ GOLD. Use 1 litre per 100 litres of water (1%). Apply only up to 21 days after fruit set. **Notes**: ***Any applications of spraying oils should be in response to a demonstrated need and be in accordance with the Crop Protection Programmes for export kiwifruit.*** Use 1000 to 1500 litres of water per hectare. The use of concentrate spraying is not advised. Avoid spray overlap. Apply when conditions favour rapid drying on the plant. **NOTE**: DO NOT MIX with Success®, Entrust®, or foliar feeds on Zespri Gold.

Use Precautions:
DO NOT spray early morning, late evening, or during the hours of darkness. These conditions may give rise to phytotoxic damage to the fruit.
DO NOT spray under slow drying conditions, i.e. high humidity, cool temperatures.
DO NOT spray plants under stress from wet or dry conditions, or disease.
DO NOT apply Serenade Max® sprays within 5 days either side of **Organic JMS Stylet-Oil.**
DO NOT use additional spray adjuvants **Organic JMS Stylet-Oil** unless advised to do so.
DO NOT spray when freezing conditions are anticipated within 48 hrs and when temperatures are above 32°C.

COMPATIBILITY:
Organic JMS Stylet-Oil is compatible with a wide range of commonly used insecticides and fungicides such as Applaud®, Manzate®, Rovral®, Lorsban®, and Tokuthion®. For other mixtures it is advisable to test compatibility on a few vines before treating the vineyard.
The following chemicals are ***not compatible*** with **Organic JMS Stylet-Oil** and should not be sprayed within 14 days either side of **Organic JMS Stylet-Oil:** Carbaryl chlorothalonil, dicofol, fenarimol, Nufilm®-17, euparen multi, tolyflunaid, Filmstar, soaps/detergents, captan, organic tin compounds, dichlofluanid and propargite.
Do not apply sulphur sprays within 10 days either side of **Organic JMS Stylet-Oil.**

Exclusion of Liability: Please Read
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

® Registered Trademark of JMS Flower Farms. Inc., USA.
®[1]Registered Trademark of UPL New Zealand Limited

Tokuthion®, Euparin® Multi are registered trademarks of Bayer AG, Germany
Shirlan® is a registered trademark of Zeneca Ltd., UK
Applaud® is a registered trademark of Nihon Nohyaku Company Limited, Japan
Manzate® is a registered trademark of Griffin LLC, USA
Rovral® is a registered trademark of FMC Corporation
Lorsban® is a trademark of Dow Agroscience, USA
Nufilm®-17 is a registered trademark of Miller Chemical and Fertilizer Corp, USA
Oil-Mate ® is a registered trademark of Elliott Chemicals Limited, NZ

See www.foodsafety.govt.nz.for registration conditions.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
31/08/2022

DOM:

Batch No.

Barcode:
942102242097 2

UN3082, ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, LIQUID N.O.S. (MINERAL OIL)
PACKING GROUP III
HAZCHEM 2X
CONTAIN SPILLAGE

24 HOUR
EMERGENCY RESPONSE
0800CHEM CALL
0800 243 622

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
31/08/2022