Prodigy™

INSECTICIDE

ACTIVE INGREDIENT: Contains 240g/L methoxyfenozide in the form of a suspension concentrate.

GROUP	18	INSECTICIDE

A target specific insecticide for the control of codling moth on pipfruit and leafrollers on avocados, blueberries, grapes, kiwifruit, stone fruit and pipfruit.

READ THIS LABEL COMPLETELY BEFORE USE

NET CONTENTS: 1 L, 5 L and 10 L

Corteva Agriscience New Zealand Limited

Private Bag 2017

New Plymouth 4342

www.corteva.co.nz

Telephone: 0800 803 939

AGRECOVERY
www.agrecovery.co.nz

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

30/03/2022

Hazardous to the aquatic environment (chronic) Category 2
Hazardous to terrestrial invertebrates

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
30/03/2022

QUALIFICATION REQUIREMENTS

Persons mixing, loading, applying, or otherwise handling Prodigy™ must meet the qualification requirements set out in the EPA Hazardous Substances (Hazardous Property Controls) Notice 2017.

RECORD KEEPING

Records of use, as described in the EPA Hazardous Substances (Hazardous Property Control) Notice 2017 and the New Zealand Standard for the Management of Agrichemicals (NZS8409), must be kept if more than 3 L of Prodigy™ is being used in a place where members of the public may be present, or where the substance may enter air or water and leave the place.

ENVIRONMENTAL HAZARDS

Very toxic to aquatic life with long lasting effects. Avoid release to the environment. DO NOT apply directly into or onto water.

Hazardous to terrestrial invertebrates. Certain insects may be killed from contact with this product.

Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

HANDLING PRECAUTIONS

EQUIPMENT

Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).

Flush equipment thoroughly with clean water after use according to Cleaning Spray Equipment instructions below. Dispose of cleaning water according to the Disposal instructions below.

STORAGE

Store in original container tightly closed and in a locked, dry, cool, well-ventilated area, away from foodstuffs. Storage must be generally in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).

Stores containing more than 100 L of Prodigy™ (or substances of the same hazard class) are subject to signage and those containing more than 1,000 L require bunding. See the Health and Safety at Work (Hazardous Substances) Regulations and the New Zealand Standard for the Management of Agrichemicals (NZS8409) for further information.

PERSONAL PROTECTION

When opening the container, preparing the product for use, and applying the product wear eye protection, cotton overalls buttoned to the neck and wrist and chemical resistant gloves. Take off

contaminated clothing and wash it before re-use.DO NOT eat, drink or smoke while using. Wash hands and face thoroughly after handling.

SPILLAGE

DO NOT touch or walk through spilled material. Wear the protective equipment recommended above. Stop leak when safe to do so. Dam area and prevent entry into waterways and drains. **Small spills/leaks:** Contain and absorb small spills with a proprietary absorbent suitable for chemical spills or inert material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. **Large spills/leaks:** Dam the area off and report them to Corteva Agriscience Emergency Services at 0800 844 455.

DISPOSAL

Triple rinse empty container and add rinsate to spray tank. Triple rinsed containers containing the Agrecovery logo on the label, are free of all residues (liquid and dry) and have an intact legible label should be taken to an Agrecovery collection site (visit www.agrecovery.co.nz or call 0800 AGRECOVERY (247 326) for information about the type and quantity of chemical acceptable for disposal). Otherwise crush and bury in a suitably approved landfill. DO NOT burn empty containers.

Dispose of unused spray mix and equipment cleaning water onto non-grazed and non-cropped ground away from edible and flowering plants. DO NOT dispose of this product; spray mix or equipment cleaning water down drains or sewers.

FIRST AID

For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

SAFETY DATA SHEET

A Safety Data Sheet for Prodigy™ is available from Corteva Agriscience on request. Call Customer Service Toll Free on 0800 803 939 or visit www.corteva.co.nz.

TRANSPORT

No restrictions on carriage of this product on a passenger service vehicle.

NOTICE

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

/03/2022

Important: You **must** read the warning and conditions given on this label. If they are not acceptable, or if you are in doubt as to the proper handling of this product (including its transport, storage, use, disposal, and requirements for approved handlers [if applicable]) then return the unopened product to the supplier at once.

Warning: This product is subject to controls and could be hazardous if handled incorrectly. The **user is responsible** for ensuring that the handling of this product is in accordance with the law (including the *Hazardous Substances and New Organisms Act 1996* and the *Agricultural Compounds and Veterinary Medicines Act 1997*) and with the directions provided. The Registrant declines all liability (whether in contract, tort (including but not limited to negligence), or otherwise) to the maximum extent permitted by law in relation to the handling of this product in a manner which is contrary to the directions given or to any applicable Acts, Regulations or Guidelines.

Further information regarding the applicable Acts, Regulations and Guidelines for the proper handling of this product may be obtained by contacting EPA New Zealand (www.epa.govt.nz) or the Ministry for Primary Industries (MPI) (www.foodsafety.govt.nz).

This product is warranted to contain the specified active ingredients within accepted analytical tolerance when packed.

Registered pursuant to the ACVM Act 1997 No. P7154
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to HSNO Act 1996, EPA Approval Code HSR000130
See www.epa.govt.nz for approval controls.

Registered to: **Corteva Agriscience New Zealand Limited** Private Bag 2017 New Plymouth 4342 Telephone 0800 803 939 www.corteva.co.nz	**EMERGENCY RESPONSE** (All hours) Ring from anywhere in New Zealand **0800 844 455** **Or Globally** **+64-6 751 2407**	**IN A TRANSPORT EMERGENCY** **DIAL 111** **FOR POLICE OR FIRE**

Shipping Name:	ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS METHOXYFENOZ~~C~~IDE) Marine Pollutant
UN Number:	3082

Date of Manufacture: Batch No. Barcode

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
30/03/2022

DIRECTIONS FOR USE

MIXING INSTRUCTIONS

Prior to pouring, shake container vigorously, then add the required quantity of Prodigy® to water in the spray vat while stirring or with agitators in motion.

APPLICATION

Use either:

- a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed; or
- a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.

COMPATIBILITY

Prodigy™ is incompatible with mineral spray oils. DO NOT mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Corteva Agriscience, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Corteva Agriscience cannot be responsible for the behaviour of such mixtures.

CLEANING THE SPRAY EQUIPMENT

After using Prodigy™ empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Discard rinse water on land already sprayed or on waste land away from desirable plants and water sources.

RAINFASTNESS

DO NOT apply if rain or overhead irrigation is expected within 6 hours.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Limits for Agricultural Compounds.

Avocados - 14 days (Consult your exporter for WHPs for export markets).
Blueberries - 7 days (Consult your exporter for WHPs for export markets).
Grapes - DO NOT apply after 80% capfall
Kiwifruit - 120 days
Pipfruit - 14 days (Consult your exporter for WHPs for export markets).
Stone fruit - 14 days (Consult your exporter for WHPs for export markets).

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
30/03/2022

Grazing restriction

The following restrictions on the use of this product **must** be observed. DO NOT feed treated crops or pasture to stock other than sheep used for leaf plucking in vineyards. Take care not to allow this product to drift onto pasture or crops that may be grazed. DO NOT graze stock in orchards which have been treated with Prodigy™.

Vineyards: DO NOT graze stock other than sheep in vineyards which have been treated with Prodigy™. Sheep may be grazed in vineyards for the purpose of leaf plucking providing they are not slaughtered for 2 months after leaving the vineyard.

RESISTANCE MANAGEMENT

GROUP	18	INSECTICIDE

Prodigy™ is a diacylhydrazine insecticide (Group 18). Some naturally occurring individual insects may become resistant when excessive use of the same insecticide occurs. To maximize the effectiveness of insecticides, products containing alternate modes of action should be used in the spray program. Consult http://resistance.nzpps.org/ for alternate modes of action and the details of resistance management strategies (if any) for the crops listed on the label.

It is an offence to use this product on animals.

Monitor treated areas closely to check for other pests. Consult field advisors to identify alternative or additive treatments when non-target pests are expected. Prodigy™ is a new generation Moult Accelerating Compound (MAC) insecticide which induces premature moulting in caterpillars. Feeding damage stops within 24 hours but insect death may take several days.

Prodigy™ can be used in Integrated Pest Management (IPM) programmes and can be applied over flowering where appropriate under the directions for use.

The rates suggested are for high volume spraying to run-off. For concentrate spraying adjust dilution rate accordingly. Thorough coverage of plants is essential for control of pests.

CROP	PEST	APPLICATION RATE	APPLICATION TIMING
Pipfruit (apples and pears)	Codling Moth, Leafrollers	For high volume spraying apply 37.5 mL per 100 litres of water and apply at 2000 litres water per hectare. For concentrate spraying adjust accordingly. DO NOT apply Prodigy™ at less than 750 mL/ha.	Apply every 14 days as part of an IPM program commencing at petal fall and re-apply when thresholds are exceeded. DO NOT apply more than four applications in any one season.
Avocado	Leafrollers (including Light Brown Apple Moth)	Apply 25 mL per 100 litres to full coverage (point of runoff) with a minimum dose of 500 mL/ha. For mature trees and water rates greater than 2000 L/ha, maintain the rates of 25 mL per 100 litres.	Apply at 3 to 4 week intervals from flowering as part of a season long leafroller spray programme. DO NOT apply more than four applications in any one season.
Blueberries		For high volume spraying apply 25 mL per 100 litres of water. The product should be applied thoroughly to ensure complete coverage using dilute spraying equipment at 1000 litres water per hectare.	Apply at not less than 7 days intervals. DO NOT apply more than three applications per season.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

30/03/2022

Grapes		Apply 15-25 mL/100 L to full coverage. DO NOT apply Prodigy™ at less than 150 mL/ha under moderate pest pressure and 250 mL/ha under severe pest pressure.	Apply a single application at 80% capfall. If pest pressure remains high at pre-bunch closure, apply an alternative product such as Success™ Naturalyte™ Insect Control.
Kiwifruit		Apply 15-25 mL/100 L to full coverage. DO NOT apply Prodigy™ at less than 300 mL/ha under moderate pest pressure and 500 mL/ha under severe pest pressure.	Apply at pre-bloom or at late flowering followed by 2 further applications at 14-day intervals.
Stone fruit		For high volume spraying apply 25 mL per 100 litres of water and apply at 2000 litres water per hectare. For concentrate spraying adjust accordingly. DO NOT apply Prodigy™ at less than 500 mL/ha.	Apply at not less than 14 days intervals commencing at petal fall. Re-apply when thresholds are exceeded. DO NOT apply more than four applications in any one season.

™ ® Trademarks of Corteva Agriscience and its affiliated companies. © 2022 Corteva.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
30/03/2022