ECOTOXIC

Keep out of reach of children

Rainbow&Brown

Granny

HERBICIDE

A non-selective herbicide for the control of many annual and perennial weeds and grasses in a wide range of situations.

ACTIVE INGREDIENT:
Contains 800g/kg glyphosate as the monoammonium salt in the form of a water soluble granule.

Contains
2.25
kg

DATE OF MANUFACTURE:

BATCH No:

GRANNY Herbicide

PLEASE READ THIS LABEL COMPLETELY BEFORE USE

BEWARE: APPLY THIS PRODUCT CAREFULLY AS SPRAYDRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS.

HAZARD CLASSIFICATIONS: Hazardous to the aquatic environment (chronic) Category 2.
HAZARDS: Toxic to aquatic life with long-lasting effects.

PRECAUTIONS:
Keep out of reach of children. Store only in the original container , tightly closed, and keep away from foodstuffs. Do not eat, drink or smoke while using. Wear waterproof gloves when handling this product. Avoid contact with skin and eyes, and inhalation of spray mist. Wash hands and exposed skin when fi nished and before meals.

Do not mix or store this product in galvanised or unlined steel (except stainless steel) containers or spray tanks. The product can react with such containers to produce potentially explosive hydrogen gas.
Container Disposal: Triple rinse empty containers into spray tank. Burn container if circumstances, especially wind direction permit. Otherwise bury in landfi ll. Avoid contamination of any water supply with chemical or empty container
Product Disposal: Use according to label or bury in landfill

FIRST AID:
If swallowed DO NOT induce vomiting. Rinse mouth. For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and fl ush skin and hair with running water. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. Immediately contact a POISON CENTRE or doctor.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/04/2024

Safety Data Sheet: Additional information is provided in the safety data sheet available from www.rainbowbrown.co.nz

STORAGE: Store at temperatures above 5°C but below 35°C.
Shelf Life: When stored appropriately, this product should show no significant degradation for up to two years from the date of manufacture. Contact Rainbow & Brown for further information about the use of any product that is older than this.

GENERAL INFORMATION: This product is a non-selective herbicide for use in agriculture and horticulture. It is water-soluble and is absorbed by green plant tissue and then translocated throughout the plant including the root system, to kill the entire plant. Target plants should be actively growing and not drought stressed for best results. Allow about 14 days for visible effects to appear. This product degrades in soil and has no residual herbicidal activity.

MIXING: Look up your target species in the Application Directions leaflet supplied with this product. Ensure all equipment is cleaned and free of residues from previous chemicals. Use only clean water. Add half the water to the tank first, then add the required amount of GRANNY Herbicide. Add the remaining water to mix thoroughly, and finally add the wetter if required. Agitate well before spraying.

EQUIPMENT
Boom: Spray up to 200L/ha using fan nozzles at pressures from 200-280kPa. Set boom height so that overlap of nozzle pattern is achieved at the top of target plants.
Aerial: Use in forest and non-crop areas only. Apply up to 100L/ha, or more in hilly terrain to ensure thorough target area contact. Do not use aerial application in intensive horticultural or cropping areas.
Handgun & Knapsack: Use a fine spray pattern to ensure thorough coverage. For handguns, use pressures of 1000-2000kPa with No 4 or No 6 spray tips.
Wiper: Premix 1 part of this product with 2 parts water and fill wiper applicator. Ensure weeds are dry and dust-free, and that wiper surface remains moistened. Make two passes in opposite directions to ensure good coverage.

COMPATIBILITY: This product is compatible with most commonly used herbicides.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

It is an offence to use this product on animals.

WITHHOLDING PERIOD
In line with good agricultural practice, a withholding period of 1 day is required.

Registered pursuant to the ACVM Act 1997, No: P7499
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, EPA Approval Code HSR007661
See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/04/2024

Conditions of Sale: This product is warranted by Rainbow & Brown Limited to conform to the chemical description and concentration specified on the label, and to be reasonably fit for the purposes stated in the directions if used in accordance with the directions. The user bears all risks of damage or failure resulting from factors beyond the control of Rainbow & Brown. Rainbow & Brown declines all liability for any damages resulting from the improper use of this product.

Registrant:
Rainbow & Brown Ltd
68a Tallyho Street, Rotorua 3015.
P O Box 100 49, Mail Centre, Rotorua 3046.

Freecall 0508 299 299.
www.RainbowBrown.co.nz

AGRECOVERY

0800 CHEMCALL 24 x 7 Chemical EMERGENCY

Rainbow&Brown

ECOTOXIC

Keep out of reach of children

Rainbow&Brown

Granny

HERBICIDE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/04/2024

A non-selective herbicide for the control of many annual and perennial weeds and grasses in a wide range of situations.

ACTIVE INGREDIENT:

Contains 800g/kg glyphosate as the monoammonium salt in the form of a water soluble granule.

Contains
4.5
kg

DATE OF MANUFACTURE:

BATCH No:

GRANNY Herbicide

PLEASE READ THIS LABEL COMPLETELY BEFORE USE

BEWARE: APPLY THIS PRODUCT CAREFULLY AS SPRAYDRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS.

HAZARD CLASSIFICATIONS: Hazardous to the aquatic environment (chronic) Category 2.

HAZARDS: Toxic to aquatic life with long-lasting effects.

PRECAUTIONS: Keep out of reach of children. Store only in the original container, tightly closed, and keep away from foodstuffs. Do not eat, drink or smoke while using. Wear waterproof gloves when handling this product. Avoid contact with skin and eyes, and inhalation of spray mist. Wash hands and exposed skin when finished and before meals.
Do not mix or store this product in galvanised or unlined steel (except stainless steel) containers or spray tanks. The product can react with such containers to produce potentially explosive hydrogen gas.

Container Disposal: Triple rinse empty containers into spray tank. Burn container if circumstances, especially wind direction permit. Otherwise bury in landfill. Avoid contamination of any water supply with chemical or empty container. **Product Disposal:** Use according to label or bury in landfill.

FIRST AID:
If swallowed DO NOT induce vomiting. Rinse mouth. or advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately . If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. Immediately contact a POISON CENTRE or doctor.

Safety Data Sheet: Additional information is provided in the safety data sheet available from www.rainbowbrown.co.nz

STORAGE: Store at temperatures above 5°C but below 35°C. Shelf Life: When stored appropriately, this product should show no significant degradation for up to two years from the date of manufacture. Contact Rainbow & Brown for further information about the use of any product that is older than this.

GENERAL INFORMATION: This product is a non-selective herbicide for use in agriculture and horticulture. It is water-soluble and is absorbed by green plant tissue and then translocated throughout the plant including the root system, to kill the entire plant. Target plants should be actively growing and not drought stressed for best results. Allow about 14 days for visible effects to appear. This product degrades in soil and has no residual herbicidal activity.

MIXING: Look up your target species in the Application Directions leaflet supplied with this product. Ensure all equipment is cleaned and free of residues from previous chemicals. Use only clean water. Add half the water to the tank first, then add the required amount of GRANNY Herbicide. Add the remaining water to mix thoroughly, and finally add the wetter if required. Agitate well before spraying.

EQUIPMENT

Boom: Spray up to 200L/ha using fan nozzles at pressures from 200-280kPa. Set boom height so that overlap of nozzle pattern is achieved at the top of target plants.

Aerial: Use in forest and non-crop areas only. Apply up to 100L/ha, or more in hilly terrain to ensure thorough target area contact. Do not use aerial application in intensive horticultural or cropping areas.

Handgun & Knapsack: Use a fine spray pattern to ensure thorough coverage. For handguns, use pressures of 1000-2000kPa with No 4 or No 6 spray tips.

Wiper: Premix 1 part of this product with 2 parts water and fi ll wiper applicator. Ensure weeds are dry and dust-free, and that wiper surface remains moistened. Make two passes in opposite directions to ensure good coverage.

COMPATIBILITY: This product is compatible with most commonly used herbicides.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

It is an offence to use this product on animals.

WITHHOLDING PERIOD:
In line with good agricultural practice, a withholding period of 1 day is required

Registered pursuant to the ACVM Act 1997, No: P7499
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, EPA Approval Code HSR007661
See www.epa.govt.nz for approval controls.

Conditions of Sale:
This product is warranted by Rainbow & Brown Limited to conform to the chemical description and concentration specified on the label, and to be reasonably fit for the purposes stated in the directions if used in accordance with the directions. The user bears all risks of damage or failure resulting from factors beyond the control of Rainbow & Brown. Rainbow & Brown declines all liability for any damages resulting from the improper use of this product.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/04/2024

Registrant:
Rainbow & Brown Ltd
68a Tallyho Street, Rotorua 3015.
P O Box 100 49, Mail Centre, Rotorua 3046.
Freecall 0508 299 299.
www.RainbowBrown.co.nz

AGRECOVERY

0800 CHEMCALL 243 622 Chemical EMERGENCY

Rainbow&Brown

Weed	Boom/ Aerial per Ha	Handgun g/100L	Knapsack g/15L	Mistblower g/10L	Critical Comments
Willow	3.8kg	425	-	-	Spray before leaf fall begins (Jan-Apr) Total spray coverage is critical for best results.
Yarrow	1.3-2.6kg	-	-	-	Apply to plant with 5-10cm leaf. For best results cultivate 10 days after treatment. Use the low rate when a crop to be drilled is competitive to yarrow and higher rate when crop is a poor competitor.

* When spraying species marked * an organosilicone penetrant must be used at 0.1-0.25% (100-200ml per 100 litres of spray solution)

Directions for Use

Agricultural Areas: May be used for the control of annual and perennial weeds as indicated in the Weeds Controlled table, prior to sowing any crop.

Minimum Tillage: Use for total weed control in a reduced tillage system. Weeds should be actively growing and not under stress from frost, drought or waterlogging. In pastures where only annual weeds are present, use 350g/ha to control seedling annual weeds and 1kg/ha to control mature annual weeds. For perennial weeds use 1kg/ha plus 50ml/100L SprayWetter (a non-ionic surfactant). Cultivation may start from 7 days after spraying perennial and 3 days after spraying annual weeds. Cultivation should occur within 21 days after spraying.

Pasture: Use for spot application to a wide range of specific weeds. This product is non-selective, and may kill any plant directly sprayed.

Orchards/vines: Citrus, apple, pear, stonefruit, kiwifruit, grapes. Apply as directed with shielded spray. For long term control of annual weeds, this product may be mixed with an approved residual herbicide. Do not apply to unprotected trees under 3 years old. Do not spray or allow spray drift onto any part of stonefruit trees, or onto green bark, suckers, buds, foliage or fruit of other trees. Spray crops before bud, sucker or bloom.

Industrial & non-agricultural areas: Treat specific weeds as directed in the Weeds Controlled Table. Do not allow spray drift from target site.

Recreational areas: Lawns, turf, parks, reserves & ornamentals. Apply before sowing/planting of these areas. In mown areas allow sufficient regrowth before spraying. May be sprayed around base of established trees and shrubs as directed, or with shielded spray or wiper. Avoid spray drift or wiper contact onto any part of the tree or shrub.

Drains & waterways: Do not spray this product directly onto water surface, nor allow runoff into the water.

Forestry: Use for control of grasses and weeds listed in the Weeds Controlled Table. Can be applied prior to burning or planting of forest or nursery, or around established trees. Do not allow spray or spray drift to contact foliage or green bark of desirable trees.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/04/2024

Rainbow & Brown

Rainbow & Brown Ltd
68A Tallyho St, Rotorua. Phone (0508) 299 299

Rainbow & Brown

GRANNY

APPLICATION DIRECTIONS

Weeds Controlled Table

Weed	Boom/ Aerial per Ha	Handgun g/100L	Knapsack g/15L	Mistblower g/10L	Critical Comments
Annual weeds	425-850g	85	15	25	Use higher rate for larger weeds.
Australian Sedge	2.5-3.8kg	425	65	125	Apply Nov-Mar. Aerial: use higher rate. Where rank growth is present, the area should be burnt 12-18 months prior to application to ensure adequate spray coverage.
Barberry	-	425	65	125	Apply from flowering to late fruit stage (Jan-Apr). Regrowth should be 1.5m tall before spraying.
Barley Grass	425g	85	15	25	Apply at early seedhead stage (Sep-Dec)
Blackberry	3.8-8.5kg	425-640	65	125	Apply from flowering-fruiting (Dec-Apr) before frosts occur. For hard-to-kill blackberry, or for aerial application or if no follow-up treatment Is planned, use the higher rate. Allow 6-10 weeks before slashing or burning.
Boxthorn	-	600	-	-	Apply when actively growing (Nov-Apr). Ensure complete coverage of dense thickets. Do not add extra surfactant or treat salt damaged bushes.
Bracken*	3.8kg	425	65	-	Apply after fronds have unfurled (Jan-Jun) but before frosts occur. Dense virgin bracken with high trash content and low leaf to root ratio should be burnt in spring and the unfurled frond regrowth sprayed the following Mar-Jun.
Broom*	3.8kg	425	65	125	Aerial – apply when plants are in full leaf. Handgun – apply all year round.
Browntop	1.3kg	210	35	65	Apply when 5-10cm high. For taller plants, graze or burn & allow regrowth to 5-10cm before spraying.

Weed	Boom/ Aerial per Ha	Handgun g/100L	Knapsack g/15L	Mistblower g/10L	Critical Comments
Buddleia	3.8kg	425	65	-	Aerial: apply to plants up to 2m high in Feb-Apr using 200-300L spray/ha. Ground: spray anytime when actively growing.
Californian thistle	1.7kg	425	65	125	Apply after flower bud development but before autumn browning. To ensure maximum shoot emergence, do not cultivate for 6 months prior to treatment. Wiper– a double pass at low speed is recommended in heavy infestations. Mowing thistles 10 days after application may assist control. Retreatment may be necessary in the next season.
Cleavers	640g	210	35	65	Apply to actively growing plants after most reach early seedhead stage
Couch	1.3-2.5kg	425	65	125	For suppression apply 900g/ha in spring and repeat in autumn. For best results, apply 1.2-1.5kg/ha in autumn to active growth of at least 10cm where no previous cultivation has occurred for at least 6 months.
Docks	1.7-3.8kg	425	65	-	Spray in spring & summer when actively growing and before seedheads have formed. Use lower rate for seedlings & high rate for mature plants. In pasture, boom spray prior to re-seeding or establishing crops to restrict seedling re-establishment.
Floating sweet grass	2.5kg	425	65	125	Spray actively growing plants after most reach early seedhead. Spraying plants under drought stress will reduce results
Indian doab	3.8kg	425	65	-	Apply at early seedhead, before frosts.
Gorse*	-	425	65	-	Apply anytime to plants not stressed by drought, grazing or previous herbicide treatment. Ensure complete coverage.
Kikuyu	2.5kg	425	65	125	Apply to actively growing plants. Under dense growth conditions, silorate and allow regrowth before treatment.
Mercer Grass	2.5-3.8kg	425-640	65-125	125-190	Use the higher rate for aquatic Mercer grass.
Nodding thistle	-	425	65	-	Apply to actively growing plants (Oct-Apr). Wiper: ensure at least 10cm of plant can be [illegible] double pass and follow-up with spot-spraying.

Weed	Boom/ Aerial per Ha	Handgun g/100L	Knapsack g/15L	Mistblower g/10L	Critical Comments
Old Man's Beard	-	425-850	125	-	Apply Nov-Mar prior to first leaf browning. Use higher rates on stems over 1cm dia. If complete coverage is not possible, cut plants at ground level in winter and spray the regrowth at the higher rate when plants are at least 50cm long (March).
Pampas grass / toetoe	-	425	65	-	Spray spring-autumn. Reduced control may result if plants are in flower at the time of spraying. Ensure complete coverage.
Paspalum	1.7-2.6kg	425	65	125	Use low rate if plants are at early seed head stage. Use 1.8kg/Ha on mature plants, before onset of frosts.
Potato (self set)	1.3kg	210	35	65	Spray actively growing plants after most reach early seedhead.
Prairie grass	640g	-	-	-	Apply at early seedhead stage.
Ragwort	2.6kg	425	65	-	Spray when actively growing, ensuring there is sufficient leaf area for uptake of herbicide. Wiper: apply to plants 10-15cm above pasture. Caution: Ragwort becomes more palatable after treatment. Keep stock away until the plants brown off.
Ratstail	2.1kg	-	-	-	Apply in autumn using 200l spray/ha. In dry conditions, delay spraying until sufficient green foliage appears.
Rautahi (Cutty grass)	2.6kg	425	65	125	Spray actively growing plants Jan-Mar. Burning or slashing may begin 30 days after treatment following brownoff.
Rushes	2.6kg	425	65	125	Burning or slashing may begin 30 days after treatment following brownoff. Wiper: apply Nov-May.
Ring fern	425g-1.3kg	125	20	-	Use 900g/ha for effective control. For pasture manipulation apply 300-600g/ha from Dec-Apr. Use high rates for dense infestations. Burn 4-6 weeks after spraying.
Ryegrass*	640g-1.3kg	210	35	_	Spray actively growing plants 3-10cm high. Use lower rate on perennial from Jan-Jul when cultivation is to follow spraying, and higher rate Aug-Dec.
Sweet brier*	5kg	425	65	-	Spray from late flowering to early leaf yellowing. Ensure complete coverage.
Tall fescue	2.6kg	425	65	125	Spray active growth Jan-Mar.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/04/2024