WARNING

KEEP OUT OF REACH OF CHILDREN

MANZATE® EVOLUTION

Fungicide

For the control of certain diseases in fruit, vegetable and ornamental crops

Active ingredient
Contains 750 g/kg MANCOZEB in the form of a water dispersible granule.

GROUP	M3	FUNGICIDE

Net Contents: 10 kg

Bar code to be inserted

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

30/07/2025

Public

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.
Any person purchasing, handling or disposing of this product must be suitably qualified.

WARNING
Causes serious eye irritation.
May cause an allergic skin reaction.
May cause damage to organs through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS
PREVENTION
Wash hands and face thoroughly after handling.
Wear protective gloves, protective clothing and eye protection.
Contaminated clothing should not be allowed out of the workplace.
Do not breathe spray.
Do not apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

RESPONSE
Do not eat, drink, or smoke when using this product.
Wash hands and face thoroughly after handling.
Take off contaminated clothing and wash it before re-use.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice / attention.
IF ON SKIN: Wash with plenty of soap and water. If skin irritation persists, get medical advice / attention. DO NOT scrub the skin.
Get medical advice if you feel unwell.
Collect spillage.
Records of use must be kept.

STORAGE
Store securely in original container, tightly closed. Do NOT store with Class 1, 2, 3.2, 4 or 5 substances or feed, seeds or foodstuffs.

IMPORTANT: Never allow **MANZATE EVOLUTION FUNGICIDE** to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of **MANZATE EVOLUTION FUNGICIDE** and create vapours which may be flammable.

DISPOSAL
Dispose of contents/container in accordance with local regulations.
Container disposal: Ensure container is completely empty. Bury in an approved landfill. Do NOT reuse container.
Product disposal: Use according to this label, otherwise bury in approved landfill. Always follow local authority requirements.

FIRST AID
If medical advice is needed, have the product container or label at hand.
IF SWALLOWED: Do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if feeling unwell. **IF INHALED**: Remove victim to fresh air. Get medical attention if person feels unwell. If not breathing, give artificial respiration preferably mouth to mouth and get immediate medical assistance.

SPILLAGE
Wear appropriate protective clothing. Shovel or sweep up and dispose of waste safely in an approved landfill. Never return to the container for reuse. Do NOT flush to surface water or sewer system.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at **www.uplcorp.com/nz**.

EXCLUSION OF LIABILITY: PLEASE READ
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to:
UPL New Zealand Ltd
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
www.uplcorp.com/nz

Registered pursuant to the ACVM Act 1997, No. P007514
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval Code: HSR000808
See www.epa.govt.nz for approval controls.

© Copyright UPL New Zealand Limited 2020, all rights reserved.
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited.

Manzate is a registered trademark of UPL Ltd, India.

Batch number:	
Date of Manufacture:	
Expiry date:	

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622)

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (MANCOZEB)
MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2X



MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

Public

GENERAL INFORMATION

MANZATE EVOLUTION FUNGICIDE is applied as a spray for the control of many important plant diseases. **MANZATE EVOLUTION FUNGICIDE** has good adhesive properties on plants, high fungicidal efficiency and is easy to use. It is particularly effective against many of the fungus diseases of certain vegetable, fruit and ornamental crops. **MANZATE EVOLUTION FUNGICIDE** should be used only in accordance with recommendations on this label.

DIRECTIONS FOR USE

The rates expressed are high volume spraying to run-off. For concentrate spraying, adjust dilution rate accordingly. Apply with well maintained, accurately calibrated spray equipment.

CROP	DISEASE	RATE	COMMENTS
FRUIT			
Apples, pears	Black spot, glomerella, ripe spot, sooty blotch	125 – 150 g/100 L water	Commence at green tip and repeating at 7 to 14-day intervals throughout the season. Use the higher rate and shorter interval early in the season or under severe disease conditions. Maintain rates of 150g/100 L for control of glomerella. Note: Early season use under frosty conditions may cause russet.
Brambles, canefruit, bushfruit	Downy mildew (dryberry), leaf spot, rust	200 g/100 L water	Begin at green tip and repeat at 10 to 14-day intervals until start of flowering.
Citrus	Brown rot	210 g/100 L water	Apply in May. If infection is serious begin applications in April and continue at 4 to 6-week intervals until late winter.
	Alternaria		Apply from October to April every 14 to 28 days.
	Verrucosis (Citrus scab)		Apply at petal fall.
	Citrus rust mite		Apply at the first sign of mite activity around December to January. Reapply after 21 to 28 days if necessary.
Grapes	Anthracnose (black spot), downy mildew	200 g/100 L water	Apply pre-bloom, post bloom and repeat every 10 to 14-day intervals as required.
	Dead arm disease	210 g/100 L water	Apply 210 g per 100 litres of water every 10 to 14 days commencing at bud burst, or in accordance with local recommendations.
Nectarines, peaches	Rust	150 g/100 L water	Apply early November. Apply 2 to 3 sprays at 2 to 3-week intervals. To avoid spray deposit on fruit do not spray within 6 weeks of harvest. Note: Do NOT allow drift onto plums as foliage damage and defoliation may occur.
Strawberries	Mycosphaerella leaf spot (frog's eye)	200 g/100 L water or 2 kg/ha	Apply every 10 to 14 days beginning in mid June. During wet weather spray every 7 days.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
30/07/2025

Public

ORNAMENTALS			
Roses	Black spot, Rust	150 to 200 g/100 L water	Begin spraying at bud burst and continue at 7 to 14-day intervals throughout the growing season.
Other ornamentals	Flower blight, rust, Septoria leaf spot	200 g/100 L water	Apply on appearance of disease and as required until disease is under control.
VEGETABLES			
Beans	Anthracnose, Rust	160 to 210 g/ 100 L water	Apply at 7 to 10-day intervals following first signs of disease.
Broccoli **Brussels sprouts** **Cabbage** **Cauliflower**	Downy mildew	150 to 200 g/ 100 L water	Plant beds. Apply every 3 to 4 days in wet weather, otherwise every 7 to 10 days.
Celery	Septoria leaf spot (late blight)	(a) 150 to 200 g/ 100 L water (b) 2 to 3 kg/ha	Plant beds. Apply every 3 to 5 days in wet weather, otherwise every 7 to 10 days. Field. Apply every 7 to 10 days from first appearance of disease. Remove residues by stripping, trimming and washing.
Carrots	Cercospora leaf spot	2 to 3 kg/ha	Apply when plants are six weeks old or when disease threatens. Apply every 7 to 10 days or as required.
Cucumber **Melons Pumpkin** **Squash** **(Cucurbits)**	Anthracnose, alternaria, cercospora leaf spot, downy mildew	2 to 3 kg/ha or 150 – 200 g/ 100 L water (high volume)	Apply as soon as plants begin to run or when disease threatens. Repeat at 7 to 10-day intervals as needed. Under severe conditions, shorten intervals to 3 to 5 days.
Lettuce	Downy mildew, leaf spot (pleasora spot), ring spot	200 g/100 L water (up to 1600 L/ha) or 2 kg/ha	Apply when disease appears or threatens. Repeat at 7 to 10-day intervals as needed. Under severe conditions, shorten intervals to 3 to 5 days. Remove residues from head by stripping and trimming and from leaf lettuce by washing or other effective means.
Maize **Sweetcorn**	Northern leaf blight	2kg/ha	As an aerial treatment in sufficient water to cover. Repeat as necessary.
Onions	Botrytis, downy mildew	2 to 3 kg/ha plus spreader sticker at 25 mL/100 L spray	Begin spraying at the 2-leaf stage and repeat at 7 to 10-day intervals as necessary.
Potatoes	Early blight (target spot), late blight	1.5 kg/ha (crop emergence) to 2 kg/ha (as vines size)	In wet weather apply every 3 to 4 days otherwise every 7 to 10 days.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

Public

Tomatoes	Anthracnose, cladosporium leaf mould, early blight, late blight, septoria leaf spot	150 to 200 g/ 100 litres water (high volume) or 1 kg/ha (planting) to 3 kg/ha (as vines size)	Apply when fruit of first clusters is well-formed. Apply at 7 to 10-day intervals or more frequently in wet weather. Do NOT use on young tomato seedlings in glass houses or tunnel houses as injury may occur. When weather conditions favour disease, more frequent applications may be necessary starting when plants are set in fields. Tomato plants treated with **MANZATE EVOLUTION FUNGICIDE** tend to develop heavy foliage and water rates should be increased accordingly to assure complete coverage. In areas where Anthracnose is a serious problem on canning tomatoes, apply 3 kg per hectare.

It is an offence to use this product on animals.
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MIXING
Always shake the container before use. Pour the required amount into the partly filled spray tank with agitation running. Complete filling the tank.

RESISTANCE MANAGEMENT
MANZATE EVOLUTION FUNGICIDE is a dithio carbamate fungicide (Group M3). Some naturally occurring individual fungi may become resistant when excessive use of the same fungicide occurs. To maximize the effectiveness of fungicides, products containing alternate modes of action should be used in the spray program. Consult http://resistance.nzpps.org/ for alternate modes of action and the details of resistance management strategies (if any) for the crops listed on the label.

APPLICATION
The rates expressed are high volume spraying to run-off. For concentrate spraying, adjust dilution rate accordingly. Apply with well maintained, accurately calibrated spray equipment.

Sprays: Use **MANZATE EVOLUTION FUNGICIDE** at rates stated. Wash required amount of product through the filler screen while tank is filling with water.
For small quantities, 3- to 4 level teaspoonfuls of **MANZATE EVOLUTION FUNGICIDE** per 5 L of water is equivalent to 150 – 200 g/ per 100 L

COMPATIBILITY
MANZATE EVOLUTION FUNGICIDE is compatible with most insecticides. Do NOT use with Bordeaux mixture or lime- sulphur.

Leaf plucking in vineyards: Sheep must not be sent for slaughter or milked for 6 months after being removed from the vineyard (and placed on clean feed).

WITHHOLDING PERIODS

Apples, pears	14 days
Brambles, canefruit, bushfruit	DO NOT use after start of flowering.
Citrus, grapes, nectarines, peaches	14 days
Strawberries:	DO NOT use after start of flowering.

Public

Roses & other ornamentals	Not applicable
Beans:	14 days
Broccoli, brussels sprouts, cabbage, cauliflower:	Seedlings only
Celery, carrots	14 days
Cucumber, melons, pumpkin, squash (cucurbits):	14 days (field), 2 days (indoors)
Lettuce, maize, sweetcorn	14 days
Maize, sweetcorn, onions	14 days
Potatoes:	7 days
Tomatoes:	7 days (field), 2 days (indoors)

RESTRICTED ENTRY INTERVALS

The following restricted entry intervals (REI) have been set for this product. Anyone entering the application area during the REI must use appropriate PPE.

– Apples, pears 5 days
– Stonefruit 6 days
– Grapes 8 days
– Onions 24 hrs
– Broccoli, brussels sprouts, cabbage, cauliflower 10 days

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
30/07/2025

Public