WARNING

Keep out of reach of children

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FUNGICIDE

Pinnacle®

For the control of a wide range of diseases in grapes, field tomatoes, potatoes, and vegetable brassicas.

ACTIVE INGREDIENT: Contains 500 g/L fluazinam in the form of a suspension concentrate.

Registered to:
Lonza NZ Limited
Hudson Road, New Plymouth, NZ.
Tel: 06-755-9234 Fax: 06-755-1174

Net Contents: 1, 5, 10, 20 L

Batch:
D.O.M.:

®Registered Trademark of Lonza NZ Limited

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
08/12/2021

READ LABEL BEFORE USE. SHAKE OR STIR WELL BEFORE USE.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

08/12/2021

HSNO Classifications: 6.1D, 6.4A, 6.5B, 6.8B, 6.9B, 9.1A.

TOXICITY

Warning - Harmful if swallowed, by skin contact or if inhaled. May cause mild eye irritation and an allergic skin reaction. Suspected of damaging fertility or the unborn child and may cause liver damage from repeated oral exposure.

ECOTOXICITY

Very toxic to aquatic organisms. Do not apply into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Avoid contamination of any water supply with product or empty container.

QUALIFICATION REQUIREMENTS: Persons mixing, loading, applying, or otherwise handling PINNACLE must meet qualification requirements set out in the EPA Hazardous Substances (Hazardous Property Controls) Notice 2017.

PRECAUTIONS: Do not handle until all safety precautions have been read and understood.

PERSONAL PROTECTION

Avoid skin and eye contact and inhalation of spray mist. When mixing or applying wear eye protection, waterproof gloves, boots and overalls. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

STORAGE

Store in original container tightly closed and in a locked dry cool area away from foodstuffs, seeds and fertilizers. Store in accordance with NZS 8409 Management of Agrichemicals. Stores containing 100 litres are subject to signage and secondary containment, and more than 100 litres require emergency response plans.

EQUIPMENT

Apply using well maintained and accurately calibrated spray equipment

DISPOSAL

Triple rinse container and add rinsate to the spray tank. Recycle empty container, discard cap and deliver clean container to an Agrecovery depot, otherwise crush and bury in a suitable landfill. Disposal of product only by using according to the label, or at an appropriate landfill.

FIRST AID: For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If exposed or concerned, get medical advice. If swallowed do NOT induce vomiting. Rinse mouth. If inhaled move the victim to fresh air immediately and kept at rest in a position comfortable for breathing. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If irritation or rash occurs, get medical advice. If splashed in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical

advice.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

TRANSPORT: Do not carry this product on a passenger service vehicle.

MIXING: Invert container and shake well before use. Add the required amount of PINNACLE to the partly filled spray tank with agitator running and complete filling the tank with water.

COMPATIBILITY: Compatible with other commonly used fungicides and insecticides with the exception of alkaline materials.

GENERAL INFORMATION:
PINNACLE is a protectant fungicide for use as a preventative treatment against botrytis, downy mildew and phomopsis (dead arm) in grapes, botrytis, sclerotinia and late blight in field tomatoes and early blight, late blight and sclerotinia in potatoes. PINNACLE also controls powdery scab on potatoes and clubroot in vegetable brassicas when applied to the soil surface and incorporated prior to planting.

DIRECTIONS FOR USE:

CROP	DISEASE	NOTES
Grapes	Botrytis, Downy Mildew and Phomopsis (dead arm)	Use 100ml/100 litres. Target the full canopy or bunch-line and ensure complete coverage of the bunches and surrounding canopy. Apply at early and late flowering; just prior to bunch closure and again at the beginning of berry ripening (veraison), but no later than 28 days before harvest. Use alternative fungicides, particularly for phomopsis control, in the critical pre-flower period and for botrytis control during the pre-harvest period. Add a non ionic surfactant at a rate of 25ml/100 litres. The rate expressed is for high volume spraying to run-off. For concentrate spraying adjust dilution rate accordingly.
Field tomatoes	Botrytis, Sclerotinia and Late Blight	Use 750ml/ha at early flowering and again approximately 14 days later, towards the end of flowering. Apply in a minimum of 500 litres water per hectare and increase if necessary to ensure complete coverage. The period of sclerotinia control is four to eight weeks, and after the second application of PINNACLE, alternative approved fungicides may be required in high risk situations. For the control of late blight after flowering, use alternative approved fungicides for the remaining duration of the crop. Add a non ionic surfactant at a rate of 25ml/100 litres.
Potatoes	Early Blight, Late Blight and Sclerotinia	Use 250ml/ha at 7-10 day intervals for late blight and sclerotinia control or use 200ml/ha plus 1kg Mancozeb (80% ai)/ha if early blight is also likely. Use the shorter interval under severe disease

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
08/12/2021

		pressure conditions. Apply when the crop has fully emerged but before any sign of disease, using sufficient water per hectare to obtain complete coverage. Add a non ionic surfactant at a rate of 25ml/100 litres.
	Powdery Scab	Apply 4 litres/ha in 500 litres of water to cultivated ground prior to planting. Within 2 hours of application, incorporate thoroughly into the soil to a depth of 15cm, using either a rotary hoe or power harrow.
Vegetable Brassicas	Clubroot	Apply 4 litres/ha in 500 litres of water to cultivated ground prior to planting. Ensure thorough incorporation to transplant root depth (not more than 10cm) within 2 hours of applying, using a rotary hoe or power harrow. Soil should be moist at application and the crop transplanted within 2 days. At least 20mm of overhead irrigation should be applied soon after planting to further assist the distribution of PINNACLE throughout the upper soil profile. Disease control may be reduced if longer transplant or irrigation intervals occur.

RAINFALL:

Do not apply if rainfall is likely within 6 hours following application. Heavy rain following treatment will remove much of the fungicide from treated plant tissue and if disease pressure remains high, earlier re-treatment will be necessary.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

It is an offence to use this product on animals.

WITHHOLDING PERIODS:

Grapes: 28 days
Field Tomatoes: 42 days
Potatoes: 14 days
Vegetable Brassicas: Do not apply after planting

RESISTANCE MANAGEMENT: **GROUP 29 FUNGICIDE** Pinnacle contains a 2,6-dinitro-aniline fungicide with an uncoupling of mitochondrial oxidative phosphorylation mode of action. Though considered of low risk naturally occurring resistant strains of disease causing fungi could still develop from over use of this fungicide. To minimise this risk use Pinnacle strictly in accordance with the label instructions. See www.resistance.nzpps.org

Registered pursuant to the ACVM Act 1997, P007697. See foodsafety.govt.nz for registration conditions
Approved under the HSNO Act 1996, HSR100383. See www.epa.govt.nz for approval conditions

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

08/12/2021

NOTICE TO BUYER: The manufacturer and vendors of Pinnacle warrant the formulation to contain the specified active ingredients within accepted tolerance when packed but make no warranty of any kind on the use of this material and accept no responsibility for results, whether the material has been used in accordance with directions or not.

EMERGENCY PHONE: 0800 CHEMCALL 0800 243 6225

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

08/12/2021