CROPSTAR™ GLYPHOSATE 680 GRANULAR HERBICIDE

HERBICIDE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/05/2025

For weed control in agriculture, horticulture, drains, aquatic areas, forestry and other situations including general weed control

ACTIVE INGREDIENT: 680 g/Kg glyphosate as the mono-ammonium salt in the form of a water soluble granule

WARNING
KEEP OUT OF REACH OF CHILDREN

Net contents: X Kg

READ LABEL BEFORE USE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/05/2025

HAZARD CLASSIFICATIONS:
Hazardous to the Aquatic Environment Chronic Category 2
WARNING
Toxic to aquatic organisms with long lasting effects.
PRECAUTIONS
PERSONAL PROTECTION
Read carefully and follow all instructions. Avoid contact with eyes and skin. Do not breathe in the dust or spray mist. When mixing or applying, wear overalls, boots, gloves, and eye protection. Wear a respirator if dust or spray mist could be inhaled.
Wash hands thoroughly after use.
STORAGE
Store in the original tightly closed container in a secure area. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

DISPOSAL
Dispose of this product only by using in accordance with this label. Do not burn. Dispose of packaging to a suitable landfill. Do not burn. Do not use packaging for any other purpose.
ENVIRONMENTAL
Take all reasonable steps to ensure that this product does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water. Collect spillage.

FIRST AID
If medical advice is needed, have product container or label at hand.
For advice, call the Poisons Centre on 0800 POISON (0800 764 766) or a Doctor.
EMERGENCY RESPONSE
In a transport emergency call 111, Police or Fire Brigade. In case of fire, call 111 Fire Brigade. For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 6225) (24 hrs).
SPILLAGE
Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Moisten to prevent dusting and sweep up. Place in a waste container. Dispose of safely to a suitable landfill
SAFETY DATA SHEET
Additional information is provided in the safety data sheet. Freephone 0800 ORIONAS (0800 674 6627) or visit www.orionagriscience.co.nz

GENERAL INFORMATION
Cropstar Glyphosate 680 Granular Herbicide is a broad spectrum, non-selective, non- residual herbicide that is non-volatile and may be used in agriculture, horticulture and for other uses including general weed control. Cropstar Glyphosate 680 Granular Herbicide is recommended prior to sowing pasture and crops. The herbicide is absorbed by plant foliage and green stems. It is inactivated immediately upon contact with soil and does not provide residual weed control. Cropstar Glyphosate 680 Granular Herbicide moves from the point of foliage or stem contact into the root system, controlling the plant above and below the ground. Visible effects may take up to 14 days or even longer depending on growing conditions. These effects are a gradual wilting and yellowing of the foliage which advances to deterioration of above and below ground parts. Avoid contact with foliage or green stems of desirable plants and trees since severe injury or destruction may result. Avoid application in winds which allow drift onto desirable plants. Apply Cropstar Glyphosate 680 Granular Herbicide only to healthy, actively growing plants with adequate leaves for herbicide uptake. Avoid treating plants that are drought affected, frost damaged, silt laden or suffering the effects of previous herbicide application. Rainfall occurring up to 2 hours after application may reduce effectiveness. The rainfast period may be reduced to 30 minutes if used in conjunction with an organo-silicone penetrant such as Silmaxx™. If Cropstar Glyphosate 680 Granular Herbicide is applied to wet plants, some efficacy may be reduced due to run off.

MIXING INSTRUCTIONS
Cropstar Glyphosate 680 Granular Herbicide mixes readily with water. Reduced results may occur if dirty water is used (e.g. water from ponds, or unlined ditches) or if hard water containing calcium salts is used. Ensure the spray tank is free of any residue of previous spray materials.
PROCEDURE
Pre-mix Cropstar Glyphosate 680 Granular Herbicide with water into a slurry prior to adding to a half full spray tank. If adding other herbicides add prior to adding the herbicide to the tank. Use tank mixtures promptly. If using a penetrant add last to avoid foaming.

USE RECOMMENDATIONS
Apply with well-maintained and calibrated spray equipment. This product must only be handled, mixed and applied by a suitably qualified person as defined under the HPC notice 2017 Part 4C if applied onto or into water (otherwise not required). This product should only be mixed, contained in or sprayed equipment made from stainless steel, fibreglass, plastic, aluminium, brass or copper. A highly flammable gas may be formed from the contact of this product with galvanised or unlined steel. All spray equipment including pumps, spray tanks, lines, nozzles, and landing gear (aircraft) should be thoroughly rinsed with water after each day of spraying.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/05/2025

WITHHOLDING PERIODS
Wheat, Barley, and Oat grains – 7 days
Animal Safety Poisonous plants, such as ragwort, may become more palatable to stock after spraying, and livestock should be kept away until plants have died down. Otherwise, withholding is not required for stock grazing. However, it is recommended that grazing is delayed for -3 days for annual weeds, and 7 days for perennial weeds to ensure absorption and translocation.
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

Refer to "Weeds controlled" section for rates of application.

- **General Weed Control**
  Handgun Mix 0.5 kg of Cropstar Glyphosate 680 Granular Herbicide with 100 litres of water.
  Knapsack Mix 75 g of Cropstar Glyphosate 680 Granular Herbicide with 15 litres of water.
  Apply to completely wet the foliage, but not to runoff.
- **Brushweed Control**
  Cropstar Glyphosate 680 Granular Herbicide may be used to control a wide range of brushweeds and woody plants.
- **Agricultural Areas**
  Cropstar Glyphosate 680 Granular Herbicide may be used for the control of annual and perennial weeds as directed in agricultural land prior to sowing any edible or non-edible crop or pasture. Where white clover control is required, tank mix with a suitable white clover herbicide and observe recommended waiting periods prior to planting legumes. Cultivation may start from 3 days for annual weeds and 7 days after spraying perennial weeds.
- **Pasture Direct (Spot) Application**
  Cropstar Glyphosate 680 Granular Herbicide is non-selective and may damage or kill any plant in the sprayed area. Retreatment and/or pasture improvement may be necessary to restrict seedling re-establishment.
- **Selective Wiper Application**
  For the control of erect weeds in pastures.
- **Pasture Renovation**
  Cropstar Glyphosate 680 Granular Herbicide may be used to control existing pasture species prior to re- seeding pasture or establishing other crops.
  Preharvest Weed Control: Wheat, barley and oats - Apply up to 2.1 kg/ha depending on specific weeds present 7 - 12 days prior to harvest when the grain moisture is less than 30%. Can be used to control weeds prior to cultivation or prior to harvest. Do not harvest within 7 days after spraying. Do not spray undersown crops. Do not spray crops that will be harvested for seed. Avoid treating plants under drought stress.
- **Orchards/Vines -** Citrus, Apple, Pear, Stonefruit, Kiwifruit, Grapevines.
  Apply as a directed or shielded spray or with wiper equipment. For long term control of annual weeds tank mix with a suitable residual herbicide. **DO NOT** apply near unprotected trees less than 3 years old. **DO NOT** allow spray or spray drift to contact any part of stone fruit trees or green bark, suckers, buds, foliage or fruit of other trees. **DO NOT** allow wiper surfaces to contact any part of the trees. Weed foliage coated in copper or calcium based compounds may reduce the activity of Cropstar Glyphosate 680 Granular Herbicide.
- **Market Gardens**
  Cropstar Glyphosate 680 Granular Herbicide can be applied before or after seeding and prior to crop emergence. Application can also be made prior to transplanting or as a post harvest clean- up. For longer term control of seedling weeds the herbicide may be tank mixed with other suitable residual herbicides. Use in accordance with the label recommendations for these products.
- **Asparagus**
  Cropstar Glyphosate 680 Granular Herbicide can be applied:

  Following the final harvest and immediately after removal of all spears but before crop re-growth occurs.
  Prior to planting or crop emergence. **DO NOT** treat emerging spears.
  In established ferns use as a directed or shielded spray or with wiper equipment. **DO NOT** allow spray, spray drift or the wiper surface to contact any part of the asparagus plants.

- **Industrial and Non- Agricultural Areas**
  Cropstar Glyphosate 680 Granular Herbicide can be used for the control of annual and perennial weeds as directed. For residual control, the herbicide can be tank mixed with suitable residual herbicides. Exercise caution when using residual herbicides prior to planting trees, shrubs or other ornamental plants.
- **Turf Grass and Recreational Areas**
  Apply before establishing lawns, parks, reserves and ornamentals. Cropstar Glyphosate 680 Granular Herbicide can also be applied around the base of established trees as a directed or shielded spray or wiper equipment. DO NOT allow spray or spray drift or the wiper surface to contact any part of the tree. In mown areas, allow sufficient re-growth before spraying.
- **Aquatic Areas -Drains and Waterways**
  Ensure that 75% of the weed is above water level or efficacy may be reduced. Following spray application, **DO NOT** not submerge the treated plants as this will cause the herbicide to be washed off. Avoid treating silt laden plants.
- **Forestry**
  Cropstar Glyphosate 680 Granular Herbicide can be applied for the control of grasses and a wide range of brushweeds. The addition of a penetrant is required for the control of bracken and tobacco. **NOT** allow spray, spray drift or the wiper surface to contact the foliage or green bark of desirable trees as severe injury can occur.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/05/2025

APPLICATION METHODS
Ensure the equipment is in good working order and correctly calibrated. Wash through with water before use and ensure filters and nozzles are free of pesticide residue. Avoid spray drift. Care must be taken when applying this product to prevent injuring desirable plants and crops. To clean equipment after use, flush thoroughly with water.

- **Boom Application**

Water volumes up to 200 litres/ha of clean water are desirable. For broad acre boom application, flat fan, wide angle nozzles (e.g. 110°) using pressures of 200 – 280 kPa are preferred. Set the boom height to ensure a double overlap of the spray pattern at the top of the weed canopy. For orchard application wide angle, even spray nozzles are preferred.

- **Knapsack and Handgun Application**

Adjust the equipment to achieve a fine spray. Use the minimum spray volume necessary to achieve uniform and complete coverage without runoff. Handgun spraying pressures of 1,000 – 2,000 kPa and No. 4 to 6 spray tips are recommended.

- **Wiper Application**

Pre- mix 1 part of Cropstar Glyphosate 680 Granular Herbicide and 4 parts of clean water. Mix only enough herbicide for immediate needs and allow the wiping area to become fully saturated before commencing. The wiping area must remain moist and clean otherwise reduced results will occur. In heavy weed infestations or when treating perennial weeds such as Californian thistle or multi- crown ragwort, a double pass in opposite directions on the same day will give the best results. As weed density increases, decrease ground speed. Drain and wash after use.

- **Aircraft Application**

Cropstar Glyphosate 680 Granular Herbicide may be applied by aircraft in forestry and non- cropping areas only. Aerial application of Cropstar Glyphosate 680 Granular Herbicide is not recommended on cropping land or near desirable plant species as even slight drift may cause severe injury or destruction. **DO NOT** apply in gusty or windy conditions, or in conditions conducive to drift e.g. still air and inversion conditions.

- **Mistblower Application**

Use a solution of 150g in 10 litres of water. For the micron attachment, increase to 1 kg in 10 litres of water. Provide complete coverage and spray to wet but not to run-off.

ADDITIVES
Apply an organo-silicone penetrant such as Silmaxx where indicated under recommendations for specific weeds.

RESISTANCE MANAGEMENT

GROUP	9	HERBICIDE

Cropstar Glyphosate 680 Granular Herbicide belongs to the glycine chemical group. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different

mode of action. Consult http://resistance.nzpps.org/index.php for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

WARRANTY
This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P7999
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR007661
See www.epa.govt.nz for approval controls.

™ CROPSTAR and SILMAXX are Trade Marks of Orion Agriscience Limited

Registrant: Agsin Pte Ltd, Zagro Global Hub, 5 Woodlands Terrace, Singapore 738430

Distributor: Orion AgriScience Ltd, Unit 1, 15 Sir Gil Simpson Drive, Harewood, Christchurch 8053
0800 674 6627
www.orionagriscience.co.nz

Bar code

Batch No.	
Date of Manufacture	

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S.
(GLYPHOSATE MONOAMMONIUM 75%)
HAZARD CLASS 9
PACKING GROUP III
HAZCHEM CODE 2Z

PLEASE READ THE LABEL COMPLETELY BEFORE USE

WEEDS CONTROLLED TABLE

Weed	Boom/ Aerial Kg/ha	Handgun Kg/100 litres	Knapsack g/15 litres	Mistblower g/10 litres	CRITICAL COMMENTS
ANNUAL WEEDS (General)	0.5 - 1.0	0.1	15	30	Use higher rate for larger weeds.
AUSTRALIAN SEDGE	3 Aerial 4.5	0.5	75	150	Apply from November to March. Where rank growth is present the area should be burnt 12- 18 months prior to aerial application to ensure adequate spray coverage.
BARLEY GRASS	0.5	0.1	15	30	
BARBERRY	-	0.5	75	150	Apply January to April from flowering to late fruit. Where bushes have been cut, regrowth should be 1.5 metres tall before spraying.
BLACKBERRY	4.5 - 10	0.5 - 0.75	75	150	Apply to healthy plants from blossom to ripe fruit stage prior to autumn frosts (December to April). Only bushes with a well developed canopy of leaves should be treated. Early treatment (December to January) may be preferred in drought prone areas. Leave sprayed blackberry for 40 - 60 days before burning or slashing and oversow the area. Where treatment has been carried out under marginal conditions some regrowth can be expected within 1 - 2 years. For aerial application of hard to kill blackberry or where no follow up pasture management is anticipated, use the higher rate.
BOXTHORN	-	0.7	-	-	Apply November to April to actively growing plants. Avoid salt damaged plants. Spray to ensure complete coverage of dense thickets. **DO NOT** use additional surfactant as poor results may occur.
BRACKEN	4.5 + Penetrant 250mL/100L	0.5 + Penetrant 200mL/100L	75 + Penetrant 30mL/15L	-	Apply when fronds are unfurled (mid January to June) but prior to frost browning. Dense virgin bracken with high trash content and low leaf to root ratio should be burnt in spring and the unfurled frond regrowth sprayed the following March to June.
BROOM	4.5 + penetrant 250mL/100L	0.5 + penetrant 200mL/100L	75 +penetrant 30mL/15L	150 +penetrant 20mL/10L	Aerial application - Best results are obtained when plants are in full leaf. Handgun - Apply all year round. Spray to ensure full coverage.
BROWNTOP	1.5	0.25	40	75	Apply to 5 - 10 cm high actively growing plants.
BUDDLEJA	4.5	0.5	75	-	Handgun - If actively growing treat all year round. Aerial - Treat up to 2 metres high during February to April.
CALIFORNIAN THISTLE	2	0.5	75	150	Apply after flower bud development but before autumn browning. To ensure maximum shoot emergence **DO NOT** cultivate for 6 months prior to treatment. Wiper - A double pass at low speed is recommended in heavy infestations. Mowing 10 days after treatment may assist control. Retreatment may be necessary in the following season.
CLEAVERS	0.75	0.25	40	75	
COUCH	1.5 - 3	0.5	75	150	For suppression use 1.5kg/ha, for control use 2 - 3 kg/ha when growing conditions are satisfactory. Use the high rate in late spring. Apply to plants with at least 4 - 5 leaves and active shoot and root growth. **DO NOT** cultivate for 6 months prior to application to ensure maximum shoot emergence. In areas of dense infestation, a repeat application may be necessary.
DOCK	2 - 4.5	0.5	75	-	Treat plants in spring or summer when actively growing and in full leaf. Use low rate for seedlings or young plants, the high rate for control of established plants.
FLOATING SWEET GRASS REED SWEET GRASS	3	0.5	75	150	Apply to actively growing plants when most have reached early seedhead.
GORSE	-	0.5 + penetrant 200mL/100L	-	-	Apply all year round. Spray to ensure complete coverage.
INDIAN DOAB	4.5	0.5	75	-	Apply when most plants have reached early seedhead.
JOHNSON GRASS	3	0.5	75	150	Apply to actively growing plants.
KIKUYU	3	0.5	75	150	Apply to actively growing plants. Under dense growth conditions silorate and allow regrowth before treatment to improve control.
MERCER GRASS					Apply to actively growing plants. Under dense growth conditions silorate and allow regrowth before treatment to improve

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/05/2025

• Aquatic • Non-aquatic	4.5 3	0.75 0.5	100 75	225 150	control.
NODDING THISTLE	-	0.5	75	-	Wiper application - Apply when actively growing when at least 10cm of the weed can be wiped. In heavy infestations a double pass is recommended.
OLD MANS BEARD	-	0.5 - 1	150	-	Apply November to March prior to leaf senescence. Vines too high for complete coverage should be cut at ground level in winter or spring and the regrowth sprayed in March when plants are at least 0.5 metres long. Follow up treatment will be necessary to control seedlings. Use the low rate for small vines.
PAMPAS GRASS / TOETOE	-	0.5	75	-	Apply spring - autumn. Ensure complete coverage. Best results are achieved when applied after flowering.
PASPALUM	2 - 3	0.5	75	150	Use the low rate from January to April (early seedhead) prior to frosts. Use 3 kg/ha on vegetative plants.
POTATO (self set)	1.5	0.25	40	75	
PRAIRIE GRASS	0.75	-	-	-	
RAGWORT	3	0.5	75	-	Apply when actively growing. Wiper application - Apply when plants are 10 - 15 cm above the pasture. In heavy infestations a double pass is recommended. (Caution: Ragwort plants may become more palatable after spraying and stock should be removed until the treated plants brown out.)
RATSTAIL	2.5	-	-	-	Apply to actively growing plants in autumn.
RAUTAHI (Cutty grass)	3	0.5	75	150	Apply to actively growing plants from January to March. Burning or slashing may take place from 30 days following brownout.
RING FERN	0.5 - 1.5	0.15	25	-	Control: Apply 1.5kg/ha in January to April. Pasture Manipulation: Apply 0.5-1 kg/ha December to April. Use the higher rate for dense infestations. Burn 4 - 6 weeks after treatment and oversow with pasture.
RUSHES	3	0.5	75	150	Apply to actively growing plants from January to March. Burning or slashing may take place from 30 days after treatment following brownout. Wiper - apply November to May.
RYEGRASS PERENNIAL	0.75 – 1.5 + penetrant 100mL/100L	0.25 + penetrant 100mL/100L	40 + penetrant 15mL/15L	-	Apply any time plants are actively growing and 3 – 10 cm high. Use low rate January to July when cultivation is to follow spraying. Use 1.5 Kg/ha August to December.
SWEET BRIAR	6 + penetrant 250mL/100L	0.5 + penetrant 200mL/100L	75 +penetrant 30mL/15L	-	Apply from late flowering to early leaf yellowing. Best results achieved with full coverage.
TALL FESCUE	3	0.5	75	150	Apply to actively growing plants January to March.
WILLOW	4.5	0.5	-	-	Apply from January to April before leaf fall begins. Total spray coverage is essential for best results.
YARROW	1.5 - 3	-	-	-	Apply to plants with 5 - 10 cm of leaf. Cultivate with a minimum of 2 grubbings from 10 days after treatment. Use the 1.5 kg/ha rate when a competitive crop to yarrow is to be drilled. Use the higher rate when the crop to be drilled is a poor competitor.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/05/2025