READ LABEL CAREFULLY BEFORE USING

Contego B Sub

BIOLOGICAL SOLUTIONS

MINISTRY OF PRIMARY INDUSTRIES
NEW ZEALAND
09/12/2024

A bio-fungicide to aid in the control of Botrytis, Powdery Mildew and other fungal pathogens on grapes, fruits, potatoes, ornamentals and vegetables.

CERTIFIED INPUT FOR ORGANICS
bio gro NZ CERT
5279

Contains: 250ml of *Bacillus subtilis* (ATCC 6051 strain) /kg
Contains 1 x 10 cfu/g in the form of a wettable powder

NET CONTENTS

1kg

biological SOLUTIONS

Registered pursuant to the ACVM Act 1997: P8116
Approved pursuant to the HSNO Act 1996: HSR100667
See www.mpi.govt.nz for registration conditions

Contego B Sub
BIOLOGICAL SOLUTIONS

READ LABEL CAREFULLY BEFORE USING

A bio-fungicide to aid in the control of Botrytis, Powdery Mildew and other fungal pathogens on grapes, fruits, potatoes, ornamentals & vegetables.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
09/12/2024

Contains: 250ml of *Bacillus subtilis* (ATCC 6051 strain) /kg
Contains 1 x 10^9 cfu/g in the form of a wettable powder

CONTEGO B SUB has been developed from naturally occurring strains of Bacillus Subtilis. It is a broad-spectrum preventative fungicide recommended for the control of powdery mildew and grey mould (Botrytis sp.) and Sour Rot in grapes during periods of low disease infection pressure. It acts by preventing spore germination, germ tube elongation and host penetration.

This product can be integrated into an overall disease and pest management strategy wherever fungicide use is necessary in an IPM programme. Thorough coverage is essential for optimum disease control. It is useful for integration into resistance management strategies and should be applied prior to or in the early stages of disease development.

This product is harmful to aquatic organisms. It is an offence to use this product on animals. Do not apply to plants in flower while bees are present.

CERTIFIED INPUT FOR ORGANICS
bio gro
5279

PRECAUTIONARY STATEMENTS
The manufacturer makes no warranty, expressed or implied, regarding the performance of this product in the absence of control over how it is used. Keep out of reach of children.

APPLICATION PROCEDURE:
Thorough coverage is essential for optimum disease control, to achieve good coverage is is critical to use the correct spray pressure, water volume nozzle-size and spacing with the correct ground speed. CONTEGO B SUB is intended for use as a dilute spray in water for application through conventional spray equipment.

MIXING DIRECTIONS:
Partially fill the spray tank with clean water and start agitation. Add the specified amount for the area to be treated to the spray tank. Complete filling the tank to the required volume. Maintain agitation until spraying is complete. Do not allow the spray mixture to stand overnight or for any prolonged periods. Spray pH must be not less than pH 4.5 and not greater pH 8.0.

COMPATIBILITY:
Do not combine CONTEGO B SUB in the spray tank with pesticides, surfactants, copper based products or fertilisers unless prior experience has shown the combination to be compatible. Any tank combinations must also be tested for effectiveness on disease and selectivity to plants before use on commercial areas.

If the use of any tank mixture is desirable, a small volume of the mixture should first be tested in a small measured volume, for physical compatibility at the dilution rates required. If this test is positive, a field test of the mixture should then be carried out on a few plants to observe effectiveness and plant safety.

APPLICATION RATES:
Use rate for all crops 1.5 – 3.0kg /ha or 150 – 350 g /100 litres in sufficient water to ensure complete coverage of plants. Actual application rates will be determined by infection and disease pressure. Standard or normal infection and disease pressure the lower application levels are recommended.

GRAPES / FRUIT: Use all season to increase protection against fungal plant pathogens, specific botrytis fungicides should be targeted at key grape botrytis timings. Can be used later in season as it has no residue and no withholding period restrictions. Two strategically timed applications are recommended at 4-5 weeks from harvest and again at 1 week from harvest. Apply in enough water to ensure thorough coverage of all parts of the plants.

ORNAMENTALS: Apply at first sign of disease and repeat at 7-10 day intervals.

VEGETABLES / POTATOES: Can be used all season to increase protection against fungal plant pathogens but specific fungicides should be targeted at key timings .Can also be used later in the season as it has nil residues and no withholding period restrictions.

HSNO CLASS: 9.1D, 6.4A, 6.5B

WARNING: *Causes eye irritation. May cause an allergic skin reaction. Harmful to aquatic life. Precautions: Read label before use. Avoid breathing dust or spray. Wash hands and face thoroughly after handling. Contaminated work clothing should not be allowed out of the work place. Avoid release into the environment. Wear protective gloves/protective clothing/eye protection/face protection as appropriate for the product.*

FIRST AID: *IF IN EYES: Rinse cautiously with water for 15 minutes. Remove contact lenses if present and easy to do. Continue rinsing. IF ON SKIN: wash with plenty of soap and water. If skin or eye irritation persists, or rash occurs: get medical advice/attention. If swallowed DO NOT induce vomiting. For advice call the National Poisons Centre 0800 POISON (0800 764 766).*

RESTRICTIONS ON USE: *This substance must not be applied onto or into water.*

PRODUCT AND PACKAGING DISPOSAL: *Dispose unused, expired or damaged product through normal solid landfill rubbish disposal systems. STORAGE: Store in a locked, dry, cool, well ventilated area. Storage must be in accordance with the current version of NZS8409 management of agrichemicals. See the safety data sheet for further information. When stored appropriately this product should show no significant degradation for one year from date of manufacture. Contact your supplier for further information about the use of any product that is older than this.*

Leaf plucking in vineyards: Sheep must not be sent for slaughter or milked for 6 months after being removed from the vineyard (and placed on clean feed)

Registered to:
Biological Solutions Ltd
4 Cargill Place, Richmond, Nelson 7020
P: 0800 10 60 91
admin@biosolutions.co.nz
www.biosolutions.co.nz

Batch No:
Date of manufacture:
Expiry date:

NET CONTENTS
1kg