KEEP OUT OF REACH OF CHILDREN

UNISHIELD

Fungicide

For the control of certain fungal diseases in apples, grapes, stonefruit, tamarillos, vegetables and ornamentals

ACTIVE INGREDIENT
800 g/kg SULPHUR in the form of a water dispersible granule.

GROUP	M2	FUNGICIDE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/07/2025

CERTIFIED INPUT FOR ORGANICS
bio gro NZ CERT
4429

Net Contents: 15 kg
Bar code to be inserted

PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/07/2025

Designed for biocidal action.

PRECAUTIONARY STATEMENTS
PREVENTION
When handling wear eye protection and gloves.
Wash hands thoroughly after handling.
Do not eat drink or smoke when using this product.
Do not apply into or onto water.
RESPONSE
Collect spillage.
STORAGE
Keep out of reach of children. Store in accordance with NZS8409 Management of Agrichemicals.
When stored appropriately this product should show no significant degradation for 3 years. Contact your supplier for further information about the use of any product that is older than this.
DISPOSAL
Ensure container is completely empty. Bury in a suitable landfill. Dispose of product only by using according to the label, or at an appropriate landfill.

FIRST AID
If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre (0800 POISON) 0800 764 766 or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If in the eyes, wash out immediately with water.

SPILLAGE
Wear appropriate protective clothing and prevent material from entering waterways. Sweep up solids without generating dust. Contain liquid spills and absorb with inert material. Dispose of waste safely in an appropriate landfill.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at https://www.uplcorp.com/nz/

EXCLUSION OF LIABILITY: PLEASE READ
Please read these conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened.

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to:
UPL New Zealand Limited
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
https://www.uplcorp.com/nz/

Registered pursuant to the ACVM Act 1997, No. P008337
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval Code HSR000816.
See www.epa.govt.nz for approval controls

Batch number:	
Date of Manufacture:	

In a transport emergency dial 111, Police or Fire Brigade.
***For specialist advice in an emergency only, call* 0800 CHEMCALL (0800 243 622)**

UN-Free

DIRECTIONS FOR USE:
It is an offence to use this product on animals.

CROP	DISEASE	RATE	NOTES
Apples (Do NOT apply to Pears)	Powdery mildew	100-200g/100 litres water	Apply from bud movement until harvest at 10-14 day intervals. Do NOT apply in combination with captan on sulphur sensitive varieties such as Kidd's Orange Red, Cox's Orange Pippin or Red Delicious. Fruit finish may be affected on some apple varieties. Do NOT use on apples post-blossom unless one sulphur containing spray has been used pre-blossom. Do NOT apply in temperatures above 25^0C.
Grapes	Powdery mildew, Erineum mite	150-300g/100 litres water	Apply at 10-14 day intervals from bud-swell until 6 weeks before harvest. Use the higher rate when conditions favour disease.
Stonefruit (Do NOT apply to apricots)	Rust	100-200g/100 litres water	Apply at 21 day intervals from fruit set until harvest. Use the higher rate when conditions favour the disease.
Tamarillos	Powdery mildew	100-200g/100 litres water	Apply at 21 day intervals throughout the growing season.
Vegetables (Do NOT apply to Cucumbers, Rockmelons & Tomatoes)	Powdery mildew, Rust	100-200g/100 litres water	Apply every 10-14 day intervals when disease infection is likely.
Ornamentals	Powdery mildew Rust	100-200g/100 litres water	Apply every 10-14 day intervals when disease infection is likely.

MIXING:
The rates suggested are for high volume spraying to just before run-off. For concentrate spraying adjust dilution rate accordingly. Use the higher rates and/or shorter spray intervals when disease pressure is high.
The recommended quantity of **UNISHIELD** may be added directly to the water in the spray tank with the agitation running.
If the leaves are hairy and/or waxy, or where the water is hard, add a suitable wetting agent at recommended rates.

MINISTER FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/07/2025

RESISTANCE MANAGEMENT
UNISHIELD contains the group M2 fungicide, sulphur. To minimise risk maintain the label dose rate. Use as part of a spray program in combination with other fungicides with different modes of action where relevant.

COMPATIBILITY:
UNISHIELD should not be used in combination oils or formulations containing oils, or used within 21 days of an oil application. The product is compatible with most other formulations of fungicides and insecticides.

EQUIPMENT:
UNISHIELD is suitable for use in most agricultural sprayers. Do not use strainers finer than 300 micrometres. Calibrate equipment to give the desired application rate and full coverage of foliage. Fine droplets are more effective than coarse (large) droplets.

WITHHOLDING PERIOD:
All crops: 1 day

Leaf plucking in vineyards: Sheep must not be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/07/2025