10kg Paper Bag

31"

11” FRONT | 4.5” | 11” FRONT | 4.5”

22”

Hazard Classification
6.3A, 9.1D

WARNING : Causes skin irritation.
Ecotoxic: Biocidal action.

PRECAUTIONS
Keep out of reach of children. Do not apply directly into / onto water.
Take all reasonable steps to ensure the substance does not cause any significant adverse effects to the environment beyond the application area.

PROTECTIVE CLOTHING
When mixing and applying spray, wear eye protection, cotton overalls, suitable gloves and footwear. Avoid skin contact and inhalation of spray mist. Remove protective clothing and wash hands and exposed skin thoroughly with soap and water before meals and after work.

STORAGE
Store in the original container, tightly closed and away from foodstuffs and animal feeds.

APPLICATION
Apply with well calibrated spray equipment. Do not allow spray drift outside the target area to occur.

FIRST AID
If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre on 0800 POISONS (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. Do NOT scrub skin. If splashed in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do so. Continue rinsing. If eye or skin irritation persists seek medical advice.

SPILLAGE
In event of spillage, prevent further spillage and contain spill to prevent product from entering waterways. Wear protective clothing and sweep up spill and place in sealed container for disposal. Avoid dust formation. Wash affected area with water and detergent. Absorb with inert material and place used absorbent materials in disposal container. Dispose of waste in at an approved landfill.

CONTAINER DISPOSAL
Crush and bury empty container in an approved landfill.

Directions for Use:
COSAVET DF is a protectant fungicide and for best results should be applied before disease infection occurs. The rates suggested are for high volume spraying to run-off. For concentrate spraying adjust dilution rates accordingly.

It is an offence to use this product on animals.

Crop	*Disease*	*Use Rate / 100L*	*Critical Comments*
Apples	Powdery Mildew	100-200 g	Apply from bud movement until harvest at 14 day intervals. Do NOT apply within 5 days of Captan application. Do NOT apply to sulphur sensitive varieties, such as Kidds Orange, Cox's Orange Pippin or Red Delicious. Do NOT apply to pears. Do NOT apply in temperatures above 25 degrees Celsius.
Grapes	Powdery Mildew	150-300 g	Apply from bud swell at 10-14 day intervals. Use the higher rate when disease is present.
Peaches Plums Nectarines	Rust	100-200 g	Apply at fruit set and repeat at 21 day intervals until harvest. Use the higher rate when conditions favour infection. Do NOT apply to apricots.
Ornamentals	Powdery Mildew and Rust	100-200 g	Apply at 10-14 day intervals when conditions favour infection.
Tamarillos	Powdery Mildew	100-200 g	Apply at 14 day intervals throughout the growing season.
Vegetables	Powdery Mildew	100-200 g	Apply at 10-14 day intervals when conditions favour infection. Use the higher rate when disease is present. Do NOT apply to cucumbers or rockmelons.

Application
Suitable for use with all types of applicators. Good spray coverage is necessary for disease control. Clean out spray equipment after use.

Mixing
Partially fill spray tank to about one quarter capacity and start agitation. Slowly add the required amount of COSAVET DF while continuing to fill the spray tank. Maintain agitation throughout application.

Compatibility
COSAVET DF is compatible with most commonly used fungicides, insecticides and foliar fertilisers, however mixtures with Captan are not recommended. COSAVET DF is NOT COMPATIBLE with spraying oils or formulations containing mineral oils. Do NOT apply within 21 days of an oil application.

Rainfall
Rainfall within 3 hours of application may reduce efficacy.

Withholding Period
1 day

Resistance prevention or management
COSAVET DF Fungicide contains the GROUP M2 fungicide, sulphur. Do not use another product containing a fungicide in the same group in conjunction with COSAVET DF Fungicide.

Conditions of Supply
To the extent permitted by law, buyers and/or users accept that neither the manufacturer, Fruitfed Supplies (Part of PGG Wrightson Ltd) as distributor, nor any other of its distributors, will accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential, howsoever arising in connection with the supply of this product including without limitation their storage, application, handling or use.

Batch No:
Date of Manufacture:

Expiry Date: When stored appropriately this product should show no significant degradation for two years from the date of manufacture.

Registered to: Sulphur Mills Australia Pty. Ltd, 4 Holly Avenue, Ryde, NSW 2112, Australia.

COSAVET™ DF

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
06/05/2021

WARNING
KEEP OUT OF REACH OF CHILDREN

Reap®

READ TOWARDS SAFER ECO-FRIENDLY PRODUCTS
PRODUCTS USED IN OVER 80 COUNTRIES WORLD WIDE

COSAVET™ DF

FUNGICIDE

ACTIVE INGREDIENT:
Contains 800g/kg sulphur in the form of a water dispersible granule

For the control of powdery mildew and rust diseases of apples, stone fruit, grapes, tamarillos, ornamentals and vegetables.

©COSAVET DF is a registered trademark of Sulphur Mills Australia Pty Limited

Distributed by:
Fruitfed Supplies

Fruitfed Supplies (Part of PGG Wrightson Ltd)
Building 2
1 William Pickering Drive
Albany
Auckland
Phone: 09-448-0510
Fax: 09-448-0511

Registered persuant to ACVM Act 1997, No. P8409.
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, HSR000741
See www.epa.govt.nz for approval controls.

CERTIFIED INPUT FOR ORGANICS bio gro NZ CERT
No. 5436

Net contents: 10kg

WARNING
READ LABEL THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

COSAVET™ DF

Date: 01-02-2021

\\192.168.1.3\k\l\Pratap\Export\country wise\New Zealand\Cosavet DF 10 kg Sulphur 80% WG paper bag **PGG Wrightson New Zealand** 01-02-2021.cdr