DANGER

KEEP OUT OF REACH OF CHILDREN

CENTURION® XTRA

Herbicide

A selective post-emergent herbicide for grass weed control in broadleaf crops, orchards, vineyards, fodder and sugar beet, and forestry

Active ingredient
Contains 360 g/litre CLETHODIM in the form of an emulsifiable concentrate

GROUP	1	HERBICIDE

Net contents: 5 L, 20 L

Bar code to be inserted

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/06/2025

Public

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of registration.
Any person purchasing, handling or disposing of this product must be suitably qualified.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/06/2025

DANGER
Combustible liquid.
May be fatal if swallowed and enters airways.
Causes skin irritation.
Causes serious eye irritation.
May cause liver damage from prolonged or repeated exposure.
Toxic to aquatic organisms with long lasting effects.
Hazardous to soil organisms.
Hazardous to terrestrial vertebrates.

PRECAUTIONARY STATEMENTS

PREVENTION
Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.
Wear protective gloves, protective clothing, and eye protection.
Wash hands thoroughly after handling.
Do not breathe spray.
Do not apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.
Records of application must be kept.

RESPONSE:
In case of fire: Use carbon dioxide, dry chemical, foam, or fog to extinguish.
IF SWALLOWED: Immediately call the National Poisons Center 0800 POISON (0800 764 766) or a doctor.
Do NOT induce vomiting.
IF ON SKIN: Wash with plenty of soap and water.
If skin irritation occurs: Get medical advice.
Take off contaminated clothing and wash before reuse.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
If eye irritation persists: Get medical advice.
Get medical advice if you feel unwell.
Collect spillage.

STORAGE:
Store locked up in a well-ventilated place.
Keep cool

DISPOSAL:
Dispose of contents/container in accordance with local regulations.
Product Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.
Container Disposal: Triple rinse empty container and add rinsate into the spray tank. If recycling, discard cap and deliver clean container to an AgRecovery depot. Otherwise crush and bury in an approved landfill.

FIRST AID
If medical advice is needed, have the product container or label at hand. If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If spilt on skin, remove contaminated clothing and wash affected areas of skin. DO NOT SCRUB THE SKIN. If concentrate is splashed in eyes flush with running water for at least 15 minutes. Take to hospital without delay. If inhaled, remove to fresh air. If breathing is difficult, take to hospital without delay.

SPILLS
When dealing with spills wear personal protective clothing. Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at www.uplcorp.com/nz.

EXCLUSION OF LIABILITY – PLEASE READ
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to:
UPL New Zealand Ltd
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
www.uplcorp.com/nz

Registered pursuant to the ACVM Act 1997, No. P008655.
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996. Approval Code HSR100646.
See www.epa.govt.nz for approval controls.

© Copyright UPL New Zealand Limited 2020, all rights reserved.
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited.

® Registered trademark

Batch number:	
Date of Manufacture:	

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622)

UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Clethodim)
MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2X

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/06/2025

Label – Back



MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/06/2025

GENERAL INFORMATION

RESISTANCE MANAGEMENT

CENTURION XTRA contains a herbicide from the cyclohexanedione group and has the inhibition of acetyl coA carboxylase mode of action. Some naturally-occurring weed biotypes resistant to **CENTURION XTRA** and other Group 1 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by **CENTURION XTRA** or other Group 1 Herbicides. See www.nzpps.org for more information.

DIRECTIONS FOR USE

For best results apply to young actively growing grass weeds. Do not apply to plants under stress from drought or waterlogging. Use the lower rate on seedling grass weeds (2-5 leaf) or actively regenerating rhizomes, and the higher rate on tillering plants or hardy rhizomatous grasses.

On established sod-bound rhizomatous grasses and established turf ryegrass in non-cultivated crops, suppression only may be obtained.

On crops which have been grazed, remove stock and allow grasses to freshen up for a few days (5-10 cm new growth) before treating. To minimise any possible crop damage, avoid spray overlaps and apply during cool conditions in the morning or evening, especially when tank mixing with other agrichemicals. **CENTURION XTRA** is considered to be rainfast if applied at least one hour before rainfall.

Always add an emulsifiable crop mineral oil at 2.0 L/ha with all applications.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/06/2025

CROP	CRITICAL COMMENTS
White clover seed	Apply after closing up, to control annual and perennial ryegrass and other grass weeds
Legume feeds (e.g. lucerne, lupins, clover, lotus, trefoil)	To control annual and perennial ryegrass and other grass weeds.
Fodder brassicas	Apply when grass seedlings are small (typically 3-6 weeks after planting) and well before canopy closure.
Fodder beet, sugar beet	For best results apply when grass seedlings are small (typically 6-8 weeks after planting) and before canopy closure. Do not apply in tank mixture with other herbicides, fungicides or insecticides unless crop safety is known. Ensure thorough cultivation prior to sowing and do not use more than 1.3 L/ha.
Forage herbs (e.g. plantain, chicory)	To control annual and perennial ryegrass and other grass weeds. Do not apply **CENTURION XTRA** to plantain seed crops after flower heads are first visible in the base of the plants.
Orchards and vineyards	To control actively growing grass weeds. Apply as a directed spray to the base of the trees or vines to avoid fruit contact.
Vegetables	To control actively growing grass weeds. Do not apply more than 670 mL/ha. Crop safety has been demonstrated on onions, carrots and lettuce. Where crop safety is unknown, a small area should be treated before widespread application. NB: Asparagus – DO NOT apply during the cutting period.
Peas, lentils, oilseed rape	To control wild oats. Apply 170 mL/ha when the majority of oats are in the 2-5 leaf growth stage. A temporary effect on leaf colour and sheen may occur in peas. This does not affect crop vigour or yield.
Forestry	Do not exceed 670 mL/ha. **Tree release** – **Spot treatment:** Use 23 mL **CENTURION XTRA** plus 100 mL emulsifiable crop oil per ten litres of water. Apply 20 mL spray solution per square

CROP	CRITICAL COMMENTS
(radiata pine, macrocarpa, douglas fir)	metre. To control broadleaf weeds and for residual control, apply in combination with Terb 500® at recommended label rates.

Grasses controlled and recommended use rates

GRASS SPECIES	RATES/HA
Wild oats, annual and perennial ryegrass, cocksfoot, barnyard grass, summergrass, floating sweet grass, canary grass, bristle grass, volunteer cereals. Note: Turf ryegrasses and tillered cereals will only be suppressed at these rates.	170-330 mL
Annual poa, smooth witchgrass, soft brome, phalaris, tall fescue, prairie grass, ripgut brome, browntop, barley grass. Note: Chewings fescue, Yorkshire fog and hairgrass will be suppressed but not controlled at these rates.	330-670 mL
Couch, paspalum, Indian doab	1.3-2.0 L

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MIXING
Half fill the spray tank and with the agitator operating and add the correct volume of **CENTURION XTRA**. Add the remainder of the water, then add the adjuvant and agitate thoroughly before using. After standing, agitate spray solution well before reusing.

EQUIPMENT
Centurion Xtra MUST only be applied via ground-based methods.
Apply using well maintained spray equipment that is accurately calibrated to apply 100-400 L of water per hectare. Use nozzles and pressures that produce a droplet size of 150-300 microns VMD. Use the high water rate for dense grass infestations. Clean equipment thoroughly after use.

COMPATIBILITY
CENTURION XTRA is compatible with Terb 500©, clopyralid and Pycus®. Do not apply in combination with other herbicides unless experience shows that efficacy is not affected. Do not apply within 7 days of phenoxy herbicides.

WITHHOLDING PERIODS
Vegetable crops – 5 weeks.
Grapes – Do not apply after the start of flowering.
Grazing and stock feed harvest withholding periods:
Legume crops, forage herbs, fodder brassicas – 3 weeks.
Fodder beet and sugar beet – 9 weeks.
® Registered trademarks

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
16/06/2025