PruneTec®

Contains: 10 g/litre tebuconazole in the form of a ready to use liquid

Net Contents: 2 litres, 5 litres, 10 litres

Group	3	Fungicide

PruneTec is a long lasting pruning wound sealant containing fungicide for the control of European Canker in Pipfruit, Silver Leaf in pip and summer fruit, Eutypa Dieback, Botryosphaeria Dieback and Petri Disease in grapes. Also for sealing grafts and pruning wounds in grapes, kiwifruit, ornamental trees and pip and summer fruit.

Registered pursuant to the ACVM Act 1997, No. P8878
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to HSNO Act 1996, No. HSR000519
See www.epa.govt.nz for approval conditions.

Do not store in freezing conditions as product will harden.

HSNO Classifications:

Reproductive toxicity Category 2
Hazardous to the aquatic environment chronic Category 3

WARNING
Hazards: Suspected of damaging fertility or the unborn child. Harmful to aquatic life with long lasting effects.
PRECAUTIONS:
Read label before use. Keep out of reach of children. When mixing or applying wear protective clothing. Do not eat, drink or smoke while using. Wash hands and face before meals and after work.
STORAGE:
Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from foodstuffs. Avoid storage under cold/frost conditions. Do not store in freezing conditions as product will harden.
SPILLS:
Ensure suitable personal protection during removal of spillages. Contain and collect spillage. Avoid release to the environment.
DISPOSAL:
Dispose of this product only at an approved landfill. Container Disposal: Triple rinse container and dispose at an approved landfill.
FIRST AID:
After contact with skin, wash immediately with water and soap. Contamination of the eyes must be treated by thorough irrigation with water. Inhalation, move to fresh air and keep person warm and at rest. Ingestion, seek medical advice if feeling unwell. If skin or eye irritation persists or person feels unwell get medical advice. For advice contact the National Poisons Centre 0800 764 766.

DIRECTIONS FOR USE:
Stir thoroughly before use. Do not dilute **PruneTec** with water as the seal integrity will be reduced.

Application: Apply **PruneTec** with a paint brush or applicator to a clean surface, in a continuous uniform layer over and beyond (minimum 1 cm) the edges of the cut surface.

Healthy Wood: Apply **PruneTec** directly after the cut is made.

Infected Wood: All infected, stained wood must be cut out, or alternatively the branch should be cut back to healthy tissue before applying **PruneTec**.

Protection of Grafts: Apply **PruneTec** after grafting and before or after taping. Do not apply before taping if the union is not perfect, otherwise **PruneTec** may penetrate the union and prevent the graft from taking. Where large wounds have been made check and repaint if cracking appears after the sealant has dried.

Brushes should be washed out in water after use.

PruneTec should only be applied to plants during the dormant period.

It is an offence to use this product on animals.

Sheep grazing withholding period in vineyards: 1-day

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/12/2024

Resistance Management: PruneTec contains tebuconazole, a group 3 fungicide with a demethylation inhibitor mode of action. Resistance to **PruneTec** and related products could develop in some disease situations from repeated use of the product. To minimise this risk, use strictly in accordance with label instructions.

CONDITIONS OF SALE & NOTICE TO BUYER:
This product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label when used in accordance with directions. No acceptance is given for liability or responsibility for any loss, damage or injury whether direct or indirect arising in connection with these goods, including their storage, application, handling or use. All such risks shall be assumed by the buyer. To the extent permitted at law, no warranties of merchantable quality or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Date of Manufacture:

Batch No:

PruneTec® is a registered trademark.

Registered to:
Max Stanton Moore
69B Rossall Street, Merivale, Christchurch, 8014
Ph: 021426528

EMERGENCY RESPONSE: New Zealand National Poisons Centre Dial 0800 764 766
Transport Emergency Dial 111

LIABILITY: This product must be used strictly as directed. Yates will not be liable for any loss or damage arising from a failure to strictly follow the directions for use.

FOR MORE GARDENING ADVICE
You can talk to Yates (NZ) 0800 693 297
yates.co.nz

®PruneTec is a registered trademark.
®Yates is a registered trademark. ™The Squares Device is a registered trademark. The front panel of this label is owned under copyright laws by Yates New Zealand.

Registered pursuant to the ACVM Act 1997, No. P8878
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, No. HSR000519
See www.epa.govt.nz for approval conditions

Registered to:
Max Stanton Moore
69B Rossall St, Merivale,
Christchurch, 8014

Distributed by:
DuluxGroup (New Zealand) Pty Ltd
The Gate, 373 Neilson St, Onehunga, Auckland 1061

BN&DOM: Printed on bottle

9 310428 550980

Yates®

PruneTec®

PRUNING & GRAFTING SEALANT
WITH APPLICATOR BRUSH

Fungicidal sealant for pruning cuts in ornamentals, pipfruit, stonefruit & vines

200 mL

Yates PruneTec® is a long-lasting pruning wound treatment and grafting sealant containing fungicide for the control of European Canker in pipfruit, Silver Leaf in pip & summerfruit, Eutypa Dieback, Botryosphaeria Dieback & Petri Disease in grapes.

- Encourages beneficial callusing to form on pruning wounds
- Dries quickly
- The seal remains elastic, and stretches as the plant grows without cracking or splitting
- Natural bark colour, to visually blend into the garden

IMPORTANT: PLEASE READ THE LABEL COMPLETELY BEFORE USE

ACTIVE INGREDIENT: CONTAINS 10G/LITRE
TEBUCONAZOLE IN THE FORM OF A READY TO USE LIQUID

DIRECTIONS: Apply evenly over a clean cut surface, covering at least 1cm beyond the edges of the cut. For healthy wood, apply directly after the cut is made. For infected wood, cut back to healthy tissue before applying. For grafting, apply after grafting and after taping. Do not dilute product with water.

Yates PruneTec® should only be applied to plants during their dormant period. It is an offense to use this product on animals.

PRECAUTIONS: Keep out of reach of children. When applying wear waterproof gloves. Do not eat, drink or smoke while using. Wash hands after use.

STORAGE & DISPOSAL: Store in original container, tightly closed, safely away from children and foodstuffs. Do not store in freezing conditions as product will harden. Dispose of this product only at an approved landfill. Avoid contamination of any water supply with contents or empty container.

FIRST AID: For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

FOR EMERGENCIES ONLY CALL: 0800 734 607 (ALL HOURS)

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/12/2024

PruneTec®

Contains: 10 g/litre tebuconazole in the form of a ready to use liquid

Net Contents: 500ml, 1 litre

Group	3	Fungicide

PruneTec is a long lasting pruning wound sealant containing fungicide for the control of European Canker in Pipfruit, Silver Leaf in pip and summer fruit, Eutypa Dieback, Botryosphaeria Dieback and Petri Disease in grapes. Also for sealing grafts and pruning wounds in grapes, kiwifruit, ornamental trees and pip and summer fruit.

Registered pursuant to the ACVM Act 1997, No. P8878
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to HSNO Act 1996, No. HSR000519
See www.epa.govt.nz for approval conditions.

Do not store in freezing conditions as product will harden.

HSNO Classifications:

Reproductive toxicity Category 2
Hazardous to the aquatic environment chronic Category 3

WARNING
Hazards: Suspected of damaging fertility or the unborn child. Harmful to aquatic life with long lasting effects.
PRECAUTIONS:
Read label before use. Keep out of reach of children. When mixing or applying wear protective clothing. Do not eat, drink or smoke while using. Wash hands and face before meals and after work.
STORAGE:
Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from foodstuffs. Avoid storage under cold/frost conditions. Do not store in freezing conditions as product will harden.
SPILLS:
Ensure suitable personal protection during removal of spillages. Contain and collect spillage. Avoid release to the environment.
DISPOSAL
Dispose of this product only at an approved landfill. Container Disposal: Triple rinse container and dispose at an approved landfill.
FIRST AID
After contact with skin, wash immediately with water and soap. Contamination of the eyes must be treated by thorough irrigation with water. Inhalation, move to fresh air and keep person warm and at rest. Ingestion, seek medical advice if feeling unwell. If skin or eye irritation persists or person feels unwell get medical advice. For advice contact the National Poisons Centre 0800 764 766.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/12/2024

DIRECTIONS FOR USE:
Stir thoroughly before use. Do not dilute **PruneTec** with water as the seal integrity will be reduced.

Application: Apply **PruneTec** with a paint brush or applicator to a clean surface, in a continuous uniform layer over and beyond (minimum 1 cm) the edges of the cut surface.

Healthy Wood: Apply **PruneTec** directly after the cut is made.

Infected Wood: All infected, stained wood must be cut out, or alternatively the branch should be cut back to healthy tissue before applying **PruneTec**.

Protection of Grafts: Apply **PruneTec** after grafting and before or after taping. Do not apply before taping if the union is not perfect, otherwise **PruneTec** may penetrate the union and prevent the graft from taking. Where large wounds have been made check and repaint if cracking appears after the sealant has dried.

Brushes should be washed out in water after use.

PruneTec should only be applied to plants during the dormant period.

It is an offence to use this product on animals.

Resistance Management: PruneTec contains tebuconazole, a group 3 fungicide with a demethylation inhibitor mode of action. Resistance to **PruneTec** and related products could develop in some disease situations from repeated use of the product. To minimise this risk, use strictly in accordance with label instructions.

CONDITIONS OF SALE & NOTICE TO BUYER:
This product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label when used in accordance with directions. No acceptance is given for liability or responsibility for any loss, damage or injury whether direct or indirect arising in connection with these goods, including their storage, application, handling or use. All such risks shall be assumed by the buyer. To the extent permitted at law, no warranties of merchantable quality or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Date of Manufacture:

Batch No:

PruneTec® is a registered trademark.

Registered to:
Max Stanton Moore
69B Rossall Street, Merivale, Christchurch, 8014
Ph: 021426528

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
12/12/2024

EMERGENCY RESPONSE: New Zealand National Poisons Centre Dial 0800 764 766
Transport Emergency Dial 111