WARNING
Keep out of reach of children
Ecotoxic

PUNCH UP™

FUNGICIDE

For the control of Botrytis in Grapes and Onions, Botrytis, Anthracnose and Rhizopus in Strawberries and Botrytis and Anthracnose in Blueberries

Active Ingredient: 375 g/kg CYPRODINIL and 250 g/kg FLUDIOXONIL in the form of a water dispersible granule

Net contents: 500g, 1 kg, 5 kg, 10 kg, 20 kg, 25 kg

READ THE LABEL COMPLETELY BEFORE USE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
22/03/2023

HAZARD CLASSIFICATIONS: Specific target organic toxicity repeated exposure Category 2, Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1, Hazardous to terrestrial vertebrates.

TOXICITY
May cause organ damage from repeated ingestion at high doses.

ECOTOXICITY
Very toxic to aquatic organisms and with long-lasting effects. Harmful to terrestrial vertebrates.
Do not apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

PRECAUTIONS
The person mixing or applying this product must be a “Qualified Contractor” or a “Qualified Person”. The spray mixture may also be prepared by a “Qualified Loader”.

RECORD KEEPING
Records of use, as described in the relevant HSNO regulations and the New Zealand Standard for the Management of Agrichemicals (NZS 8409), must be kept if using more than 3kg of Punch Up within 24 hours.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure place away from foodstuffs. Storage must be in accordance with the current version of the NZS 8409 Management of Agrichemicals. Stores containing more than 100 L require signage , secondary containment, and emergency response plan.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of dust or spray mist. When mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.
USE: Apply with well maintained and calibrated spray equipment.
DISPOSAL: Dispose of this product only by using in accordance with this label, or at a suitable landfill. Triple rinse container and add rinsate to the spray tank. Dispose of container to a suitable landfill or agricultural container recycling scheme. Do not burn. Do not use container for any other purpose.

First Aid
If medical advice is needed, have product container or label at hand. If swallowed do not induce vomiting. Rinse mouth. Get medical advice if you feel unwell. If splashed in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Get medical advice if irritation persists.
If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately for advice.

Emergency Response
In a transport emergency call **111, Police or Fire Brigade**. In a fire, evacuate and call **111 Fire Brigade.**

Spillage
Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. If granules have spilt, moisten to prevent dusting and sweep up. If liquid, absorb spill with inert material such as soil, sand, etc. Place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfill.

Safety Data Sheet
Additional information is provided in the safety data sheet.

GENERAL INFORMATION
Punch Up is a mixture of anilinopyrimidine and phenylpyrrole fungicides, which in combination have protectant and systemic activity against *Botrytis* (grey mould) infections in grapes and bulb onions, *Botrytis* (grey mould), *Colletotrichum* (anthracnose) and *Rhizopus* (leak) infections in strawberry, and Botrytis (grey mould) and Colletotrichum (anthracnose) in blueberries. Punch Up has multi-site action, inhibiting spore germination, penetration of the germ tube and mycelial growth. It is rainfast within 2 hours following application.

DIRECTIONS FOR USE

It is an offence to use this product on animals.

Crop	Disease	Rate	Remarks
Blueberries	Grey Mould (*Botrytis*), Anthracnose (*Colletotrichum*)	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	A maximum of 2 applications of Punch Up can be made per season. Apply in sufficient water to ensure thorough and uniform coverage without runoff (Approx. 1000 L/ha). The first application should be made at full flower, with the second application no later than 14 days after the completion of flowering.
Grapes	*Botrytis*	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	The rate suggested is for high volume spraying to runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage of the bunch zone is achieved. The addition of Synergy ™ Wetter Sticker at label rates will assist coverage. A maximum of 2 applications per season can be made. Punch Up can be applied at either early flowering (5% capfall), late flowering (80% capfall), pre bunch closure or in the period from the beginning of berry ripening up until 28 days before harvest. Apply an approved fungicide from a different chemical group for the remainder of the season.
Bulb Onions		800 g/ha	A maximum of 2 applications of Punch Up can be made per season at key timings for *Botrytis* control. Apply in sufficient water to ensure thorough and uniform coverage. Applications MUST be made with the addition of Synergy Wetter Sticker at label rates.
Strawberries	Grey Mould (*Botrytis*), Anthracnose (*Colletotrichum*) and leak (*Rhizopus*)	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	The rate suggested is for high volume spraying to runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved. The addition of Synergy Wetter Sticker at label rates will assist coverage. Punch Up can be applied at 7 to 14 day intervals from flowering onwards. The closer interval is recommended when conditions are conducive to disease infection. Punch Up applications may be applied in a block or in alternation with other non anilinopyrimidine fungicides registered for botrytis control. See resistance management for the maximum number of applications.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
22/03/2023

MIXING
Add the required amount of Punch Up into the mixing chamber of the partly filled spray tank and add the remainder of the water, ensuring complete mixing before spraying. Continue agitation during spraying.
COMPATIBILITY
Punch Up is compatible with other commonly used fungicides and insecticides.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Blueberries	Last application no later than 14 days after flowering and at least 28 days before harvest
Strawberries	3 days
Bulb onions	7 days
Grapes	28 days

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
22/03/2023

RESISTANCE MANAGEMENT

GROUP	9 + 12	FUNGICIDE

Punch Up contains a Group 9 fungicide (cyprodinil, belonging to the anilinopyrimidine class of chemistry) and Group 12 fungicide (fludioxonil, belonging to the phenylpyrrole class of chemistry). Resistance to this fungicide and related anilinopyrimidine products could develop in some situations from repeated use. To minimise this risk, use the product strictly in accordance with the label instructions.
Blueberries, Grapes and Bulb Onions: Do NOT use more than 2 applications of Punch Up or any other anilinopyrimidine fungicide on the same crop during the season.
Strawberries: A maximum of 3 applications of Punch Up or any other anilinopyrimidine fungicide per season can be made if more than six sprays in total are made for Botrytis control. If 6 or less Botrytis control sprays are made then do NOT apply more than 2 applications of Punch Up or any other anilinopyrimidine fungicide per season.
Consult http://resistance.nzpps.org/index.php for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P8964
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR100937
See www.epa.govt.nz for approval conditions.

™ PUNCH UP and SYNERGY are Trade Marks of Orion Agriscience Ltd

Registrant: Shandong Rainbow International Co. Ltd,
30th Floor, Building A5-5, Hanyu Financial Centre, No. 7000 Jingshi Road, Jinan Innovation Zone, Jinan, Shandong, China

NZ Address: 6 Horlicks Place, Randwick Park, Auckland 2105 Tel: (021) 02886306

Bar code

Batch No.	
Date of Manufacture	

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (CYPRODINIL 38%, FLUDIOXONIL 25%)
PACKING GROUP III
HAZCHEM CODE 2Z