WARNING

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

SULPHUR 800 WG

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/07/2025

FUNGICIDE/MITICIDE

ACTIVE INGREDIENT: 800 g/kg SULPHUR in the form of a water dispersible granule

GROUP M2 FUNGICIDE

For the control of certain diseases of Apples, Grapes, Stonefruit, Tamarillos, Vegetables and Ornamentals.

KENSO NZ
Together we grow

Registered pursuant to the ACVM Act 1997, No. P008992
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval No. HSR000741
See www.epa.govt.nz for approval controls

NET CONTENTS: 10 KG

Registrant:
Kenso Corporation (M) Sdn Bhd
Lot 29, Jalan Sungai Pinang 5/4, Taman Perindustrian Pulau Indah, 42920 Pelabuhan Klang, Selangor, Malaysia.

NZ Agent:
Kenso NZ Limited
2 Bond Crescent, Forrest Hill, Auckland 0620 New Zealand.
Ph: 0800 536 766

SULPHUR 800 WG FUNGICIDE

HAZARD CLASSIFICATION
Skin irritation Category 2, Designed for biocidal action

WARNING
Causes skin irritation.

HANDLING PRECAUTIONS
STORAGE
Keep out of reach of children. Store in the original packaging in a cool, dry, well ventilated place and protect from sunlight, away from human and animal foodstuffs, medicines and remedies, seeds and fertilisers. Storage of certain quantities of this product trigger specific storage requirements (Refer to SDS). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION
Avoid contact with eyes and skin. Avoid inhalation of dust and spray mist. When mixing or applying, wear chemical resistant: coveralls, footwear and gloves, and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

DISPOSAL
Dispose of this product only by using in accordance with this label, or at a suitable waste receival facility. Dispose of packaging to a suitable waste receival facilityl. Do not burn packaging. Do not use packaging for any other purpose.

SPILLAGE
Exclude non-essential people from the area. Wear protective clothing. Moisten to prevent dusting and sweep up. Place in a waste container. Dispose of safely to an approved waste or hazardous substances receival facility.

FIRST AID
If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor immediately for advice.

SAFETY DATA SHEET
A Safety Data Sheet is available from the Kenso NZ website: www.kenso.co.nz

CONDITIONS OF SALE
Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

FREE RECYCLING
AGRECOVERY
agrecovery.co.nz

EMERGENCY RESPONSE
Ring 0800 CHEMCALL, or
0800 243 622 (24 hours)

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/07/2025

Product of India

9 421028 660495

KENSO NZ
Together we grow

Batch No.:
Date of Manufacture:

WARNING

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

SULPHUR 800 WG

FUNGICIDE/MITICIDE

ACTIVE INGREDIENT: 800 g/kg SULPHUR in the form of a water dispersible granule

GROUP M2 FUNGICIDE

For the control of certain diseases of Apples, Grapes, Stonefruit, Tamarillos, Vegetables and Ornamentals.

Registered pursuant to the ACVM Act 1997, No. P008992
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval No. HSR00741
See www.epa.govt.nz for approval controls

Kenso Corporation (M) Sdn Bhd
2 Bond Crescent, Forrest Hill,
Auckland 0620 NZ
Phone: 0800 536 766

FREE RECYCLING
AgRecovery
agrecovery.co.nz

HAZARD CLASSIFICATION
Skin irritation Category 2, Designed for biocidal action

WARNING
Causes skin irritation.

HANDLING PRECAUTIONS
STORAGE
Keep out of reach of children. Store in the original packaging in a cool, dry, well ventilated place and protect from sunlight, away from human and animal foodstuffs, medicines and remedies, seeds and fertilisers. Storage of certain quantities of this product trigger specific storage requirements (Refer to SDS). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION
Avoid contact with eyes and skin. Avoid inhalation of dust and spray mist. When mixing or applying, wear chemical resistant: coveralls, footwear and gloves, and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT
Apply with well-maintained and calibrated spray equipment.

CONDITION OF SALE
Kenso Corporation (M) Sdn. Bhd. will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Kenso Corporation (M) Sdn. Bhd. is limited to the replacement of the goods or (at the option of Kenso Corporation (M) Sdn. Bhd.) the refund of the price paid and where possible sufficient part of the goods to enable proper examination being returned to Kenso Corporation (M) Sdn. Bhd. within thirty days of delivery.

Registered pursuant to the ACVM Act 1997, No. P008992
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996, Approval No. HSR000741
See www.epa.govt.nz for approval controls

Batch No. :

Date of Manufacture :

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/07/2025

DISPOSAL
Dispose of this product only by using in accordance with this label, or at a suitable waste receival facility. Dispose of packaging to a suitable waste receival facility. Do not burn packaging. Do not use packaging for any other purpose.

ENVIRONMENTAL
Use this product carefully. Do not contaminate aquatic environments with product, spray drift or packaging. Do not allow spray drift outside the target area to occur.

SPILLAGE
Exclude non-essential people from the area. Wear protective clothing. Moisten to prevent dusting and sweep up. Place in a waste container. Dispose of safely to an approved waste or hazardous substances receival facility.

GENERAL INFORMATION
Sulphur 800 WG Fungicide/Miticide is a protectant fungicide and must be applied before disease infection occurs. For best results, even spray coverage is necessary.

CAUTION
Some fruits and crops are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use on any crop unless local experience has proven sulphur to be safe in that locality. Do NOT apply Sulphur 800 WG Fungicide/Miticide when the temperature is above 25°C, or under conditions of extreme humidity, as crop damage can occur.
Cucurbits and other crops grown under glass can be affected by an application of Sulphur 800 WG Fungicide/Miticide during periods of rapid growth.

Do NOT apply to sulphur shy apples such as Cox's Orange.
Do NOT apply to apricots.
Do NOT apply to pears.
Do NOT apply to apples with Captan or within 5 days of a Captan treatment.
Do NOT apply with crop oil or within 21 days of a crop oil application.

APPLICATION METHODS
Suitable for use with all applicators. The rates expressed are for high volume spraying to run off. For concentrated spraying, adjust dilution rate accordingly. Sulphur is corrosive so all spray equipment should be thoroughly flushed with clean water after use.

COMPATIBILITY
Sulphur 800 WG Fungicide/Miticide is compatible with most insecticides, fungicides and foliar fertilisers. When using with untried emulsifiable concentrate formulations, test the compatibility of both products by mixing small quantities before final mixing. Do NOT use with spraying oils or any products containing mineral oil.

RAINFALL
Rain within 3 hours of treatment may reduce efficacy.

It is an offence to use to use this product on animals.
DIRECTIONS FOR USE

Crop	Pest /Disease	Rate	Remarks
Apples	Powdery mildew	100-200 g per 100 litres	Apply from bud movement until harvest, however, use before December may affect fruit finish. DO NOT apply in temperatures above 25°C. DO NOT apply in combination with captan. DO NOT apply to sulphur sensitive varieties such as Kidds Orange Red, Cox's Orange Pippin or Red Delicious. Fruit finish may be affected on some apple varieties. DO NOT apply on pears.
Grapes	Powdery mildew	150-300 g per 100 litres	Apply at 10 to 14 day intervals from bud swell until 6 weeks before harvest. Use the higher rates when disease is evident. Erineum mites will also be controlled.
Peas	Powdery mildew	100-200 g per 100 litres	Apply when conditions favour disease and repeat as necessary. Use the higher rates where infection has already occurred.
Peaches Plums Nectarines	Rust	100-200 g per 100 litres	Apply at fruit set then at 21 day intervals until harvest. Use the higher rate during conditions of high pressure. DO NOT apply to apricot.
Roses	Powdery mildew	200-300 g per 100 litres	Apply on a 14 day schedule using the higher rate during periods of severe disease pressure. DO NOT use in mixture with maldison.
Tamarillos	Powdery mildew	100-200 g per 100 litres	Apply at 10 to 14 day intervals throughout the growing season.
Vegetables	Powdery mildew	100-300 g per 100 litres or 2-3kg/ha	Use at regular 10 - 14 day intervals during periods of infection. Use the higher rate where infection has already occurred.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/07/2025

WITHHOLDING PERIODS
Nil.

RESISTANCE MANAGEMENT GROUP **M2** FUNGICIDE

Sulphur 800 WG Fungicide is an inorganic (M2) group of fungicides. Some naturally occurring individual fungi may become resistant when excessive use of the same fungicide occurs. To minimize this risk use strictly in accordance with the label instructions and resistance management strategies. Consult http://resistance.nzpps.org/ for alternate modes of action and the details of resistance management strategies (if any) for the crops listed on the label.

FIRST AID
If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor immediately for advice.

SAFETY DATA SHEET
A Safety Data Sheet is available from the Kenso NZ website: www.kenso.co.nz