DANGER
KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE

ChampION++ ®

Active Ingredient: 300g/kg copper as copper hydroxide in the form of a water dispersible granule.

A protective copper fungicide/bactericide for the control of many diseases of fruit, vegetables and forestry.

QR Code

Net: 10 kg

Barcode: 9414614257810

Label codes: 04-23//PS/02578-10KNZ

Nufarm

BioGro logo

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
05/07/2023

ChampION++ ®

GHS CLASSIFICATION:

Acute oral toxicity Category 4, Serious eye damage Category 1, Skin sensitisation Category 1, Specific target organ toxicity (repeated exposure) Category 2,
Hazardous to terrestrial vertebrates,
Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1

This product must not be used for any purpose, or in any manner, contrary to this label unless authorized under appropriate legislation.

DANGER

- Harmful if swallowed.
- May cause an allergic skin reaction.
- May cause damage to lungs through prolonged or repeated exposure.
- Causes serious eye damage.
- Very toxic to aquatic life with long lasting effects. Avoid contamination of any water supply with product or empty container. Do not apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.
- Harmful to terrestrial vertebrates

HANDLING PRECAUTIONS

The person mixing or applying this product must be a “Qualified Contractor” or a “Qualified Person”. The spray mixture may also be prepared by a “Qualified Loader”.

STORAGE:

- Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs. Store in accordance with NZS8409 Management of Agrichemicals.
- Stores containing 100kg ChampION^{++} are subject to signage and more than 100kg require emergency response plans.

PERSONAL PROTECTION:

- Do not get in eyes. Avoid skin contact and inhalation of spray mist or dust.
- Do not eat, drink or smoke while using this product.
- When mixing or applying wear protective clothing such as impervious gloves, eye and face protection, waterproof hat, coat, trousers and boots with socks.
- Remove protective clothing and wash hands and face thoroughly before meals and after work.
- Wash contaminated clothing before reuse.
- Contaminated clothing should not be allowed to leave the workplace.

EQUIPMENT:

- Apply using well maintained and accurately calibrated spray equipment. Clean equipment thoroughly after use.

DISPOSAL:

- Ensure container is completely empty. Burn if circumstances, especially wind direction permit. Otherwise crush and bury in a suitable landfill.
- Dispose of product only by using according to the label or at an approved treatment facility.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

05/07/2023

First Aid: If medical advice is needed, have product container or label at hand.
IF SWALLOWED: do NOT induce vomiting. Rinse mouth. Call the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell. **IF ON SKIN:** wash with plenty of soap and water. If skin irritation or rash occurs: get medical advice / attention. **IF INHALED:** move victim to fresh air immediately. If not breathing, give artificial respiration and seek medical attention. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. Get medical advice if you feel unwell.

Spillage: Collect spillage. Wear appropriate protective clothing and prevent material from entering waterways. Sweep up solids without generating dust. Contain liquid spills and absorb with inert material. Dispose of waste safely in an approved landfill.

Transport: Do not carry ChampION^{++} on a passenger service vehicle.

Record Keeping: Records of use as described in NZS8409 Management of Agrichemicals, must be kept if 3kg or more of ChampION^{++} is applied within 24 hours. Records must be kept for a minimum of 3 years from application date.

Mixing: Add two thirds of the required quantity of water to the spray tank, and with the agitation system going, slowly add the ChampION^{++}. Continue agitation while adding the remainder of the water. The addition of insecticides, spreader-stickers etc should be made last, after the ChampION has been thoroughly mixed. Agitate thoroughly before and during spraying and before recommencing after a stoppage.

Compatibility: ChampION^{++} is compatible with most commonly used fungicides and insecticides. Do not mix with Attack® for use on glasshouse crops. Do not mix with Diazinon or with strongly alkaline or acidic materials. If compatibility is unknown conduct a compatibility jar test using proposed use rates before mixing and applying a whole tank. Do not apply this product in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.

Resistance Management: ChampION^{++} is a member of the inorganic copper group of fungicides with multi-site activity. ChampION^{++} is a GROUP M1 FUNGICIDE Some naturally occurring fungi and bacteria resistant to ChampION^{++} and other Group M1 fungicides may exist through normal genetic variability in any population. These will not be controlled by ChampION^{++} or other Group M1 fungicides. The resistant individuals can eventually dominate the population if copper products are used repeatedly. Since the occurrence of resistance is difficult to detect prior to use, Nufarm Limited accepts no liability for any losses that may result from the failure of ChampION^{++} to control resistant fungi or bacteria. To prevent or delay the development of resistance use ChampION^{++} in tank mixtures where appropriate, and/or in rotation with fungicides or bacteriocides having a different mode of action. Refer to www.resistance.nzpps.org/ for further information.

Directions for use:

It is an offence to use this product on animals.

ChampION^{++} is a form of cupric hydroxide with extremely small particle and crystallite size. It is a free flowing and easy to pour formulation which goes easily into suspension to give a spray with excellent tenacity on plant surfaces. ChampION^{++} is highly active against a range of plant diseases.

ChampION^{++} may be applied through standard ground and aerial spray equipment. The recommended dilution rates in the table below are for high volume spraying to runoff. The volume of water applied per hectare will depend on the crop being sprayed and where necessary should increase during the season to ensure complete coverage of all stages of the crop. For concentrate spraying apply the same volume of ChampION^{++} per hectare as for dilute sprays.

In conditions of severe disease pressure use the higher recommended rate and reduce the spray interval.

Crop Caution: Soft new growth of some plant species may be hardened or injured by copper sprays.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

05/07/2023

CROP	DISEASE	RATE		COMMENTS
		Per 100L Water	Per Hectare	
FRUIT CROPS				
Apples Pears (including Nashi)	Black spot	110g	2.2kg minimum	Apply as a full cover spray. Make application between silver tip and green tip. **CAUTION:** Late application may cause phytotoxicity. Discontinue use when green tip is 1cm.
Avocados	Anthracnose	70-90g	2kg minimum	Apply at monthly intervals from flowering to harvest. During prolonged wet weather reduce application intervals to 14 days.
Bushfruit	Cane wilt Leaf spot Cane spot Spur blight	70-90g	0.65kg minimum	Apply at bud burst and green tip and continue at 10-14 day intervals
Citrus	Melanose Verrucosis Brown rot	45-90g 90-110g	1.75kg (mature trees) 1.75kg (mature trees)	Apply during pre and post bloom periods. Apply in the autumn and continue as necessary. Spray the skirts of trees to at least 1 metre high and also treat the ground out to 0.5metres beyond the skirt. Copper marking may occur on sensitive varieties or under slow drying conditions.
Feijoas	Leaf spot *(Glomerella sphaceloma)*	79-90g	1kg minimum	Apply and repeat as necessary.
Grapes	Black spot Downy mildew	65-130g	0.9kg minimum	Begin at budbreak and continue during the pre and post bloom periods at 10-14 day intervals. Discontinue use if signs of injury appear.
Kiwifruit	Leaf spot (*Phomopsis* sp & others). *Botryosphaeria*	70-90g	1.3kg minimum	Apply as a dormant season spray only. Apply after harvest but before leaf fall, after pruning and as two sprays one month apart before bud burst.
	Psa (*Pseudomonas syringae* pathovar *actinidiae*). MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND 05/07/2023	70-90g		Apply after harvest but before leaf fall, after pruning and as two sprays one month apart before budburst. Pre- and Post-flowering. DO NOT apply to open flowers. Apply as a protectant treatment prior to a significant rain (infection) event or immediately after when the crop is dry. Use a full dilute water rate, appropriate for the size of canopy, to give complete coverage. WARNING: Higher rates, multiple applications, poor drying conditions and/or tank mixtures may result in leaf drop, leaf marking and marginal leaf burn. Fruit marking may occur from any application post flowering. Apply alone post-flowering to minimize the risk of crop injury.
Passionfruit	Septoria spot Brown spot Grease spot	45g	0.44kg minimum	Apply at 2-4 week intervals alternating with mancozeb from November to March. Where grease spot is a problem 2 week intervals may be necessary.

CROP	DISEASE	RATE		COMMENTS
		Per 100L Water	Per Hectare	
Pears (including Nashi)	Fire blight *(Erwinia amylovora)*	25-45g	1.4kg minimum	***Bearing Trees:*** Make one application between the silver leaf and green tip period. On smooth skinned varieties do not apply between green tip and harvest as russeting may result. ***Non Bearing Trees:*** Maintain a protective cover of regular sprays at 14-21 day intervals until active vegetative growth ceases. Reduce interval between sprays if wet weather prevails. ***Post Harvest:*** Apply at 21 day intervals from harvest to leaf fall. Reduce interval between sprays if wet weather occurs and vegetative growth is active.
Stonefruit	Leaf curl Shot hole Bladder plum	110g	2.2kg minimum	Apply at bud movement but before and within one week of buds opening. Make a second application at 70-90g/100 litres 10-14 days later.
	Bacterial blast	312g	6.25kg minimum	Apply at leaf fall and repeat at monthly intervals throughout the winter until bud movement, then treat as for leaf curl.
Strawberries	Leaf spot	70-90g	0.65kg minimum	Apply when plants are established and continue on a weekly schedule throughout the season. Use the higher rate early season or under high disease pressure. Discontinue use if signs of crop injury appear.
Tamarillos	Leaf spot	70g	0.9kg minimum	Apply as necessary.
VEGETABLES				
Asparagus	Stemphylium leaf spot	250-375g	2.5kg minimum	Commence spraying at first appearance of disease after closing up and repeat at 14 day intervals to maintain disease control. Ensure good plant coverage. Under high disease pressure a programme of up to 6 applications may be necessary.
Beans	Bacterial blight (Halo & common)	250-310g	1.36kg minimum	For protective sprays apply at 7-14 day intervals from when plants are 15cm high.
Brassicas: Cabbage Cauliflower Broccoli Brussels sprouts	Downy mildew	70-90g	0.3kg minimum	Apply after crop emergence and/or transplanting at 10-14 day intervals as necessary.
Celery	Septoria leaf spot	90g	0.44kg minimum	Apply as soon as plants are established in the field then at 5-7 day intervals. An approved mineral crop oil may be added at 1%.
Cucurbits	Angular leaf spot Downy mildew	70-130g	0.65kg minimum	Apply at 7 day intervals after plants have started to vine.
Lettuce	Downy mildew	70-110g	0.33kg minimum	Begin treatment from crop emergence and repeat every 7-10 days as required to suppress the disease.
Onions	Downy mildew Bacterial blight	90g	1kg minimum	Apply on first appearance of disease and repeat every 10-14 days while conditions favour disease development.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
05/07/2023

CROP	DISEASE	RATE		COMMENTS
		Per 100L Water	Per Hectare	
Peas	Downy mildew Leaf spot	70-90g	0.66kg minimum	Begin treatment from crop emergence and repeat at 7-14 day intervals depending on disease severity.
Potatoes	Early blight Late blight	130-200g	1.3kg minimum	Apply every 7-10 days from when plants are 15cm high until 2 weeks before harvest.
Tomatoes	Early blight Late blight Bacterial speck Bacterial spot	70-130g	0.8kg increasing to 1.75kg	Apply at 7-10 day intervals. Reduce spray interval if conditions are favourable for development of the disease. Ensure spray volumes of 1500 litre/hectare as crop reaches maximum growth and ensure full coverage of upper and lower leaf surfaces. Repeat application after more than 13mm rain or irrigation.
FORESTRY				
Pine Trees	Pine needle blight *(Dothistroma pini)*		2.5- 4.4kg	**Aerial application to plantations:** Apply in 60-100 litres of water per hectare. Make one application in November or early December. Repeat at yearly intervals if necessary.
			1.25-3.1 kg	**Ground operated equipment in nurseries:** Apply with 1% summer oil in 600-1100 litres of water. Commence spraying in October or November and repeat at monthly intervals until planting out.

Frost Injury Protection: Bacterial Ice Nucleation Inhibitor: Application of ChampION^{++} made to the crops listed on this label, at the rates and stages of growth indicated, just prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae, Erwinia herbicola* and *Pseudomonas fluorescens*) and may therefore provide some protection against light frost. Do not rely solely on ChampION^{++} for frost protection, especially where weather conditions favour severe frost.

Registered pursuant to the ACVM Act 1997, No P9228
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code HSR000739.
See www.epa.govt.nz for approval controls.

® ChampION++ is a registered trademark of Nufarm Americas Inc.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

05/07/2023

CONDITIONS OF SALE:

CAUTION: READ THIS NOTICE BEFORE OPENING THE CONTAINER

If you open this container you will be deemed to have accepted the conditions below.

The results from use of this product may be affected by factors beyond Nufarm's control, including mixing, use, climatic conditions, time of application, crop or crop stage and the possible development of resistance to the active ingredient, even though the product is reasonably fit for the purposes stated in this label, and even though label directions are followed. Nufarm will replace, free of charge, any product which is found to have a manufacturing defect. To the extent permitted by law, all other warranties, guarantees, representations and conditions, whether expressed or implied by law, trade, custom or otherwise are expressly excluded.
When you use this product, you agree that neither Nufarm Limited nor any manufacturer of any component of the product will be liable for any damage, injury or loss including consequential loss which arises in any way from the storage, handling, or use of the product. This exclusion of liability includes loss caused by negligent acts or omissions of Nufarm Limited or any supplier or manufacturer of the product or any of their employees and agents.

UN 3077
ENVIRONMENTALLY
HAZARDOUS SUBSTANCE,
SOLID, N.O.S. (copper)
Class 9, PG III, Marine Pollutant.

Marketed by:

Nufarm Limited
Central Park
Building 3, Level 4
666 Great South Road
Ellerslie
Auckland, 1051 NZ.
Telephone 0-9-270 4157
www.nufarm.co.nz

Nufarm emergency phonelink: 0800-651 911 toll free 24 hours

Batch No.:
D.O.M.: