WARNING

KEEP OUT OF REACH OF CHILDREN

K-POW ®

FUNGICIDE AND ADJUVANT

ACTIVE INGREDIENT: K-POW contains: 200g/L POTASSIUM BICARBONATE (bicarbonate and carbonate) and 130g/L FATTY ACIDS (potassium salts) in the form of a soluble concentrate.

For the control of Powdery Mildew on Grapes and to improve coverage of fungicides on grapes.

GROUP	NC	FUNGICIDE

AGRECOVERY
Rural Recycling Programme

NET CONTENTS 20, **100, 200, 1000 litre**s (**Bold** = Main Commercial pack sizes)

Registered pursuant to the ACVM Act 1997. No.P009371
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996. *Approval Code* ***HSR101188***

CERTIFIED biogro NEW ZEALAND INPUT for ORGANICS
4429

Registered to: ***UPL New Zealand Limited***
PO Box 51584
Pakuranga 2140
Phone: 0800 100 325
www.upl-ltd.com/nz

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL **(0800 243 622)**

® Registered Trade Mark of UPL New Zealand Limited

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

01/03/2021

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.

HSNO Classification: 6.3B, 6.4A, 9.3C
WARNING
May cause skin and eye irritation.
Harmful to terrestrial vertebrates.

PRECAUTIONS:
Store in original container, tightly closed, away from foodstuffs and under lock and key. Avoid contact with seeds, fertilisers and other pesticides.
Wear overalls, gloves and goggles while mixing and spraying.
Avoid contact with skin and eyes, and inhalation of spray mist.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking and after work.
Wash protective clothing after use.
Apply with well maintained and calibrated spray equipment.
Avoid contamination of any water supply with chemical or empty container.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.

CONTAINER DISPOSAL: 20 litre HDPE: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. Avoid contamination of any water supply with chemical or empty container.
200 L & 1000 L Tote Tanks: Triple rinse with clean water. Drain rinsate onto waist ground eg under hedges. Do not allow rinsate to enter any waterways. Once cleaned, container may be reused.
PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID
If swallowed DO NOT induce vomiting. For advice, contact the National poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If splashed in eyes, wash out immediately with water. If inhaled move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SPILLAGE
Wear impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water.

GENERAL INFORMATION
K-POW is a potassium bicarbonate based fungicide and surfactant that is active against Powdery Mildew of Grapes. Control is enhanced by the addition of copper and/or sulphur products.

MIXING
K-POW is formulated to readily mix with water. Simply measure the required amount and add to the partly filled spray tank with agitation operating. Add K-POW after other products added have been fully dispersed. Complete filling and maintain agitation until spraying is complete

COMPATABILITY
K-POW is compatible with copper hydroxide and cuprous oxide based copper fungicides and with wettable sulphur. Add K-POW to the spray tank only after other products have been added and are thoroughly mixed and diluted. For other products before using, check compatibility of the mixture with a jar test and observe all label

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

01/03/2021

precautions. The addition of wetting or sticking agents is **not** recommended. **DO NOT USE if any compatibility issues are noted.**

WATER HARDNESS
K-POW is a highly alkaline product. Caution is required when mixing in hard water, water with high levels of solubilized minerals present, and water with high pH. Conduct testing before commercial use. Precipitation of the product in the spray tank and mineral deposits on fruit can occur.

DIRECTIONS FOR USE
Apply K-POW in mixture with Sulphur for preventative control of powdery mildew in the period from bud burst to veraison. Apply in mixture with a copper spray when used to eradicate any early established stage of powdery mildew infection later in the season.

CAUTION
DO NOT use during hot periods of the day. Apply morning or evening and on days when temperatures are cooler. DO NOT apply to red grapes after veraison. Maintain good agitation while spraying.

Crop	Target	Use Rate	Comments
Wine Grapes	**Powdery Mildew**	Apply 1.65 litres **K-POW** / 100 litres of water in mixture with a copper and/or sulphur product.	Apply in sufficient water to obtain full coverage of vines to the point of run off. Higher water rates will generally give better coverage. *K-Pow can be applied at any growth stage throughout the season at 10-14 day application intervals until 28 days before harvest, during the high risk period, Flowering until Veraison, keep application intervals close (7-10 days).* If heavy rainfall occurs, replace cover sprays as soon as possible. For powdery mildew control apply preventatively when conditions favour infection or as an eradicant at an early stage of infection and before powdery mildew becomes well established in the crop with the addition of sulphur and or copper. Follow label directions for the copper and sulphur products.

WITHHOLDING PERIOD: 28 days
It is an offence to use this product on animals

STORAGE
Store in a cool dry place avoiding temperatures above 35°C.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at **www.upl-ltd.com/nz**

EXCLUSION OF LIABILITY:
Please read these conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened.
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND

Batch No.
DOM

01/03/2021

® - Registered Trademark of UPL New Zealand Limited

COPY RIGHT:
© Copyright UPL New Zealand Limited (2016), all rights reserved. All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND

01/03/2021