December 2024

Front Label

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
13/01/2025

DANGER
Keep out of reach of children

GA200

For non-selective weed control against a wide range of grasses, broad-leaf weeds and clovers.

ACTIVE INGREDIENT:
Contains 200g/litre glufosinate-ammonium in the form of a soluble concentrate.

Read the directions for use before using this product.

Contains *X* Litres

Date of Manufacture:

Batch No:

December 2024

Back Label

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
13/01/2025

READ THE LABEL AND LEAFLET COMPLETELY BEFORE MIXING OR APPLYING

HAZARD CLASSIFICATIONS:
Eye irritation Category 2, Reproductive toxicity Category 2, Specific target organ toxicity (repeated exposure) Category 1, Aquatic toxicity (chronic) Category 3, Hazardous to soil organisms, Hazardous to terrestrial vertebrates.

TOXICITY
Causes serious eye irritation. Causes damage to organs from repeated oral exposure at high doses. Suspected of damaging fertility or the unborn child.

ECOTOXICITY:
Harmful to aquatic life with long lasting effects. Toxic to the soil environment. Harmful to terrestrial vertebrates.
NON-SELECTIVE HERBICIDE: Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Do not allow drift to occur outside the target area.

QUALIFIED HANDLER:
This product must be under the control of an suitably trained handler when it is applied in a wide dispersive manner or used by a commercial contractor.
This product must not be used for any purpose or in any manner contrary to this label unless authorised under appropriate legislation.

PRECAUTIONS
Obtain special instructions before use. Do not handle until all safety precautions have been read and understood.
PERSONAL PROTECTION: Avoid contact with skin, eyes and inhalation of spray mist.
When handling, wear eye protection, waterproof gloves, boots and overalls. Remove protective clothing and wash hands and face thoroughly before meals and after work.
Do not eat, drink or smoke while using. Clean protective clothing and equipment before re-use.

STORAGE:
Keep out of reach of children. Store in accordance with the current version of NZS 8409 Management of Agrichemicals. Store in the original container away from sources of ignition in a locked, dry, cool area away from foodstuffs, fertilisers and seeds. Avoid storage at temperatures below 5°C and above 35°C.

SPILLAGE: Collect spillage. Wear appropriate protective clothing and prevent material from entering waterways. Exclude non-essential people from the area. Contain liquid spills and absorb with inert material such as soil or sand and place in waste containers. Dispose of waste safely in an approved landfill.

DISPOSAL: Avoid contamination of any water supply with chemical or empty container.
Follow all local, regional and national laws and regulations regarding hazardous waste disposal.

Product Disposal: Dispose of this product only by using according to the label directions or if this is not possible alternatively contact Rainbow & Brown for advice. Do not dispose of this product down drains or sewers.
Container Disposal: Triple rinse empty containers and add residue to spray tank. Do not burn containers. Crush or puncture and bury in a suitably approved landfill or through Agrecovery recycling service.

FIRST AID: If medical advice is needed, have product container or label at hand. **Call a POISON CENTRE, TEL 0800 POISON (0800 764766) or a doctor.**
If exposed or concerned, get medical advice.
If swallowed do NOT induce vomiting. Rinse mouth.
If on skin wash with soap and water.
In in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention.
If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult and get medical assistance.

TRANSPORT: Do not carry this product on a passenger service vehicle.

RECORD KEEPING: Records of use required as described in the current NZS8409 Management of Agrichemicals.

MIXING: Add two thirds of the required quantity of water to the spray tank. With the agitation system going, add the product to the spray tank. Agitate for 3-5 minutes while adding remaining water.

COMPATIBILITY: GA200 is compatible with most commonly used residual herbicides.

GENERAL COMMENTS: GA200 is absorbed into the foliage and green stems of actively growing plants. It is inactive in soil and is not translocated as an active ingredient, so thorough coverage of target weeds is essential to obtain optimal results.
Visual effects of treatment with GA200 occur within 3-10 days. The effect will occur more rapidly under ideal growing conditions.
Rainfall within 2 hours of application may wash the product off the foliage. Rain within 6 hours will reduce efficacy to the point where it may be necessary to reapply the product.
Note: Do not spray in temperatures below 10°C. Avoid contact with green plant parts (including immature bark) of desirable plants as damage may occur. Do not mow or cultivate after treatment until weeds start to die back.

WITHHOLDING PERIOD: 28 days on all crops.

Registered pursuant to the ACVM Act 1997, No. P009422.
See www.foodsafety.govt.nz for conditions of registration.

Approved pursuant to the HSNO Act 1996, EPA Approval Code HSR007813
See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
13/01/2025

December 2024

CONDITIONS OF SALE:
This product is warranted by Rainbow & Brown Limited to conform to the chemical description and concentration specified on the label, and to be reasonably fit for the purposes stated in the directions if used in accordance with the directions. The user bears all risks of damage or failure resulting from factors beyond the control of Rainbow & Brown. Rainbow & Brown declines all liability for any damages resulting from the improper use of this product.

[R&B LOGO]
Registrant:
Rainbow & Brown Ltd
68A Tallyho Street, Rotorua 3015
PO Box 100 49, Mail Centre, Rotorua 3046
Freecall (0508) 299 299
www.RainbowBrown.co.nz

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
13/01/2025

December 2024

Leaflet

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
13/01/2025

DIRECTION FOR USE: RATES OF APPLICATION

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

USE	DOSAGE RATE	REMARKS
Berryfruit - Raspberries and Brambles Primocane control	500-750mL/100 litres applied through cone nozzles in sufficient water to ensure complete coverage	Apply as a directed high-volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of a non-ionic wetter at label rates is recommended. Knockdown takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached a height of 15 cm. Do not apply to plants under stress.
Orchard and vineyard weed control- Pipfruit, kiwifruit, stonefruit, grapes and citrus.	Use a dosage rate suitable for the weeds present. If the dilution rate is less than 1 litre /100 litres of water, add a non-ionic wetter at label rates	Apply in spring when the majority of the annual weeds have germinated. Mix with residual herbicides for long-term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water for optimum coverage. May be applied up to one day before crop emergence.
Non-crop land. General weed control.	7.5-10 litres/ha in 300-500 litres of water or 100mL/10 litre knapsack.	Apply as a drenching spray to the point of run-off. Mix with residual herbicides for long-term weed control.

WEED SUSCEPTIBILITY

The lower rates recommended are appropriate for young succulent actively growing weeds or when the weed population is sparse and good coverage and penetration is possible. Use the higher dosage rates for mature weeds or where the weed population is dense.

Note: GA200 will not completely control mature weeds with well-developed taproots or underground storage organs. Some regrowth may occur following treatment.

RESISTANCE MANAGEMENT

GA200 contains glufosinate-ammonium, a phosphinic acid herbicide with the inhibition of glutamine synthetase mode of action . Certain individual weed biotypes may become resistant. To minimise the

risk of these weeds dominating the population, use strictly in accordance with the label instructions. GA200 should be used in a programme that incorporates effective herbicide products with different modes of action. Refer to the web site www.resistance.nzpps.org > for more detailed information.

GROUP	10	HERBICIDE

Dosage Range of major weeds:

3-5 litres/ha			
Annual poa	Clover seedlings	Rough-stalked meadow grass	Subterranean clover
Asparagus seedling	Cornbind	Scotch thistle	Wild turnip
Beaked hawksbeard	Daisy	Sow thistle	Willow weed
Black nightshade	Dandelion	Staggerweed	Wireweed
Calandrina	Mouse-ear chickweed	Summer grass	
Chickweed	Musky storksbill	Spurrey	
Cleavers	Redroot	Stinking mayweed	

5-7.5 litres/ha			
Bristle grass	Clovers	Parsley piert	Storksbill
Bur medic	Cut-leaved geranium	Prairie grass	Strawberry
Catsear	Docks	Ryegrasses*	Twin cress
Chewings fescue	Fleabane	Sheeps sorrel	Yorkshire fog

7.5-10 litres/ha			
Broom	Crowfoot grass	Mallows	Wild strawberry
Buttercup's	Gorse	Paspalum	Willow herb
Couch	Hawksbeard	Tall fescue	Yarrow
Field pansy	Kikuyu grass		

10 litres plus/ha			
Bracken	Plantains	Rushes	Indian doab
Meadow rice grass	Rape	Sedges	

*Resistant weeds: Hairy buttercup and some ryegrass strains

Conditions of Sale

This product is warranted by Rainbow & Brown Limited to conform to the chemical description and concentration specified on the label, and to be reasonably fit for the purposes stated in the directions if used in accordance with the directions. The user bears all risks of damage or failure resulting from factors beyond the control of Rainbow & Brown. Rainbow & Brown declines all liability for any damages resulting from the improper use of this product.

[R&B LOGO]
Rainbow & Brown Ltd
PO Box 10049, 68A Tallyho St, Rotorua
Phone Toll Free: (0508) 299 299 email: mail@rainbowbrown.co.nz

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
13/01/2025