WARNING

KEEP OUT OF REACH OF CHILDREN

Proclaim® Opti

Insecticide

syngenta®

Active Ingredient:
PROCLAIM® Opti contains 50 g/kg EMAMECTIN BENZOATE in the form of a water dispersible granule.

GROUP	6	INSECTICIDE

For the control of Leafrollers and Codling Moth on Pipfruit and Leafrollers on Avocados, Grapes and Kiwifruit

IMPORTANT: READ THIS BOOKLET BEFORE USING THIS PRODUCT.

Net Contents: 80g, 120g, 160 g, 400g, 500g and 600 g

Registered pursuant to the ACVM Act 1997, No. P9526.
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code HSR101328.
See www.epa.govt.nz for approval controls.

Syngenta Crop Protection Limited
Level 4, 60 Parnell Road, Parnell, Auckland 1052

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
25/02/2026

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.
Any person mixing, loading or handling this product MUST be a qualified loader or under the guidance of a qualified loader.
Any person applying this product MUST be a qualified contractor/person or under the guidance of a qualified contractor/person.

GHS Classifications
Acute oral and inhalation toxicity, category 4
Specific target organ toxicity (repeated exposure), category 2
Hazardous to the aquatic environment, acute and chronic, category 1
Hazardous to terrestrial vertebrates
Hazardous to terrestrial invertebrates

WARNING
Harmful if swallowed or inhaled.
May cause neurotoxic damage through prolonged or repeated oral exposure at high doses.
Very toxic to aquatic life with long lasting effects. Higher risk of toxic effects in estuarine environments.
Harmful to terrestrial vertebrates.
Very toxic to terrestrial invertebrates.

PRECAUTIONS
Store in original container, tightly closed, away from foodstuffs.
Use only outdoors or in a well-ventilated area.
When mixing, loading and / or applying wear protective clothing, including long sleeve shirt, long trousers and closed shoes, chemical resistant gloves, face shield or goggles.
DO NOT breathe dust or spray. Avoid contact with skin and eyes.
If clothing becomes contaminated with product, remove clothing immediately.
DO NOT eat, drink or smoke while handling the product or using.
Remove protective clothing and wash hands and face thoroughly before eating, drinking or smoking and after work.
Wash protective clothing separately from other clothing, daily after use and before reuse.
Apply with well maintained and calibrated spray equipment.
Records of use MUST be kept as per Clause 48 of the Hazardous Property Controls Notice 2017.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
DO NOT carry on a passenger service vehicle.
Avoid contamination of any water supply with chemical or empty container. DO NOT apply into or onto water.
Direct contamination or spray drift to any aquatic environment MUST be strictly avoided.

Container disposal: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.
For additional information refer to the EPA Notices: Hazardous Substances (Disposal) Notice 2017.

FIRST AID
If SWALLOWED, DO NOT induce vomiting. Rinse Mouth.
If SKIN or HAIR contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin.
If INHALED, move the victim to fresh air and keep at rest in a position comfortable for breathing.
For advice or if you feel unwell, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.
If medical advice is needed, have product container or label at hand.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
25/02/2026

SPILLAGE
Collect spillage. Wear suitable protective clothing including face shield, chemical resistant gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

GENERAL INFORMATION
PROCLAIM® Opti has translaminar activity but is not systemic, therefore thorough coverage is essential. After application, PROCLAIM® Opti quickly moves into young leaves from where it is taken up by the feeding Leafrollers. PROCLAIM® Opti takes about 4 days to achieve maximum control. When PROCLAIM® Opti comes into contact with soil it readily and tightly binds to the soil. PROCLAIM® Opti that is not absorbed into plants is quickly degraded.
Note: DO NOT apply if rainfall is expected before spray would have dried, as reduced efficacy may result. The addition of a non-ionic surfactant is required in all uses. A reduction in efficacy may result if a non-ionic surfactant at label rates is not included.
Crop Monitoring: For effective Leafroller and Codling Moth control, regularly monitor crops during the season.

RESISTANCE MANAGEMENT **GROUP 6 INSECTICIDE**
PROCLAIM® Opti Insecticide is a member of the avermectin group of insecticides. PROCLAIM® Opti is a Group 6 insecticide. Resistance to this insecticide could develop from repeated use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies. Since the occurrence of resistant insects is difficult to detect prior to use, Syngenta Crop Protection Ltd accepts no liability for any losses that may result from the failure of PROCLAIM® Opti to control resistant insects. Consult your local supplier, consultant, a Syngenta representative or New Zealand Plant Protection Society (Inc), www.resistance.nzpps.org for alternate modes of action and the details of resistance management strategies for the crops listed on the label.
Avoid using this pesticide exclusively all season, and avoid unnecessary spraying. Maintain good cultural control practices.

INTEGRATED PEST MANAGEMENT
PROCLAIM® Opti is compatible with IPM programmes using the predator mites *Typhlodromus pyri*, *Phytoseiulus persimilis* and *T.occidentalis.*

DIRECTIONS FOR USE
It is an offence to use this product on animals.

The application rate is for high volume spraying to the point of runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved.

Set up and operate the sprayer to achieve even coverage throughout the crop canopy, applying sufficient water to cover the crop to just prior to the point of runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

Crop	Pest	Rates	Critical Comments
Apples, **Pears**	Leafrollers, Codling moth	2 g/100 litres as a dilute spray. Add a non-ionic wetting agent at label rates	Begin applications when local thresholds are reached. Apply PROCLAIM® Opti in a sufficient volume of water to ensure thorough coverage of the crop canopy to the point of pre-runoff. The water rate should be adjusted to the size of the canopy. Apply a maximum of 3 sprays at a minimum of 14 day intervals.
Avocados	Leafrollers	4 g/100 litres as a dilute spray. Add a non-ionic wetting agent at label rates	Begin applications when local thresholds are reached. Apply PROCLAIM® Opti in a sufficient volume of water to ensure thorough coverage of the crop canopy to the point of pre-runoff. The water rate should be adjusted to the size of the canopy. Apply a maximum of 3 sprays at a minimum of 21 day intervals.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
25/02/2026

Crop	Pest	Rates	Critical Comments
Grapes	Leafrollers	4 g/100 litres as a dilute spray. Add a non-ionic wetting agent at label rates	Begin applications when local thresholds are reached. Apply PROCLAIM® Opti in a sufficient volume of water to ensure thorough coverage of the crop canopy to the point of pre-runoff. PROCLAIM® Opti can be applied at any time from post flowering to pre bunch closure. For optimal control of Leafrollers, apply soon after egg lay when larvae are small in size and before larvae become webbed into the bunches. Eggs laid on new growth after application may not be controlled. Make no more than 2 applications of PROCLAIM® Opti per calendar year, with a minimum interval period of 14 days.
Kiwifruit	Leafrollers	2 g/100 litres as a dilute spray. Add a non-ionic wetting agent at label rates.	Apply PROCLAIM® Opti at fruit set when local thresholds are reached, with further applications at 21 day intervals if crop monitoring shows a need. Apply PROCLAIM® Opti in a sufficient volume of water to ensure thorough coverage of the crop canopy to the point of pre-runoff. Make no more than 3 applications of PROCLAIM® Opti per season. The final spray MUST NOT be made within 42 days of harvest.

EPA RESTRICTIONS

The maximum application rate of PROCLAIM® Opti on grapes is 2g emamectin benzoate/ha (40 g PROCLAIM® Opti/ha). The maximum application frequency of PROCLAIM® Opti on grapes MUST NOT be more than twice per calendar year, with a minimum interval period of 14 days.

The maximum application rate of PROCLAIM® Opti on pome fruits is 2.5 g emamectin benzoate/ha (50 g PROCLAIM® Opti /ha). The maximum application frequency of PROCLAIM® Opti on pome fruits MUST NOT be more than 3 times per calendar year, with a minimum interval period of 14 days.

The maximum application rate of PROCLAIM® Opti on avocados is 6g emamectin benzoate/ha (120 g PROCLAIM® Opti /ha). The maximum application frequency of PROCLAIM® Opti on avocados MUST NOT be more than 3 times per calendar year, with a minimum interval period of 21 days.

The maximum application rate of PROCLAIM® Opti on kiwifruit is 2g emamectin benzoate/ha (40 g PROCLAIM® Opti /ha). The maximum application frequency of PROCLAIM® Opti on kiwifruit MUST NOT be more than 3 times per calendar year, with a minimum interval period of 14 days.

BUFFER ZONE AND SPRAY DRIFT

When applying to avocados, the person in charge of the application of PROCLAIM® Opti and any person applying PROCLAIM® Opti MUST ensure that PROCLAIM® Opti is not applied within 5 metres of a downwind water body.
DO NOT apply when weather conditions favour drift from the target area.
Take all reasonable steps to ensure that the substance does NOT cause any significant adverse effect in the environment beyond the application area.

POLLINATOR SAFETY

DO NOT apply until after crops have finished flowering.
DO NOT apply PROCLAIM® Opti to plants if:

a) Bees are foraging; or
b) The plants are in flower or part flower, and are likely to be visited by non-target invertebrate pollinators (including bees).

MIXING

Add the required amount of PROCLAIM® Opti to the partly filled spray tank, then add the remainder of the water while thoroughly agitating. Continue agitation while spraying.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
25/02/2026

APPLICATION
PROCLAIM® Opti MUST only be applied with ground based equipment.
PROCLAIM® Opti MUST NOT be applied when wind speeds are less than 3 km/hr or more than 20 km/hr as measured at the application site.

COMPATIBILITY
For information on compatibility of PROCLAIM® Opti, consult your local Syngenta representative. As formulations of other manufacturer's products are beyond the control of Syngenta, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.

WITHHOLDING PERIODS
It is an offence for users of this product to cause residues exceeding the relevant MRD in the Food Notice: Maximum Residue Levels for Agricultural Compounds
Avocados ..... 3 days
Grapes ..... DO NOT apply after bunch closure
Kiwifruit ..... 42 days
Pipfruit ..... 3 days
For export crops – contact your exporter or Industry crop adviser.
Leaf plucking in vineyards:
Sheep must not be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
25/02/2026

STORAGE
Store in a cool, dry place, and protect from sunlight. Avoid temperatures below minus 5°C or above 35°C. See safety data sheet for additional storage information. Stores containing quantities of 100 kg or more require signage and secondary containment.

SAFETY DATA SHEET
If additional hazard information is required, refer to the Safety Data Sheet. For a copy, visit our website at www.syngenta.co.nz

DISCLAIMER
This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser MUST determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland 1052. Phone 0800 333 336.

Agrecovery logo

Batch Number	
DOM	

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S.
(emamectin benzoate)
MARINE POLLUTANT
PACKING GROUP III, HAZCHEM 2Z
CONTAIN SPILLAGE



In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours).