Blue AG

Stinger 510 SL

HERBICIDE

WARNING

KEEP OUT OF REACH OF CHILDREN

MINISTER FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

GROUP 9 HERBICIDE

ACTIVE INGREDIENT

510 g/litre Glyphosate

as the isopropylamine salt in the form of a soluble concentrate.

A non-selective herbicide for the control of many annual and perennial weeds in a wide range of situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS)
0800 CHEMCALLOR 0800 243 622 OR
PGG WRIGHTSON LTD 0800 10 22 76

20 L
NET CONTENTS

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

PEEL OFF

HAZARD CLASSIFICATION: Skin irritation Category 2, Hazardous to the aquatic environment (chronic) Category 2.
TOXICITY: WARNING: Causes skin irritation.
ECOTOXICITY: Toxic to some aquatic plants and with long-lasting effects.

QUALIFIED HANDLER: Required when applied onto or into water and the water has the potential to leave the place containing the application area.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure cool, dry place out of direct sunlight and away from seed, fertilizers, feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT: Apply with well-maintained and calibrated spray equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409:2004).

DISPOSAL INFORMATION
CONTAINER DISPOSAL: Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

ENVIRONMENT: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur. If accidental spraying occurs, immediately wash spray off foliage with plenty of water to minimise damage.

SPILLAGE: Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil or sand, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfil.

FIRST AID: If swallowed do not induce vomiting. Rinse mouth. For advice, call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor. If in eyes rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a POISON CENTRE or doctor if irritation persists. If skin or hair contact occurs, remove contaminated clothing and wash with running water. Get advice if irritation persists.

COMPATIBILITY: Stinger 510SL is compatible with commonly used herbicides.

GENERAL INFORMATION
Stinger 510SL is a non-selective glyphosate herbicide for use in agriculture and horticulture.

- Before planting crops and trees
- In pasture renovation and reduced tillage systems
- In vineyards, orchards and market gardens
- For pre-harvest weed control
- On turf and recreational areas
- In drains and waterways.

Stinger 510SL does not have adverse ecotoxic effects on fish or crustacea. Stinger 510SL is water soluble and is absorbed by green plant tissue. It is then translocated throughout the plant, including the root system, to kill the entire plant. Plants should be actively growing, and not under drought stress, with clean foliage at the time of spraying. The product degrades in the soil and has no residual weed control activity. Rainfall within 2 hours of application may reduce effectiveness. Application to foliage which is already wet will increase the time required for rainfastness. Visible effects may not be apparent for 14 days depending on weather conditions. Some perennial weeds may not show effects for 2 - 3 weeks.

DIRECTIONS FOR USE
APPLICATION METHODS: Stinger 510SL must only be placed in equipment made from stainless steel, fibreglass, plastic, aluminium, brass or copper. Other types of container may cause a reaction in which flammable hydrogen gas is generated. All spray equipment including pumps, spray tanks, lines and nozzles, and landing gear of aircraft should be thoroughly washed with water after each spraying day.

BOOM EQUIPMENT: Application can be made in spray volumes up to 200 litres/ha using fan nozzles at pressures in the range of 200-280kPa. Boom height should be set so that a double overlap of nozzle pattern is achieved at the top of the target plants.
AERIAL EQUIPMENT: Applicable to forest and non-crop areas only. Can be applied by aircraft in spray volumes of up to 100 litres/ha. Increase spray volumes in hilly terrain to ensure maximum target contact. Do not use in intensive horticultural cropping areas.
HANDGUN AND KNAPSACK: Ensure a fine spray pattern to give complete coverage. For handguns, use pressures of 1000 2000 kPa and No. 4 and No. 6 spray tips.
WIPER EQUIPMENT: Pre-mix 1 part Stinger 510SL in 3 parts clean water to make a 25% solution. Mix only enough for immediate use. Plants must be clean and dry and the wiper fully saturated. Apply up and down slopes for effective applications.
MISTBLOWER: Refer to table.

MIXING: Fill the clean spray tank with half the required amount of clean water. Add Stinger 510SL as required and mix well before adding the remaining water. If surfactant is required, add this last to minimise foaming. Agitate well before spraying. When tank-mixing with other chemicals, add Stinger 510SL after other chemicals, except concentrates, but before suspension concentrates. Soil particles in the water containing calcium salts deactivate Stinger 510SL. If hard water cannot be avoided, the addition of a pH buffer is recommended.

PRIMARY INDUSTRIES NEW ZEALAND
19/11/2025

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS: Wheat, barley, oat grain- 7 days

ANIMAL SAFETY: Poisonous plants, such as ragwort, may become more palatable to stock after spraying, and livestock should be kept away until plants have died down. Otherwise, withholding is not required for stock grazing. However, it is recommended that grazing is delayed for 3 days for annual weeds, and 7 days for perennial weeds. This will allow **Stinger 510SL** to penetrate the weeds for maximum effectiveness.

WEED CONTROL NOTES

AGRICULTURAL AREAS: May be used for the control of annual and perennial weeds as indicated in the Weeds Controlled Table prior to sowing any crop. **ASPARAGUS:** Pre-crop emergence and post-harvest applications. Apply prior to planting or crop emergence. Do not treat emerging spears. Application can also be made after removal of all spears but before crop regrowth occurs. In established ferns, apply as a directed or shielded spray or with wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plant. **BROAD-ACRE CROPS FOR WEED CONTROL: Wheat, barley, oats.** Can be used to control weeds prior to cultivation or prior to harvest. Not to be used as preharvest desiccant. Spray 7-12 days prior to harvest when the grain moisture is less than 30%. Do not harvest within 7 days after spraying. Do not spray undersown crops. Do not spray crops that will be harvested for seed. Do not apply more than 2.8 L/ha. Avoid treating plants under drought stress. **BRUSHWEEDS:** Refer to table. **DRAINS AND WATERWAYS:** For best results ensure that 75% of the weed to be sprayed is above water level. Avoid treating silt-laden weeds. Following spraying do not submerge the weed, as this will cause the herbicide to be washed off. **FORESTRY:** For the control of grasses and weeds as listed in the table. Can be applied for weed control prior to burning or planting of forest or nursery or around already established trees. Do not allow spray drift or spray contact with foliage or green bark of desirable trees. **MARKET GARDENS:** Apply before or after seeding but prior to crop emergence. Spraying can also be made prior to transplanting or as a post-harvest clean up. **MINIMUM TILLAGE:** For total weed control prior to seeding in a reduced tillage system. Weeds should be actively growing and not under stress from frost, drought or waterlogging. For the improved control of white clover, tank mix with a dicamba or tribenuron-methyl formulation at approved label rates. Do within 28 days of applying dicamba. If using with tribenuron-methyl then do not sow crops within 14 days of application, or sow brassicas if the soil pH exceeds 6.5 or use within 2 months after applying lime. Consult partner product label for further directions. Cultivation may start from 3 days after spraying annual weeds, and 7 days after spraying perennial weeds. Cultivation should occur within 21 days after spraying. Do not sow until a satisfactory seedbed is formed.
NON-AGRICULTURAL AREAS: Treat specific problem weeds as indicated in the Weeds Controlled Table.
ORCHARDS/VINES: Crops: citrus, apple, pear, stonefruit, kiwifruit, grapes. Apply as a directed or shielded spray. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of the stonefruit tree, or green bark, suckers, buds, foliage or fruit of other trees. Spray before crops bud, sucker or bloom. **PASTURE:** May be applied as a spot application to control a wide range of weeds in pasture. It is nonselective and may damage or kill any plant in the spray area. **RECREATIONAL AREAS:** Lawns, turf, parks, reserves and ornamentals. Apply before the sowing and planting of these areas, or use as a spot application. In mown areas, allow sufficient regrowth before spraying. Can also be sprayed around the base of established trees or shrubs, as a directed or shielded spray. Avoid contact with any part of the shrub or tree.

RESISTANCE MANAGEMENT: Stinger 510SL contains the GROUP 9 herbicide glyphosate. Rotate or alternate herbicide types during or between seasons to avoid resistance in some weed species. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult http://resistance.nzpps.org for alternative modes of action and details of resistance management strategies (if any).

WEEDS CONTROLLED TABLE

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	WHEN TO APPLY
Annual weeds	0.7-1.4	140	20	40	Use the higher rate for larger weeds.
Australian sedge	4.2 - 6.3 (aerial)	700	105	210	Apply from November to March. Burn or slash from 30 days after treatment.
Barberry	-	700	105	210	Apply from flowering to late fruit stage (January - April). Regrowth should be 1,5m tall before spraying is carried out.
Barley grass	0.7	140	20	40	Apply when plants are at early seed head stage. (September - December).
Blackberry	6.3-14	1060	105	210	Apply from flowering-fruiting stage (December - April), before frosts occur. For hard to kill blackberry, aerial application, or no follow-up action planned, use the higher rate. Allow 6-10 weeks before slashing or burning.
Boxthorn	-	880	105	-	Apply to actively growing plants (November - April). Ensure complete coverage of dense thickets. Do not use extra surfactant, or treat salt damaged bushes.
Bracken/ broom	6.3 + organosili-cone penetrant 250 mL/ 100 L	700 + organosili-cone penetrant 250 mL/ 100 L	105 + organosili-cone penetrant 20 mL	210 + organosili-cone penetrant 20 mL	Apply after bracken fronds have unfurled (mid-Jan-June), but before frosts occur. For ground application to broom anytime, but for aerial application when broom is in full leaf.
Browntop	4.2	700	105	210	Apply when plants are 5-10cm high. For taller plants, graze or burn and allow regrowth of 5-10 cm height before treatment.
Buddleia	6.3	700	105	-	Apply using handgun anytime. Aerialtreat buddleja up to 2 metres high (February-April) using spray volume of 200-300 litres/ha.
Californian thistle	2.8	700	105	210	Apply after flower bud development. Retreatment and/or cropping or pasture improvement may be necessary to restrict reestablishment from seed. Mowing thistles 10 days after spraying may assist control.
Cleavers	1	350	50	105	Spray actively growing plants after most reach early seedhead.
Couch	2.1-4.2	700	105	210	For suppression, apply 2.1 litres/ha in spring and repeat in autumn. For best results, apply 2.8 - 4.2 litres/ha in autumn to active growth of at least 10cm where no previous cultivation has occurred for at least 6 months.
Docks	2.8-6.3	700	105	-	Spray plants in spring or summer when actively growing and before seedheads have formed. Use the lower rate for seedlings and the higher rate for mature plants. In pasture situations boom spray prior to reseeding or establishing crops to restrict seedling reestablishment. Under trees or vines may be tank-mixed with Gesatop 500FW for residual seedling control.
Floating sweet grass	4.2	700	105	210	Spray actively growing plants after most reach early seedhead. Spraying plants under drought stress may give reduced results.
Gorse	-	700 + organosili-cone penetrant 200 mL/ 100 L	-	-	Apply anytime ensuring complete coverage of plants not stressed by drought, grazing or previous herbicide treatment.
Indian doab	6.3	700	105	-	Apply at early seedhead stage before frosts occur.
Johnson grass	4.2	700	105	-	Apply to actively growing plants.
Kikuyu/ Mercer grass	4.2	700	105	210	Apply at early seedhead stage. On dense infestations, cut and spray the regrowth.
Nodding thistle	-	700	105	-	When applying by wiper ensure plants are actively growing (October - April) when at least 10cm of the plant can be wiped. A double pass is recommended and follow-up with spot treatments if necessary.
Old man's beard	3.5	700-1400	210	-	Apply Nov-March prior to first leaf browning. Use higher rates on stems over 1cm in diameter. If complete coverage is not possible plants should be cut at ground level in winter and the regrowth sprayed with the higher rate when plants are at least 50cm lonq (March).
Pampas grass/toetoe	-	700	105	-	Spray Spring-Autumn. Reduced control may result if plants are in flower at the time of spraying. Ensure complete coverage.
Paspalum	2.8-4.2	700	105	210	Spray at low rate if plants are at early seedhead stage. Use 4.2 litres/ha on mature vegetative plants, before onset of frosts.
Potato (self set)	2.1	350	55	105	Spray actively growing plants after most reach early seedhead.
Prairie grass	1	140	20	-	Apply at early seedhead stage.
Ragwort	4.2	700	105	-	Spray when actively growing and ensure there is sufficient leaf area for herbicide uptake. When applying by wiper use on plants 10-15cm above the pasture. For best results, a double pass is recommended with a follow-up spot treatment if necessary. Caution: Ragwort becomes more palatable after treatment. Keep stock away from treated areas until plants brown-off.
Ratstail	3.5	700	105	-	Apply at a spray volume of 135 litres/ ha in autumn. In dry conditions, delay sprayinq until sufficient qreen foliage is present.
Rautahi (cutty grass)	4.2	700	105	210	Spray to actively growing plants from January - March. Burning or slashing may begin 30 days after treatment following brown-off.
Rushes	4.2	700	105	210	Burning or slashing may begin 30 days after treatment following brown-off. Wiper application to be carried out in November - May.
Ring fern	0.7-2.1	210	35	-	For effective control, apply 2.1 litres/ha. For pasture manipulation, apply 0.7-1.4 litres/ha from December - April. Use higher rates for dense infestations. Burn 4-6 weeks after treatment.
Ryegrass (annual Italian short rotation hybrids)					Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Established < 8 months	2.1-2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Established > 8 months	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Ryegrass (Perennial)	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20 mL		Spray to actively growing plants with at least [illegible] of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Sweet brier	8.5 + organosili-cone penetrant 250mL/ 100L	700 + organosili-cone penetrant 200mL/ 100L	105 + organosili-cone penetrant 20 mL	-	Spray when plant is in late flowering stage at early leaf flowing stage. Ensure full coverage.
Tall fescue	4.2	700	105	210	Apply January - March to actively growth.
Willow	6.3 + organosili-cone penetrant 250 mL/100L	700 + organosili-cone penetrant 250 mL/100L	-	-	Apply before leaf fall begins (January - April). Total spray coverage is essential for best results.
Yarrow	2.1 -4.2	-	-	-	Apply to plant with 5-10cm of leaf. For best results cultivate 10 days after treatment. Use the low rate where a crop to be grazed is competitive to yarrow. Use higher rate where this is a poor competitor.

Add the Slikka™ wetter and penetrant at the suppliers recommended rates.
Slikka™ is a trade name of UPL Ltd CROP SOLUTIONS.

Registered pursuant to the ACVM Act 1997, No. P9580. See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000133. See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

Blue AG

Stinger 510 SL

HERBICIDE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

WARNING
KEEP OUT OF REACH OF CHILDREN

GROUP 9 HERBICIDE

ACTIVE INGREDIENT

510 g/litre Glyphosate

as the isopropylamine salt in the form of a soluble concentrate.

A non-selective herbicide for the control of many annual and perennial weeds in a wide range of situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS) 0800 CHEMCALL OR 0800 243 622 OR PGG WRIGHTSON LTD 0800 10 22 76

200 L
NET CONTENTS

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

PEEL OFF

HAZARD CLASSIFICATION: Skin irritation Category 2,
Hazardous to the aquatic environment (chronic) Category 2.
TOXICITY: WARNING: Causes skin irritation.
ECOTOXICITY: Toxic to some aquatic plants and with long-lasting effects.

QUALIFIED HANDLER: Required when applied onto or into water and the water has the potential to leave the place containing the application area.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure cool, dry place out of direct sunlight and away from seed, fertilizers, feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT: Apply with well-maintained and calibrated spray equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409:2004).

DISPOSAL INFORMATION
CONTAINER DISPOSAL: Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

ENVIRONMENT: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur. If accidental spraying occurs, immediately wash spray off foliage with plenty of water to minimise damage.

SPILLAGE: Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil or sand, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfil.

FIRST AID: If swallowed do not induce vomiting. Rinse mouth. For advice, call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor. If in eyes rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a POISON CENTRE or doctor if irritation persists. If skin or hair contact occurs, remove contaminated clothing and wash with running water. Get advice if irritation persists.

COMPATIBILITY: Stinger 510SL is compatible with commonly used herbicides.

GENERAL INFORMATION
Stinger 510SL is a non-selective glyphosate herbicide for use in agriculture and horticulture.

- Before planting crops and trees
- In pasture renovation and reduced tillage systems
- In vineyards, orchards and market gardens
- For pre-harvest weed control
- On turf and recreational areas
- In drains and waterways.

Stinger 510SL does not have adverse ecotoxic effects on fish or crustacea. Stinger 510SL is water soluble and is absorbed by green plant tissue. It is then translocated throughout the plant, including the root system, to kill the entire plant. Plants should be actively growing, and not under drought stress, with clean foliage at the time of spraying. The product degrades in the soil and has no residual weed control activity. Rainfall within 2 hours of application may reduce effectiveness. Application to foliage which is already wet will increase the time required for rainfastness. Visible effects may not be apparent for 14 days depending on weather conditions. Some perennial weeds may not show effects for 2 - 3 weeks.

DIRECTIONS FOR USE
APPLICATION METHODS: Stinger 510SL must only be placed in equipment made from stainless steel, fibreglass, plastic, aluminium, brass or copper. Other types of container may cause a reaction in which flammable hydrogen gas is generated. All spray equipment including pumps, spray tanks, lines and nozzles, and landing gear of aircraft should be thoroughly washed with water after each spraying day.
BOOM EQUIPMENT: Application can be made in spray volumes up to 200 litres/ha using fan nozzles at pressures in the range of 200-280kPa. Boom height should be set so that a double overlap of nozzle pattern is achieved at the top of the target plants.
AERIAL EQUIPMENT: Applicable to forest and non-crop areas only. Can be applied by aircraft in spray volumes of up to 100 litres/ha. Increase spray volumes in hilly terrain to ensure maximum target contact. Do not use in intensive horticultural cropping areas.
HANDGUN AND KNAPSACK: Ensure a fine spray pattern to give complete coverage. For handguns, use pressures of 1000 2000 kPa and No. 4 and No. 6 spray tips.
WIPER EQUIPMENT: Pre-mix 1 part Stinger 510SL in 3 parts clean water to make a 25% solution. Mix only enough for immediate use. Plants must be clean and dry and the wiper fully saturated. Apply up and down slopes for effective application.
MISTBLOWER: Refer to table.

MIXING: Fill the clean spray tank with half the required amount of water. Add Stinger 510SL as required and mix well before adding the remainder of water. If surfactant is required, add this last to minimise foaming. Agitate well before spraying. When tank-mixing with other chemicals, add Stinger 510SL after other water- soluble formulations, but before suspension concentrates. Soil particles and hard water containing calcium salts deactivate Stinger 510SL. If hard water cannot be avoided, the addition of a pH buffer is recommended.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS: Wheat, barley, oat grain- 7 days

ANIMAL SAFETY: Poisonous plants, such as ragwort, may become more palatable to stock after spraying, and livestock should be kept away until plants have died down. Otherwise, withholding is not required for stock grazing. However, it is recommended that grazing is delayed for 3 days for annual weeds, and 7 days for perennial weeds. This will allow **Stinger 510SL** to penetrate the weeds for maximum effectiveness.

WEED CONTROL NOTES

AGRICULTURAL AREAS: May be used for the control of annual and perennial weeds as indicated in the Weeds Controlled Table prior to sowing any crop. **ASPARAGUS:** Pre-crop emergence and post-harvest applications. Apply prior to planting or crop emergence. Do not treat emerging spears. Application can also be made after removal of all spears but before crop regrowth occurs. In established ferns, apply as a directed or shielded spray or with wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plant. **BROAD-ACRE CROPS FOR WEED CONTROL: Wheat, barley, oats.** Can be used to control weeds prior to cultivation or prior to harvest. Not to be used as preharvest desiccant. Spray 7-12 days prior to harvest when the grain moisture is less than 30%. Do not harvest within 7 days after spraying. Do not spray undersown crops. Do not spray crops that will be harvested for seed. Do not apply more than 2.8 L./ha. Avoid treating plants under drought stress. **BRUSHWEEDS:** Refer to table. **DRAINS AND WATERWAYS:** For best results ensure that 75% of the weed to be sprayed is above water level. Avoid treating silt-laden weeds. Following spraying do not submerge the weed, as this will cause the herbicide to be washed off. **FORESTRY:** For the control of grasses and weeds as listed in the table. Can be applied for weed control prior to burning or planting of forest or nursery or around already established trees. Do not allow spray drift or spray contact with foliage or green bark of desirable trees. **MARKET GARDENS:** Apply before or after seeding but prior to crop emergence. Spraying can also be made prior to transplanting or as a post-harvest clean up. **MINIMUM TILLAGE:** For total weed control prior to seeding in a reduced tillage system. Weeds should be actively growing and not under stress from frost, drought or waterlogging. For the improved control of white clover, tank mix with a dicamba or tribenuron-methyl formulation at approved label rates. Do within 28 days of applying dicamba. If using with tribenuron-methyl then do not sow crops within 14 days of application, or sow brassicas if the soil pH exceeds 6.5 or use within 2 months after applying lime. Consult partner product label for further directions. Cultivation may start from 3 days after spraying annual weeds, and 7 days after spraying perennial weeds. Cultivation should occur within 21 days after spraying. Do not sow until a satisfactory seedbed is formed. **NON-AGRICULTURAL AREAS:** Treat specific problem weeds as indicated in the Weeds Controlled Table. **ORCHARDS/VINES:** Crops: citrus, apple, pear, stonefruit, kiwifruit, grapes. Apply as a directed or shielded spray. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of the stonefruit tree, or green bark, suckers, buds, foliage or fruit of other trees. Spray before crops bud, sucker or bloom. **PASTURE:** May be applied as a spot application to control a wide range of weeds in pasture. It is nonselective and may damage or kill any plant in the spray area. **RECREATIONAL AREAS:** Lawns, turf, parks, reserves and ornamentals. Apply before the sowing and planting of these areas, or use as a spot application. In mown areas, allow sufficient regrowth before spraying. Can also be sprayed around the base of established trees or shrubs, as a directed or shielded spray. Avoid contact with any part of the shrub or tree.

RESISTANCE MANAGEMENT: Stinger 510SL contains the GROUP 9 herbicide glyphosate. Rotate or alternate herbicide types during or between seasons to avoid resistance in some weed species. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult http://resistance.nzpps.org for alternative modes of action and details of resistance management strategies (if any).

WEEDS CONTROLLED TABLE

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	WHEN TO APPLY
Annual weeds	0.7-1.4	140	20	40	Use the higher rate for larger weeds.
Australian sedge	4.2 - 6.3 (aerial)	700	105	210	Apply from November to March. Burn or slash from 30 days after treatment.
Barberry	-	700	105	210	Apply from flowering to late fruit stage (January - April). Regrowth should be 1,5m tall before spraying is carried out.
Barley grass	0.7	140	20	40	Apply when plants are at early seed head stage. (September - December).
Blackberry	6.3-14	1060	105	210	Apply from flowering-fruiting stage (December - April), before frosts occur. For hard to kill blackberry, aerial application, or no follow-up action planned, use the higher rate. Allow 6-10 weeks before slashing or burning.
Boxthorn	-	880	105	-	Apply to actively growing plants (November - April). Ensure complete coverage of dense thickets. Do not use extra surfactant, or treat salt damaged bushes.
Bracken/ broom	6.3 + organosili-cone penetrant 250 mL/ 100 L	700 + organosili-cone penetrant 250 mL/ 100 L	105 + organosili-cone penetrant 20 mL	210 + organosili-cone penetrant 20 mL	Apply after bracken fronds have unfurled (mid-Jan-June), but before frosts occur. For ground application to broom anytime, but for aerial application when broom is in full leaf.
Browntop	4.2	700	105	210	Apply when plants are 5-10cm high. For taller plants, graze or burn and allow regrowth of 5-10 cm height before treatment.
Buddleia	6.3	700	105	-	Apply using handgun anytime. Aerialtreat buddleja up to 2 metres high (February-April) using spray volume of 200-300 litres/ha.
Californian thistle	2.8	700	105	210	Apply after flower bud development. Retreatment and/or cropping or pasture improvement may be necessary to restrict reestablishment from seed. Mowing thistles 10 days after spraying may assist control.
Cleavers	1	350	50	105	Spray actively growing plants after most reach early seedhead.
Couch	2.1-4.2	700	105	210	For suppression, apply 2.1 litres/ha in spring and repeat in autumn. For best results, apply 2.8 - 4.2 litres/ha in autumn to active growth of at least 10cm where no previous cultivation has occurred for at least 6 months.
Docks	2.8-6.3	700	105	-	Spray plants in spring or summer when actively growing and before seedheads have formed. Use the lower rate for seedlings and the higher rate for mature plants. In pasture situations boom spray prior to reseeding or establishing crops to restrict seedling reestablishment. Under trees or vines may be tank-mixed with Gesatop 500FW for residual seedling control.
Floating sweet grass	4.2	700	105	210	Spray actively growing plants after most reach early seedhead. Spraying plants under drought stress may give reduced results.
Gorse	-	700 + organosili-cone penetrant 200 mL/ 100 L	-	-	Apply anytime ensuring complete coverage of plants not stressed by drought, grazing or previous herbicide treatment.
Indian doab	6.3	700	105	-	Apply at early seedhead stage before frosts occur.
Johnson grass	4.2	700	105	-	Apply to actively growing plants.
Kikuyu/ Mercer grass	4.2	700	105	210	Apply at early seedhead stage. On dense infestations, cut and spray the regrowth.
Nodding thistle	-	700	105	-	When applying by wiper ensure plants are actively growing (October - April) when at least 10cm of the plant can be wiped. A double pass is recommended and follow-up with spot treatments if necessary.
Old man's beard	3.5	700-1400	210	-	Apply Nov-March prior to first leaf browning. Use higher rates on stems over 1cm in diameter. If complete coverage is not possible plants should be cut at ground level in winter and the regrowth sprayed with the higher rate when plants are at least 50cm long (March).
Pampas grass/toetoe	-	700	105	-	Spray Spring-Autumn. Reduced control may result if plants are in flower at the time of spraying. Ensure complete coverage.
Paspalum	2.8-4.2	700	105	210	Spray at low rate if plants are at early seedhead stage. Use 4.2 litres/ha on mature vegetative plants, before onset of frosts.
Potato (self set)	2.1	350	55	105	Spray actively growing plants after most reach early seedhead.
Prairie grass	1	140	20	-	Apply at early seedhead stage.
Ragwort	4.2	700	105	-	Spray when actively growing and ensure there is sufficient leaf area for herbicide uptake. When applying by wiper use on plants 10-15cm above the pasture. For best results, a double pass is recommended with a follow-up spot treatment if necessary. Caution: Ragwort becomes more palatable after treatment. Keep stock away from treated areas until plants brown-off.
Ratstail	3.5	700	105	-	Apply at a spray volume of 135 litres/ ha in autumn. In dry conditions, delay spraying until sufficient green foliage is present.
Rautahi (cutty grass)	4.2	700	105	210	Spray to actively growing plants from January - March. Burning or slashing may begin 30 days after treatment following brown-off.
Rushes	4.2	700	105	210	Burning or slashing may begin 30 days after treatment following brown-off. Wiper application to be carried out in November - May.
Ring fern	0.7-2.1	210	35	-	For effective control, apply 2.1 litres/ha in a pasture formulation, apply 0.7-1.4 litres/ha from December - April. Use higher rates for dense infestations. Burn 4-6 weeks after treatment.
Ryegrass (annual Italian short rotation hybrids)					Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Established < 8 months	2.1-2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Established > 8 months	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Ryegrass (Perennial)	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20 mL		Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Sweet brier	8.5 + organosili-cone penetrant 250mL/ 100L	700 + organosili-cone penetrant 200mL/ 100L	105 + organosili-cone penetrant 20 mL	-	Spray when plant is in late flowering to early leaf yellowing stage. Ensure full coverage.
Tall fescue	4.2	700	105	210	Apply January - March to active growth.
Willow	6.3 + organosili-cone penetrant 250 mL/100L	700 + organosili-cone penetrant 250 mL/100L	-	-	Apply before leaf fall begins (January - April). Total spray coverage is essential for best results.
Yarrow	2.1 -4.2	-	-	-	Apply to plant with 5-10cm of leaf. For best results cultivate 10 days after treatment. Use the low rate when a crop to be drilled is competitive to yarrow. Use higher rate when it is a poor competitor.

Add the Slikka™ wetter and penetrant at the suppliers recommended rates.
Slikka™ is a trade name of UPL Ltd CROP SOLUTIONS.

Registered pursuant to the ACVM Act 1997, No. P9580. See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000133. See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

Blue AG

Stinger 510 SL

HERBICIDE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

WARNING
KEEP OUT OF REACH OF CHILDREN

GROUP 9 HERBICIDE

ACTIVE INGREDIENT

510 g/litre Glyphosate

as the isopropylamine salt in the form of a soluble concentrate.

A non-selective herbicide for the control of many annual and perennial weeds in a wide range of situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS) 0800 CHEMCALL OR 0800 243 622 OR PGG WRIGHTSON LTD 0800 10 22 76

1000 L
NET CONTENTS

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

PEEL OFF

HAZARD CLASSIFICATION: Skin irritation Category 2, Hazardous to the aquatic environment (chronic) Category 2.
TOXICITY: WARNING: Causes skin irritation.
ECOTOXICITY: Toxic to some aquatic plants and with long-lasting effects.

QUALIFIED HANDLER: Required when applied onto or into water and the water has the potential to leave the place containing the application area.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure cool, dry place out of direct sunlight and away from seed, fertilizers, feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT: Apply with well-maintained and calibrated spray equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409:2004).

DISPOSAL INFORMATION

CONTAINER DISPOSAL: Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

ENVIRONMENT: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur. If accidental spraying occurs, immediately wash spray off foliage with plenty of water to minimise damage.

SPILLAGE: Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil or sand, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfil.

FIRST AID: If swallowed do not induce vomiting. Rinse mouth. For advice, call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor. If in eyes rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a POISON CENTRE or doctor if irritation persists. If skin or hair contact occurs, remove contaminated clothing and wash with running water. Get advice if irritation persists.

COMPATIBILITY: Stinger 510SL is compatible with commonly used herbicides.

GENERAL INFORMATION

Stinger 510SL is a non-selective glyphosate herbicide for use in agriculture and horticulture.

- Before planting crops and trees
- In pasture renovation and reduced tillage systems
- In vineyards, orchards and market gardens
- For pre-harvest weed control
- On turf and recreational areas
- In drains and waterways.

Stinger 510SL does not have adverse ecotoxic effects on fish or crustacea. Stinger 510SL is water soluble and is absorbed by green plant tissue. It is then translocated throughout the plant, including the root system, to kill the entire plant. Plants should be actively growing, and not under drought stress, with clean foliage at the time of spraying. The product degrades in the soil and has no residual weed control activity. Rainfall within 2 hours of application may reduce effectiveness. Application to foliage which is already wet will increase the time required for rainfastness. Visible effects may not be apparent for 14 days depending on weather conditions. Some perennial weeds may not show effects for 2 - 3 weeks.

DIRECTIONS FOR USE

APPLICATION METHODS: Stinger 510SL must only be placed in equipment made from stainless steel, fibreglass, plastic, aluminium, brass or copper. Other types of container may cause a reaction in which flammable hydrogen gas is generated. All spray equipment including pumps, spray tanks, lines and nozzles, and landing gear of aircraft should be thoroughly washed with water after each spraying day.
BOOM EQUIPMENT: Application can be made in spray volumes up to 200 litres/ha using fan nozzles at pressures in the range of 200-280kPa. Boom height should be set so that a double overlap of nozzle pattern is achieved at the top of the target plants.
AERIAL EQUIPMENT: Applicable to forest and non-crop areas only. Can be applied by aircraft in spray volumes of up to 100 litres/ha. Increase spray volumes in hilly terrain to ensure maximum target contact. Do not use in intensive horticultural cropping areas.
HANDGUN AND KNAPSACK: Ensure a fine spray pattern to give complete coverage. For handguns, use pressures of 1000 2000 kPa and No. 4 and No. 6 spray tips.
WIPER EQUIPMENT: Pre-mix 1 part Stinger 510SL in 3 parts clean water to make a 25% solution. Mix only enough for immediate use. Plants must be clean and dry and the wiper fully saturated. Apply up and down slopes for effective applications.
MISTBLOWER: Refer to table.

MIXING: Fill the clean spray tank with half the required amount of clean water. Add Stinger 510SL as required and mix well before adding the remaining water. If surfactant is required, add this last to minimise foaming. Agitate well before spraying. When tank-mixing with other chemicals, add Stinger 510SL after other water soluble products but before suspension concentrates. Soil particles and hard water contaminants or salts deactivate Stinger 510SL. If hard water cannot be avoided, the addition of a pH buffer is recommended.

MINISTRY PRIMARY INDUSTRIES NEW ZEALAND
19/11/2025

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS: Wheat, barley, oat grain- 7 days

ANIMAL SAFETY: Poisonous plants, such as ragwort, may become more palatable to stock after spraying, and livestock should be kept away until plants have died down. Otherwise, withholding is not required for stock grazing. However, it is recommended that grazing is delayed for 3 days for annual weeds, and 7 days for perennial weeds. This will allow **Stinger 510SLSL** to penetrate the weeds for maximum effectiveness.

WEED CONTROL NOTES

AGRICULTURAL AREAS: May be used for the control of annual and perennial weeds as indicated in the Weeds Controlled Table prior to sowing any crop. **ASPARAGUS:** Pre-crop emergence and post-harvest applications. Apply prior to planting or crop emergence. Do not treat emerging spears. Application can also be made after removal of all spears but before crop regrowth occurs. In established ferns, apply as a directed or shielded spray or with wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plant. **BROAD-ACRE CROPS FOR WEED CONTROL:** Wheat, barley, oats. Can be used to control weeds prior to cultivation or prior to harvest. Not to be used as preharvest desiccant. Spray 7-12 days prior to harvest when the grain moisture is less than 30%. Do not harvest within 7 days after spraying. Do not spray undersown crops. Do not spray crops that will be harvested for seed. Do not apply more than 2.8 L./ha. Avoid treating plants under drought stress. **BRUSHWEEDS:** Refer to table. **DRAINS AND WATERWAYS:** For best results ensure that 75% of the weed to be sprayed is above water level. Avoid treating silt-laden weeds. Following spraying do not submerge the weed, as this will cause the herbicide to be washed off. **FORESTRY:** For the control of grasses and weeds as listed in the table. Can be applied for weed control prior to burning or planting of forest or nursery or around already established trees. Do not allow spray drift or spray contact with foliage or green bark of desirable trees. **MARKET GARDENS:** Apply before or after seeding but prior to crop emergence. Spraying can also be made prior to transplanting or as a post-harvest clean up. **MINIMUM TILLAGE:** For total weed control prior to seeding in a reduced tillage system. Weeds should be actively growing and not under stress from frost, drought or waterlogging. For the improved control of white clover, tank mix with a dicamba or tribenuron-methyl formulation at approved label rates. Do within 28 days of applying dicamba. If using with tribenuron-methyl then do not sow crops within 14 days of application, or sow brassicas if the soil pH exceeds 6.5 or use within 2 months after applying lime. Consult partner product label for further directions. Cultivation may start from 3 days after spraying annual weeds, and 7 days after spraying perennial weeds. Cultivation should occur within 21 days after spraying. Do not sow until a satisfactory seedbed is formed. **NON-AGRICULTURAL AREAS:** Treat specific problem weeds as indicated in the Weeds Controlled Table. **ORCHARDS/VINES:** Crops: citrus, apple, pear, stonefruit, kiwifruit, grapes. Apply as a directed or shielded spray. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of the stonefruit tree, or green bark, suckers, buds, foliage or fruit of other trees. Spray before crops bud, sucker or bloom. **PASTURE:** May be applied as a spot application to control a wide range of weeds in pasture. It is nonselective and may damage or kill any plant in the spray area. **RECREATIONAL AREAS:** Lawns, turf, parks, reserves and ornamentals. Apply before the sowing and planting of these areas, or use as a spot application. In mown areas, allow sufficient regrowth before spraying. Can also be sprayed around the base of established trees or shrubs, as a directed or shielded spray. Avoid contact with any part of the shrub or tree. **RESISTANCE MANAGEMENT: Stinger 510SLSL** contains the GROUP 9 herbicide glyphosate. Rotate or alternate herbicide types during or between seasons to avoid resistance in some weed species. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult http://resistance.nzpps.org for alternative modes of action and details of resistance management strategies (if any).

WEEDS CONTROLLED TABLE

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	WHEN TO APPLY
Annual weeds	0.7-1.4	140	20	40	Use the higher rate for larger weeds.
Australian sedge	4.2 - 6.3 (aerial)	700	105	210	Apply from November to March. Burn or slash from 30 days after treatment.
Barberry	-	700	105	210	Apply from flowering to late fruit stage (January - April). Regrowth should be 1,5m tall before spraying is carried out.
Barley grass	0.7	140	20	40	Apply when plants are at early seed head stage. (September - December).
Blackberry	6.3-14	1060	105	210	Apply from flowering-fruiting stage (December - April), before frosts occur. For hard to kill blackberry, aerial application, or no follow-up action planned, use the higher rate. Allow 6-10 weeks before slashing or burning.
Boxthorn	-	880	105	-	Apply to actively growing plants (November - April). Ensure complete coverage of dense thickets. Do not use extra surfactant, or treat salt damaged bushes.
Bracken/ broom	6.3 + organosili-cone penetrant 250 mL/ 100 L	700 + organosili-cone penetrant 250 mL/ 100 L	105 + organosili-cone penetrant 20 mL	210 + organosili-cone penetrant 20 mL	Apply after bracken fronds have unfurled (mid-Jan-June), but before frosts occur. For ground application to broom anytime, but for aerial application when broom is in full leaf.
Browntop	4.2	700	105	210	Apply when plants are 5-10cm high. For taller plants, graze or burn and allow regrowth of 5-10 cm height before treatment.
Buddleia	6.3	700	105	-	Apply using handgun anytime. Aerialtreat buddleja up to 2 metres high (February-April) using spray volume of 200-300 litres/ha.
Californian thistle	2.8	700	105	210	Apply after flower bud development. Retreatment and/or cropping or pasture improvement may be necessary to restrict reestablishment from seed. Mowing thistles 10 days after spraying may assist control.
Cleavers	1	350	50	105	Spray actively growing plants after most reach early seedhead.
Couch	2.1-4.2	700	105	210	For suppression, apply 2.1 litres/ha in spring and repeat in autumn. For best results, apply 2.8 - 4.2 litres/ha in autumn to active growth of at least 10cm where no previous cultivation has occurred for at least 6 months.
Docks	2.8-6.3	700	105	-	Spray plants in spring or summer when actively growing and before seedheads have formed. Use the lower rate for seedlings and the higher rate for mature plants. In pasture situations boom spray prior to reseeding or establishing crops to restrict seedling reestablishment. Under trees or vines may be tank-mixed with Gesatop 500FW for residual seedling control.
Floating sweet grass	4.2	700	105	210	Spray actively growing plants after most reach early seedhead. Spraying plants under drought stress may give reduced results.
Gorse	-	700 + organosili-cone penetrant 200 mL/ 100 L	-	-	Apply anytime ensuring complete coverage of plants not stressed by drought, grazing or previous herbicide treatment.
Indian doab	6.3	700	105	-	Apply at early seedhead stage before frosts occur.
Johnson grass	4.2	700	105	-	Apply to actively growing plants.
Kikuyu/ Mercer grass	4.2	700	105	210	Apply at early seedhead stage. On dense infestations, cut and spray the regrowth.
Nodding thistle	-	700	105	-	When applying by wiper ensure plants are actively growing (October - April) when at least 10cm of the plant can be wiped. A double pass is recommended and follow-up with spot treatments if necessary.
Old man's beard	3.5	700-1400	210	-	Apply Nov-March prior to first leaf browning. Use higher rates on stems over 1cm in diameter. If complete coverage is not possible plants should be cut at ground level in winter and the regrowth sprayed with the higher rate when plants are at least 50cm long (March).
Pampas grass/toetoe	-	700	105	-	Spray Spring-Autumn. Reduced control may result if plants are in flower at the time of spraying. Ensure complete coverage.
Paspalum	2.8-4.2	700	105	210	Spray at low rate if plants are at early seedhead stage. Use 4.2 litres/ha on mature vegetative plants, before onset of frosts.
Potato (self set)	2.1	350	55	105	Spray actively growing plants after most reach early seedhead.
Prairie grass	1	140	20	-	Apply at early seedhead stage.
Ragwort	4.2	700	105	-	Spray when actively growing and ensure there is sufficient leaf area for herbicide uptake. When applying by wiper use on plants 10-15cm above the pasture. For best results, a double pass is recommended with a follow-up spot treatment if necessary. Caution: Ragwort becomes more palatable after treatment. Keep stock away from treated areas until plants brown-off.
Ratstail	3.5	700	105	-	Apply at a spray volume of 135 litres/ ha in autumn. In dry conditions, delay spraying until sufficient green foliage is present.
Rautahi (cutty grass)	4.2	700	105	210	Spray to actively growing plants from January - March. Burning or slashing may begin 30 days after treatment following brown-off.
Rushes	4.2	700	105	210	Burning or slashing may begin 30 days after treatment following brown-off. Wiper application to be carried out in November - May.
Ring fern	0.7-2.1	210	35	-	For effective control, apply 2.1 litres/ha. For [illegible] apply 0.7-1.4 litres/ha from December - April. Use higher rates for dense infestations. Burn 4-6 weeks after treatment.
Ryegrass (annual Italian short rotation hybrids)					Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Established < 8 months	2.1-2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Established > 8 months	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Ryegrass (Perennial)	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20 mL		Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Sweet brier	8.5 + organosili-cone penetrant 250mL/ 100L	700 + organosili-cone penetrant 200mL/ 100L	105 + organosili-cone penetrant 20 mL	-	Spray when plant is in late flowering to early leaf yellowing stage. Ensure full coverage.
Tall fescue	4.2	700	105	210	Apply January - March to active growth.
Willow	6.3 + organosili-cone penetrant 250 mL/100L	700 + organosili-cone penetrant 250 mL/100L	-	-	Apply before leaf fall begins (January - April). Total spray coverage is essential for best results.
Yarrow	2.1 -4.2	-	-	-	Apply to plant with 5-10cm of leaf. For best results cultivate 10 days after treatment. Use the low rate when a crop to be drilled is competitive to yarrow. Use higher rate when it is a poor competitor.

Add the Slikka™ wetter and penetrant at the suppliers recommended rates.
Slikka™ is a trade name of UPL Ltd CROP SOLUTIONS.

Registered pursuant to the ACVM Act 1997, No. P9580. See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000133. See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

150 X 240 (20 L)

Pantone 7485C

Pantone 368C

ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Blue AG

Stinger 510 SL

HERBICIDE

GROUP **9** HERBICIDE

ACTIVE INGREDIENT
510 g/litre Glyphosate
as the isopropylamine salt in the form of a soluble concentrate.

A non-selective herbicide for the control of many annual and perennial weeds in a wide range of situations.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS)
0800 CHEMCALL OR 0800 243 622 OR
PGG WRIGHTSON LTD 0800 10 22 76

20 L
NET CONTENTS

150 X 240 (20 L) | Pantone 425C | BLACK C:60 M:40 Y:40 K:100 | Pantone 368C | Pantone 485 C

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

PEEL OFF

HAZARD CLASSIFICATION: Hazardous to the aquatic environment (chronic) Category 2.

ECOTOXICITY: Toxic to some aquatic plants and with long-lasting effects.

QUALIFIED HANDLER: Required when applied onto or into water and the water has the potential to leave the place containing the application area.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure cool, dry place out of direct sunlight and away from seed, fertilizers, feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT: Apply with well-maintained and calibrated spray equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409:2004)

DISPOSAL INFORMATION

CONTAINER DISPOSAL: Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

ENVIRONMENT: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur. If accidental spraying occurs, immediately wash spray off foliage with plenty of water to minimise damage.

SPILLAGE: Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil or sand, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfil.

FIRST AID: If swallowed do not induce vomiting. Rinse mouth. For advice, call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor. If in eyes rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a POISON CENTRE or doctor if irritation persists. If skin or hair contact occurs, remove contaminated clothing and wash with running water. Get advice if irritation persists.

COMPATIBILITY: STINGER 510SL is compatible with commonly used herbicides.

GENERAL INFORMATION

STINGER 510SL is a non-selective glyphosate herbicide for use in agriculture and horticulture.

- Before planting crops and trees
- In pasture renovation and reduced tillage systems
- For pre-harvest weed control
- In vineyards, orchards and market gardens
- On turf and recreational areas
- In drains and waterways.

STINGER 510SL does not have adverse ecotoxic effects on fish or crustacea. **STINGER 510SL** is water soluble and is absorbed by green plant tissue. It is then translocated throughout the plant, including the root system, to kill the entire plant. Plants should be actively growing, and not under drought stress, with clean foliage at the time of spraying. The product degrades in the soil and has no residual weed control activity. Rainfall within 2 hours of application may reduce effectiveness. Application to foliage which is already wet will increase the time required for rainfastness. Visible effects may not be apparent for 14 days depending on weather conditions. Some perennial weeds may not show effects for 2 - 3 weeks.

DIRECTIONS FOR USE

APPLICATION METHODS: STINGER 510SL must only be placed in equipment made from stainless steel, fibreglass, plastic, aluminium, brass or copper. Other types of container may cause a reaction in which flammable hydrogen gas is generated. All spray equipment including pumps, spray tanks, lines and nozzles, and landing gear of aircraft should be thoroughly washed with water after each spraying day.

BOOM EQUIPMENT: Application can be made in spray volumes up to 200 litres/ha using fan nozzles at pressures in the range of 200-280kPa. Boom height should be set so that a double overlap of nozzle pattern is achieved at the top of the target plants.

AERIAL EQUIPMENT: Applicable to forest and non-crop areas only. Can be applied by aircraft in spray volumes of up to 100 litres/ha. Increase spray volumes in hilly terrain to ensure maximum target contact. Do not use in intensive horticultural cropping areas.

HANDGUN AND KNAPSACK: Ensure a fine spray pattern to give complete coverage. For handguns, use pressures of 1000 2000 kPa and No. 4 and No. 6 spray tips.

WIPER EQUIPMENT: Pre-mix 1 part STINGER 510SL in 3 parts clean water to make a 25% solution. Mix only enough for immediate use. Plants must be clean and dry and the wiper fully saturated. Apply up and down slopes for effective application.

MISTBLOWER: Refer to tab[illegible]

MIXING: Fill the clean spray tank [illegible] required amount of clean water. Add **STINGER 510SL** [illegible] required and mix well before adding [illegible] water. If surfactant is required add it [illegible] to minimise foaming. Agitate well before [illegible] tank-mixing with other chemicals [illegible] **STINGER 510SL** after other water-soluble concentrates, but before suspension concentrates. Avoid particles and hard water containing calcium salts deactivate **STINGER 510SL**. If hard water cannot be avoided, the addition of a pH buffer is recommended.

19/11/2025

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -
D.O.M. -

9 415251 004621 >

150 X 240 (20 L)

Pantone 425C

GBD STD MULTIFOLD®

MINISTRY FOR PRIMARY INDUSTRIES

NEW ZEALAND

19/11/2025

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS: Wheat, barley, oat grain- 7 days

ANIMAL SAFETY: Poisonous plants, such as ragwort, may become more palatable to stock after spraying, and livestock should be kept away until plants have died down. Otherwise, withholding is not required for stock grazing. However, it is recommended that grazing is delayed for 3 days for annual weeds, and 7 days for perennial weeds. This will allow **STINGER 510SL** to penetrate the weeds for maximum effectiveness.

WEED CONTROL NOTES
AGRICULTURAL AREAS: May be used for the control of annual and perennial weeds as indicated in the Weeds Controlled Table prior to sowing any crop. **ASPARAGUS:** Pre-crop emergence and post-harvest applications. Apply prior to planting or crop emergence. Do not treat emerging spears. Application can also be made after removal of all spears but before crop regrowth occurs. In established ferns, apply as a directed or shielded spray or with wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plant.
BROAD-ACRE CROPS FOR WEED CONTROL: Wheat, barley, oats. Can be used to control weeds prior to cultivation or prior to harvest. Not to be used as preharvest desiccant. Spray 7-12 days prior to harvest when the grain moisture is less than 30%. Do not harvest within 7 days after spraying. Do not spray undersown crops. Do not spray crops that will be harvested for seed. Do not apply more than 2.8 L./ha. Avoid treating plants under drought stress. **BRUSHWEEDS:** Refer to table. **DRAINS AND WATERWAYS:** For best results ensure that 75% of the weed to be sprayed is above water level. Avoid treating silt-laden weeds. Following spraying do not submerge the weeds as this will cause the herbicide to be washed off. **FORESTRY:** For the control of grasses and weeds as listed in the table. Can be applied for weed control prior to burning or planting of forest or nursery or around already established trees. Do not allow spray drift or spray contact with foliage or green bark of desirable trees. **MARKET GARDENS:** Apply before or after seeding but prior to crop emergence. Spraying can also be made prior to transplanting or as a post-harvest clean up. **MINIMUM TILLAGE:** For total weed control prior to seeding in a reduced tillage system. Weeds should be actively growing and not under stress from frost, drought or waterlogging. For the improved control of white clover, tank mix with a dicamba or tribenuron-methyl formulation at approved label rates. Do within 28 days of applying dicamba. If using with tribenuron-methyl then do not sow crops within 14 days of application, or sow brassicas if the soil pH exceeds 6.5 or use within 2 months after applying lime. Consult partner product label for further directions. Cultivation may start from 3 days after spraying annual weeds, and 7 days after spraying perennial weeds. Cultivation should occur within 21 days after spraying. Do not sow until a satisfactory seedbed is formed. **NON-AGRICULTURAL AREAS:** Treat specific problem weeds as indicated in the Weeds Controlled Table. **ORCHARDS/VINES:** Crops: citrus, apple, pear, stonefruit, kiwifruit, grapes. Apply as a directed or shielded spray. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of the stonefruit tree, or green bark, suckers, buds, foliage or fruit of other trees. Spray before crops bud, sucker or bloom. **PASTURE:** May be applied as a spot application to control a wide range of weeds in pasture. It is nonselective and may damage or kill any plant in the spray area. **RECREATIONAL AREAS:** Lawns, turf, parks, reserves and ornamentals. Apply before the sowing and planting of these areas, or use as a spot application. In mown areas, allow sufficient regrowth before spraying. Can also be sprayed around the base of established trees or shrubs, as a directed or shielded spray. Avoid contact with any part of the shrub or tree. **RESISTANCE MANAGEMENT: STINGER 510SL** contains the GROUP 9 herbicide glyphosate. Rotate or alternate herbicide types during or between seasons to avoid resistance in targeted species. Some naturally occurring weeds may become resistant by the excessive use of these herbicides. To prevent the development of resistance within the weeds and/or rotate herbicides with a different mode of action, consult http://resistance.nz.org for alternative modes of action and details of resistance management strategies (if any).

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
19/11/2025

WEEDS CONTROLLED TABLE

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	[illegible] TO APPLY
Annual weeds	0.7-1.4	140	20	40	Use the higher rate for larger weeds
Australian sedge	4.2 - 6.3 (aerial)	700	105	210	Apply from November to March. Burn or slash from 30 days after treatment.
Barberry	-	700	105	210	Apply from flowering to late fruit stage (January - April). Regrowth should be 1,5m tall before spraying is carried out.
Barley grass	0.7	140	20	40	Apply when plants are at early seed head stage. (September - December).
Blackberry	6.3-14	1060	105	210	Apply from flowering-fruiting stage (December - April), before frosts occur. For hard to kill blackberry, aerial application, or no follow-up action planned, use the higher rate. Allow 6-10 weeks before slashing or burning.

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	WHEN TO APPLY
Boxthorn	-	880	105	-	Apply to actively growing plants (November - April). Ensure complete coverage of dense thickets. Do not use extra surfactant, or treat salt damaged bushes.
Bracken/ broom	6.3 + organosili-cone penetrant 250 mL/ 100 L	700 + organosili-cone penetrant 250 mL/ 100 L	105 + organosili-cone penetrant 20 mL	210 + organosili-cone penetrant 20 mL	Apply after bracken fronds have unfurled (mid-Jan-June), but before frosts occur. For ground application to broom anytime, but for aerial application when broom is in full leaf.
Browntop	4.2	700	105	210	Apply when plants are 5-10cm high. For taller plants, graze or burn and allow regrowth of 5 - 10 cm height before treatment.
Buddleia	6.3	700	105	-	Apply using handgun anytime. Aerialtreat buddleja up to 2 metres high (February-April) using spray volume of 200-300 litres/ha.
Californian thistle	2.8	700	105	210	Apply after flower bud development. Retreatment and/or cropping or pasture improvement may be necessary to restrict reestablishment from seed. Mowing thistles 10 days after spraying may assist control.
Cleavers	1	350	50	105	Spray actively growing plants after most reach early seedhead.
Couch	2.1-4.2	700	105	210	For suppression, apply 2.1 litres/ha in spring and repeat in autumn. For best results, apply 2.8 - 4.2 litres/ha in autumn to active growth of at least 10cm where no previous cultivation has occurred for at least 6 months.
Docks	2.8-6.3	700	105	-	Spray plants in spring or summer when actively growing and before seedheads have formed. Use the lower rate for seedlings and the higher rate for mature plants. In pasture situations boom spray prior to reseeding or establishing crops to restrict seedling reestablishment. Under trees or vines may be tank-mixed with Gesatop 500FW for residual seedling control.
Floating sweet grass	4.2	700	105	210	Spray actively growing plants after most reach early seedhead. Spraying plants under drought stress may give reduced results.
Gorse	-	700 + organosili-cone penetrant 200 mL/ 100 L	-	-	Apply anytime ensuring complete coverage of plants not stressed by drought, grazing or previous herbicide treatment.
Indian doab	6.3	700	105	-	Apply at early seedhead stage before frosts occur.
Johnson grass	4.2	700	105	-	Apply to actively growing plants.
Kikuyu/ Mercer grass	4.2	700	105	210	Apply at early seedhead stage. On dense infestations, cut and spray the regrowth.
Nodding thistle	-	700	105	-	When applying by wiper ensure plants are actively growing (October - April) when at least 10cm of the plant can be wiped. A double pass is recommended and follow-up with spot treatments if necessary.
Old man's beard	3.5	700-1400	210	-	Apply Nov-March prior to first leaf browning. Use higher rates on stems over 1cm in diameter. If complete coverage is not possible plants should be cut at ground level in winter and the regrowth sprayed with the higher rate when plants are at least 50cm long (March).
Pampas grass/toetoe	-	700	[illegible]	[illegible]	Spray Spring-Autumn. Reduced control may result if plants are in flower at the time of spraying. Ensure complete coverage.
Paspalum	2.8-4.2	700	105	210	Spray at low rate if plants are at early seedhead stage. Use 4.2 litres/ha on mature vegetative plants, before onset of frosts.
Potato (self set)	[illegible]	350	55	105	Spray actively growing plants after most reach early seedhead.
Prairie grass	[illegible]	140	20	-	Apply at early seedhead stage.
Ragwort	[illegible]	700	[illegible]	[illegible]	Spray when actively growing and ensure there is sufficient leaf area for herbicide uptake. When applying by wiper use on plants 10-15cm above the pasture. For best results, a double pass is recommended with a follow-up spot treatment if necessary. Caution: Ragwort becomes more palatable after treatment. Keep stock away from treated areas until plants brown off.
Ratstail	3.5	700	105	-	Apply at a spray volume of 135 litres/ ha in autumn. In dry conditions, delay spraying until sufficient green foliage is present.
Rautahi (cutty grass)	4.[illegible]	700	105	210	Spray to actively growing plants from January - March. Burning or slashing may begin 30 days after treatment following brown-off.
Rushes	4.2	700	105	210	Burning or slashing may begin 30 days after treatment following brown-off. Wiper application to be carried out in November - May.
Ring fern	0.7-2.1	210	35	[illegible]	For effective control, apply 2.1 litres/ha. For pasture manipulation, apply 0.7-1.4 litres/ha from December - April. Use higher rates for dense infestations. Burn 4-6 weeks after treatment.
Ryegrass (annual Italian short rotation hybrids)	[illegible]	[illegible]	[illegible]	[illegible]	Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	WHEN TO APPLY
Established < 8 months	2.1-2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Established > 8 months	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Ryegrass (Perennial)	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20 mL		Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Sweet brier	8.5 + organosili-cone penetrant 250mL/ 100L	700 + organosili-cone penetrant 200mL/ 100L	105 + organosili-cone penetrant 20 mL	-	Spray when plant is in late flowering to early leaf yellowing stage. Ensure full coverage.
Tall fescue	4.2	700	105	210	Apply January - March to active growth.
Willow	6.3 + organosili-cone penetrant 250 mL/100L	700 + organosili-cone penetrant 250 mL/100L	-	-	Apply before leaf fall begins (January - April). Total spray coverage is essential for best results.
Yarrow	2.1 -4.2	-	-	-	Apply to plant with 5-10cm of leaf. For best results cultivate 10 days after treatment. Use the low rate when a crop to be drilled is competitive to yarrow. Use higher rate when it is a poor competitor.

Add the Slikka™ wetter and penetrant at the suppliers recommended rates.
Slikka™ is a trade name of UPL Ltd CROP SOLUTIONS.

Registered pursuant to the ACVM Act 1997, No. P9580. See www.foodsafety.qovt.nz for registration conditions. Approved pursuant to the HSNO Act 1996, No. HSR101103. See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

300X200 (200 L)

Pantone 7485C

Pantone 368C

ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Blue AG

Stinger 510 SL

HERBICIDE

GROUP 9 HERBICIDE

ACTIVE INGREDIENT

510 g/litre Glyphosate

as the isopropylamine salt in the form of a soluble concentrate.

A non-selective herbicide for the control of many annual and perennial weeds in a wide range of situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS) 0800 CHEMCALL
OR 0800 243 622 OR PGG WRIGHTSON LTD 0800 10 22 76

200 L
NET CONTENTS

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

300X200 (200 L)

Pantone 425C

BLACK
C:60 M:40 Y:40 K:100

Pantone 368C

Pantone 485 C

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

PEEL OFF

HAZARD CLASSIFICATION: Hazardous to the aquatic environment (chronic) Category 2.

ECOTOXICITY: Toxic to some aquatic plants and with long-lasting effects.

QUALIFIED HANDLER: Required when applied onto or into water and the water has the potential to leave the place containing the application area.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure cool, dry place out of direct sunlight and away from seed, fertilizers, feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT: Apply with well-maintained and calibrated spray equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409:2004)

DISPOSAL INFORMATION

CONTAINER DISPOSAL: Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

ENVIRONMENT: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur. If accidental spraying occurs, immediately wash spray off foliage with plenty of water to minimise damage.

SPILLAGE: Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil or sand, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfil.

FIRST AID: If swallowed do not induce vomiting. Rinse mouth. For advice, call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor. If in eyes rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a POISON CENTRE or doctor if irritation persists. If skin or hair contact occurs, remove contaminated clothing and wash with running water. Get advice if irritation persists.

COMPATIBILITY: STINGER 510SL is compatible with commonly used herbicides.

GENERAL INFORMATION

STINGER 510SL is a non-selective glyphosate herbicide for use in agriculture and horticulture.

- Before planting crops and trees
- In pasture renovation and reduced tillage systems
- In vineyards, orchards and market gardens
- For pre-harvest weed control
- On turf and recreational areas
- In drains and waterways.

STINGER 510SL does not have adverse ecotoxic effects on fish or crustacea. **STINGER 510SL** is water soluble and is absorbed by green plant tissue. It is then translocated throughout the plant, including the root system, to kill the entire plant. Plants should be actively growing, and not under drought stress, with clean foliage at the time of spraying. The product degrades in the soil and has no residual weed control activity. Rainfall within 2 hours of application may reduce effectiveness. Application to foliage which is already wet will increase the time required for rainfastness. Visible effects may not be apparent for 14 days depending on weather conditions. Some perennial weeds may not show effects for 2 - 3 weeks.

DIRECTIONS FOR USE

APPLICATION METHODS: STINGER 510SL must only be placed in equipment made from stainless steel, fibreglass, plastic, aluminium, brass or copper. Other types of container may cause a reaction in which flammable hydrogen gas is generated. All spray equipment including pumps, spray tanks, lines and nozzles, and landing gear of aircraft should be thoroughly washed with water after each spraying day.

BOOM EQUIPMENT: Application can be made in spray volumes up to 200 litres/ha using fan nozzles at pressures in the range of 200-280kPa. Boom height should be set so that a double overlap of nozzle pattern is achieved at the top of the target plants.

AERIAL EQUIPMENT: Applicable to forest and non-crop areas only. Can be applied by aircraft in spray volumes of up to 100 litres/ha. Increase spray volumes in hilly terrain to ensure maximum target contact. Do not use in intensive horticultural cropping areas.

HANDGUN AND KNAPSACK: Ensure a fine spray pattern to give complete coverage. For handguns, use pressures of 1000 2000 kPa and No. 4 and No. 6 spray tips.

WIPER EQUIPMENT: Pre-mix 1 part STINGER 510SL in 3 parts clean water to make a 25% solution. Mix only enough for immediate use. Plants must be clean and dry and wiped fully saturated. Apply up and down slopes for effective applications.

MISTBLOWER: Refer to table.

MIXING: Fill the clean spray tank with half the required amount of clean water, add **STINGER 510SL**as required and mix well before adding the remaining water. If adjuvant is required, add this last to minimise foaming. Agitate well before spraying. When mixing with other chemicals, add **STINGER 510SL** after other water- soluble concentrates, but before suspension concentrates. Soil particles and hard water containing calcium salts deactivate **STINGER 510SL.** If hard water cannot be avoided, the addition of a pH buffer is recommended.

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

MINISTRY PRIMARY INDUSTRIES NEW ZEALAND
19/11/2025

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

9 415251 004638 >

300X200 (200 L)

Pantone 425C

MULTIFOLD®

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS: Wheat, barley, oat grain- 7 days

ANIMAL SAFETY: Poisonous plants, such as ragwort, may become more palatable to stock after spraying, and livestock should be kept away until plants have died down. Otherwise, withholding is not required for stock grazing. However, it is recommended that grazing is delayed for 3 days for annual weeds, and 7 days for perennial weeds. This will allow **STINGER 510SL** to penetrate the weeds for maximum effectiveness.

WEED CONTROL NOTES

AGRICULTURAL AREAS: May be used for the control of annual and perennial weeds as indicated in the Weeds Controlled Table prior to sowing any crop. **ASPARAGUS:** Pre-crop emergence and post-harvest applications. Apply prior to planting or crop emergence. Do not treat emerging spears. Application can also be made after removal of all spears but before crop regrowth occurs. In established ferns, apply as a directed or shielded spray or with wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plant. **BROAD-ACRE CROPS FOR WEED CONTROL:** Wheat, barley, oats. Can be used to control weeds prior to cultivation or prior to harvest. Not to be used as preharvest desiccant. Spray 7-12 days prior to harvest when the grain moisture is less than 30%. Do not harvest within 7 days after spraying. Do not spray undersown crops. Do not spray crops that will be harvested for seed. Do not apply more than 2.8 L./ha. Avoid treating plants under drought stress. **BRUSHWEEDS:** Refer to table. **DRAINS AND WATERWAYS:** For best results ensure that 75% of the weed to be sprayed is above water level. Avoid treating silt-laden weeds. Following spraying do not submerge the weed, as this will cause the herbicide to be washed off. **FORESTRY:** For the control of grasses and weeds as listed in the table. Can be applied for weed control prior to burning or planting of forest or nursery or around already established trees. Do not allow spray drift or spray contact with foliage or green bark of desirable trees. **MARKET GARDENS:** Apply before or after seeding but prior to crop emergence. Spraying can also be made prior to transplanting or as a post-harvest clean up. **MINIMUM TILLAGE:** For total weed control prior to seeding in a reduced tillage system. Weeds should be actively growing and not under stress from frost, drought or waterlogging. For the improved control of white clover, tank mix with a dicamba or tribenuron-methyl formulation at approved label rates. Do within 28 days of applying dicamba. If using with tribenuron-methyl then do not sow crops within 14 days of application, or sow brassicas if the soil pH exceeds 6.5 or use within 2 months after applying lime. Consult partner product label for further directions. Cultivation may start from 3 days after spraying annual weeds, and 7 days after spraying perennial weeds. Cultivation should occur within 21 days after spraying. Do not sow until a satisfactory seedbed is formed. **NON-AGRICULTURAL AREAS:** Treat specific problem weeds as indicated in the Weeds Controlled Table. **ORCHARDS/VINES:** Crops: citrus, apple, pear, stonefruit, kiwifruit, grapes. Apply as a directed or shielded spray. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of the stonefruit tree, or green bark, suckers, buds, foliage or fruit of other trees. Spray before crops bud, sucker or bloom. **PASTURE:** May be applied as a spot application to control a wide range of weeds in pasture. It is nonselective and may damage or kill any plant in the spray area. **RECREATIONAL AREAS:** Lawns, turf, parks, reserves and ornamentals. Apply before the sowing and planting of these areas, or use as a spot application. In mown areas, allow sufficient regrowth before spraying. Can also be sprayed around the base of established trees or shrubs, as a directed or shielded spray. Avoid contact with any part of the shrub or tree. **RESISTANCE MANAGEMENT: STINGER 510SL** contains the GROUP 9 herbicide glyphosate. Rotate or alternate herbicide types during or between seasons to avoid resistance in some weed species. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult http://resistance.nzpps.org for alternative modes of action and details of resistance management strategies (if any).

WEEDS CONTROLLED TABLE

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	WHEN TO APPLY
Annual weeds	0.7-1.4	140	20	40	Use the higher rate for larger weeds.
Australian sedge	4.2 - 6.3 (aerial)	700	105	210	Apply from November to March. Burn or slash from 30 days after treatment.
Barberry	-	700	105	210	Apply from flowering to late fruit stage (January - April). Regrowth should be 1,5m tall before spraying is carried out.
Barley grass	0.7	140	20	40	Apply when plants are at early seed head stage. (September - December).
Blackberry	6.3-14	1060	105	210	Apply from flowering-fruiting stage (December - April), before frosts occur. For hard to kill blackberry, aerial application, or no follow-up action planned, use the higher rate. Allow 6-10 weeks before slashing or burning.
Boxthorn	-	880	105	-	Apply to actively growing plants (November - April). Ensure complete coverage of dense thickets. Do not use extra surfactant, or treat salt damaged bushes.
Bracken/ broom	6.3 + organosili-cone penetrant 250 mL/ 100 L	700 + organosili-cone penetrant 250 mL/ 100 L	105 + organosili-cone penetrant 20 mL	210 + organosili-cone penetrant 20 mL	Apply after bracken fronds have unfurled (mid-Jan-June), but before frosts occur. For ground application to broom anytime, but for aerial application when broom is in full leaf.
Browntop	4.2	700	105	210	Apply when plants are 5-10cm high. For taller plants, qraze or burn and allow regrowth of 5-10 cm height before treatment.
Buddleia	6.3	700	105	-	Apply using handgun anytime. Aerialtreat buddleja up to 2 metres high (February-April) using spray volume of 200-300 litres/ha.
Californian thistle	2.8	700	105	210	Apply after flower bud development. Retreatment and/or cropping or pasture improvement may be necessary to restrict reestablishment from seed. Mowing thistles 10 days after spraying may assist control.
Cleavers	1	350	50	105	Spray actively growing plants after most reach early seedhead.
Couch	2.1-4.2	700	105	210	For suppression, apply 2.1 litres/ha in spring and repeat in autumn. For best results, apply 2.8 - 4.2 litres/ha in autumn to active growth of at least 10cm where no previous cultivation has occurred for at least 6 months.
Docks	2.8-6.3	700	105	-	Spray plants in spring or summer when actively growing and before seedheads have formed. Use the lower rate for seedlings and the higher rate for mature plants. In pasture situations boom spray prior to reseeding or establishing crops to restrict seedling reestablishment. Under trees or vines may be tank-mixed with Gesatop 500FW for residual seedling control.
Floating sweet grass	4.2	700	105	210	Spray actively growing plants after most reach early seedhead. Spraying plants under drought stress may give reduced results.
Gorse	-	700 + organosili-cone penetrant 200 mL/ 100 L	-	-	Apply anytime ensuring complete coverage of plants not stressed by drought, grazing or previous herbicide treatment.
Indian doab	6.3	700	105	-	Apply at early seedhead stage before frosts occur.
Johnson grass	4.2	700	105	-	Apply to actively growing plants.
Kikuyu/ Mercer grass	4.2	700	105	210	Apply at early seedhead stage. On dense infestations, cut and spray the regrowth.
Nodding thistle	-	700	105	-	When applying by wiper ensure plants are actively growing (October - April) when at least 10cm of the plant can be wiped. A double pass is recommended and follow-up with spot treatments if necessary.
Old man's beard	3.5	700-1400	210	-	Apply Nov-March prior to first leaf browning. Use higher rates on stems over 1cm in diameter. If complete coverage is not possible plants should be cut at ground level in winter and the regrowth sprayed with the higher rate when plants are at least 50cm lonq (March).
Pampas grass/toetoe	-	700	105	-	Spray Spring-Autumn. Reduced control may result if plants are in flower at the time of spraying. Ensure complete coverage.
Paspalum	2.8-4.2	700	105	210	Spray at low rate if plants are at early seedhead stage. Use 4.2 litres/ha on mature vegetative plants, before onset of frosts.
Potato (self set)	2.1	350	55	105	Spray actively growing plants after most reach early seedhead.
Prairie grass	1	140	20	-	Apply at early seedhead stage.
Ragwort	4.2	700	105	-	Spray when actively growing and ensure there is sufficient leaf area for herbicide uptake. When applying by wiper use on plants 10-15cm above the pasture. For best results, a double pass is recommended with a follow-up spot treatment if necessary. Caution: Ragwort becomes more palatable after treatment. Keep stock away from treated areas until plants brown-off.
Ratstail	3.5	700	105	-	Apply at a spray volume of 135 litres/ ha in autumn. In dry conditions, delay sprayinq until sufficient qreen foliage is present.
Rautahi (cutty grass)	4.2	700	105	210	Spray to actively growing plants from January - March. Burning or slashing may begin 30 days after treatment following brown-off.
Rushes	4.2	700	105	210	Burning or slashing may begin 30 days after treatment following brown-off. Wiper application to be carried out in November - May.
Ring fern	0.7-2.1	210	35	-	For effective control, apply 2.1 litres/ha. For pasture manipulation, apply 0.7-1.4 litres/ha from December - April. Use higher rates for dense infestations. Burn 4-6 weeks after treatment.
Ryegrass (annual Italian short rotation hybrids)					Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Established < 8 months	2.1-2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Established > 8 months	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Ryegrass (Perennial)	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20 mL		Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Sweet brier	8.5 + organosili-cone penetrant 250mL/ 100L	700 + organosili-cone penetrant 200mL/ 100L	105 + organosili-cone penetrant 20 mL	-	Spray when plant is in late flowering to early leaf yellowing stage. Ensure full coverage.
Tall fescue	4.2	700	105	210	Apply January - March to active growth.
Willow	6.3 + organosili-cone penetrant 250 mL/100L	700 + organosili-cone penetrant 250 mL/100L	-	-	Apply before leaf fall begins (January - April). Total spray coverage is essential for best results.
Yarrow	2.1 -4.2	-	-	-	Apply to plant with 5-10cm of leaf. For best results cultivate 10 days after treatment. Use the low rate when a crop to be drilled is competitive to yarrow. Use higher rate when it is a poor competitor.

Add the Slikka™ wetter and penetrant at the suppliers recommended rates.
Slikka™ is a trade name of UPL Ltd CROP SOLUTIONS.

Registered pursuant to the ACVM Act 1997, No. P9580. See www.foodsafety.qovt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR101103. See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

300X200 (1000 L)

Pantone 7485C

Pantone 368C

ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Blue AG

Stinger 510 SL

HERBICIDE

GROUP 9 HERBICIDE

ACTIVE INGREDIENT

510 g/litre Glyphosate

as the isopropylamine salt in the form of a soluble concentrate.

A non-selective herbicide for the control of many annual and perennial weeds in a wide range of situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS) 0800 CHEMCALL
OR 0800 243 622 OR PGG WRIGHTSON LTD 0800 10 22 76

1000 L
NET CONTENTS

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

300X200 (1000 L)

Pantone 425C

BLACK
C:60 M:40 Y:40 K:100

Pantone 368C

Pantone 485 C

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

PEEL OFF

HAZARD CLASSIFICATION: Hazardous to the aquatic environment (chronic) Category 2.

ECOTOXICITY: Toxic to some aquatic plants and with long-lasting effects.

QUALIFIED HANDLER: Required when applied onto or into water and the water has the potential to leave the place containing the application area.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure cool, dry place out of direct sunlight and away from seed, fertilizers, feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

EQUIPMENT: Apply with well-maintained and calibrated spray equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409:2004)

DISPOSAL INFORMATION
CONTAINER DISPOSAL: Triple rinse empty container and add residue to spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise, crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

ENVIRONMENT: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur. If accidental spraying occurs, immediately wash spray off foliage with plenty of water to minimise damage.

SPILLAGE: Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil or sand, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfil.

FIRST AID: If swallowed do not induce vomiting. Rinse mouth. For advice, call the National Poisons Centre 0800 POISON (0800 764 766) (24 hrs) or a doctor. If in eyes rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a POISON CENTRE or doctor if irritation persists. If skin or hair contact occurs, remove contaminated clothing and wash with running water. Get advice if irritation persists.

COMPATIBILITY: STINGER 510SL is compatible with commonly used herbicides.

GENERAL INFORMATION
STINGER 510SL is a non-selective glyphosate herbicide for use in agriculture and horticulture.
- Before planting crops and trees
- In pasture renovation and reduced tillage systems
- In vineyards, orchards and market gardens

STINGER 510SL does not have adverse ecotoxic effects on fish or crustacea. **STINGER 510SL** is water soluble and is absorbed by green plant tissue. It is then translocated throughout the plant, including the root system, to kill the entire plant. Plants should be actively growing, and not under drought stress, with clean foliage at the time of spraying. The product degrades in the soil and has no residual weed control activity. Rainfall within 2 hours of application may reduce effectiveness. Application to foliage which is already wet will increase the time required for rainfastness. Visible effects may not be apparent for 14 days depending on weather conditions. Some perennial weeds may not show effects for 2 - 3 weeks.

DIRECTIONS FOR USE
APPLICATION METHODS: STINGER 510SL must only be placed in equipment made from stainless steel, fibreglass, plastic, aluminium, brass or copper. Other types of container may cause a reaction in which flammable hydrogen gas is generated. All spray equipment including pumps, spray tanks, lines and nozzles, and landing gear of aircraft should be thoroughly washed with water after each spraying day.
BOOM EQUIPMENT: Application can be made in spray volumes up to 200 litres/ha using fan nozzles at pressures in the range of 200-280kPa. Boom height should be set so that a double overlap of nozzle pattern is achieved at the top of the target plants.
AERIAL EQUIPMENT: Applicable to forest and non-crop areas only. Can be applied by aircraft in spray volumes of up to 100 litres/ha. Increase spray volumes in hilly terrain to ensure maximum target contact. Do not use in intensive horticultural cropping areas.
HANDGUN AND KNAPSACK: Ensure a fine spray pattern to give complete coverage. For handguns, use pressures of 1000 2000 kPa and No. 4 and No. 6 spray tips.
WIPER EQUIPMENT: Pre-mix 1 part STINGER 510SL in 3 parts clean water to make a 25% solution. Mix only enough for immediate use. Plants must be clean and dry and the wiper fully saturated. Apply up and down slopes for effective applications.
MISTBLOWER: Refer to table.

MIXING: Fill the clean spray tank with half the required amount of clean water. Add **STINGER 510SL**as required and mix well before adding the remaining water. If surfactant is required, add this last to minimise foaming. Agitate well before spraying. When tank-mixing with other chemicals, add **STINGER 510SL** after other water- soluble concentrates, but before suspension concentrates. Soil particles and hard water containing calcium salts deactivate **STINGER 510SL.** If hard water cannot be avoided, the addition of a pH buffer is recommended.

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

9 415251 005

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025

300X200 (1000 L)

Pantone 425C

MULTIFOLD®

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS: Wheat, barley, oat grain- 7 days

ANIMAL SAFETY: Poisonous plants, such as ragwort, may become more palatable to stock after spraying, and livestock should be kept away until plants have died down. Otherwise, withholding is not required for stock grazing. However, it is recommended that grazing is delayed for 3 days for annual weeds, and 7 days for perennial weeds. This will allow **STINGER 510SL** to penetrate the weeds for maximum effectiveness.

WEED CONTROL NOTES

AGRICULTURAL AREAS: May be used for the control of annual and perennial weeds as indicated in the Weeds Controlled Table prior to sowing any crop. **ASPARAGUS:** Pre-crop emergence and post-harvest applications. Apply prior to planting or crop emergence. Do not treat emerging spears. Application can also be made after removal of all spears but before crop regrowth occurs. In established ferns, apply as a directed or shielded spray or with wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plant. **BROAD-ACRE CROPS FOR WEED CONTROL:** Wheat, barley, oats. Can be used to control weeds prior to cultivation or prior to harvest. Not to be used as preharvest desiccant. Spray 7-12 days prior to harvest when the grain moisture is less than 30%. Do not harvest within 7 days after spraying. Do not spray undersown crops. Do not spray crops that will be harvested for seed. Do not apply more than 2.8 L./ha. Avoid treating plants under drought stress. **BRUSHWEEDS:** Refer to table. **DRAINS AND WATERWAYS:** For best results ensure that 75% of the weed to be sprayed is above water level. Avoid treating silt-laden weeds. Following spraying do not submerge the weed, as this will cause the herbicide to be washed off. **FORESTRY:** For the control of grasses and weeds as listed in the table. Can be applied for weed control prior to burning or planting of forest or nursery or around already established trees. Do not allow spray drift or spray contact with foliage or green bark of desirable trees. **MARKET GARDENS:** Apply before or after seeding but prior to crop emergence. Spraying can also be made prior to transplanting or as a post-harvest clean up. **MINIMUM TILLAGE:** For total weed control prior to seeding in a reduced tillage system. Weeds should be actively growing and not under stress from frost, drought or waterlogging. For the improved control of white clover, tank mix with a dicamba or tribenuron-methyl formulation at approved label rates. Do within 28 days of applying dicamba. If using with tribenuron-methyl then do not sow crops within 14 days of application, or sow brassicas if the soil pH exceeds 6.5 or use within 2 months after applying lime. Consult partner product label for further directions. Cultivation may start from 3 days after spraying annual weeds, and 7 days after spraying perennial weeds. Cultivation should occur within 21 days after spraying. Do not sow until a satisfactory seedbed is formed. **NON-AGRICULTURAL AREAS:** Treat specific problem weeds as indicated in the Weeds Controlled Table. **ORCHARDS/VINES:** Crops: citrus, apple, pear, stonefruit, kiwifruit, grapes. Apply as a directed or shielded spray. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of the stonefruit tree, or green bark, suckers, buds, foliage or fruit of other trees. Spray before crops bud, sucker or bloom. **PASTURE:** May be applied as a spot application to control a wide range of weeds in pasture. It is nonselective and may damage or kill any plant in the spray area. **RECREATIONAL AREAS:** Lawns, turf, parks, reserves and ornamentals. Apply before the sowing and planting of these areas, or use as a spot application. In mown areas, allow sufficient regrowth before spraying. Can also be sprayed around the base of established trees or shrubs, as a directed or shielded spray. Avoid contact with any part of the shrub or tree. **RESISTANCE MANAGEMENT: STINGER 510SL** contains the GROUP 9 herbicide glyphosate. Rotate or alternate herbicide types during or between seasons to avoid resistance in some weed species. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs. To prevent or delay the development of resistance, use in tank mixes and/or rotate use with herbicides with a different mode of action. Consult http://resistance.nzpps.org for alternative modes of action and details of resistance management strategies (if any).

WEEDS CONTROLLED TABLE

WEED	BOOM/ AERIAL L/HA	HANDGUN MLS/ 100 LITRES	KNAPSACK MLS/15 LITRES	MISTBLOWER MLS/10 LITRES	WHEN TO APPLY
Annual weeds	0.7-1.4	140	20	40	Use the higher rate for larger weeds.
Australian sedge	4.2 - 6.3 (aerial)	700	105	210	Apply from November to March. Burn or slash from 30 days after treatment.
Barberry	-	700	105	210	Apply from flowering to late fruit stage (January - April). Regrowth should be 1,5m tall before spraying is carried out.
Barley grass	0.7	140	20	40	Apply when plants are at early seed head stage. (September - December).
Blackberry	6.3-14	1060	105	210	Apply from flowering-fruiting stage (December - April), before frosts occur. For hard to kill blackberry, aerial application, or no follow-up action planned, use the higher rate. Allow 6-10 weeks before slashing or burning.
Boxthorn	-	880	105	-	Apply to actively growing plants (November - April). Ensure complete coverage of dense thickets. Do not use extra surfactant, or treat salt damaged bushes.
Bracken/ broom	6.3 + organosili-cone penetrant 250 mL/ 100 L	700 + organosili-cone penetrant 250 mL/ 100 L	105 + organosili-cone penetrant 20 mL	210 + organosili-cone penetrant 20 mL	Apply after bracken fronds have unfurled (mid-Jan-June), but before frosts occur. For ground application to broom anytime, but for aerial application when broom is in full leaf.
Browntop	4.2	700	105	210	Apply when plants are 5-10cm high. For taller plants, qraze or burn and allow regrowth of 5-10 cm height before treatment.
Buddleia	6.3	700	105	-	Apply using handgun anytime. Aerialtreat buddleja up to 2 metres high (February-April) using spray volume of 200-300 litres/ha.
Californian thistle	2.8	700	105	210	Apply after flower bud development. Retreatment and/or cropping or pasture improvement may be necessary to restrict reestablishment from seed. Mowing thistles 10 days after spraying may assist control.
Cleavers	1	350	50	105	Spray actively growing plants after most reach early seedhead.
Couch	2.1-4.2	700	105	210	For suppression, apply 2.1 litres/ha in spring and repeat in autumn. For best results, apply 2.8 - 4.2 litres/ha in autumn to active growth of at least 10cm where no previous cultivation has occurred for at least 6 months.
Docks	2.8-6.3	700	105	-	Spray plants in spring or summer when actively growing and before seedheads have formed. Use the lower rate for seedlings and the higher rate for mature plants. In pasture situations boom spray prior to reseeding or establishing crops to restrict seedling reestablishment. Under trees or vines may be tank-mixed with Gesatop 500FW for residual seedling control.
Floating sweet grass	4.2	700	105	210	Spray actively growing plants after most reach early seedhead. Spraying plants under drought stress may give reduced results.
Gorse	-	700 + organosili-cone penetrant 200 mL/ 100 L	-	-	Apply anytime ensuring complete coverage of plants not stressed by drought, grazing or previous herbicide treatment.
Indian doab	6.3	700	105	-	Apply at early seedhead stage before frosts occur.
Johnson grass	4.2	700	105	-	Apply to actively growing plants.
Kikuyu/ Mercer grass	4.2	700	105	210	Apply at early seedhead stage. On dense infestations, cut and spray the regrowth.
Nodding thistle	-	700	105	-	When applying by wiper ensure plants are actively growing (October - April) when at least 10cm of the plant can be wiped. A double pass is recommended and follow-up with spot treatments if necessary.
Old man's beard	3.5	700-1400	210	-	Apply Nov-March prior to first leaf browning. Use higher rates on stems over 1cm in diameter. If complete coverage is not possible plants should be cut at ground level in winter and the regrowth sprayed with the higher rate when plants are at least 50cm lonq (March).
Pampas grass/toetoe	-	700	105	-	Spray Spring-Autumn. Reduced control may result if plants are in flower at the time of spraying. Ensure complete coverage.
Paspalum	2.8-4.2	700	105	210	Spray at low rate if plants are at early seedhead stage. Use 4.2 litres/ha on mature vegetative plants, before onset of frosts.
Potato (self set)	2.1	350	55	105	Spray actively growing plants after most reach early seedhead.
Prairie grass	1	140	20	-	Apply at early seedhead stage.
Ragwort	4.2	700	105	-	Spray when actively growing and ensure there is sufficient leaf area for herbicide uptake. When applying by wiper use on plants 10-15cm above the pasture. For best results, a double pass is recommended with a follow-up spot treatment if necessary. Caution: Ragwort becomes more palatable after treatment. Keep stock away from treated areas until plants brown-off.
Ratstail	3.5	700	105	-	Apply at a spray volume of 135 litres/ ha in autumn. In dry conditions, delay sprayinq until sufficient qreen foliage is present.
Rautahi (cutty grass)	4.2	700	105	210	Spray to actively growing plants from January - March. Burning or slashing may begin 30 days after treatment following brown-off.
Rushes	4.2	700	105	210	Burning or slashing may begin 30 days after treatment following brown-off. Wiper application to be carried out in November - May.
Ring fern	0.7-2.1	210	35	-	For effective control, apply 2.1 litres/ha. For pasture manipulation, apply 0.7-1.4 litres/ha from December - April. Use higher rates for dense infestations. Burn 4-6 weeks after treatment.
Ryegrass (annual Italian short rotation hybrids)					Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Established < 8 months	2.1-2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Established > 8 months	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20mL/ 100L	210 + organosili-cone penetrant 20mL/ 100L	
Ryegrass (Perennial)	2.8 + organosili-cone penetrant 100mL/ 100L	700 + organosili-cone penetrant 100mL/ 100L	105 + organosili-cone penetrant 20 mL		Spray to actively growing plants with at least 5 -10cm of green leaf. A higher rate may be necessary that reflects the weed spectrum present.
Sweet brier	8.5 + organosili-cone penetrant 250mL/ 100L	700 + organosili-cone penetrant 200mL/ 100L	105 + organosili-cone penetrant 20 mL	-	Spray when plant is in late flowering to early leaf yellowing stage. Ensure full coverage.
Tall fescue	4.2	700	105	210	Apply January - March to active growth.
Willow	6.3 + organosili-cone penetrant 250 mL/100L	700 + organosili-cone penetrant 250 mL/100L	-	-	Apply before leaf fall begins (January - April). Total spray coverage is essential for best results.
Yarrow	2.1 -4.2	-	-	-	Apply to plant with 5-10cm of leaf. For best results cultivate 10 days after treatment. Use the low rate when a crop to be drilled is competitive to yarrow. Use higher rate when it is a poor competitor.

Add the Slikka™ wetter and penetrant at the suppliers recommended rates.
Slikka™ is a trade name of UPL Ltd CROP SOLUTIONS.

Registered pursuant to the ACVM Act 1997, No. P9580. See www.foodsafety.qovt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR101103. See www.epa.govt.nz for approval controls.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
19/11/2025