DANGER

KEEP OUT OF REACH OF CHILDREN

BAMMER®

HERBICIDE

For broad spectrum weed control in orchards, vineyards, waste areas and for seedbed preparation pre-sowing of crops

Active ingredient
Contains 150 g/L GLUFOSINATE AMMONIUM and
4.5 g/L CARFENTRAZONE ETHYL in the form of an oil in water emulsion

GROUP	10 & 14	HERBICIDE

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
09/09/2025

AGRECOVERY FOUNDING BRAND

Net contents: 1 L, 4 L, 5 L, 10 L, 20 L

Bar code to be inserted

Public

PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of registration.

Any person purchasing, handling or disposing of this product must be suitably qualified.

DANGER
Causes serious eye damage.
Suspected of damaging fertility or the unborn child.
Causes damage to liver from repeated oral exposure at high doses.
Harmful to aquatic life with long lasting effects.
Hazardous to soil organisms.
Hazardous to terrestrial vertebrates.

PRECAUTIONARY STATEMENTS
PREVENTION
Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
Wear protective gloves, protective clothing and eye protection.
Do not breathe spray.
Do not apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.
RESPONSE
Wash hands and face thoroughly after handling.
Do not eat, drink, or smoke when using this product.
Records of use must be kept.
STORAGE
Store locked up.
Store in a cool dry place avoiding temperatures above 35°C.
DISPOSAL
Dispose of contents/container in accordance with local regulations.
Container disposal: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery® (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID
If medical advice is needed, have the product container or label at hand. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Immediately call a doctor or the National Poisons Centre 0800 POISON (0800 764 766).
IF exposed or concerned: Get medical advice.
Get medical advice if you feel unwell.

SPILLAGE
Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

Label – Back

SHELF LIFE
When stored appropriately, this product should show no significant degradation for two years from the date of manufacture. When using this product beyond this shelf life, contact UPL New Zealand Limited for further information.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at **www.uplcorp.com/nz/**

EXCLUSION OF LIABILITY: PLEASE READ
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered to and distributed by:
UPL New Zealand Ltd
PO Box 51584
Pakuranga 2140
Auckland
Phone: 0800 100 325
www.uplcorp.com/nz/
Registered pursuant to the ACVM Act 1997, No. P009722.
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval Code HSR101379.
See www.epa.govt.nz for approval controls.

© Copyright UPL New Zealand Limited 2020, all rights reserved.
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited.
® Registered trademark

Batch number:	
Date of Manufacture:	

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622)

UN-Free

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
09/09/2025

GENERAL INFORMATION

BAMMER is an herbicide that contains two active ingredients: glufosinate ammonium and carfentrazone-ethyl.

Glufosinate ammonium is taken into plants through foliage and green stems. Once inside the plant it prevents the plant from metabolising ammonia that is a by-product of photosynthesis. Ammonia accumulates in the plant and eventually causes death. Plants are more susceptible when growing actively.

Carfentrazone-ethyl is a fast-acting contact herbicide that desiccates and controls weeds through a process of membrane disruption. It is rapidly absorbed through the foliage and green stems of plants.

Neither active ingredient provides residual weed control. Regrowth of more robust weed growth is likely to occur after some weeks following treatment. An alternative herbicide or mixture may be required for repeat treatment. The addition of a drift control agent like **LI-1000®** or **UNISON** is recommended.

RESISTANCE MANAGEMENT

This product contains glufosinate-ammonium, a **Group 10** herbicide, that has a mode of action that inhibits glutamine synthetase and carfentrazone-ethyl, a **Group 14** herbicide, that causes inhibition of protoporphyrinogen oxidase (PPO).

Continuous use of this product may eventually lead to a build-up of weeds resistant to it. This can be prevented or delayed by alternating with herbicides from other groups. Use strictly in accordance with label instructions.

RATE PER HA	WEEDS CONTROLLED
4–5 L	Annual poa, asparagus seedlings, beaked hawksbeard, black nightshade, calandrinia, chickweed, cleavers, clover seedlings, cornbind, daisy, dandelion, mouse-ear chickweed, musky storksbill, redroot, rough-stalked meadow grass, scotch thistle, sow thistle, spurrey, staggerweed, stinking mayweed, subterranean clover, summer grass, wild turnip, willow weed, wireweed
5–7.5 L	Bristle grass, bur medick, catsear, chewings fescue, clovers, cut-leaved geranium, docks, fleabane, parsley piert, prairie grass, ryegrasses*, sheep's sorrel, storksbill, strawberry, twin cress, yorkshire fog
7.5–9 L	Broom, buttercups*, couch, crowfoot grass, field pansy, gorse, hawksbeard, kikuyu grass, mallows, paspalum, tall fescue, wild strawberry, willow herb, yarrow
9 L	Bracken, plantains, rape, rushes, sedges

*Some ryegrasses and hairy buttercup are resistant.

IMPORTANT

Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Toxic weeds may become more palatable after spraying with partner products and **BAMMER** may concentrate this effect.

BAMMER MUST be applied using ground-based methods only.

BAMMER MUST NOT be applied when wind speeds are less than 3 km/hr or more than 20 km/hr.

When **BAMMER** is applied with a boom sprayer, the boom height should be kept as low as practicable to avoid spray drift, with a nozzle set to provide medium-coarse sized droplets. Small sized droplets are not approved.

The maximum application rate for **BAMMER** is 9 L/ha, with a maximum of 3 applications per calendar year and a minimum interval period of 10 days.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
09/09/2025

DIRECTIONS FOR USE

SITUATION	RATE	COMMENTS
Stale seed bed	4–5 L/ha in 100-200 L of water	Apply when weeds are small and before sowing crop. Allow 7 days before sowing shallow seeded crops such as onion, carrot and parsnip.
Non-crop land Boom spraying Hand gun application and motorised knapsack Airblast knapsack	4–9 L/ha in 300-400 L of water 1–2.5 L/100 L 100–250 mL/10 L 400–750 mL/ 10 L	Use the lower rate on weeds that are young, succulent and actively growing. Use the high rate on weeds that are mature or when the weed population is very dense and more difficult to obtain coverage on. Do not use airblast knapsack sprayers near desirable crops.
Orchards* Boom spraying Hand gun application and motorised knapsack	4–9 L/ha in 300-400 L of water 1– 2.5 L/100 L 100–250 mL/10 L	Apply rates as above relative to weed growth state.
	* May be used under pomefruit, citrus, grapes, vineyards, kiwifruit, and summerfruit.	

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

TANK CLEANING

BAMMER is a very sticky product that will adhere to tank walls and may therefore cause crop damage if not fully removed before using sprayers for selective weed control or the application of fungicides, insecticides etc.

After spraying with **BAMMER** add 500 mL **ALL-CLEAR®** per 100 L of water and circulate through the spray tank and all spray lines. Allow to stand for 15 min, then empty the tank by spraying out through the boom or gun.

Should small quantities of **BAMMER** remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to sensitive crops and other vegetation. To ensure residue and deposit removal allow **ALL-CLEAR** mixture to stand overnight in lines and tank. This applies particularly to aged and perished spray hoses and lines. This treatment can remove any residues of **BAMMER** in the spray rig.
Do not apply sprayer cleaning solutions or rinsate to sensitive crops.

WITHHOLDING PERIODS

All crops 28 days

® Registered trademarks

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
09/09/2025