WARNING

KEEP OUT OF REACH OF CHILDREN

Miravis® Prime

ADEPIDYN® technology Fungicide

syngenta®

Active Ingredients:
MIRAVIS® Prime contains 150 g/litre PYDIFLUMETOFEN (ADEPIDYN®) and 250 g/litre FLUDIOXONIL in the form of a suspension concentrate.

GROUP	7	12	FUNGICIDE

For the control of *Botrytis* in Grapes.

Net contents: 5, 10, 15 and 20 litres

Registered pursuant to the ACVM Act 1997 No. P9724
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code HSR101506.
See www.epa.govt.nz for approval controls

Syngenta Crop Protection Limited
Level 4, 60 Parnell Road, Parnell, Auckland 1052

[placeholder for internal codes]

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
02/12/2024

PLEASE READ THE LABEL COMPLETELY BEFORE USE
This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.
Any person mixing, loading or handling this product MUST be a qualified loader or under the guidance of a qualified loader.
Any person applying this product MUST be a qualified contractor/person or under the guidance of a qualified contractor/person.

GHS Classifications:
Specific Target Organ Toxicity- (repeated exposure) Category 2
Hazardous to the aquatic environment acute Category 1
Hazardous to the aquatic environment chronic Category 1
WARNING
May cause damage to liver through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONS
Store in original container, tightly closed, away from foodstuffs.
When mixing, loading or applying this product wear protective clothing including overalls, chemical resistant gloves, waterproof hat.
DO NOT breathe spray mist or vapours.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking and after work.
Wash protective clothing after use.
Apply with well maintained and calibrated spray equipment.
Remove protective clothing and wash hands and face thoroughly before meals and after using.
Records of use MUST be kept as per Clause 48 of the Hazardous Property Controls Notice 2017.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
DO NOT carry on a passenger service vehicle.
Avoid contamination of any water supply with chemical or empty container. This product MUST NOT be applied directly into or onto water.

Container Disposal: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.
Product Disposal: Disposal of this product only by using according to the label, or at an approved landfill or other approved facility.
For additional information refer to EPA Notice: Hazardous Substances (Disposal) Notice 2017

FIRST AID
Get medical advice or attention if you feel unwell.
For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

SPILLAGE
Collect spillage. Wear suitable protective clothing including face shield, chemical resistant gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
02/12/2024

GENERAL INFORMATION
MIRAVIS® Prime is a combination of two broad spectrum fungicides with protectant properties. MIRAVIS® Prime is rainfast within 1 hour.
MIRAVIS® Prime should not affect beneficial species including Parasitic Wasps and Predatory Mites.

RESISTANCE MANAGEMENT **GROUP 7 12 FUNGICIDE**
MIRAVIS® Prime contains the active ingredients Pydiflumetofen (ADEPIDYN®), from the SDHI (Succinate DeHydrogenase Inhibitor mode of action group (GROUP 7 FUNGICIDE) and Fludioxonil from the phenylpyrrole mode of action group (GROUP 12 FUNGICIDE).
Resistance to this fungicide could develop from repeated use. To minimise this risk use strictly in accordance with the label instructions and resistance management strategies. DO NOT use in conjunction with other products containing a GROUP 7 fungicide when adhering to the fungicide mixture and alternation guidelines for grapes. Some naturally occurring individual fungi strains resistant to MIRAVIS® Prime and other Group 7 and Group 12 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by MIRAVIS® Prime and other Group 7 and Group 12 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Crop Protection Ltd, New Zealand accepts no liability for any losses that may result from the failure of MIRAVIS® Prime to control resistant fungi. Refer to the NZ Plant Protection Society website (www.resistance.nzpps.org) for further details on resistance management.

GRAPES
Use a maximum of 1 application of MIRAVIS® Prime per season. MIRAVIS® Prime MUST be applied in alternation with another approved, non-cross resistant, fungicide recommended for the control of the target disease. DO NOT apply more than 2 Group 7 fungicides per season per crop.

DIRECTIONS FOR USE
Shake well before use
It is an offence to use this product on animals.

Crop	Disease	Rate	Critical Comments
Grapes	Botrytis bunch rot (Grey mould) (*Botrytis cinerea*) Application for Botrytis will also result in control of Powdery mildew (*Erysiphe nectar*)	100 mL/100L of water as a dilute spray	Use the industry accepted rate calculator to determine the hectare dosage. Apply MIRAVIS® Prime as part of a regular Botrytis program up to the start of 80% capfall (EL 25). Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of vine growth, vigour, and row spacing. The rate suggested is for high volume spraying to runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage of the inflorescences is achieved. MIRAVIS® Prime MUST NOT be applied to the same area more than once per season. If only powdery mildew is being targeted, an alternative product should be used at the recommended label rates and timings. **Note:** For export wine pre-harvest intervals consult your exporter or refer to the latest New Zealand Winegrowers Vineyard Spray Schedule.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
02/12/2024

EPA RESTRICTIONS

Application Rate
The maximum application rate of this substance is 1.3L of MIRAVIS® Prime/ha (200 g pydiflumetofen/ha and 333 g fludioxonil/ha), with a maximum of one application per year.

Application Method
DO NOT apply when wind speeds are less than 3 km/hr or more than 20km/hr as measured at the application site.

Apply with ground based equipment and, at a minimum, medium droplets, as defined by the American Society of Agricultural and Biological Engineers ASABE Standard (S572) or the British Crop Production Council guideline.

SPRAY DRIFT
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

POLLINATOR SAFETY
DO NOT apply this substance to plants if:
(a) Bees are foraging; or
(b) The plants are in flower or part flower and are likely to be visited by non-target invertebrate pollinators including bees.

MIXING
Shake well before use
Half-fill the spray tank with clean water and start agitation. Add the required amount of MIRAVIS® Prime and allow to disperse. Complete the filling of the spray tank with the rest of the water while continuing to agitate the mixture thoroughly. Maintain agitation until spraying is complete. DO NOT leave the spray mix in the sprayer overnight.

APPLICATION
Apply by ground-based methods only.
Thorough coverage is essential for effective results.

Application in vine crops
The application rate is for high volume spraying to the point of runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved.
Set up and operate the sprayer to achieve even coverage throughout the crop canopy, applying sufficient water to cover the crop to just prior to the point of runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

DO NOT apply in a dilute water volume greater than 1300L/ha.

COMPATIBILITY
MIRAVIS® Prime is compatible with commonly used fungicides and insecticides. Before using any tank mixture, consult and comply with the recommendations of the partner product. Each product should be added separately to the bulk of the water in the spray tank whilst filling and agitator working.

WITHHOLDING PERIODS
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds
For export crops – contact your exporter or Industry crop adviser.
Grapes ..... DO NOT apply after 80% capfall (EL 25)
Leaf plucking in vineyards
Sheep MUST NOT be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

MINISTER FOR PRIMARY INDUSTRIES
NEW ZEALAND
02/12/2024

STORAGE
Store in a dry, shaded area, avoiding temperatures below minus 10°C and above 35°C. See safety data sheet for additional storage information. Stores containing quantities of 100 litres or more require signage and secondary containment.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to Safety Data Sheet. For a copy visit our website at www.syngenta.co.nz

DISCLAIMER
This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Distributed by and registered to Syngenta Crop Protection Limited, Level 4, 60 Parnell Road, Parnell, Auckland 1052. Phone 0800 333 336.

Bar code
Agrecovery logo

Batch No.	
Date of Manufacture	

UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.
(fludioxonil and pydiflumetofen) MARINE POLLUTANT.
PACKING GROUP III, HAZCHEM 3Z
CONTAIN SPILLAGE

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only, call 0800 734 607 (24 hours).

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
02/12/2024