ECOTOXIC

WYNCA® SUNPHOSATE 540K HERBICIDE

A non-residual, non-selective herbicide for weed control prior to planting crops and pasture, prior to harvesting some crops and for general weed control in horticulture, agriculture, and forestry.

Active Ingredient: 540g/litre glyphosate as the potassium salt in the form of a soluble concentrate.

Net contents: 1- 1000 litres

(XINAN CHEMICAL LOGO)

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/07/2024

PLEASE READ THE LABEL COMPLETLY BEFORE USE

HAZARD CLASSIFICATION: Hazardous to the aquatic environment (chronic) Category 2.
ECOTOXIC: Toxic to aquatic life with long-lasting effects.
ENVIRONMENT: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Non-selective herbicide - toxic to plants. Avoid contact with foliage or green stems of fruit or crops, desirable plants and trees, since severe injury or death may occur.
Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Do not allow drift to occur outside the target area.
If accidental spraying occurs, immediately wash spray off foliage with plenty of water to minimise damage.
Do not apply concentrate product directly onto or into water. Follow label instructions.
QUALIFIED HANDLER: This product must be under the control of a suitably qualified handler when applied onto or into water and the water has the potential to leave the place containing the application area.
PRECAUTIONS:
PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.
STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure place away from feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements (Refer to Safety Note). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.
USE: Apply with well-maintained and calibrated spray equipment.
DISPOSAL: Dispose of this product only by using in accordance with this label, or at a suitable landfill. Triple rinse container and add rinsate to the spray tank. Dispose of container to a suitable landfill or agricultural recycling scheme. Do not burn. Do not use container for any other purpose.

First Aid
If swallowed do NOT induce vomiting. Rinse mouth. Get medical advice if you feel unwell.
If skin or hair contact occurs, remove contaminated clothing and wash with soap and water. Get medical advice if irritation persists.
If splashed in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention.
For advice, call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/07/2024

Emergency Response
In a transport emergency **call 111, Police or Fire Brigade**.

Spillage
Collect spill. Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil, sand, etc, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfill.

GENERAL INFORMATION
WYNCA SUNPHOSATE 540 K Herbicide is a broad spectrum, non-selective, non-residual herbicide that is not volatile and may be used in agriculture, horticulture, drains, aquatic areas, forestry and for other uses including general weed control. WYNCA SUNPHOSATE 540 K Herbicide is recommended prior to sowing pasture, crops and for the control of a wide range of brushweeds.
WYNCA SUNPHOSATE 540 K Herbicide is absorbed by plant foliage and green stems. It is inactivated immediately upon contact with the soil and does not provide residual weed control. WYNCA SUNPHOSATE 540 K Herbicide moves from the point of foliage or stem contact into the root system controlling the plant above and below the ground. Visible effects may take up to 14 days or even longer depending on growing conditions. These effects are a gradual wilting and yellowing of the foliage which advances to deterioration of above and below ground parts.

Apply WYNCA SUNPHOSATE 540 K Herbicide only to healthy, actively growing plants with adequate leaves for herbicide uptake. Avoid treating plants that are silt laden or affected by drought, frost, water logging, grazing or previous herbicide application.
Rainfall occurring up to 2 hours after application may reduce effectiveness. When Organosilicone is added the rainfree period may be reduced to 30 minutes. If WYNCA SUNPHOSATE 540 K Herbicide is applied to wet plants, efficacy may be reduced due to runoff.
With brushweeds wait for stem dehydration to occur prior to burning or slashing.

MIXING AND COMPATIBILITY:
WYNCA SUNPHOSATE 540 K Herbicide mixes readily with clean water. Reduced results may occur if dirty water (e.g. from ponds), hard water containing calcium salts, or water containing metal ions (e.g. iron) is used. Ensure the spray tank is free of any residue of previous spray materials. Half fill the spray tank with water. Add the required amount of WYNCA SUNPHOSATE 540 K Herbicide, agitate and continue filling the tank. Avoid foaming by keeping the filler nozzle below water level. Do not store mixed solutions for longer than 5 days. Do not use with 2,4-D Amine. If adding other herbicides, ensure these are fully dispersed before adding the WYNCA SUNPHOSATE 540 K Herbicide. Use tank mixtures promptly.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Withholding period: Wheat, barley, oat grain – 7 days

Animal safety: After application of WYNCA SUNPHOSATE 540 K Herbicide a withholding period for grazing stock is not required except where ragwort and other poisonous weeds are present.
Stock should be withheld from the treated area until these plants brown out. In all other situations it is recommended that grazing or cultivating is delayed for 1 day after treatment of annual weeds or 3 days if perennial weeds are present to ensure absorption of WYNCA SUNPHOSATE 540 K Herbicide.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/07/2024

USE RECOMMENDATIONS:

Refer to “Weeds Controlled” section for rates of application.

General Weed Control
Handgun - mix 660 mL of WYNCA SUNPHOSATE 540 K Herbicide with 100 litres of water (0.66% solution).
Knapsack - mix 100 mL of WYNCA SUNPHOSATE 540 K Herbicide with 15 litres of water.
Small sprayers - mix 6.5 mL of WYNCA SUNPHOSATE 540 K Herbicide with every 1 litre of water.
Apply to completely wet the foliage but not to run-off.

Brushweed Control
WYNCA SUNPHOSATE 540 K Herbicide may be used to control a wide range of brushweeds and woody plants.

Agricultural Areas
WYNCA SUNPHOSATE 540 K Herbicide may be used for the control of annual and perennial weeds as directed in agricultural land prior to sowing any edible or non-edible crop or pasture.
Where white clover control is required, tank mix with a suitable white clover herbicide and observe recommended waiting periods prior to planting legumes. Cultivation may start from 3 days after spraying perennial weeds and 1 day for annual weeds.

Pasture - Direct (Spot) Application
WYNCA SUNPHOSATE 540 K Herbicide is non-selective and may damage or kill any plant in the sprayed area. Apply as a directed low pressure spray. Retreatment and/or pasture improvement may be necessary to restrict seedling re-establishment.

Pasture - Selective Wiper Application
For the control of erect weeds in pastures. Pre-mix 1 part WYNCA SUNPHOSATE 540 K Herbicide in 3 parts clean water to make a 25% solution. Mix only enough for immediate use. Plants must be clean and dry and the wiper fully saturated. Apply up and down slopes for effective applications
.

Pasture Renewal
WYNCA SUNPHOSATE 540 K Herbicide may be used to control existing pasture species prior to re-seeding pasture or establishing crops.

Pasture Seedhead Suppression (Unimproved Pasture)
Use in unimproved pasture to delay onset of flowering and to maintain pasture grasses in a vegetative growth phase. Treat spring pasture at early grass seedhead emergence with 180–200ml WYNCA SUNPHOSATE 540 K

Herbicide /ha. Temporary pasture brownout will occur. Do not use Organosilicone or any other surfactant. Accurate spray calibration is important. Grazing can be carried out at any time after 3 days from treatment, but for optimum results graze from 21 days after treatment.

DO NOT USE ON RYEGRASS OR OTHER IMPROVED PASTURES AS YIELD SUPPRESSION MAY OCCUR. IT IS ADVISABLE NOT TO TREAT MORE THAN 10-15% OF THE FARM IN ANY ONE YEAR. DO NOT USE WHEN SOIL MOISTURE LEVELS ARE INADEQUATE FOR PASTURE GROWTH FOLLOWING APPLICATION.

Spraying of Silage Before Cutting

Where paddocks are to be used for cropping after silage, WYNCA SUNPHOSATE 540 K Herbicide may be applied before the silage is cut. Use 2.0-2.7 L/ha plus Organosilicone 100mL/100 litres of water in 100-200 litres of water per hectare. Use the high rate on perennial ryegrass or if the silage is rank or lodged. Refer to higher label rates on other specific weeds. After spraying wait at least 4 days before cutting. Grass quality is maintained for 7-10 days after the initial planned cutting date, even into the early stages of brown-out.

Spraying of Silage or Hay Paddocks After Cutting

Before spraying ensure that there is at least 5-10 cm of actively growing green leaf on all tillers. Typically this takes three weeks or more after cutting. Refer to application rates in the table. Always use Organosilicone.

Pre-Harvest Weed Control: Wheat, Barley, Oats

For wheat, barley and oats - apply 7-12 days prior to harvest when the grain moisture content is less than 30%. For preharvest weed control only. Do NOT use as a preharvest desiccant. Do not apply to undersown crops including clovers as they may be severely damaged. Do not apply to crops which will be harvested for seed. Use rates up to 2.67 litres/ha (1.44 kg ai/ha) depending on the specific weeds to be treated. Avoid treating drought stressed plants.

Orchards/Vines: Citrus, Apples, Pears, Stonefruit, Kiwifruit, Grapevines

Apply as a directed or shielded spray or with wiper equipment. For long term control of annual weeds WYNCA SUNPHOSATE 540 K Herbicide may be tank mixed with a suitable residual herbicide. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of stonefruit trees, or green bark, suckers, buds, foliage or fruit of other trees. Do not allow wiper surfaces to contact any part of the trees. Weed foliage coated in copper or calcium based compounds may reduce the activity of WYNCA SUNPHOSATE 540 K Herbicide.

Market Gardens

WYNCA SUNPHOSATE 540 K Herbicide can be applied before or after seeding and prior to crop emergence. WYNCA SUNPHOSATE 540 K Herbicide can be applied prior to transplanting or as a post harvest clean-up for longer term control of seedling weeds. WYNCA SUNPHOSATE 540 K Herbicide may be tank mixed with suitable residual herbicides. Use in accordance with the label recommendations for these products.

Asparagus

WYNCA SUNPHOSATE 540 K Herbicide can be applied:
Following the final harvest and immediately after removal of all spears but before crop regrowth occurs.
Prior to planting or crop emergence. Do not treat emerging spears.
In established ferns as a directed or shielded spray or with wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plants.

Industrial and Non-agricultural Areas

WYNCA SUNPHOSATE 540 K Herbicide can be used for the control of annual and perennial weeds as directed.
For residual control
WYNCA SUNPHOSATE 540 K Herbicide can be tank mixed with Simazine or another suitable residual herbicide. Exercise caution when using residual herbicides prior to planting trees, shrubs or other ornamental plants.

Turfgrass and Recreational Areas

Apply before establishing lawns, parks, reserves and ornamentals. WYNCA SUNPHOSATE 540 K Herbicide can be applied around the base of established trees as a directed or shielded spray or by wiper equipment. Do not allow spray, spray drift or the wiper surface to contact any part of the tree. In mown areas, allow sufficient re-growth before spraying.

Aquatic Areas: Drains and Waterways

Ensure that 75% of the weed to be treated is above water level or efficacy may be reduced. Following spray application do not submerge the treated plant as this will cause the herbicide to be washed off. Avoid treating silt laden plants.

Forestry

WYNCA SUNPHOSATE 540 K Herbicide can be applied for the control of grasses and a wide range of brushweeds. The addition of Organosilicone is required for the control of bracken and gorse.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/07/2024

Do not allow spray, spray drift or the wiper surface to contact the foliage or green bark of desirable trees as severe injury can occur.

WEEDS CONTROLLED - BRUSHWEEDS

Weed	Boom or Aerial L/Ha	Handgun L/100L	Knapsack mL/10L	Mistblower mL/10L	Critical Comment
Barberry	-	0.66	66	200	Apply Jan. to April from flowering to late fruit. Where bushes have been cut, regrowth should be 1.5 meters tall before spraying.
Blackberry	6-12.3	0.66 - 1	66	200	Apply to healthy plants from blossom to ripe fruit stage prior to Autumn frosts (Dec. to Apr.). Only bushes with a well developed canopy of leaves should be treated. Early treatment (Dec. to Jan.) may be preferred in drought prone areas. Leave sprayed blackberry for 40-60 days before burning or slashing and oversow the area. Where treatment has been carried out under marginal conditions some regrowth can be expected within 1-2 years. For aerial application of hard to kill blackberry or where no follow-up pasture management is anticipated, use the higher rate.
Boxthorn	-	0.83	-	-	Apply Nov. to Apr. to actively growing plants. Avoid salt damaged plants. Spray to ensure complete coverage of dense thickets. DO NOT use additional surfactant as poor results may occur.
Bracken	6 + 250mL Organosilicone per 100L (Min 500mL/Ha)	0.66 + 200mL Organosilicone per 100L	66 + 20mL Organosilicone	-	Apply when fronds are unfurled (mid Jan. to June) but prior to frost browning. DENSE VIRGIN BRACKEN WITH HIGH TRASH CONTENT AND LOW LEAF TO ROOT RATIO SHOULD BE BURNT IN SPRING AND THE UNFURLED FROND REGROWTH SPRAYED THE FOLLOWING MARCH TO JUNE.
Broom	6 + 250mL Organosilicone per 100L (Min 500mL/Ha)	0.66 + 200mL Organosilicone per 100L	66 + 20mL Organosilicone	200 + 20mL Organosilicone	Aerial Application - Best results are obtained when plants are in full leaf. Handgun Application - Apply all year round. Ensure full coverage.
Buddleja	6	0.66	66	-	Handgun - If actively growing treat all year. Aerial - Treat buddleja up to 2 meters high during Feb. to Apr.
Gorse	-	0.66 + 200mL Organosilicone per 100L	-	-	Apply all year round. **Spray to ensure complete coverage.**
Old Man's Beard	-	0.66-1.3	130	-	Apply Nov. to Mar. prior to leaf senescence. Vines too high for complete coverage should be cut at ground level in winter or spring and the regrowth sprayed in Mar. when plants are at least 0.5 meters long. Follow-up treatment will be necessary to control seedlings. Use the low rate for small vines.
Pampas Grass / Toetoe	-	0.66	66	-	Apply Spring-Autumn. Ensure complete coverage. Best results are achieved when applied after flowering.
Ring Fern	0.66-2	0.2	23	-	Control: Apply 2 litres/ha in Jan. to Apr. Pasture Manipulation: Apply 700ml - 1.4 litres/ha Dec. to Apr. Use the high rate for dense infestations. Burn 4-6 weeks after treatment and oversow with pasture.
Sweet Brier	8 + 250mL Organosilicone per 100L (Min 500mL/Ha)	0.66 + 200mL Organosilicone per 100L	66 + 20mL Organosilicone	-	Apply from late flowering to early leaf yellowing. Best results achieved with full coverage.
Willow	9-12 + 250mL Organosilicone per 100L (Min 500mL/Ha)	0.66+ 200mL Organosilicone per 100L	-	-	Apply from Jan. to Apr. before leaf fall begins. Total spray coverage is essential for best results. Use the high rate for pussy willow control.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/07/2024

WEEDS CONTROLLED – ANNUAL WEEDS

Weed	Boom or Aerial L/Ha	Handgun L/100L	Knapsack mL/10L	Mistblower mL/10L	Critical Comment
Annual Weeds	0.66-1.3	0.13	13	40	Use higher rate for larger weeds.
Barley Grass	0.66	0.13	13	40	
Cleavers	1	0.33	33	100	
Ryegrass Short rotation (Annual, Italian, Hybrid) Established <8 months Established >8 months	2-2.7 + 100mL Organosilicone per 100L 2.7 + 100mL Organosilicone per 100L	0.66 + 100mL Organosilicone per 100L 0.66 + 100mL Organosilicone per 100L	66 + 10mL Organosilicone 66 + 10mL Organosilicone	200 + 10mL Organosilicone 200 + 10mL Organosilicone	Ensure plants are actively growing and have at least 5-10 cm of green leaf. Add Organosilcone at all times.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND 26/07/2024

WEEDS CONTROLLED – PERENNIAL WEEDS

Weed	Boom or Aerial L/Ha	Handgun L/100L	Knapsack mL/10L	Mistblower mL/10L	Critical Comment
Australian Sedge	4 Aerial 6	0.66	66	200	Apply from Nov. to Mar. Where rank growth is present the area should be burnt 12-18 months prior to aerial application to ensure adequate spray coverage.
Browntop	4	0.66	66	200	Apply to 5-10cm high actively growing plants. Some regrowth may occur. Longterm control requires a pasture renewal programme utilizing annual crops and repeat applications of WYNCA SUNPHOSATE 540 K Herbicide before re-establishing perennial pasture.
Californian Thistle	2.7	0.66	66	200	Apply after flower bud development but before autumn browning. To ensure maximum shoot emergence do not cultivate for 6 months prior to treatment. Wiper - a double pass at low speed is recommended in heavy infestations. Mowing thistles 10 days after treatment may assist control. Retreatment may be necessary in the following season.
Couch	2-4	0.66	66	200	For suppression use 2 litres/ha, for control use 2.7 - 4 litres/ha when growing conditions are satisfactory. Autumn treatment is preferable. Use the high rate in late spring. Apply to plants with at least 4-5 leaves (at least 10-15 cm leaf length) and active shoot and root growth. Do not cultivate for 6 months prior to application to ensure maximum shoot emergence. Some regrowth may occur. Long term control requires a pasture renewal programme utilizing annual crops and repeat applications of WYNCA SUNPHOSATE 540 K Herbicide before re-establishing perennial pasture. See the Use Recommendations section for control of couch in a pre-harvest situation.
Chewings Fescue Red Fescue	6 + 100mL Organosilicone per 100L	0.66 + 100mL Organosilicone per 100L	-	-	Apply to healthy actively growing plants with 10-15 cm leaf length. Spring treatment gives best results. Some regrowth may occur. Long term control requires a pasture renewal programme utilizing annual crops and repeat applications of WYNCA SUNPHOSATE 540 K Herbicide before re-establishing perennial pasture.
Dock	2.7-6	0.66	66	-	Treat plants in spring or summer when actively growing and in full leaf. Use low rate for seedlings and young plants. Use high rate for control of established plants.
Floating Sweet Grass Reed Sweet Grass	4	0.66	66	200	Apply to actively growing plants when most have reached early seedhead.
Indian Doab	6	0.66	66	-	Apply when most plants have reached early seedhead.

Weed	Boom or Aerial L/Ha	Handgun L/100L	Knapsack mL/10L	Mistblower mL/10L	Critical Comment
Johnson Grass	4	0.66	66	200	Apply to actively growing plants.
Kikuyu	4	0.66	66	200	Apply to actively growing plants. Under dense growth conditions mulch and allow at least 10-15 cm of regrowth before treatment. Long term control requires a pasture renewal programme utilizing annual crops and repeat applications of WYNCA SUNPHOSATE 540 K Herbicide before re-establishing perennial pastures.
Mercer Grass Aquatic Non-aquatic	 6 4	 1 0.66	 100 66	 300 200	
Nodding Thistle	4	0.66	66	-	Wiper Application (see Application Instructions). Apply when actively growing when at least 10cm of the weed can be wiped. In heavy infestations a double pass is recommended.
Paspalum	2.7-4	0.66	66	200	Use the low rate from Jan. to Apr. (early seedhead) prior to frosts. Use 4 litres/ha on vegetative plants.
Potato (Self Set)	2	0.33	33	100	
Prairie Grass	1	-	-	-	
Ragwort	4	0.66	66	-	Apply when actively growing. Wiper - apply when plants are 10-15cm above the pasture. In heavy infestations a double pass is recommended. **Caution:** Ragwort plants may become more palatable following spraying and stock should be removed until the treated plants brown out.
Ratstail	3.3	-	-	-	Apply to actively growing plants in Autumn.
Rautahi (Cutty Grass)	4	0.66	66	200	Apply to actively growing plants from Jan. to Mar. Burning or slashing may take place from 30 days after treatment following brownout. Wiper (Rushes only). Apply Nov. to May.
Rushes	4	0.66	66	200	
Ryegrass Perennial	2.7 + 100mL Organosilicone per 100L	0.66 + 100mL Organosilicone per 100L	66 + 10mL Organo silicone	-	Ensure plants are actively growing and have at least 5 – 10 cm of green leaf. Add Organosilicone at all times.
Tall Fescue	4	0.66	66	200	Apply to actively growing plants Jan. to Mar.
Yarrow	2-4	-	-	-	Apply to plants with 5-10cm of leaf. Cultivate with a minimum of 2 grubbings from 10 days after treatment. Use the 2 litre rate when a competitive crop to yarrow is to be drilled. Use the 4 litre rate when the crop to be drilled is a poor competitor.

For wiper see application instructions in directions

RESISTANCE MANAGEMENT

GROUP	9	HERBICIDE

WYNCA SUNPHOSATE 540 K Herbicide contains the active ingredient glyphosate, which belongs to the glycine chemical group. Some naturally occurring weeds may become resistant when excessive use of the same herbicide occurs.
Consult http://resistance.nzpps.org/index.php for alternative modes of action and details of resistance management strategies (if any) for the weeds or crops listed on the label.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/07/2024

Registered pursuant to the ACVM Act 1997, No. P9745
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR100587
See www.epa.govt.nz for approval conditions.

Registrant: Zhejiang Xinan Chemical Industrial Group Co., Ltd.
93 Daqiao Road, Xinanjiang,
Jiande District, Zhejiang, China

NZ Agent:
AR & JA DRYSDALE LTD
PO Box 72 275
Papakura 2244
Telephone: (021) 939 728

Bar code

Batch No.	
Date of Manufacture	

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
26/07/2024