ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

DISRUPTOR™
540 KALIUM

A non-residual, non-selective herbicide for weed control prior to planting crops and pasture, prior to harvesting some crops and for general weed control in horticulture, agriculture, and forestry.

Contains 540 g/litre glyphosate as the potassium salt in the form of a soluble concentrate.

Registrant:
Evossentials
PO Box 1173
Pukekohe 2340
Tel 0800 382 5423

www.evossentials.co.nz

FREE RECYCLING
AGRECOVERY
agrecovery.co.nz

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

20 L

™ Trademark of Evoessentials Ltd

DISRUPTOR 540 Kalium **READ ENTIRE BOOKLET BEFORE OPENING OR USING**

TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

KEEP OUT OF REACH OF CHILDREN.
READ LABEL BEFORE USE.

To avoid risks to man and the environment, comply with instructions for use.

OPERATOR PROTECTION: Avoid contact with skin and eyes and avoid breathing spray mist. When mixing and applying spray wear appropriate protective clothing, suitable waterproof gloves, suitable boots, and eye and face protection (face shield). Remove protective clothing and wash hands and face thoroughly before meals and after work.
ENVIRONMENTAL PROTECTION: Prevent spray drift as it may cause severe damage to non-target plants. Where the substance is applied onto or into water and the water has the potential to leave the place containing the application area, clauses 62, 63 and 64 of the Hazardous Property Controls Notice 2017 apply.
EQUIPMENT: Apply using appropriate, well maintained spray equipment that is accurately calibrated. Flush equipment thoroughly with clean water after use. Do not contaminate water when disposing of equipment wash waters.
TRANSPORT: Do not carry more than 1 litre of DISRUPTOR 540 Kalium on a passenger service vehicle.
STORAGE: Store in original container tightly closed in a locked, dry, cool area away from foodstuffs, stockfeeds, seeds and fertilisers. Store in accordance with the current version of the New Zealand Standard for the Management of Agrichemicals NZS8409. Solutions of DISRUPTOR 540 Kalium should be mixed, stored and applied only in stainless steel, aluminum, brass, copper, fibreglass, plastic or plastic lined containers. Do not mix or store this product in galvanized or unlined steel (except stainless steel) containers or spray tanks. DISRUPTOR 540 Kalium or solutions of DISRUPTOR 540 Kalium react with such containers to produce hydrogen gas which may form a highly combustible gas mixture which can flash or explode if ignited. Stores containing 1000 litres or more of DISRUPTOR 540 Kalium require secondary containment systems, emergency response plans and signage.
SPILLAGE: Wear appropriate protective clothing and prevent further spillage and product from entering waterways. COLLECT SPILLAGE. Contain and absorb spills with inert material and place in suitable waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely.
DISPOSAL: Avoid contamination of any water supply with product or empty container.
Container: Triple rinse empty container and add rinsate to the spray tank. Recycle the rinsed container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill.
Product: Dispose of product only by using according to the label, or at an approved landfill. Do not store this product for disposal in galvanized or unlined steel (except stainless steel) containers.

FIRST AID
If swallowed, do not induce vomiting. Rinse mouth. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water.

Conditions of Sale
The use of DISRUPTOR 540 Kalium herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Evoessentials Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Evoessentials Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Registered pursuant to the ACVM Act 1997 No. P9876 See www.foodsafety.govt.nz for registration conditions. Approved pursuant to the HSNO Act 1996 Approval Code HSR100587. See www.epa.govt.nz for approval controls.	Batch No. & D.O.M sticker Emergency Phonelink: For 24 hour specialist emergency advice call 0800 734 607

GENERAL INFORMATION

DISRUPTOR 540 Kalium is a broad spectrum, non-selective, non-residual herbicide that is not volatile and may be used in agriculture, horticulture, drains, aquatic areas, forestry and for other uses including general weed control. DISRUPTOR 540 Kalium is recommended prior to sowing pasture, crops and for the control of a wide range of brushweeds.

DISRUPTOR 540 Kalium is absorbed by plant foliage and green stems. It is inactivated immediately upon contact with the soil and does not provide residual weed control. DISRUPTOR 540 Kalium moves from the point of foliage or stem contact into the root system controlling the plant above and below the ground. Visible effects may take up to 14 days or even longer depending on growing conditions. These effects are a gradual wilting and yellowing of the foliage which advances to deterioration of above and below ground parts.

Apply DISRUPTOR 540 Kalium only to healthy, actively growing plants with adequate leaves for herbicide uptake. Avoid treating plants that are silt laden or affected by drought, frost, water logging, grazing or previous herbicide application.

Rainfall occurring up to 2 hours after application may reduce effectiveness. When an organosilicone surfactant is added the rainfree period may be reduced to 30 minutes. If DISRUPTOR 540 Kalium is applied to wet plants, efficacy may be reduced due to runoff.

With brushweeds wait for stem dehydration to occur prior to burning or slashing.

Animal safety
After application of DISRUPTOR 540 Kalium a withholding period for grazing stock is not required (except where ragwort is present). However, it is recommended that grazing or cultivating is delayed for 1 day after treatment of annual weeds or 3 days if perennial weeds are present to ensure absorption of DISRUPTOR 540 Kalium.

RESISTANCE MANAGEMENT

DISRUPTOR 540 Kalium contains glyphosate, a herbicide from the glycine chemical family with inhibition of EPSP synthase mode of action, GROUP 9 HERBICIDE. Weeds resistant to glyphosate and other Group 9 herbicides could eventually dominate from excessive use of these herbicides. To minimise this risk use strictly in accordance with label instructions. DISRUPTOR 540 Kalium should be used in a program that incorporates effective herbicide products with different modes of action. Refer to the NZCPR web site (www.nzpps.org/resistance) for more detailed information.

DIRECTIONS FOR USE

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Withholding Period
Wheat, barley, oat grain – 7 days

Refer to “Weeds Controlled” section for rates of application.

General Weed Control
Handgun – mix 660ml of DISRUPTOR 540 Kalium with 100 litres of water (0.66% solution).
Knapsack - mix 100ml of DISRUPTOR 540 Kalium with 15 litres of water.
Small sprayers – mix 6.5ml of DISRUPTOR 540 Kalium with every 1 litre of water.
Apply to completely wet the foliage but not to run-off.

Brushweed Control
DISRUPTOR 540 Kalium may be used to control a wide range of brushweeds and woody plants.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

Agricultural Areas
DISRUPTOR 540 Kalium may be used for the control of annual and perennial weeds as directed in agricultural land prior to sowing any edible or non-edible crop or pasture.
Where white clover control is required, tank mix with a suitable white clover herbicide and observe recommended waiting periods prior to planting legumes. Cultivation may start from 3 days after spraying perennial weeds and 1 day for annual weeds.

Pasture – Direct (Spot) Application
DISRUPTOR 540 Kalium is non-selective and may damage or kill any plant in the sprayed area. Apply as a directed low pressure spray. Retreatment and/or pasture improvement may be necessary to restrict seedling re-establishment.

Pasture – Selective Wiper Application
For the control of erect weeds in pastures.

Pasture Renewal
DISRUPTOR 540 Kalium may be used to control existing pasture species prior to re-seeding pasture or establishing crops.

Pasture Seedhead Suppression (Unimproved Pasture)
Use in unimproved pasture to delay onset of flowering and to maintain pasture grasses in a vegetative growth phase.
Treat spring pasture at early grass seedhead emergence with 180-200 ml DISRUPTOR 540 Kalium /ha. Temporary pasture brownout will occur. Do not use an organosilicone surfactant or any other surfactant. Accurate spray calibration is important. Grazing can be carried out at any time after 3 days from treatment, but for optimum results graze from 21 days after treatment.

DO NOT USE ON RYEGRASS OR OTHER IMPROVED PASTURES AS YIELD SUPPRESSION MAY OCCUR. IT IS ADVISABLE NOT TO TREAT MORE THAN 10-15% OF THE FARM IN ANY ONE YEAR. DO NOT USE WHEN SOIL MOISTURE LEVELS ARE INADEQUATE FOR PASTURE GROWTH FOLLOWING APPLICATION.

Spraying of Silage Before Cutting
Where paddocks are to be used for cropping after silage, DISRUPTOR 540 Kalium may be applied before the silage is cut. Use 2.0-2.7 L/ha plus an organosilicon surfactant 100mL/100 litres of water in 100-200 litres of water per hectare. Use the high rate on perennial ryegrass or if the silage is rank or lodged. Refer to higher label rates on other specific weeds. After spraying wait at least 4 days before cutting. Grass quality is maintained for 7-10 days after the initial planned cutting date, even into the early stages of brown-out.

Spraying of Silage or Hay Paddocks After Cutting
Before spraying ensure that there is at least 5-10 cm of actively growing green leaf on all tillers. Typically, this takes three weeks or more after cutting. Refer to application rates in the table. Always use an organosilicone surfactant.

Pre-Harvest weed control: Wheat, Barley, Oats
Wheat, barley and oats: apply 7-12 days prior to harvest when the grain moisture content is less than 30%.
Do not use as a pre-harvest desiccant.
Do not apply to undersown crops including clovers as they may be severely damaged.
Do not apply to crops which will be harvested for seed. Use rates up to 2.67 litres/ha depending on the specific weeds to be treated. Avoid treating drought stressed plants.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

Orchards/Vines: Citrus, Apples, Pears, Stonefruit, Kiwifruit, Grapevines
Apply as a directed or shielded spray or with wiper equipment. For long term control of annual weeds DISRUPTOR 540 Kalium may be tank mixed with a suitable residual herbicide. Do not apply near unprotected trees less than 3 years old. Do not allow spray or spray drift to contact any part of stone fruit trees, or green bark, suckers, buds, foliage or fruit of other trees. Do not allow wiper surfaces to contact any part of the trees. Weed foliage coated in copper or calcium based compounds may reduce the activity of DISRUPTOR 540 Kalium.

Market Gardens
DISRUPTOR 540 Kalium can be applied before or after seeding and prior to crop emergence. DISRUPTOR 540 Kalium can be applied prior to transplanting or as a post-harvest clean-up for longer term control of seedling weeds. DISRUPTOR 540 Kalium may be tank mixed with suitable residual herbicides. Use in accordance with the label recommendations for these products.

Asparagus
DISRUPTOR 540 Kalium can be applied:

- Following the final harvest and immediately after removal of all spears but before crop regrowth occurs.
- Prior to planting or crop emergence. Do not treat emerging spears. In established ferns as a directed or shielded spray or with wiper equipment.

Do not allow spray, spray drift or the wiper surface to contact any part of the asparagus plants.

Industrial and Non-Agricultural Areas
DISRUPTOR 540 Kalium can be used for the control of annual and perennial weeds as directed. For residual control, DISRUPTOR 540 Kalium can be tank mixed with simazine or another suitable residual herbicide. Exercise caution when using residual herbicides prior to planting trees, shrubs or other ornamental plants.

Turfgrass and Recreational Areas
Apply before establishing lawns, parks, reserves and ornamentals. DISRUPTOR 540 Kalium can be applied around the base of established trees as a directed or shielded spray or by wiper equipment. Do not allow spray or spray drift or the wiper surface to contact any part of the tree. In mown areas, allow sufficient re-growth before spraying.

Aquatic Areas: Drains and Waterways
Ensure that 75% of the weed to be treated is above water level or efficacy may be reduced. Following spray application do not submerge the treated plant as this will cause the herbicide to be washed off. Avoid treating silt laden plants.

Forestry
DISRUPTOR 540 Kalium can be applied for the control of grasses and a wide range of brushweeds. The addition of an organo-silicone surfactant is required for the control of bracken and gorse.
Do not allow spray, spray drift or the wiper surface to contact the foliage or green bark of desirable trees as severe injury can occur.

MIXING AND COMPATIBILITY

DISRUPTOR 540 Kalium mixes readily with clean water. Reduced results may occur if dirty water (e.g. from ponds), hard water containing calcium salts, or water containing metal ions (e.g. iron) is used.

Do not mix, store or apply this product in galvanized steel or unlined steel containers or spray tanks, since a highly flammable gas mixture may be formed. Use stainless steel, aluminium, brass, copper, fibreglass, plastic or plastic lined containers or spray tanks. Spray tanks, pumps, lines and nozzles should be thoroughly rinsed with clean water following application.

Ensure the spray tank is free of any residue of previous spray materials prior to filling. Half fill the spray tank with water. Add the required amount of DISRUPTOR 540 Kalium, agitate and continue filling the tank. Avoid foaming by keeping the filler nozzle below water level. Do not store mixed solutions for longer than 5 days. Do not use with 2,4-D Amine. If adding other herbicides, ensure these are fully dispersed before adding the DISRUPTOR 540 Kalium. Add surfactant, if required, near the end of the filling process to minimise foaming. Always maintain adequate agitation during application and use the tank mix promptly as a gradual loss of activity may occur over a period of days following spray preparation. Clean all equipment after use by washing thoroughly with clean water.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

APPLICATION INSTRUCTIONS

Beware: Spray drift hazard. Apply this product carefully. Prevent spray drift as it may cause severe damage to non-target plants. Non-selective herbicide toxic to plants and algae. Avoid contact with foliage or green stems of fruit or crops, desirable plants and trees, since severe injury or death may occur.

Broadcast Boom Application. The use of flat fan, wide angle nozzles (e.g. 110º) using pressures of 200-280kPa are preferred. Set boom height to ensure a double overlap of the spray pattern at the top of the weed canopy. Apply in 100-200 litres of clean water per hectare. For orchard application wide angle, even spray nozzles are preferred.

Knapsack or Handgun Applications. Adjust equipment to apply a fine spray. Use the minimum spray volume necessary to achieve uniform and complete coverage. Handgun spraying pressures of 1,000-2,000 kPa and No. 4 to No. 6 spray tips are recommended.

Wiper Application. Wiper equipment (e.g. Ropewick, canvas, felt or carpet applicators) may be used to apply DISRUPTOR 540 Kalium. Pre-mix 1 part of DISRUPTOR 540 Kalium and 2 parts of clean water to prepare a 33% solution for use through the wiper. Mix only enough herbicide for immediate needs and allow the wiping area to become fully saturated before commencing. The wiping area must remain moist and clean otherwise reduced results will occur. In heavy weed infestations or when treating perennial weeds such as Californian thistle or multi-crown ragwort, a double pass in opposite directions on the same day will give the best results. As weed density increases, decrease ground speed.

Aircraft Application. DISRUPTOR 540 Kalium may be applied by aircraft in forestry and non-cropping areas only. Aerial application of DISRUPTOR 540 Kalium is not recommended on cropping land or near desirable plant species as even slight drift may cause severe injury or destruction. Do not apply in gusty or windy conditions or in other conditions conducive to drift, e.g. still air and inversion conditions.

Mistblower Application. Use a solution of 200ml of DISRUPTOR 540 Kalium in 10 litres of water. For the micron attachment increase to 1.3 litres of DISRUPTOR 540 Kalium in 10 litres of water. Provide complete coverage and spray to wet but not to runoff.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

WEEDS CONTROLLED

Weed	Boom or Aerial litres/ha	Handgun mL/ 100 litres	Knapsack mL/10 litres	Mistblower mL/10 litres	Critical Comments
WEEDS CONTROLLED – BRUSHWEEDS					
BARBERRY *(Berberis glaucocarpa)*	-	660	66	200	Apply Jan. to April from flowering to late fruit. Where bushes have been cut, regrowth should be 1.5 metres tall before spraying
BLACKBERRY *(Rubus fruticosus)*	6-12.3	660 mL to 1 litre	66	200	Apply to healthy plants from blossom to ripe fruit stage prior to Autumn frosts (Dec. to Apr). Only bushes with a well developed canopy of leaves should be treated. Early treatment (Dec. to Jan.) may be preferred in drought prone areas. Leave sprayed blackberry for 40-60 days before burning or slashing and oversow the area. Where treatment has been carried out under marginal conditions some regrowth can be expected within 1-2 years. For aerial application of hard to kill blackberry or where no follow- up pasture management is anticipated, use the higher rate.
BOXTHORN (*Lycium ferocissimum*)	-	830	-	-	Apply Nov. to Apr to actively growing plants. Avoid salt damaged plants. Spray to ensure complete coverage of dense thickets. Do not use additional surfactant as poor results may occur.
BRACKEN *(Pteridium aquilinum)* AN ORGANOSILICONE SURFACTANT MUST BE USED	6 + Organosilicone at label rate	660 + Organosilicone at label rate	66 + Organosilicone at label rate	-	Apply when fronds are unfurled (mid Jan. to June.) but prior to frost browning. DENSE VIRGIN BRACKEN WITH HIGH TRASH CONTENT AND LOW LEAF TO ROOT RATIO SHOULD BE BURNT IN SPRING AND THE UNFURLED FROND REGROWTH SPRAYED THE FOLLOWING MARCH TO JUNE
BROOM *(Cytisus scoparius)* AN ORGANOSILICONE SURFACTANT MUST BE USED	6 + Organosilicone at label rate	660 + Organosilicone at label rate	66 + Organosilicone at label rate	200 + Organosilicone at label rate	Aerial Application – Best results are obtained when plants are in full leaf. Handgun Application – Apply all year round. Ensure full coverage
BUDDLEJA *(Buddleja davidii)*	6	660	66	-	Handgun – If actively growing treat all year. Aerial – Treat buddleja up to 2 metres high during Feb. to Apr.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND 04/08/2024

Weed	Boom or Aerial litres/ha	Handgun mL/ 100 litres	Knapsack mL/10 litres	Mistblower mL/10 litres	Critical Comments
GORSE *(Ulex europaeus L)* AN ORGANOSILICONE SURFACTANT MUST BE USED	-	660 + Organosilicone at label rate	-	-	Apply all year round. **Spray to ensure complete coverage**.
OLD MAN'S BEARD *(Clematis vitalba)*	-	660 mL to1.3 litres	130	-	Apply Nov. to Mar. prior to leaf senescence. Vines too high for complete coverage should be cut at ground level in winter or spring and the regrowth sprayed in Mar. when plants are at least 0.5 metres long. Follow-up treatment will be necessary to control seedlings. Use the low rate for small vines.
PAMPAS GRASS / TOETOE *(Cortaderia* spp.*)*	-	660	66	-	Apply Spring-Autumn. Ensure complete coverage. Best results are achieved when applied after flowering.
RING FERN *(Paesia scaberula)*	0.66 to 2	200	23	-	Control: Apply 2 litres/ha in Jan. to Apr. Pasture Manipulation: Apply 700ml – 1.4 litres/ha Dec. to Apr. Use the high rate for dense infestations. Burn 4-6 weeks after treatment and oversow with pasture.
SWEET BRIER *(Rosa rubiginosa)* AN ORGANOSILICONE SURFACTANT MUST BE USED	8 + Organosilicone at label rate	660 + Organosilicone at label rate	66 + Organosilicone at label rate	-	Apply from late flowering to early leaf yellowing. Best results achieved with full coverage.
WILLOW (*Salix* species)	9 to12 + Organosilicone at label rate	660 + Organosilicone at label rate	-	-	Apply from Jan. to Apr. before leaf fall begins. Total spray coverage is essential for best results. Use the high rate for pussy willow control.
WEEDS CONTROLLED - ANNUAL WEEDS					
ANNUAL WEEDS (Seedlings)	0.66 to 1.3	130	13	40	
BARLEY GRASS *(Hordeum murinum)*	0.66	130	13	40	
CLEAVERS *(Galium aparine)*	1	330	33	100	
RYEGRASS - short rotation (Annual, Italian, Hybrid)					Ensure plants are actively growing and have at least 5-10 cm of green leaf. Add an organosilicone surfactant at all times.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

Weed	Boom or Aerial litres/ha	Handgun mL/ 100 litres	Knapsack mL/10 litres	Mistblower mL/10 litres	Critical Comments
Established < 8 months Established > 8 months	2 to 2.7 + Organosilicone at label rate 2.7 + Organosilicone at label rate	660 + Organosilicone at label rate 660 + Organosilicone at label rate	66 + Organosilicone at label rate 66 + Organosilicone at label rate	200 + Organosilicone at label rate 200 + Organosilicone at label rate	
WEEDS CONTROLLED - PERENNIAL WEEDS					
AUSTRALIAN SEDGE *(Carex longebrachiata)*	4 Aerial 6	660	66	200	Apply from Nov. to Mar. Where rank growth is present the area should be burnt 12-18 months prior to aerial application to ensure adequate spray coverage.
BROWNTOP *(Agrostis capillaris)*	4	660	66	200	Apply to 5-10cm high actively growing plants. Some regrowth may occur. Longterm control requires a pasture renewal programme utilizing annual crops and repeat applications of DISRUPTOR 540 Kalium before re-establishing perennial pasture.
CALIFORNIAN THISTLE *(Cirsium arvense)*	2.7	660	66	200	Apply after flower bud development but before autumn browning. To ensure maximum shoot emergence do not cultivate for 6 months prior to treatment. Wiper – a double pass at low speed is recommended in heavy infestations. Mowing thistles 10 days after treatment may assist control. Retreatment may be necessary in the following season.
COUCH *(Elytrigia repens)*	2-4	660	66	200	For suppression use 2 litres/ha, for control use 2.7 – 4 litres/ha when growing conditions are satisfactory. Autumn treatment is preferable. Use the high rate in late spring. Apply to plants with at least 4-5 leaves (at least 10 – 15 cm leaf length) and active shoot and root growth. Do not cultivate for 6 months prior to application to ensure maximum shoot emergence. Some regrowth may occur. Longterm control requires a pasture renewal programme utilizing annual crops and repeat applications of DISRUPTOR 540 Kalium before re-establishing perennial pasture. See the Use Recommendations section for control of couch in a pre- harvest situation.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
04/08/2024

Weed	Boom or Aerial litres/ha	Handgun mL/ 100 litres	Knapsack mL/10 litres	Mistblower mL/10 litres	Critical Comments
CHEWINGS FESCUE *(Festuca rubra* subsp. *commutata)* RED FESCUE *(Festuca rubra* subsp. *rubra*)	6 + Organosilicone at label rate	660 + Organosilicone at label rate	66	-	Apply to healthy actively growing plants with 10 – 15 cm leaf length. Spring treatment gives best results. Some regrowth may occur. Longterm control requires a pasture renewal programme utilizing annual crops and repeat applications of DISRUPTOR 540 Kalium before re-establishing perennial pasture.
DOCK *(Rumex obtusifolius)*	2.7-6	660	66	-	Treat plants in spring or summer when actively growing and in full leaf. Use low rate for seedlings and young plants. Use high rate for control of established plants.
FLOATING SWEET GRASS *(Glyceria fluitans)* REED SWEET GRASS *(Glyceria maxima)*	4	660	66	200	Apply to actively growing plants when most have reached early seedhead.
INDIAN DOAB *(Cynodon dactylon)*	6	660	66	-	Apply when most plants have reached early seedhead.
JOHNSON GRASS *(Sorghum halepense)*	4	660	66	200	Apply to actively growing plants.
KIKUYU *(Pennisetum clandestinum)*	4	660	66	200	Apply to actively growing plants. Under dense growth conditions mulch and allow at least 10 – 15 cm of regrowth before treatment. Longterm control requires a pasture renewal programme utilizing annual crops and repeat applications of DISRUPTOR 540 Kalium before re-establishing perennial pastures.
MERCER GRASS *(Paspalum distichum)* Aquatic Non-Aquatic	 6 4	 1 litre 660 mL	 100 66	 300 200	
NODDING THISTLE *(Carduus nutans)*	4	660	66	-	Wiper Application (see Application Instructions). Apply when actively growing when at least 10cm of the weed can be wiped. In heavy infestations a double pass is recommended.
PASPALUM *(Paspalum dilatatum*	2.7-4	660	66	200	Use the low rate from Jan. to Apr. (early seedhead) prior to frosts. Use 4 litres/ha on vegetative plants.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024

Weed	Boom or Aerial litres/ha	Handgun mL/ 100 litres	Knapsack mL/10 litres	Mistblower mL/10 litres	Critical Comments
POTATO (SELF SET) *(Solanum tuberosum)*	2	330	33	100	
PRAIRIE GRASS *(Bromus willdenowii)*	1	-	-	-	
RAGWORT *(Scenecio jacobaea)*	4	660	66	-	Apply when actively growing. Wiper – apply when plants are 10-15cm above the pasture. In heavy infestations a double pass is recommended. **Caution**: Ragwort plants may become more palatable following spraying and stock should be removed until the treated plants brown out.
RATSTAIL *(Sporobolus africanus)*	3.3	-	-	-	Apply to actively growing plants in Autumn.
RAUTAHI (CUTTY GRASS) *(Carex coriacea)*	4	660	66	200	Apply to actively growing plants from Jan. to Mar. Burning or slashing may take place from 30 days after treatment following brownout. Wiper (Rushes only). Apply Nov. to May.
RUSHES *(Juncus* spp)	4	660	66	200	
RYEGRASS PERENNIAL *(Lolium perenne)*	2.7 + Organosilicone at label rate	660 mL + Organosilicone at label rate	66 + Organosilicone at label rate	-	Ensure plants are actively growing and have at least 5 – 10 cm of green leaf. Add Organosilicone at all times.
TALL FESCUE *(Festuca arundinacea*	4	660	66	200	Apply to actively growing plants Jan. to Mar.
YARROW *(Achillea millefolium)*	2 to 4	-	-	-	Apply to plants with 5-10cm of leaf. Cultivate with a minimum of 2 grubbings from 10 days after treatment. Use the 2 litre rate when a competitive crop to yarrow is to be drilled. Use the 4 litre rate when the crop to be drilled is a poor competitor.

*Organosilicone surfactant

For wiper see application instructions in directions.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
04/08/2024