150 X 240 (20 L)

Pantone 7485C

Pantone 368C

DANGER • ECOTOXIC

KEEP OUT OF REACH OF CHILDREN

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

Blue AG

Patriot 1®

NON-SELECTIVE CONTACT HERBICIDE

GROUP **10** HERBICIDE

ACTIVE INGREDIENT

200 g/litre
Glufosinate-ammonium
in the form of a soluble concentrate.

A non-selective contact herbicide for the control of a wide range of grasses, broad leaf weeds and clovers in a variety of crop land and non-crop land situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS)
0800 CHEMCALL OR 0800 243 622 OR
PGG WRIGHTSON LTD 0800 10 22 76

20 L
NET CONTENTS

150 X 240 (20 L) Pantone 425C BLACK C:60 M:40 Y:40 K:100 Pantone 368C Pantone 485 C

PEEL OFF

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

HAZARD CLASSIFICATIONS

Skin irritation Category 2, Eye irritation Category 2, Reproductive toxicity Category 2, Specific target organ toxicity (repeated exposure) Category 1, Hazardous to soil organisms, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment chronic Category 3.

TOXICITY

DANGER: Causes damage to kidneys and nervous system through prolonged or repeated exposure via any route. Causes eye irritation and skin irritation. Suspected of damaging fertility or the unborn child through repeated exposure at high doses.

ECOTOXIC

Harmful to aquatic life. This substance must not be applied onto or into water. Toxic to soil organisms. Harmful to terrestrial vertebrates. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

HANDLING PRECAUTIONS: Do NOT handle this product until all safety precautions have been read and understood. Keep out of reach of children. Avoid inhalation of spray mist. Avoid contact with eyes and skin. Do not eat, drink or smoke while using. Use only outdoors in a well-ventilated area.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS 8409). Flush equipment thoroughly with clean water after use.

STORAGE: Store in the original container locked up in a cool well ventilated place, out of direct sunlight and away stockfeed or foodstuff. Store in accordance with the current version of NZS 8409 Management of Agrichemicals.

PROTECTIVE CLOTHING: When mixing or applying wear appropriate protective clothing including impervious gloves, chemical resistant boots, eye protection (goggles with side pieces or a face shield) and cotton overalls buttoned to the neck and wrist. Remove protective clothing and wash hands, arms and face thoroughly with soap and water before meals and after work.

SPILLAGE: Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways; bund if necessary. Wear appropriate protective clothing as in the Precautions box of this label to clean up spills. Absorb spillage with inert absorbent material, such as vermiculite, sand or clay, and place in waste containers. Wash contaminated area with water and detergent, absorb and add to waste for disposal. Dispose of the waste safely in an approved landfill.

DISPOSAL INFORMATION

PRODUCT DISPOSAL: Dispose of product only by using in accordance with label directions, through the Agrecovery® Chemical Recovery service, or at an approved landfill.
CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Take empty container to an Agrecovery® collection site. If this is not possible bury in an approved landfill. Do not contaminate aquatic environment with product or empty container.

FIRST AID

If exposed or concerned get medical advice/attention.
If swallowed, rinse mouth. Do NOT induce vomiting. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell.
If inhaled, Remove patient to fresh air and maintain in a position comfortable for breathing. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor for advice if person feels unwell.
If in eyes, Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists get medical advice/attention.
If on skin, wash with plenty of soap and water. If skin irritation persists, get medical advice/attention. Wash contaminated clothing before reuse.

FORT PRIMARY INDUSTRIES NEW ZEALAND 10/10/2025

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use.
PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

9 415251 004843 >

150 X 240 (20 L)

Pantone 425C

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

DIRECTIONS FOR USE
BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS. Apply using ground-based application only. The maximum application rate is 10L/ha (2.03kg ai/ha) with a maximum of two applications per year and minmum application interval of 90 days.

RESISTANCE MANAGEMENT
Patriot 1 contains the GROUP 10 herbicide, Glufosinate-ammonium, a glutamine synthetase inhibitor. Certain weeds may develop resistance to this herbicide through prolonged and repeated use. Herbicides from different groups should either be used alternately or mixed together to prevent the numbers of resistant individuals building up. To minimize this risk use strictly in accordance with label instructions and strictly in accordance with label instructions and resistance management strategies at www.resistance.nzpps.org

MIXING
Add the required amount of Patriot 1 to a partly filled spray tank with the agitation system operating. Complete filling the tank with water and apply, maintaining agitation.

COMPATIBILITY
Patriot 1 is compatible with most commonly used residual herbicides.

RAINFALL
Rainfall within 2 hours of application may wash Patriot 1 off the foliage, and within 6 hours will reduce effectiveness to the degree that re-treatment may be necessary.

USING THE SPRAY
IMPORTANT: In order to select the appropriate nozzles for the required spray quality and to operate the equipment to minimise spray drift, this product must be applied in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).

WEED SUSCEPTIBILITY
The lower rates indicated are appropriate when weeds are young, succulent and actively growing, or when the weed population is sparse and good coverage and penetration is possible. Use the higher rates when weeds are mature or the population is dense.

NB: Patriot 1 will not completely control mature weeds with well-developed tap roots or underground storage organs: some re-growth may occur following treatment.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
10/10/2025

RECOMMENDATIONS FOR USE

USE	RATE	REMARKS
Berryfruit – Raspberries and Brambles Primocane control	500-750ml/100 litres applied through cone nozzles in sufficient water to ensure complete coverage of primocanes.	Apply as a directed high volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of a non-ionic wetter at label rate is recommended. "Knockdown" takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached 15cm in height. Do not apply to plants under stress.
Orchard and vineyard weed control. Pipfruit, Kiwifruit, Stonefruit, Grapes and Citrus	Use a rate appropriate to the weeds present. If dilution rate is less than 1 litre/100 litres of water add a non-ionic wetter at label rates.	Apply in spring when the majority of the annual weeds have germinated. Residual herbicides may be applied in mixture for long-term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water to ensure complete coverage. May be applied up to one day before crop emergence.
Non-cropland. General weed control.	7.5-10 litres/ha in 300-500 litres of water or 100ml/ 10 litre knapsack.	Apply in a drenching spray to the point of run-off. May be mixed with residual herbicides for long-term weed control.

INDICATED RATE RANGE FOR VARIOUS WEED SPECIES

3 - 5 Litres/ha

Annual Poa	Asparagus seedlings	Beaked Hawksbeard	Black Nightshade	Calandrinia
Chickweed	Cleavers	Clover seedlings	Cornbind	Daisy
Dandelion	Mouse-ear chickweed	Musky Storksbill	Redroot	Rough-stalked meadow grass
Scotch Thistle	Sow Thistle	Staggerweed	Summer grass	Spurrey
Stinking Mayweed	Subterranean clover	Wild turnip	Willow weed	Wireweed

5 – 7.5 Litres/ha

Bristle grass	Bur medick	Catsear	Chewing's Fescue	Clovers
Cut-leaved geranium	Docks	Fleabane	Parsley Piert	Prairie grass
Ryegrasses*	Sheep's sorrel	Storksbill	Twin cress	Yorkshire fog

7.5 – 10 Litres/Ha

Broom	Buttercups*	Couch	Crowfoot grass	Field pansy
Gorse	Hawksbeard	Kikuyu grass	Mallows	Paspalum
Tall Fescue	Wild strawberry	Willow herb	Yarrow	

10 Litres/ha

Bracken	Indian doab	Meadow rice grass	Plantains	Rape
Rushes	Sedges			

*Resistant weeds: Hairy buttercup, some ryegrass strains.

WITHHOLDING PERIODS: 28 days for all crops. It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.
ADDITIONAL INFORMATION: Patriot 1 is absorbed into the foliage and green stems of actively growing plants. It is inactive in soil and is not translocated as an active herbicide, so thorough coverage of target weeds is essential for good results. Visual effects of treatment will become evident more rapidly under ideal growing conditions, generally within 3-10 days. Do not spray in temperatures below 10°C. Avoid contact on green parts (including immature bark) of desirable plants, as damage may occur. Do not mow or cultivate after treatment until weeds start to die off.
Registered pursuant to the ACVM Act 1997, No. P9924 See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval No. HSR100830
See www.epa.govt.nz for approval controls

MULTIFOLD®

300X200 (200 L)

Pantone 7485C

Pantone 368C

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

DANGER • ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Blue AG

Patriot 1®

NON-SELECTIVE CONTACT HERBICIDE

GROUP 10 HERBICIDE

ACTIVE INGREDIENT

200 g/litre Glufosinate-ammonium

in the form of a soluble concentrate.

A non-selective contact herbicide for the control of a wide range of grasses, broad leaf weeds and clovers in a variety of crop land and non-crop land situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS) 0800 CHEMCALL OR 0800 243 622 OR PGG WRIGHTSON LTD 0800 10 22 76

200 L
NET CONTENTS

300X200 (200 L)

Pantone 425C

BLACK
C:60 M:40 Y:40 K:100

Pantone 368C

Pantone 485 C

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

PEEL OFF

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

HAZARD CLASSIFICATIONS
Skin irritation Category 2, Eye irritation Category 2, Reproductive toxicity Category 2, Specific target organ toxicity (repeated exposure) Category 1, Hazardous to soil organisms, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment chronic Category 3.

TOXICITY
DANGER: Causes damage to kidneys and nervous system through prolonged or repeated exposure via any route. Causes eye irritation and skin irritation. Suspected of damaging fertility or the unborn child through repeated exposure at high doses.

ECOTOXIC
Harmful to aquatic life. This substance must not be applied onto or into water. Toxic to soil organisms. Harmful to terrestrial vertebrates. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

HANDLING PRECAUTIONS: Do NOT handle this product until all safety precautions have been read and understood. Keep out of reach of children. Avoid inhalation of spray mist. Avoid contact with eyes and skin. Do not eat, drink or smoke while using. Use only outdoors in a well-ventilated area.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS 8409). Flush equipment thoroughly with clean water after use.

STORAGE: Store in the original container locked up in a cool well ventilated place, out of direct sunlight and away stockfeed or foodstuff. Store in accordance with the current version of NZS 8409 Management of Agrichemicals.

PROTECTIVE CLOTHING: When mixing or applying wear appropriate protective clothing including impervious gloves, chemical resistant boots, eye protection (goggles with side pieces or a face shield) and cotton overalls buttoned to the neck and wrist. Remove protective clothing and wash hands, arms and face thoroughly with soap and water before meals and after work.

SPILLAGE: Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways; bund if necessary. Wear appropriate protective clothing as in the Precautions box of this label to clean up spills. Absorb spillage with inert absorbent material, such as vermiculite, sand or clay, and place in waste containers. Wash contaminated area with water and detergent, absorb and add to waste for disposal. Dispose of the waste safely in an approved landfill.

DISPOSAL INFORMATION
PRODUCT DISPOSAL: Dispose of product only by using in accordance with label directions, through the Agrecovery® Chemical Recovery service, or at an approved landfill.
CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Take empty container to an Agrecovery® collection site. If this is not possible bury in an approved landfill. Do not contaminate aquatic environment with product or empty container.

FIRST AID
If exposed or concerned get medical advice/attention.
If swallowed, rinse mouth. Do NOT induce vomiting. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell.
If inhaled, Remove patient to fresh air and maintain in a position comfortable for breathing. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor for advice if person feels unwell.
If in eyes, Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical advice/attention.
If on skin, wash with plenty of soap and water. If skin irritation persists, get medical advice/attention. Wash contaminated clothing before reuse.

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

9 415251 005185 >

300X200 (200 L)

Pantone 425C

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

MULTIFOLD®

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

DIRECTIONS FOR USE
BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS. Apply using ground-based application only. The maximum application rate is 10L/ha (2.03kg ai/ha) with a maximum of two applications per year and minmum application interval of 90 days.

RESISTANCE MANAGEMENT
Patriot 1 contains GROUP 10 herbicide, Glufosinate-ammonium, a glutamine synthetase inhibitor. Certain weeds may develop resistance to this herbicide through prolonged and repeated use. Herbicides from different groups should either be used alternately or mixed together to prevent the numbers of resistant individuals building up. To minimize this risk use strictly in accordance with label instructions and resistance management strategies at www.resistance.nzpps.org

MIXING
Add the required amount of Patriot 1 to a partly filled spray tank with the agitation system operating. Complete filling the tank with water and apply, maintaining agitation.

COMPATIBILITY
Patriot 1 is compatible with most commonly used residual herbicides.

RAINFALL
Rainfall within 2 hours of application may wash Patriot 1 off the foliage, and within 6 hours will reduce effectiveness to the degree that re-treatment may be necessary.

USING THE SPRAY
IMPORTANT: In order to select the appropriate nozzles for the required spray quality and to operate the equipment to minimise spray drift, this product must be applied in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).

WEED SUSCEPTIBILITY
The lower rates indicated are appropriate when weeds are young, succulent and actively growing, or when the weed population is sparse and good coverage and penetration is possible. Use the higher rates when weeds are mature or the population is dense.

NB: Patriot 1 will not completely control mature weeds with well-developed tap roots or underground storage organs: some re-growth may occur following treatment.

RECOMMENDATIONS FOR USE

USE	RATE	REMARKS
Berryfruit – Raspberries and Brambles Primocane control	500-750ml/100 litres applied through cone nozzles in sufficient water to ensure complete coverage of primocanes.	Apply as a directed high volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of a non-ionic wetter at label rate is recommended. "Knockdown" takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached 15cm in height. Do not apply to plants under stress.
Orchard and vineyard weed control. Pipfruit, Kiwifruit, Stonefruit, Grapes and Citrus	Use a rate appropriate to the weeds present. If dilution rate is less than 1 litre/100 litres of water add a non-ionic wetter at label rates.	Apply in spring when the majority of the annual weeds have germinated. Residual herbicides may be applied in mixture for long-term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water to ensure complete coverage. May be applied up to one day before crop emergence.
Non-cropland. General weed control.	7.5-10 litres/ha in 300-500 litres of water or 100ml/10 litre knapsack.	Apply in a drenching spray to the point of run-off. May be mixed with residual herbicides for long-term weed control.

INDICATED RATE RANGE FOR VARIOUS WEED SPECIES

3 - 5 Litres/ha

Annual Poa	Asparagus seedlings	Beaked Hawksbeard	Black Nightshade	Calandrinia
Chickweed	Cleavers	Clover seedlings	Cornbind	Daisy
Dandelion	Mouse-ear chickweed	Musky Storksbill	Redroot	Rough-stalked meadow grass
Scotch Thistle	Sow Thistle	Staggerweed	Summer grass	Spurrey
Stinking Mayweed	Subterranean clover	Wild turnip	Willow weed	Wireweed

5 – 7.5 Litres/ha

Bristle grass	Bur medick	Catsear	Chewing's Fescue	Clovers
Cut-leaved geranium	Docks	Fleabane	Parsley Piert	Prairie grass
Ryegrasses*	Sheep's sorrel	Storksbill	Twin cress	Yorkshire fog

7.5 – 10 Litres/ha

Broom	Buttercups*	Couch	Crowfoot grass	Field pansy
Gorse	Hawksbeard	Kikuyu grass	Mallows	Paspalum
Tall Fescue	Wild strawberry	Willow herb	Yarrow	

10 Litres/ha

Bracken	Indian doab	Meadow rice grass	Plantains	Rape
Rushes	Sedges			

*Resistant weeds: Hairy buttercup, some ryegrass strains.

WITHHOLDING PERIODS: 28 days for all crops
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.
ADDITIONAL INFORMATION
Patriot 1 is absorbed into the foliage and green stems of actively growing plants. It is inactive in soil and is not translocated as an active herbicide, so thorough coverage of target weeds is essential for good results. Visual effects of treatment will become evident more rapidly under ideal growing conditions, generally within 3-10 days. Do not spray in temperatures below 10°C. Avoid contact on green parts (including immature bark) of desirable plants, as damage may occur. Do not mow or cultivate after treatment until weeds start to die off.
Registered pursuant to the ACVM Act 1997, No. P9924
See www.foodsafety.govt.nz for registration conditionsz
Approved pursuant to the HSNO Act 1996, Approval No. HSR100830
See www.epa.govt.nz for approval controls

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

300X200 (1000 L)

Pantone 7485C

Pantone 368C

DANGER • ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Blue AG

Patriot 1®

NON-SELECTIVE CONTACT HERBICIDE

GROUP 10 HERBICIDE

ACTIVE INGREDIENT

200 g/litre Glufosinate-ammonium

in the form of a soluble concentrate.

A non-selective contact herbicide for the control of a wide range of grasses, broad leaf weeds and clovers in a variety of crop land and non-crop land situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS) 0800 CHEMCALL OR 0800 243 622 OR PGG WRIGHTSON LTD 0800 10 22 76

1000 L
NET CONTENTS

300X200 (1000 L)

Pantone 425C

BLACK
C:60 M:40 Y:40 K:100

Pantone 368C

Pantone 485 C

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

PEEL OFF

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

HAZARD CLASSIFICATIONS
Skin irritation Category 2, Eye irritation Category 2, Reproductive toxicity Category 2, Specific target organ toxicity (repeated exposure) Category 1, Hazardous to soil organisms, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment chronic Category 3.

TOXICITY
DANGER: Causes damage to kidneys and nervous system through prolonged or repeated exposure via any route. Causes eye irritation and skin irritation. Suspected of damaging fertility or the unborn child through repeated exposure at high doses.

ECOTOXIC
Harmful to aquatic life. This substance must not be applied onto or into water. Toxic to soil organisms. Harmful to terrestrial vertebrates. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

HANDLING PRECAUTIONS: Do NOT handle this product until all safety precautions have been read and understood. Keep out of reach of children. Avoid inhalation of spray mist. Avoid contact with eyes and skin. Do not eat, drink or smoke while using. Use only outdoors in a well-ventilated area.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS 8409). Flush equipment thoroughly with clean water after use.

STORAGE: Store in the original container locked up in a cool well ventilated place, out of direct sunlight and away stockfeed or foodstuff. Store in accordance with the current version of NZS 8409 Management of Agrichemicals.

PROTECTIVE CLOTHING: When mixing or applying wear appropriate protective clothing including impervious gloves, chemical resistant boots, eye protection (goggles with side pieces or a face shield) and cotton overalls buttoned to the neck and wrist. Remove protective clothing and wash hands, arms and face thoroughly with soap and water before meals and after work.

SPILLAGE: Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways; bund if necessary. Wear appropriate protective clothing as in the Precautions box of this label to clean up spills. Absorb spillage with inert absorbent material, such as vermiculite, sand or clay, and place in waste containers. Wash contaminated area with water and detergent, absorb and add to waste for disposal. Dispose of the waste safely in an approved landfill.

DISPOSAL INFORMATION
PRODUCT DISPOSAL: Dispose of product only by using in accordance with label directions, through the Agrecovery® Chemical Recovery service, or at an approved landfill.
CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Take empty container to an Agrecovery® collection site. If this is not possible bury in an approved landfill. Do not contaminate aquatic environment with product or empty container.

FIRST AID
If exposed or concerned get medical advice/attention.
If swallowed, rinse mouth. Do NOT induce vomiting. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell.
If inhaled, Remove patient to fresh air and maintain in a position comfortable for breathing. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor for advice if person feels unwell.
If in eyes, Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical advice/attention.
If on skin, wash with plenty of soap and water. If skin irritation persists, get medical advice/attention. Wash contaminated clothing before reuse.

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

9 415251 004867 >

300X200 (1000 L)

Pantone 425C

MULTIFOLD®

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

DIRECTIONS FOR USE
BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS. Apply using ground-based application only. The maximum application rate is 10L/ha (2.03kg ai/ha) with a maximum of two applications per year and minmum application interval of 90 days.

RESISTANCE MANAGEMENT
Patriot 1 contains GROUP 10 herbicide, Glufosinate-ammonium, a glutamine synthetase inhibitor. Certain weeds may develop resistance to this herbicide through prolonged and repeated use. Herbicides from different groups should either be used alternately or mixed together to prevent the numbers of resistant individuals building up. To minimize this risk use strictly in accordance with label instructions and resistance management strategies at www.resistance.nzpps.org

MIXING
Add the required amount of Patriot 1 to a partly filled spray tank with the agitation system operating. Complete filling the tank with water and apply, maintaining agitation.

COMPATIBILITY
Patriot 1 is compatible with most commonly used residual herbicides.

RAINFALL
Rainfall within 2 hours of application may wash Patriot 1 off the foliage, and within 6 hours will reduce effectiveness to the degree that re-treatment may be necessary.

USING THE SPRAY
IMPORTANT: In order to select the appropriate nozzles for the required spray quality and to operate the equipment to minimise spray drift, this product must be applied in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).

WEED SUSCEPTIBILITY
The lower rates indicated are appropriate when weeds are young, succulent and actively growing, or when the weed population is sparse and good coverage and penetration is possible. Use the higher rates when weeds are mature or the population is dense.

NB: Patriot 1 will not completely control mature weeds with well-developed tap roots or underground storage organs: some re-growth may occur following treatment.

RECOMMENDATIONS FOR USE

USE	RATE	REMARKS
Berryfruit – Raspberries and Brambles Primocane control	500-750ml/100 litres applied through cone nozzles in sufficient water to ensure complete coverage of primocanes.	Apply as a directed high volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of a non-ionic wetter at label rate is recommended. "Knockdown" takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached 15cm in height. Do not apply to plants under stress.
Orchard and vineyard weed control. Pipfruit, Kiwifruit, Stonefruit, Grapes and Citrus	Use a rate appropriate to the weeds present. If dilution rate is less than 1 litre/100 litres of water add a non-ionic wetter at label rates.	Apply in spring when the majority of the annual weeds have germinated. Residual herbicides may be applied in mixture for long-term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water to ensure complete coverage. May be applied up to one day before crop emergence.
Non-cropland. General weed control.	7.5-10 litres/ha in 300-500 litres of water or 100ml/10 litre knapsack.	Apply in a drenching spray to the point of run-off. May be mixed with residual herbicides for long-term weed control.

INDICATED RATE RANGE FOR VARIOUS WEED SPECIES

3 - 5 Litres/ha

Annual Poa	Asparagus seedlings	Beaked Hawksbeard	Black Nightshade	Calandrinia
Chickweed	Cleavers	Clover seedlings	Cornbind	Daisy
Dandelion	Mouse-ear chickweed	Musky Storksbill	Redroot	Rough-stalked meadow grass
Scotch Thistle	Sow Thistle	Staggerweed	Summer grass	Spurrey
Stinking Mayweed	Subterranean clover	Wild turnip	Willow weed	Wireweed

5 – 7.5 Litres/ha

Bristle grass	Bur medick	Catsear	Chewing's Fescue	Clovers
Cut-leaved geranium	Docks	Fleabane	Parsley Piert	Prairie grass
Ryegrasses*	Sheep's sorrel	Storksbill	Twin cress	Yorkshire fog

7.5 – 10 Litres/ha

Broom	Buttercups*	Couch	Crowfoot grass	Field pansy
Gorse	Hawksbeard	Kikuyu grass	Mallows	Paspalum
Tall Fescue	Wild strawberry	Willow herb	Yarrow	

10 Litres/ha

Bracken	Indian doab	Meadow rice grass	Plantains	Rape
Rushes	Sedges			

*Resistant weeds: Hairy buttercup, some ryegrass strains.

WITHHOLDING PERIODS: 28 days for all crops
It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.
ADDITIONAL INFORMATION
Patriot 1 is absorbed into the foliage and green stems of actively growing plants. It is inactive in soil and is not translocated as an active herbicide, so thorough coverage of target weeds is essential for good results. Visual effects of treatment will become evident more rapidly under ideal growing conditions, generally within 3-10 days. Do not spray in temperatures below 10°C. Avoid contact on green parts (including immature bark) of desirable plants, as damage may occur. Do not mow or cultivate after treatment until weeds start to die off.
Registered pursuant to the ACVM Act 1997, No. P9924
See www.foodsafety.govt.nz for registration conditionsz
Approved pursuant to the HSNO Act 1996, Approval No. HSR100830
See www.epa.govt.nz for approval controls

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

150 X 240 (20 L)

Pantone 7485C

Pantone 368C

DANGER - POISON

KEEP OUT OF REACH OF CHILDREN

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

Shaleen Narayan
Manager Approvals

Blue AG

Patriot 1®

NON-SELECTIVE CONTACT HERBICIDE

GROUP **10** HERBICIDE

ACTIVE INGREDIENT

200 g/litre Glufosinate-ammonium

in the form of a soluble concentrate.

A non-selective contact herbicide for the control of a wide range of grasses, broad leaf weeds and clovers in a variety of crop land and non-crop land situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS)
0800 CHEMCALL OR 0800 243 622 OR
PGG WRIGHTSON LTD 0800 10 22 76

1 L
NET CONTENTS

150 X 240 (20 L)

Pantone 425C | BLACK C:60 M:40 Y:40 K:100 | Pantone 368C | Pantone 485 C

PEEL OFF

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

HAZARD CLASSIFICATIONS

Skin irritation Category 2, Eye irritation Category 2, Reproductive toxicity Category 2, Specific target organ toxicity (repeated exposure) Category 1, Hazardous to soil organisms, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment chronic Category 3.

TOXICITY

DANGER: Causes damage to kidneys and nervous system through prolonged or repeated exposure via any route. Causes eye irritation and skin irritation. Suspected of damaging fertility or the unborn child through repeated exposure at high doses.

ECOTOXIC

Harmful to aquatic life. This substance must not be applied onto or into water. Toxic to soil organisms. Harmful to terrestrial vertebrates. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

HANDLING PRECAUTIONS: Do NOT handle this product until all safety precautions have been read and understood. Keep out of reach of children. Avoid inhalation of spray mist. Avoid contact with eyes and skin. Do not eat, drink or smoke while using. Use only outdoors in a well-ventilated area.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS 8409). Flush equipment thoroughly with clean water after use.

STORAGE: Store in the original container locked up in a cool well ventilated place, out of direct sunlight and away stockfeed or foodstuff. Store in accordance with the current version of NZS 8409 Management of Agrichemicals.

PROTECTIVE CLOTHING: When mixing or applying wear appropriate protective clothing including impervious gloves, chemical resistant boots, eye protection (goggles with side pieces or a face shield) and cotton overalls buttoned to the neck and wrist. Remove protective clothing and wash hands, arms and face thoroughly with soap and water before meals and after work.

SPILLAGE: Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways, bunds, drains. Wear appropriate protective clothing as above. Precautions based on this label apply to spills. Absorb spillage with inert absorbent material such as vermiculite, sand or [illegible] waste containers. Wash contaminated areas with water and detergent, absorb and add to waste for disposal. Dispose of the waste safely in an approved landfill.

DISPOSAL INFORMATION

PRODUCT DISPOSAL: Dispose of product only by using in accordance with label directions, through the Agrecovery® Chemical Recovery service, or at an approved landfill.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Take empty container to an Agrecovery® collection site. If this is not possible bury in an approved landfill. Do not contaminate aquatic environment with product or empty container.

FIRST AID

If exposed or concerned get medical advice/attention.

If swallowed, rinse mouth. Do NOT induce vomiting. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell.

If inhaled, Remove patient to fresh air and maintain in a position comfortable for breathing. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor for advice if person feels unwell.

If in eyes, Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical advice/attention.

If on skin, wash with plenty of soap and water. If skin irritation persists, get medical advice/attention. Wash contaminated clothing before reuse.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
10/10/2025

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use.
PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

150 X 240 (20 L)

Pantone 425C

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

DIRECTIONS FOR USE
BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS. Apply using ground-based application only. The maximum application rate is 10L/ha (2.03kg ai/ha) with a maximum of two applications per year and minmum application interval of 90 days.

RESISTANCE MANAGEMENT
Patriot 1 contains the GROUP 10 herbicide, Glufosinate-ammonium, a glutamine synthetase inhibitor. Certain weeds may develop resistance to this herbicide through prolonged and repeated use. Herbicides from different groups should either be used alternately or mixed together to prevent the numbers of resistant individuals building up. To minimize this risk use strictly in accordance with label instructions and strictly in accordance with label instructions and resistance management strategies at www.resistance.nzpps.org

MIXING
Add the required amount of Patriot 1 to a partly filled spray tank with the agitation system operating. Complete filling the tank with water and apply, maintaining agitation.

COMPATIBILITY
Patriot 1 is compatible with most commonly used residual herbicides.

RAINFALL
Rainfall within 2 hours of application may wash the product off the foliage, and within 6 hours will reduce the effectiveness to the degree that re-treatment may be necessary.

USING THE SPRAY
IMPORTANT: In order to select the appropriate nozzle for the required spray quality and to operate the equipment to minimise spray drift, the product must be applied in accordance with the New Zealand standard for the Management of Agrichemicals (NZS8409).

WEED SUSCEPTIBILITY
The lower rates indicated are appropriate when weeds are young, succulent and actively growing, or when the weed population is sparse and good coverage and penetration is possible. Use the higher rates when weeds are mature or the population is dense.

NB: Patriot 1 will not completely control mature weeds with well-developed tap roots or underground storage organs: some re-growth may occur following treatment.

RECOMMENDATIONS FOR USE

USE	RATE	REMARKS
Berryfruit – Raspberries and Brambles Primocane control	500-750ml/100 litres applied through cone nozzles in sufficient water to ensure complete coverage of primocanes.	Apply as a directed high volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of a non-ionic wetter at label rate is recommended. "Knockdown" takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached 15cm in height. Do not apply to plants under stress.
Orchard and vineyard weed control. Pipfruit, Kiwifruit, Stonefruit, Grapes and Citrus	Use a rate appropriate to the weeds present. If dilution rate is less than 1 litre/100 litres of water add a non-ionic wetter at label rates.	Apply in spring when the majority of the annual weeds have germinated. Residual herbicides may be applied in mixture for long-term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water to ensure complete coverage. May be applied up to one day before crop emergence.
Non-cropland. General weed control.	7.5-10 litres/ha in 300-500 litres of water or 100ml/ 10 litre knapsack.	Apply in a drenching spray to the point of run-off. May be mixed with residual herbicides for long-term weed control.

INDICATED RATE RANGE FOR VARIOUS WEED SPECIES

3 - 5 Litres/ha

Annual Poa	Asparagus seedlings	Beaked Hawksbeard	Black Nightshade	Calandrinia
Chickweed	Cleavers	Clover seedlings	Cornbind	Daisy
Dandelion	Mouse-ear chickweed	Musky Storksbill	Redroot	Rough-stalked meadow grass
Scotch Thistle	Sow Thistle	Staggerweed	Summer grass	Spurrey
Stinking Mayweed	Subterranean clover	Wild turnip	Willow weed	Wireweed

5 – 7.5 Litres/ha

Bristle grass	Bur medick	Catsear	Chewing's Fescue	Clovers
Cut-leaved geranium	Docks	Fleabane	Parsley Piert	Prairie grass
Ryegrasses*	Sheep's sorrel	Storksbill	Twin cress	Yorkshire fog

7.5 – 10 Litres/Ha

Broom	Buttercups*	Couch	Crowfoot grass	Field pansy
Gorse	Hawksbeard	Kikuyu grass	Mallows	Paspalum
Tall Fescue	Wild strawberry	Willow herb	Yarrow	

10 Litres/ha

Bracken	Indian doab	Meadow rice grass	Plantains	Rape
Rushes	Sedges			

*Resistant weeds: Hairy buttercup, some ryegrass strains.

WITHHOLDING PERIODS: 28 days for all crops. It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.
ADDITIONAL INFORMATION: Patriot 1 is absorbed into the foliage and green stems of actively growing plants. It is inactive in soil and is not translocated as an active herbicide, so thorough coverage of target weeds is essential for good results. Visual effects of treatment will become evident more rapidly under ideal growing conditions, generally within 3-10 days. Do not spray in temperatures below 10°C. Avoid contact on green parts (including immature bark) of desirable plants, as damage may occur. Do not mow or cultivate after treatment until weeds start to die off.
Registered pursuant to the ACVM Act 1997, No. P9924 See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval No. HSR100830
See www.epa.govt.nz for approval controls

MULTIFOLD®

150 X 240 (20 L)

Pantone 7485C

Pantone 368C

DANGER • ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/10/2025

Blue AG

Patriot 1®

NON-SELECTIVE CONTACT HERBICIDE

GROUP 10 HERBICIDE

ACTIVE INGREDIENT

200 g/litre Glufosinate-ammonium

in the form of a soluble concentrate.

A non-selective contact herbicide for the control of a wide range of grasses, broad leaf weeds and clovers in a variety of crop land and non-crop land situations.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS)
0800 CHEMCALL OR 0800 243 622 OR
PGG WRIGHTSON LTD 0800 10 22 76

10 L
NET CONTENTS

150 X 240 (20 L) | Pantone 425C | BLACK C:60 M:40 Y:40 K:100 | Pantone 368C | Pantone 485 C

PEEL OFF

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

HAZARD CLASSIFICATIONS

Skin irritation Category 2, Eye irritation Category 2, Reproductive toxicity Category 2, Specific target organ toxicity (repeated exposure) Category 1, Hazardous to soil organisms, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment chronic Category 3.

TOXICITY

DANGER: Causes damage to kidneys and nervous system through prolonged or repeated exposure via any route. Causes eye irritation and skin irritation. Suspected of damaging fertility or the unborn child through repeated exposure at high doses.

ECOTOXIC

Harmful to aquatic life. This substance must not be applied onto or into water. Toxic to soil organisms. Harmful to terrestrial vertebrates. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

HANDLING PRECAUTIONS: Do NOT handle this product until all safety precautions have been read and understood. Keep out of reach of children. Avoid inhalation of spray mist. Avoid contact with eyes and skin. Do not eat, drink or smoke while using. Use only outdoors in a well-ventilated area.

EQUIPMENT: Apply using accurately calibrated and maintained equipment in accordance with The New Zealand Standard for the Management of Agrichemicals (NZS 8409). Flush equipment thoroughly with clean water after use.

STORAGE: Store in the original container locked up in a cool well ventilated place, out of direct sunlight and away stockfeed or foodstuff. Store in accordance with the current version of NZS 8409 Management of Agrichemicals.

PROTECTIVE CLOTHING: When mixing or applying wear appropriate protective clothing including impervious gloves, chemical resistant boots, eye protection (goggles with side pieces or a face shield) and cotton overalls buttoned to the neck and wrist. Remove protective clothing and wash hands, arms and face thoroughly with soap and water before meals and after work.

SPILLAGE: Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways, bunds if possible. Wear appropriate protective clothing as per Precautions box of this label. Contain spills. Absorb spillage with inert absorbent material such as vermiculite, sand or [illegible] in waste containers. Wash contaminated area with water and detergent, absorb and add to waste for disposal. Dispose of the waste safely in an approved landfill.

DISPOSAL INFORMATION

PRODUCT DISPOSAL: Dispose of product only by using in accordance with label directions, through the Agrecovery® Chemical Recovery service, or at an approved landfill.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Take empty container to an Agrecovery® collection site. If this is not possible bury in an approved landfill. Do not contaminate aquatic environment with product or empty container.

FIRST AID

If exposed or concerned get medical advice/attention.

If swallowed, rinse mouth. Do NOT induce vomiting. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell.

If inhaled, Remove patient to fresh air and maintain in a position comfortable for breathing. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor for advice if person feels unwell.

If in eyes, Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists get medical advice/attention.

If on skin, wash with plenty of soap and water. If skin irritation persists, get medical advice/attention. Wash contaminated clothing before reuse.

INSPECTED FOR PRIMARY INDUSTRIES NEW ZEALAND
10/10/2025

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.

This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use.
PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

150 X 240 (20 L)

Pantone 425C

READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING

DIRECTIONS FOR USE
BEWARE: APPLY THIS PRODUCT CAREFULLY. SPRAY DRIFT MAY CAUSE SERIOUS DAMAGE TO OTHER DESIRABLE PLANTS. Apply using ground-based application only. The maximum application rate is 10L/ha (2.03kg ai/ha) with a maximum of two applications per year and minmum application interval of 90 days.

RESISTANCE MANAGEMENT
Patriot 1 contains the GROUP 10 herbicide, Glufosinate-ammonium, a glutamine synthetase inhibitor. Certain weeds may develop resistance to this herbicide through prolonged and repeated use. Herbicides from different groups should either be used alternately or mixed together to prevent the numbers of resistant individuals building up. To minimize this risk use strictly in accordance with label instructions and strictly in accordance with label instructions and resistance management strategies at www.resistance.nzpps.org

MIXING
Add the required amount of Patriot 1 to a partly filled spray tank with the agitation system operating. Complete filling the tank with water and apply, maintaining agitation.

COMPATIBILITY
Patriot 1 is compatible with most commonly used residual herbicides.

RAINFALL
Rainfall within 2 hours of application may wash the product off the foliage, and within 6 hours will reduce the effectiveness to the degree that re-treatment may be necessary.

USING THE SPRAY
IMPORTANT: In order to select the appropriate nozzle for the required spray quality and to operate the equipment to minimise spray drift, this product must be applied in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409).

WEED SUSCEPTIBILITY
The lower rates indicated are appropriate when weeds are young, succulent and actively growing, or when the weed population is sparse and good coverage and penetration is possible. Use the higher rates when weeds are mature or the population is dense.

NB: Patriot 1 will not completely control mature weeds with well-developed tap roots or underground storage organs: some re-growth may occur following treatment.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
10/10/2025

RECOMMENDATIONS FOR USE

USE	RATE	REMARKS
Berryfruit – Raspberries and Brambles Primocane control	500-750ml/100 litres applied through cone nozzles in sufficient water to ensure complete coverage of primocanes.	Apply as a directed high volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of a non-ionic wetter at label rate is recommended. "Knockdown" takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached 15cm in height. Do not apply to plants under stress.
Orchard and vineyard weed control. Pipfruit, Kiwifruit, Stonefruit, Grapes and Citrus	Use a rate appropriate to the weeds present. If dilution rate is less than 1 litre/100 litres of water add a non-ionic wetter at label rates.	Apply in spring when the majority of the annual weeds have germinated. Residual herbicides may be applied in mixture for long-term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water to ensure complete coverage. May be applied up to one day before crop emergence.
Non-cropland. General weed control.	7.5-10 litres/ha in 300-500 litres of water or 100ml/ 10 litre knapsack.	Apply in a drenching spray to the point of run-off. May be mixed with residual herbicides for long-term weed control.

INDICATED RATE RANGE FOR VARIOUS WEED SPECIES

3 - 5 Litres/ha

Annual Poa	Asparagus seedlings	Beaked Hawksbeard	Black Nightshade	Calandrinia
Chickweed	Cleavers	Clover seedlings	Cornbind	Daisy
Dandelion	Mouse-ear chickweed	Musky Storksbill	Redroot	Rough-stalked meadow grass
Scotch Thistle	Sow Thistle	Staggerweed	Summer grass	Spurrey
Stinking Mayweed	Subterranean clover	Wild turnip	Willow weed	Wireweed

5 – 7.5 Litres/ha

Bristle grass	Bur medick	Catsear	Chewing's Fescue	Clovers
Cut-leaved geranium	Docks	Fleabane	Parsley Piert	Prairie grass
Ryegrasses*	Sheep's sorrel	Storksbill	Twin cress	Yorkshire fog

7.5 – 10 Litres/Ha

Broom	Buttercups*	Couch	Crowfoot grass	Field pansy
Gorse	Hawksbeard	Kikuyu grass	Mallows	Paspalum
Tall Fescue	Wild strawberry	Willow herb	Yarrow	

10 Litres/ha

Bracken	Indian doab	Meadow rice grass	Plantains	Rape
Rushes	Sedges			

*Resistant weeds: Hairy buttercup, some ryegrass strains.

WITHHOLDING PERIODS: 28 days for all crops. It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.
ADDITIONAL INFORMATION: Patriot 1 is absorbed into the foliage and green stems of actively growing plants. It is inactive in soil and is not translocated as an active herbicide, so thorough coverage of target weeds is essential for good results. Visual effects of treatment will become evident more rapidly under ideal growing conditions, generally within 3-10 days. Do not spray in temperatures below 10°C. Avoid contact on green parts (including immature bark) of desirable plants, as damage may occur. Do not mow or cultivate after treatment until weeds start to die off.
Registered pursuant to the ACVM Act 1997, No. P9924 See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval No. HSR100830
See www.epa.govt.nz for approval controls

MULTIFOLD®