WARNING
Keep out of reach of children

ECOTOXIC

Swap 375 + 250 WG™

FUNGICIDE

For the control of Botrytis in Grapes and Onions, Botrytis, Anthracnose and Rhizopus in Strawberries
and Botrytis and Anthracnose in Blueberries

Active Ingredient: 375 g/kg CYPRODINIL and 250 g/kg FLUDIOXONIL in the form of a water dispersible granule

Distributed by: Farmlands Co-operative Society Limited
535 Wairakei Road, Burnside, Christchurch 8053
TEL: 0800 200 600

Net contents: 500g, 1 kg, 5 kg, 10 kg, 20 kg, 25 kg

Batch:
D.O.M.:

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
29/04/2024

PLEASE READ THE LABEL COMPLETELY BEFORE USE

HAZARD CLASSIFICATIONS: Specific target organ toxicity (repeated exposure) Category 2, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1.

TOXICITY: May cause organ damage from repeated ingestion at high doses.

ECOTOXICITY:
Very toxic to aquatic life and with long-lasting effects. Harmful to terrestrial vertebrates.
Do not apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

TRAINED HANDLER: This product must be under the control of a trained handler during mixing, loading and use.
RECORD-KEEPING: Records of use are required when 3 kg or more is applied within a 24 hour period in an application area where the substance is likely to enter air or water and leave the application area.

PRECAUTIONS
Avoid contact with eyes and skin. Avoid inhalation of dust or spray mist. When mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before rest breaks, meals and after work.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure place away from foodstuffs. Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals. Storage of 100kg or more requires signage, secondary containment and an emergency response plan.
USE: Apply with well maintained and calibrated spray equipment.
DISPOSAL:
Do not contaminate environment with product or container.
Product disposal - If possible dispose of by using according to the label, otherwise dispose of in an approved landfill.
Container disposal - Ensure container is completely empty. Dispose of to an agricultural recycling service or bury in suitable landfill. Do not burn. Do not use container for any other purpose.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
29/04/2024

FIRST AID
Call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor for advice.
If swallowed do NOT induce vomiting. Rinse mouth. Get medical advice if person feels unwell. If splashed in eyes, wash out immediately with running water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention. If skin or hair contact occurs, remove contaminated clothing and wash affected area with soap and running water. If skin irritation persists, get medical attention. If inhaled, immediately move to fresh air and rest. Begin artificial respiration if the victim is not breathing. Use mouth to nose rather than mouth to mouth. Get immediate medical assistance.

Emergency Response
In a transport emergency call **111, Police or Fire Brigade**. In a fire, evacuate and call **111 Fire Brigade.**

SPILLAGE
Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. If granules have spilt, moisten to prevent dusting and sweep up. If liquid, absorb spill with inert material such as soil, sand, etc. Place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfill.

GENERAL INFORMATION

Swap 375 + 250 WG is a mixture of anilinopyrimidine and phenylpyrrole fungicides, which in combination have protectant and systemic activity against *Botrytis* (grey mould) infections in grapes and bulb onions, *Botrytis* (grey mould), *Colletotrichum* (anthracnose) and *Rhizopus* (leak) infections in strawberry, and Botrytis (grey mould) and Colletotrichum (anthracnose) in blueberries. **Swap 375 + 250 WG** has multi-site action, inhibiting spore germination, penetration of the germ tube and mycelial growth. It is rainfast within 2 hours following application.

MIXING
Add the required amount of **Swap 375 + 250 WG** into the mixing chamber of the partly filled spray tank and add the remainder of the water, ensuring complete mixing before spraying. Continue agitation during spraying.

COMPATIBILITY
Swap 375 + 250 WG is compatible with other commonly used fungicides and insecticides.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
29/04/2024

DIRECTIONS FOR USE

It is an offence to use this product on animals.

Apply only by ground-based application methods.

Crop	Disease	Rate	Remarks
Blueberries	Grey Mould (*Botrytis*), Anthracnose (*Colletotrichum*)	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	A maximum of 2 applications of **Swap 375 + 250 WG** can be made per season. Apply in sufficient water to ensure thorough and uniform coverage without runoff (Approx. 1000 L/ha). The first application should be made at full flower, with the second application no later than 14 days after the completion of flowering.
Grapes	*Botrytis*	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	The rate suggested is for high volume spraying to runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage of the bunch zone is achieved. The addition of Synergy ™ Wetter Sticker at label rates will assist coverage. A maximum of 2 applications per season can be made. **Swap 375 + 250 WG** can be applied at either early flowering (5% capfall), late flowering (80% capfall), pre bunch closure or in the period from the beginning of berry ripening up until 28 days before harvest. Apply an approved fungicide from a different chemical group for the remainder of the season.
Bulb Onions		800 g/ha	A maximum of 2 applications of **Swap 375 + 250 WG** can be made per season at key timings for *Botrytis* control. Apply in sufficient water to ensure thorough and uniform coverage. Applications MUST be made with the addition of Synergy Wetter Sticker at label rates.
Strawberries	Grey Mould (*Botrytis*), Anthracnose (*Colletotrichum*) and leak (*Rhizopus*)	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	The rate suggested is for high volume spraying to runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved. The addition of Synergy Wetter Sticker at label rates will assist coverage. **Swap 375 + 250 WG** can be applied at 7 to 14 day intervals from flowering onwards. The closer interval is recommended when conditions are conducive to disease infection. **Swap 35 + 250 WG** applications may be applied in a block or in alternation with other non anilinopyrimidine fungicides registered for botrytis control. See resistance management for the maximum number of applications.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

Blueberries	Last application no later than 14 days after flowering and at least 28 days before harvest
Strawberries	3 days
Bulb onions	7 days
Grapes	28 days

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
29/04/2024

RESISTANCE MANAGEMENT

GROUP	9 + 12	FUNGICIDE

Swap 375 + 250 WG contains a Group 9 fungicide (cyprodinil, belonging to the anilinopyrimidine class of chemistry) and Group 12 fungicide (fludioxonil, belonging to the phenylpyrrole class of chemistry). Resistance to this fungicide and related anilinopyrimidine products could develop in some situations from repeated use. To minimise this risk, use the product strictly in accordance with the label instructions.
Blueberries, Grapes and Bulb Onions: Do NOT use more than 2 applications of **Swap 375 + 250 WG** or any other anilinopyrimidine fungicide on the same crop during the season.
Strawberries: A maximum of 3 applications of **Swap 375 + 250 WG** or any other anilinopyrimidine fungicide per season can be made if more than six sprays in total are made for Botrytis control. If 6 or less Botrytis control sprays are made then do NOT apply more than 2 applications of **Swap 375 + 250 WG** or any other anilinopyrimidine fungicide per season.
Consult http://resistance.nzpps.org/index.php for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P10008
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR100937
See www.epa.govt.nz for approval controls.

Swap 375 + 250 WG™ is a tradename of AGROBEATS TECH CO., LTD

Registrant:
AGROBEATS TECH CO., LTD
C-47,No.69, Shuangfeng Road, Pukou Economic Development Zone, Pukou District, Nanjing, P.R. China

NZ Agent:
AgStar New Zealand Pty Ltd
4 Putakitaki Street, Lincoln, 7608
Tel 027 483 9756

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (CYPRODINIL 38%, FLUDIOXONIL 25%)
PACKING GROUP III
HAZCHEM CODE 2Z
IN A TRANSPORT EMERGENCY, DIAL 111 POLICE OR FIRE BRIGADE.

WARNING
Keep out of reach of children

ECOTOXIC

Swap 375 + 250 WG™

FUNGICIDE

For the control of Botrytis in Grapes and Onions, Botrytis, Anthracnose and Rhizopus in Strawberries
and Botrytis and Anthracnose in Blueberries

Active Ingredient: 375 g/kg CYPRODINIL and 250 g/kg FLUDIOXONIL in the form of a water dispersible granule

Net contents: 500g, 1 kg, 5 kg, 10 kg, 20 kg, 25 kg

Batch:
D.O.M.:

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
29/04/2024

PLEASE READ THE LABEL COMPLETELY BEFORE USE

HAZARD CLASSIFICATIONS: Specific target organ toxicity (repeated exposure) Category 2, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1.

TOXICITY: May cause organ damage from repeated ingestion at high doses.

ECOTOXICITY:
Very toxic to aquatic life and with long-lasting effects. Harmful to terrestrial vertebrates.
Do not apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

TRAINED HANDLER: This product must be under the control of a trained handler during mixing, loading and use.
RECORD-KEEPING: Records of use are required when 3 kg or more is applied within a 24 hour period in an application area where the substance is likely to enter air or water and leave the application area.

PRECAUTIONS
Avoid contact with eyes and skin. Avoid inhalation of dust or spray mist. When mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before rest breaks, meals and after work.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure place away from foodstuffs. Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals. Storage of 100kg or more requires signage, secondary containment and an emergency response plan.
USE: Apply with well maintained and calibrated spray equipment.
DISPOSAL:
Do not contaminate environment with product or container.
Product disposal - If possible dispose of by using according to the label, otherwise dispose of in an approved landfill.
Container disposal - Ensure container is completely empty. Dispose of to an agricultural recycling service or bury in suitable landfill. Do not burn. Do not use container for any other purpose.

FIRST AID
Call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor for advice.
If swallowed do NOT induce vomiting. Rinse mouth. Get medical advice if person feels unwell. If splashed in eyes, wash out immediately with running water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention. If skin or hair contact occurs, remove contaminated clothing and wash affected area with soap and running water. If skin irritation persists, get medical attention. If inhaled, immediately move to fresh air and rest. Begin artificial respiration if the victim is not breathing. Use mouth to nose rather than mouth to mouth. Get immediate medical assistance.

Emergency Response
In a transport emergency call **111, Police or Fire Brigade**. In a fire, evacuate and call **111 Fire Brigade.**

MINISTRY FOR PRIMARY INDUSTRIES
29/04/2024

SPILLAGE
Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. If granules have spilt, moisten to prevent dusting and sweep up. If liquid, absorb spill with inert material such as soil, sand, etc. Place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfill.

GENERAL INFORMATION

Swap 375 + 250 WG is a mixture of anilinopyrimidine and phenylpyrrole fungicides, which in combination have protectant and systemic activity against *Botrytis* (grey mould) infections in grapes and bulb onions, *Botrytis* (grey mould), *Colletotrichum* (anthracnose) and *Rhizopus* (leak) infections in strawberry, and Botrytis (grey mould) and Colletotrichum (anthracnose) in blueberries. **Swap 375 + 250 WG** has multi-site action, inhibiting spore germination, penetration of the germ tube and mycelial growth. It is rainfast within 2 hours following application.

MIXING
Add the required amount of **Swap 375 + 250 WG** into the mixing chamber of the partly filled spray tank and add the remainder of the water, ensuring complete mixing before spraying. Continue agitation during spraying.

COMPATIBILITY
Swap 375 + 250 WG is compatible with other commonly used fungicides and insecticides.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
29/04/2024

DIRECTIONS FOR USE

It is an offence to use this product on animals.

Apply only by ground-based application methods.

Crop	Disease	Rate	Remarks
Blueberries	Grey Mould (*Botrytis*), Anthracnose (*Colletotrichum*)	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	A maximum of 2 applications of **Swap 375 + 250 WG** can be made per season. Apply in sufficient water to ensure thorough and uniform coverage without runoff (Approx. 1000 L/ha). The first application should be made at full flower, with the second application no later than 14 days after the completion of flowering.
Grapes	*Botrytis*	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	The rate suggested is for high volume spraying to runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage of the bunch zone is achieved. The addition of Synergy ™ Wetter Sticker at label rates will assist coverage. A maximum of 2 applications per season can be made. **Swap 375 + 250 WG** can be applied at either early flowering (5% capfall), late flowering (80% capfall), pre bunch closure or in the period from the beginning of berry ripening up until 28 days before harvest. Apply an approved fungicide from a different chemical group for the remainder of the season.
Bulb Onions		800 g/ha	A maximum of 2 applications of **Swap 375 + 250 WG** can be made per season at key timings for *Botrytis* control. Apply in sufficient water to ensure thorough and uniform coverage. Applications MUST be made with the addition of Synergy Wetter Sticker at label rates.
Strawberries	Grey Mould (*Botrytis*), Anthracnose (*Colletotrichum*) and leak (*Rhizopus*)	80 g/100 litres of water, applying sufficient mixture to ensure 800 g product /ha	The rate suggested is for high volume spraying to runoff. For concentrate spraying, adjust dilution rate accordingly. It is essential that good coverage is achieved. The addition of Synergy Wetter Sticker at label rates will assist coverage. **Swap 375 + 250 WG** can be applied at 7 to 14 day intervals from flowering onwards. The closer interval is recommended when conditions are conducive to disease infection. **Swap 375 + 250 WG** applications may be applied in a block or in alternation with other non anilinopyrimidine fungicides registered for botrytis control. See resistance management for the maximum number of applications.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

Blueberries	Last application no later than 14 days after flowering and at least 28 days before harvest
Strawberries	3 days
Bulb onions	7 days
Grapes	28 days

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
29/04/2024

RESISTANCE MANAGEMENT

GROUP	9 + 12	FUNGICIDE

Swap 375 + 250 WG contains a Group 9 fungicide (cyprodinil, belonging to the anilinopyrimidine class of chemistry) and Group 12 fungicide (fludioxonil, belonging to the phenylpyrrole class of chemistry). Resistance to this fungicide and related anilinopyrimidine products could develop in some situations from repeated use. To minimise this risk, use the product strictly in accordance with the label instructions.
Blueberries, Grapes and Bulb Onions: Do NOT use more than 2 applications of **Swap 375 + 250 WG** or any other anilinopyrimidine fungicide on the same crop during the season.
Strawberries: A maximum of 3 applications of **Swap 375 + 250 WG** or any other anilinopyrimidine fungicide per season can be made if more than six sprays in total are made for Botrytis control. If 6 or less Botrytis control sprays are made then do NOT apply more than 2 applications of **Swap 375 + 250 WG** or any other anilinopyrimidine fungicide per season.
Consult http://resistance.nzpps.org/index.php for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P10008
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR100937
See www.epa.govt.nz for approval controls.

Swap 375 + 250 WG™ is a tradename of AGROBEATS TECH CO., LTD

Registrant:
AGROBEATS TECH CO., LTD
C-47,No.69, Shuangfeng Road, Pukou Economic Development Zone, Pukou District, Nanjing, P.R. China

NZ Agent:
AgStar New Zealand Pty Ltd
4 Putakitaki Street, Lincoln, 7608
Tel 027 483 9756

UN 3077
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (CYPRODINIL 38%, FLUDIOXONIL 25%)
PACKING GROUP III
HAZCHEM CODE 2Z
IN A TRANSPORT EMERGENCY, DIAL 111 POLICE OR FIRE BRIGADE.