WARNING – KEEP OUT OF REACH OF CHILDREN

SPIROSEC 100SC

A systemic insecticide for the control of apple leafcurling midge and armoured scales in apples, pear leafcurling midges in pears, mealy bug in grapes, armoured scales in kiwifruit, Kelly's citrus thrips, Australian Citrus Whitefly and armoured scales in citrus, and thrips and mealy bug in blueberries.

CONTAINS: 100 g/L Spirotetramat in the form of a suspension concentrate

Net Contents: 1 / 5 / 10/ 20 litre

Batch No.:

Date of manufacture:

Registered pursuant to the ACVM Act 1997, No. P010020. See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval Code: HSR100545. See www.epa.govt.nz for approval controls.

May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child.
Harmful to aquatic life with long lasting effects.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
21/03/2025

PRECAUTIONS:
Obtain special instructions before use
Do not handle until all safety precautions have been read and understood.
Avoid breathing fumes, mists, vapours or spray.
Do not get in eyes, on skin, or on clothing.
Contaminated work clothing should not be allowed out of the workplace.
Avoid release to the environment.
Wear protective gloves, protective clothing and eye or face protection.
use personal protective equipment as required.
Wash contaminated clothing before reuse.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
21/03/2025

STORAGE:
Store locked up.
Protect from sunlight.
Store in a dry place. Store in a closed container.
Store in a well-ventilated place. Keep cool.

SHELF LIFE:
When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

DISPOSAL:
Triple rinse empty container and add rinsate to the spray tank.
Return empty container to Agrecovery collection point for recycling
Avoid contamination of any water supply with chemical or empty container.

SPILLAGE:
Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste container. Wash area with water and absorb with further inert material. Dispose of waste safely (such as a suitable landfill).

FIRST AID: If medical advice is needed, have product container or label at hand.
IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.
IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice.
CALL A POISON CENTRE OR DOCTOR IF EFFECTS PERSIST OR YOU FEEL UNWELL.
24-HOUR EMERGENCY TELEPHONE NUMBER – FOR ADVICE CONTACT THE NATIONAL POISONS CENTRE – 0800 POISON (0800 764 766) – OR A DOCTOR, IMMEDIATELY. HAVE PRODUCT CONTAINER OR LABEL AT HAND.
FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY, CALL 0800 CHEMCALL (24HRS) OR 0800 243 6225. IN A TRANSPORT EMERGENCY DIAL 111

Spirosec 100SC is a systemic (xylem and phloem mobile) insecticide belonging to the chemical class of ketoenols for the control of apple leaf curling midge and armoured scales in apples, pear leaf curling midges in pears, mealy bug in grapes, armoured scales in kiwifruit, Kelly's citrus thrips, Australian Citrus Whitefly and armoured scales in citrus, and thrips and mealy bug in blueberries.

It is an offence to use this product on animals.

DIRECTIONS FOR USE:
Mixing: Pour **Spirosec 100SC** directly into the spray tank with agitators running.

RESISTANCE MANAGEMENT: **Group 23 Insecticide**

Spirosec 100SC contains spirotetramat from the inhibitors of acetyl CoA carboxylase Lipid synthesis, growth regulation mode of action (Group 23). Resistance to this insecticide could develop in some situations from repeated use. To minimize this risk, use strictly in accordance with label instructions. Do not apply more than 3 applications per season.

RECOMMENDATION:
The rates suggested for use in apples is for high volume spraying of a dilute spray to the point of run-off. For concentrate spraying adjust dilution rate accordingly.

Crop	Pest	Application Rate	Comments
Apples	Apple leaf curling midge (ALCM)	48 ml/100L water	Apply product to coincide with peak ALCM egg laying periods. Apply from mid-November and mid-December to control the second generation and from mid-January to early February to control the third generation
	Armoured Scales	48 ml/100L water	Apply product to coincide with scale crawler release periods. Apply from mid-November to early December to control the first generation of San Jose Scale. Apply as necessary from late January to early Febreaury to control the second generation. In the South Island only, apply from mid-December to mid-January to control Oystershell Scale.
Blueberries	Thrips Mealy bug	150 ml/100L water	Apply at 14 – 21 day intervals a maximum of 3 times per season. Apply in sufficient water to obtain good coverage.
Citrus	Kelly's citrus thrips	96 ml/100L water	Apply post-flowering when pest threshholds are reached. Repeat if needed before calyx closure and pre-harvest. Apply a maximum of 3 applications per season, with at least 14 days between the first two applications. No more than 2 applications should be made within 90 days of harvest.
	Australian citrus whitefly, Armoured Scales	48 ml/100L water	
Grapes	Mealy bug	150 ml/100L water	Apply a maximum of 2 applications. Apply when new season's shoots are from 20 – 50cm long, just

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
21/03/2025

			prior to flowering or immediately post flowering. Do NOT apply more than 10 days post flowering. Do NOT apply in a water rate that will result in run-off.
Kiwifruit	Armoured Scales	48 ml/100L water	Apply two applications, the first at green tip and the second pre-flowering. Apply in sufficient water to obtain good coverage.
Pears	Pear leaf curling midge (PLCM)	48 ml/100L water	Apply during periods of peak PLCM egg laying, either immediately prior to flowering or immediately post-flowering. Apply in sufficient water to obtain good coverage.

COMPATIBILITY:
Spirosec 100SC is compatible with most fungicides.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD:
Apples- 14 days.
Blueberries – 7 days.
Citrus – 21 days
Grapes - Do not apply after 10 days post-flowering.
Kiwifruit – Apply only pre-flowering.
Pears – 35 days.

GRAZING:
Livestock that have been grazing or fed treated crops should be placed on clean feed for 2 months prior to slaughter.

Leaf plucking in vineyards - Sheep must not be sent for slaughter or milked for 6 months after being removed from the vineyard (and placed on clean feed)

CONDITIONS OF SALE:
Grochem®, a brand of ADAMA New Zealand Limited (Grochem) will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by the Statutes applicable to the sale and supply of these goods. To the extent allowed by such Statutes the liability of Grochem is limited to the replacement of the goods or (at the option of Grochem) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Grochem within thirty days of delivery.

REGISTERED TO: ADAMA New Zealand Ltd.
DISTRIBUTED BY:
: Grochem, a brand of ADAMA New Zealand Limited

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
21/03/2025

Level 1, 93 Bolt Road
Tahunanui, Nelson 7011
PO Box 1799, Nelson 7040
New Zealand
Telephone 0800 447 624
Email: grochem@grochem.com
www.grochem.com

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
21/03/2025