DANGER

Keep out of reach of children

FUNGICIDE

Mancostar 750

For the control of fungal diseases in a wide range of crops

ACTIVE INGREDIENT: Contains 750 g/kg mancozeb in the form of a water dispersible granule

Net Contents: 500 g – 25 kg

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/12/2025

READ THE LABEL COMPLETELY BEFORE MIXING AND APPLYING

HAZARD CLASSIFICATIONS: Skin irritation Category 2, Eye irritation Category 2, Respiratory sensitization Category 1, Skin sensitization Category 1, Specific target organ toxicity (repeated) Category 2, Hazardous to terrestrial vertebrates, Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1.

This product must not be used for any purpose or in any manner contrary to this label unless authorized under appropriate legislation.

TOXICITY

DANGER - May cause allergy or asthma symptoms or breathing difficulties if inhaled.
Causes skin and serious eye irritation. May cause an allergic skin reaction. May cause organ damage from repeated or prolonged exposure.

ECOTOXICITY

Very toxic to aquatic organisms and with long-lasting effects. Harmful to terrestrial vertebrates.
Do not apply directly into or onto water.
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

QUALIFIED HANDLER: This product must be under the personal control of a suitably qualified handler when mixed, loaded or applied.

RECORD KEEPING: Records of use must be kept if using more than 3 kg of **Mancostar 750** within 24 hours in a place where the product is likely to enter air or water and leave the application area.

PRECAUTIONS

Read label before use.

PERSONAL PROTECTION

Do not inhale dust or spray mist. Avoid contact with eyes and skin. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. In case of inadequate ventilation, wear respiratory protection with cartridge fitted with suitable filter for particulates. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Contaminated work clothing should not be allowed outside the workplace. Clean protective clothing and equipment daily after work.

STORAGE

Keep out of reach of children. Store in the original packaging in a cool, dry, secure and well- ventilated area away from foodstuffs. Storage must be in accordance with the current version of NZS 8409.

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
11/12/2025

Management of Agrichemicals. Stores containing quantities of more than 100 kg require emergency response plans, signage and secondary containment.

USE

Apply with well-maintained and calibrated spray equipment. Workers entering an area within 48-hours after the product has been applied must wear gloves. The public must not be permitted to access any area where this product has been sprayed within the first 48-hours after application.

SPILLAGE

Collect spill. Exclude all bystanders from the vicinity of the spill. Do not touch or walk through spilled material. Wear facemask or goggles, chemical resistant overalls (buttoned to neck and wrists), gloves and boots. If granules have spilt, moisten to prevent dust and sweep up. If liquid spill , absorb spill with inert material such as soil, sand, etc. Place in waste container. Wash area with water, and absorb with further inert material. Contain liquid spills with earth or non-combustible material, and prevent entry into waterways and drains. Dispose of safely at an approved landfill.

DISPOSAL

Avoid contamination of any water supply with product or empty container.

Product disposal: Dispose of this product only by using in accordance with this label, or to an approved landfill, or other approved facility. Do not burn.

Container disposal: Dispose of empty packaging to an approved landfill. Do not use packaging for any other purpose.

FIRST AID

For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label to hand.

If swallowed, do NOT induce vomiting. Rinse mouth. Get medical advice.

If skin or hair contact occurs, remove contaminated clothing and wash skin and hair with plenty of soap and water. Take off contaminated clothing and wash before reuse. If irritation or rash occurs, get medical attention.

If in eyes, wash out immediately with running water and keep rinsing for 15 minutes. Remove contact lenses if present and easy to do. Get medical attention if irritation persists.

If inhaled, immediately move to fresh air and keep at rest until recovered. If experiencing respiratory symptoms, call a POISON CENTRE or doctor.

SAFETY DATA SHEET

Additional information is provided in the safety data sheet available from the supplier, www.agstar.co.nz .

In a transport emergency call 111, Police or Fire Brigade

GENERAL INFORMATION

Mancostar 750 is a fungicide with protective properties for the control of fungal diseases in range of fruit, vegetable and ornamental crops.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/12/2025

DIRECTIONS FOR USE

RATES AND TIMING OF APPLICATION:

Mancostar 750 must not be applied at rates exceeding 3.2 kg mancozeb/ha and must not be applied to the same area more than six times per year. The following rates are for a full coverage spray to run-off. For more concentrated spraying with lower volume applications, adjust the dilution rate accordingly.

It is an offence to use this product on animals.

A) FRUIT CROPS			
CROP	**DISEASE**	**RATE**	**CRITICAL COMMENTS**
Apples Pears	Black spot *Glomerella* Ripe spot Sooty blotch	110 – 160 g / 100 litres water	Apply at green tip, and repeat at 10 - 14 day intervals. Use highest rate and shortest interval early in season or where severe disease exists. Maintain high rate for *Glomerella*. NOTE: Early season use under frosty conditions may cause russet.
Boysenberries Loganberries Raspberries	Downy mildew (Dryberry) Leaf spot Rust	210 g / 100 litres water	Apply at green tip, and repeat at 10 - 14 day intervals.
Citrus	Brown rot	210 g / 100 litres water	Apply in May. Where severe disease exists, apply at 4 - 6 week intervals from April to late winter.
	Alternaria		Apply from October to April at 14 - 28 day intervals.
	Verrucosis		Apply at petal fall
	Citrus rust mite		Apply at first signs of mite activity (Dec/Jan). Apply at 21 - 28 day intervals if necessary.
Currants (Red and Black) Gooseberries	Leaf spot	210 g / 100 litres water	Apply at green tip, pre-flowering, and repeat at 14 - 21 day intervals.
Grapes	*Anthracnose* (Black spot) Downy mildew	210 g / 100 litres water	Apply at bud burst, and repeat at 10 -14 day intervals as required.
Nectarines Peaches	Rust	210 g / 100 litres water	Apply in early November. Repeat application if disease is severe. NOTE: To avoid spray deposits, Do NOT spray within 6 weeks of harvest. Do NOT allow drift onto plums as foliage damage and defoliation may occur.
Strawberries	*Mycosphaerella* leaf spot (Frogs eye)	210 g / 100 litres water or 2 kg / ha	Apply from mid-June. Apply at 10 - 14 day intervals. In wet weather, repeat at 7 day intervals.
B) ORNAMENTALS			
Ornamentals	Blights, Spots	210 g / 100	Apply at 7 - 14 day intervals throughout the growing

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND
11/12/2025

MINISTRY FOR PRIMARY INDUSTRIES NEW ZEALAND 11/12/2025

	and Rust	litres water	season.
C) VEGETABLES			
Beans	*Anthracnose* Rust	160 - 210 g / 100 litres water	Apply at the first signs of disease and repeat at 7 - 10 day intervals.
Broccoli Brussels sprouts Cabbage Cauliflower	Downy mildew	160 - 210 g / 100 litres water	Apply at 7 - 10 day intervals. In wet weather, apply to seedbed at 3 - 4 day intervals.
Carrots	*Cercospora* leaf spot	160 - 210 g / 100 litres water	Apply at the first signs of disease and repeat at 7 - 10 day intervals.
Celery	*Septoria* leaf spot	160 - 210 g / 100 litres water	Seedbed: Apply at 3 - 5 day intervals. Field: Apply at 7 - 10 day intervals.
Cucurbits: Cucumber Melons Pumpkin Squash	*Anthracnose* *Alternaria* *Cercospora* leaf spot Downy mildew	160 - 210 g / 100 litres water	Apply as plants begin to run or at first signs of disease. Repeat at 5 - 7 day intervals.
Lettuce	Downy mildew Leaf spot Ring spot	160 - 210 g / 100 litres water	Apply at the first signs of disease and repeat at 7 - 10 day intervals.
Maize Sweetcorn	Northern leaf blight	2.1 kg / ha	Apply as an aerial treatment at the first signs of disease. Apply in sufficient water for coverage. Repeat as necessary.
Onions	*Botrytis* Downy mildew	160 - 210 g / 100 litres water	Apply at 7 - 10 day intervals.
Potatoes	Early blight (Target spot) Late blight	1.6 – 2.1 Kg / ha	Apply lower rate as crop emerges, higher rate as vines size. Repeat at 7 - 10 day intervals. In wet weather, repeat at 3 - 4 day intervals.
Tomatoes	*Anthracnose* Early blight Late blight *Septoria* leaf spot	160 - 210 g / 100 litres water	Apply at 7 - 10 day intervals. Use highest rate and shortest interval where severe disease conditions exist. Do NOT use on young tomato seedlings in glass houses or tunnel houses as injury may occur.

APPLICATION METHODS: Mancostar 750 WDG is non-corrosive and almost non-abrasive. **Mancostar 750** can be applied with conventional spraying equipment. Do not use strainers finer than 300 micrometres. Use a finer droplet size for more uniform coverage. Ensure the underside of leaves are treated. Wash equipment thoroughly after use.

MIXING: Mancostar 750 mixes readily with water. Add the required amount of **Mancostar 750** into the partly filled tank then add the remaining water while agitating. The spray mixture should be kept agitated before and during spraying, and before spraying after a stoppage.

COMPATIBILITY: Mancostar 750 is compatible with most commonly used insecticides and fungicides. Do NOT use with lime sulphur or Bordeaux mixture. For hard to wet vegetables such as onions, the addition of a non-ionic wetting agent such as Triton X-AG will improve coverage.

Restricted Entry Interval (REI)

Apples, pears	5 days
Stonefruit	6 days
Grapes	8 days
Onions	24 hours
Broccoli, brussels sprouts cabbage, cauliflower	10 days

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

Cucurbits, Tomatoes	1 day
Apples, Beans, Boysenberries, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Citrus, Currants (Red and Black), Gooseberries, Grapes, Lettuce, Loganberries, Maize, Nectarines, Onions, Peaches, Pears, Potatoes, Raspberries, Strawberries, Sweetcorn	14 days
Export Crops	Check with your exporter or industry crop adviser

Leaf plucking in vineyards: Sheep must not be sent for laughter or milking for 6 months after being removed from the vineyard (and placed on clean feed).

RESISTANCE MANAGEMENT

GROUP	M3	FUNGICIDE

Mancostar 75 WDG is a member of the dithio carbamate group of fungicides. Resistance to this and other M3 fungicides could develop from excessive use. To minimise this risk, use strictly in accordance with label instructions and resistance management strategies that exist for any diseases listed on the label. Consult http://resistance.nzpps.org/index.php for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

Registered pursuant to the ACVM Act 1997, No. P10323
See www.foodsafety.govt.nz for registration conditions.
Approved under the HSNO Act 1996, HSR100972
See www.epa.govt.nz for approval controls

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/12/2025

.

Registrant: AgStar New Zealand Pty Ltd
4 Putakitaki Street, Lincoln, 7608

www.agstar.co.nz

Distributed by: Farmlands Co-operative Society Limited
535 Wairakei Road, Burnside, Christchurch 8053
TEL: 0800 200 600

Batch No.:
Date of Manufacture :
Expiry date:

CONDITIONS OF SALE

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING.
IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer or registrants control. The manufacturer and registrant decline all liability for damage resulting from improper storage or use of the product.

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
11/12/2025