310x550 (10 KG)

Pantone 290 C

Pantone 638 C

DANGER • ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Blue AG

Captrex

FUNGICIDE

GROUP **M4** FUNGICIDE

ACTIVE INGREDIENT
900 g/kg Captan
in the form of a water dispersible granule.

A fungicide for the control of a wide range of fungus diseases on fruit, vegetables, ornamentals, turf, greenhouse crops and for certain soil borne diseases.

AGRECOVERY

Read the label completely before use
EMERGENCY RESPONSE (ALL HOURS)
0800 CHEMCALL OR 0800 243 622 OR
PGG WRIGHTSON LTD 0800 10 22 76

10 KG
NET CONTENTS

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/11/2025

310x550 (10 KG)

Pantone 485 C | Pantone 425C | BLACK C:60 M:40 Y:40 K:100 | Pantone 638 C

READ THE LABEL COMPLETELY BEFORE OPENING AND USING

HAZARD CLASSIFICATION
Eye damage Category 1,
Skin sensitisation Category 1,
Carcinogenicity Category 2,
Hazardous to the aquatic environment (acute) Category 1.

TOXICITY
Causes serious eye damage. May cause an allergic skin reaction. Suspected of causing cancer.

ECOTOXICITY
Very toxic to aquatic life. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water.

QUALIFIED HANDLER: This product must be under the control of a suitably qualified handler with mixed, loaded and applied.

RECORD KEEPING: A record of use must be kept according to HSNO Regulations and N.Z. Standard for the Management of Agrichemicals (NZS 8409). If 3 kg or more is applied within 24 hours, in places where the product is likely to enter air or water and leave the application area, a written record is to be kept for each application.

PERSONAL PROTECTION: Avoid contact with skin, eyes and inhalation of spray mist. When handling, wear eye protection, waterproof gloves, boots and overalls. Remove protective clothing and wash hands and face thoroughly before rest breaks, meals and after work. Do not eat, drink or smoke while using. Wash protective clothing and equipment after use.

STORAGE: Store in the original container in a locked, dry, cool area away from foodstuffs, fertilisers and seeds. Avoid storage at temperatures below 5°C and above 35°C. Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals. Stores containing more than 100 kg require signage, and an emergency response plan.

EQUIPMENT: Apply using well maintained and accurately calibrated equipment. Fine droplets are more effective than coarse droplets. Ensure that undersides of leaves are treated.

SPILLAGE: Collect spill. Wear appropriate protective clothing and prevent material from entering waterways. Contain liquid spills and absorb with inert material and place in waste container. Sweep up solids without creating dust. Dispose of waste safely in a suitable landfill.

DISPOSAL INFORMATION
Avoid contamination of any water supply with chemical or empty container.
PRODUCT DISPOSAL: Dispose of this product only by using according to the label directions or if this is not possible dispose of safely at an approved landfill.
CONTAINER DISPOSAL: Dispose of through a recycling facility if available, or otherwise at an approved landfill.
Do not use container for any other purpose.

FIRST AID
If exposed or concerned, get medical advice. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If swallowed do NOT induce vomiting. Rinse mouth. If inhaled move the victim to fresh air immediately and kept at rest in a position comfortable for breathing. If skin contact occurs, remove contaminated clothing and wash with soap and running water. If irritation or rash develops, get medical advice. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Get immediate medical attention or advice.

MIXING: Captrex mixes readily with water. While agitating, slowly add the required quantity of **Captrex** through the filler screen to a half-filled spray tank and continue filling. Maintain agitation during filling and spraying and agitate thoroughly after a stoppage before recommencing spraying.

When tank mixing **Captrex** with another product, ensure that products are added in the following order with tank agitation running:
1. First add any WP or WG fungicides or insecticides to the half-filled tank (add WG & WP formulationsfollowed by SC and EW formulations).
2. If adding foliar fertilisers such as calcium chloride or calcium nitrate or a surfactant, these must always beadded last. Check for any precautions on the labels of products being mixed.
3. Do not leave mixed spray standing overnight.

COMPATIBILITY: Captrex is compatible with most commonly used pesticides. Do not mix **Captrex** with Bordeaux mixture, lime sulphur, oils or dicofol. Use with emulsifiable concentrate (EC) formulations is not recommended. If used with sulphur on Kidds Orange, Delicious and some sulphur shy varieties, some leaf spotting or leaf drop may occur. For further information see: www.resistance.nzpps.org.

RESISTANCE MANAGEMENT: GROUP M4 FUNGICIDE
Fungal diseases have not yet developed resistance to captan which belongs to the phthalimide group of fungicides. To minimise the risk of resistance and to prolong the usefulness of specific fungicides, Captrex should be used strictly in accordance with label instructions and resistance management strategies that exist for any disease listed on the label. Maintain good agricultural practice.

It is an offence to use this product on animals.

DIRECTIONS FOR USE
The rates suggested are for high volume spraying to run-off. For concentrate spraying, adjust dilution rates accordingly. Use the higher rates and/or shorter spray intervals when disease pressure is high.

FRUIT CROPS

CROPS	DISEASE	RATE	NOTES
Apples, Pears	Black spot, Glomerella, Ripe spot and Sooty blotch	65 - 110 g/ 100 litres water	Blackspot: Apply the high rate every 7 days from green tip through to petal fall and the low rate at 10-14 day intervals from petal fall through to harvest. Where glomerella, ripe spot or sooty blotch are a problem use the high rate from January to harvest. Products containing captan may increase the risk of russet on some varieties. Where this is of concern use alternative materials between full bloom and second cover. **Do not apply to Anjou Pears.**
Blueberries	Stem blight, Leaf spot and Botrytis	110 g/100 litres water	Spray before flowering when 10% of flowers are open, at petal fall and every 10-14 days until harvest.
Boysenberries, Loganberries and other Brambles	Cane spot	110 g/100 litres water	Apply at petal fall and repeat every 10-14 days until harvest.
Citrus	Brown rot	110-215 g/ 100 litres water or 110 g/100 litres water	As a skirt spray before winter rains. Thoroughly wet trees to a height of 1 metre and repeat 10 weeks later. Spray entire tree in autumn and repeat 4-6 weeks later. Do not use oil products within 14 days either side of a **Captrex** spray.
Red and Black Currants	Leaf spot and Botrytis	110 g/100 litres water	Apply every 10-14 days from green tip until harvest.
Gooseberries	Leaf spot and Botrytis	110 g/100 litres water	Apply just before flowering and repeat at petal fall, preharvest and postharvest. For botrytis flower rot, applications at 5-7 day intervals over flowering may be necessary depending on the weather.
Grapes	Botrytis, Dead arm, and Anthracnose	110 g/100 litres water	Apply every 14 days, commencing when shoots are 5-8 cm long.
	Downy Mildew	80 g/100 litres water	Apply at the start of flowering at 80 g/100 litres. (then use an alternative product or **Captrex** at 110g/100 litres to control botrytis and downy mildew over the flowering period). After flowering continue to apply 80 g/100 litres at 14 day intervals.
Raspberries	Cane spot, Spur blight and Botrytis	110 g/100 litres water	Optimum coverage is essential. **Main crop:** Apply every 10-14 days from green tip until harvest. **Autumn crop:** Apply every 10-14 days from November until harvest.
Stonefruit	Brown rot, Scab and Shothole	110 g/100 litres water	Apply at 7 day intervals from bud movement until bloom, then at 3-4 day intervals until petal fall. After petal fall apply at 14 day intervals until 3 weeks before harvest, then apply at 7 day intervals until harvest or if weather is wet.
Strawberries	Leaf spot and Botrytis	110 g/100 litres water or 2.7 kg/ha	Apply as necessary for leaf spot. For botrytis, apply at 7 day intervals from the beginning of flowering and between pickings if warm wet conditions prevail.

VEGETABLE CROPS

CROPS	DISEASE	RATE	NOTES
Peas, Beans	Root rot and Damping off	6.2 - 8 kg/ha	Apply as a broadcast spray pre-planting and work into upper 10cm of soil.
Carrots	Ring spot and Cercospora blight	2.7 kg/ha	As necessary.
Cucurbits	Anthracnose and Downy mildew	1.3-2.7 kg/ha	As necessary ensuring optimum spray coverage.
Lettuce	Leaf spot	4.0 kg/ha	Every 10-14 days from rosette stage.
Potato Seed	Seed rot, Damping off	110 g/100 litres water	Pre-planting dip within 6 hours of cutting.
Pea Seed	Seed rot, Damping off	90 g/500 ml water / 100 kg of seed	Keep stock solution well agitated and do not allow to stand overnight. **Captrex** does not contain a sticker. A suitable sticker should be added to the stock solution.Contact your local representative for further advice. Ensure treated seed is clearly labelled.

ORNAMENTALS - AMENITY TURF

CROPS	DISEASE	RATE	NOTES
Lawns	Brown patch, Leaf spot and Damping off	5.3 kg/ha or 110 g/ 100 litres water	As a soil drench using sufficient water for good soil penetration.
Ornamentals	Rose black spot. Carnation leaf spot. Chrysanthemum spot.	110 g/100 litres water	As necessary.
Soil and Glasshouse Drench	Damping off	110 g/100 litres per 100 square metres	Work into top 10cm of soil before planting.

KNAPSACK SPRAYER DOSAGE RATES:
1. **Hydraulic types:** 12g per 10 litres of water
2. **Motorised mistblower types:** Use 35g per 10 litres of water. Use middle settings on the restrictor and operate at sufficient throttle to give small droplets. Move all foliage in the airstream.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD
Boysenberries: 14 days
Blueberries: 1 day
Carrots: 14 days
Citrus: 28 days
Cucurbits: 3 days
Currants: 1 day
Gooseberries: 1 day
Grapes: 21 days
Lettuce: 7 days
Loganberries: 14 days
Other brambles: 14 days
Pipfruit: 3 days
Raspberries: 1 day
Strawberries: 1 day
Stonefruit: 7 days
Treated seed:
Do not use for human consumption, stock or poultry food

For export crops: Consult your exporter for advice about any restrictions.

Leaf plucking in vineyards: Sheep must not be sent for slaughter or milked for 2 months after being removed from the vineyard (and placed on clean feed).

Registered pursuant to the ACVM Act 1997, No. P10336
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, approval code HSR000503
See www.epa.govt.nz for approval controls.

CONDITIONS OF SUPPLY- Read before buying and using.
If these conditions are unacceptable, return at once to the retailer unopened within seven days and the purchase price will be refunded in full.
This product conforms to the chemical description on the label. However to the full extent permitted by law, all warranties of any kind, expressed or implied, with respect to the fitness for any purpose (including crop yield/performance/injury, product effectiveness or any other consequence), condition, merchantable quality or any other matter, are expressly excluded. Buyers and/or users accept that PGG Wrightson Ltd and its distributors, are not liable in any way (including in negligence, tort and equity) to you for any loss, costs, loss of income, damage or injury whether direct, indirect or consequential, howsoever arising in connection with this product including without limitation storage, application, handling or use. PGG Wrightson Ltd's liability if any is limited solely to an amount not exceeding twice the purchase price of the product. To read PGG Wrightson Ltd's terms of trade that apply to PGG Wrightson transactions, visit www.pggwrightson.co.nz.

REGISTERED TO: PGG Wrightson Ltd
1 Robin Mann Place
Christchurch Airport Christchurch 8053
PO Box 292 Christchurch 8140

Batch No -

D.O.M. -

9 41525 00524 6

MINISTRY FOR PRIMARY INDUSTRIES
NEW ZEALAND
10/11/2025