

Random cardinalities of coalescence classes in a six-state Markov grand coupling

Paul

Abstract

We construct an independent random map grand coupling of a strictly positive six-state transition matrix for which the eventual coalescence classes have cardinalities $(2, 4)$ or $(3, 3)$, each with probability $1/2$. The number of classes is always two, giving a negative answer to Open Problem 3.11 of Grimmett and Holmes, *Coalescence in Markov Chains* (2026). The transition matrix has identical rows and satisfies $P^2 = P$. A pathwise proof uses two proper colorings of the disjoint union of two three-vertex paths. We extend the construction to bipartite graphs, characterize when that construction has a random cardinality profile, and derive a second counterexample by splitting states in Grimmett and Holmes' four-state example. A self-contained proof of the classical equal stationary mass theorem explains why cardinalities can vary while stationary masses remain equal, and shows that six states are minimal among finite irreducible independent random map couplings. Finally, we distinguish the explicit construction from the established semigroup theory underlying it.

Keywords: Markov chains; grand couplings; coalescence; random mappings; transformation semigroups.

1. The question and the result

Let S be a nonempty finite set, and let μ be a probability measure on the set S^S of functions from S to itself. Let $F_1, F_2, \dots$ be independent with common law μ . We use ordinary function composition, with the rightmost function applied first, and put

$$H_0 = \text{id}_S, \quad H_t = F_t \circ \dots \circ F_1, \quad X_t^i = H_t(i). \quad (1)$$

The same function F_t updates every coordinate at time t . For each fixed i , the process X^i is a Markov chain with transition probabilities

$$P_{xy} = \mu\{f : f(x) = y\}, \quad x, y \in S. \quad (2)$$

Indeed, F_{t+1} is independent of the maps already used, so its conditional action on the current state x has the distribution in (2). We assume throughout, except where explicitly stated otherwise, that P is irreducible.

If two coordinates meet, they subsequently coincide, since future functions act identically on equal inputs. Thus

$$i \sim j \iff H_t(i) = H_t(j) \text{ for some } t \geq 0$$

is an equivalence relation. Write $\mathcal{C}_\infty$ for its random partition of S . For a partition $\mathcal{C}$, define $\text{prof}(\mathcal{C})$ to be the tuple of its block cardinalities arranged in nondecreasing order, with

repetitions retained. This tuple records the multiset of cardinalities without choosing labels for the blocks.

Grimmett and Holmes [3, Open Problem 3.11] ask whether these cardinalities must be deterministic. The relevant distinction is between the number of eventual classes and their sizes. The former is almost surely constant; an elementary minimum rank proof is recalled in Section 4. The following construction shows that the latter need not be.

Theorem 1.1. *There is a probability measure μ supported on sixteen maps of $S = \{1, 2, 3, 4, 5, 6\}$, each of probability $1/16$, consistent with*

$$P = \mathbf{1}\pi, \quad \pi = \frac{1}{8}(1, 2, 1, 1, 2, 1), \quad (3)$$

where $\mathbf{1}$ is the column vector of six ones, such that

$$\mathbb{P}\{\text{prof}(\mathcal{C}_\infty) = (2, 4)\} = \mathbb{P}\{\text{prof}(\mathcal{C}_\infty) = (3, 3)\} = \frac{1}{2}. \quad (4)$$

For every sequence of maps from this support, the eventual partition is already the partition into fibers of F_1 . In particular, there are exactly two classes on every such sequence. The matrix P is strictly positive, irreducible and aperiodic, and $P^2 = P$.

Both P and μ are fixed in this statement. The randomness in (4) occurs across realizations of one grand coupling, rather than through a choice between different coupling laws.

Earlier examples show that the eventual partition can be random even when its cardinality profile is fixed. In the four-state coupling of Grimmett and Holmes [3, Example 3.10], the two possible partitions both have profile $(2, 2)$. The two minimum rank partitions in the six-state example of Budzban and Feinsilver [1, Sections 3–4] likewise share the profile $(2, 2, 2)$. The construction above gives different cardinality profiles under a single coupling law. Corollary 6.2 shows more generally how a finite state-splitting lift converts a nonconstant eventual partition into a nonconstant cardinality profile.

The main construction and its proof appear in Section 2. Section 3 gives a graph formulation and an infinite family. Sections 4 and 5 explain the stationary mass and semigroup structure; the former also proves that six states are minimal. Section 6 derives another counterexample directly from a previously published coupling.

2. The six-state construction

Consider the simple undirected graph G with vertex set S and edge set

$$E = \{\{1, 2\}, \{2, 3\}, \{4, 5\}, \{5, 6\}\}. \quad (5)$$

It is the disjoint union of two paths on three vertices. Introduce two subsets

$$A_0 = \{1, 3, 4, 6\}, \quad A_1 = \{1, 3, 5\}. \quad (6)$$

For either $b \in \{0, 1\}$, every edge has one endpoint in A_b and one in $S \setminus A_b$. In graph terminology, $(A_b, S \setminus A_b)$ is a proper two-coloring.

Let $\vec{E}$ be the eight oriented edges obtained by giving each edge in (5) both orientations. Choose B uniformly from $\{0, 1\}$ and, independently, choose (U, V) uniformly from $\vec{E}$. Define

$$f_{b,u,v}(x) = \begin{cases} u, & x \in A_b, \\ v, & x \notin A_b. \end{cases} \quad (7)$$

The law μ is the law of $f_{B,U,V}$. The choices give sixteen distinct maps: the two values of b have different fiber partitions, and, for fixed b , the values on the two parts recover the



Figure 1: The two colorings used in the construction. Shaded vertices form A_b . Reversing the colors on the second connected component changes the cardinalities but preserves the condition that adjacent vertices have different colors..

ordered pair (u, v) . Equivalently, in the notation $(f(1), \dots, f(6))$, they have the forms

$$f_{0,u,v} = (u, v, u, u, v, u), \quad f_{1,u,v} = (u, v, u, v, u, v). \quad (8)$$

cf. the Appendix for all sixteen maps.

2.1 Verification of the single-coordinate transition matrix

The degrees of the six vertices of G are $(1, 2, 1, 1, 2, 1)$. Since every oriented edge is equally likely,

$$\mathbb{P}(U = j) = \mathbb{P}(V = j) = \frac{\deg_G(j)}{8} = \pi_j. \quad (9)$$

For fixed x and fixed b , the output $f_{b,U,V}(x)$ is either U or V , according to the membership of x in A_b . Both endpoint distributions are π , so

$$\mathbb{P}(F_t(x) = j \mid B_t = b) = \pi_j.$$

Averaging over b proves (3). In particular, every transition has positive probability and every state has a positive self-transition. Furthermore,

$$\pi P = (\pi \mathbf{1})\pi = \pi, \quad P^2 = \mathbf{1}(\pi \mathbf{1})\pi = \mathbf{1}\pi = P.$$

Each individual coordinate has distribution π after one step, and its states at times $1, 2, \dots$ are independent with common law π . This assertion concerns one coordinate; the coordinates updated by the same random maps are dependent.

2.2 The pathwise argument

Lemma 2.1. *Every map in (7) sends the two endpoints of every edge of G to the two distinct endpoints of an edge of G .*

Proof. If $\{x, y\} \in E$, then exactly one of x, y belongs to A_b . Consequently,

$$\{f_{b,u,v}(x), f_{b,u,v}(y)\} = \{u, v\} \in E.$$

Since the graph has no loops, then $u \neq v$. □

Proof of Theorem 1.1. The matrix assertions were established above. Fix any infinite sequence of allowed maps. The first map has image $\{u_1, v_1\}$, an edge of G , and fibers A_{b_1} and $S \setminus A_{b_1}$. All starting states in either fiber therefore meet at time one.

At time one the two remaining locations are the endpoints of an edge. Lemma 2.1 implies that, after the next map, they are still the endpoints of an edge and remain distinct.

Induction gives the same statement at every subsequent time. Hence states in different fibers of the first map never meet, while states in the same fiber have already met. This proves

$$\mathcal{C}_\infty = \{A_{B_1}, S \setminus A_{B_1}\}. \quad (10)$$

The cardinalities of A_0 and its complement are 4 and 2; those of A_1 and its complement are 3 and 3. Since B_1 is fair, (4) follows. $\square$

Remark 2.2 (The profile law can vary while P stays fixed). Replace the fair choice of B by $\mathbb{P}(B = 0) = \theta$, where $0 \leq \theta \leq 1$, and keep the independent uniform oriented edge choice. Equation (9) still gives the same matrix P . The probabilities of the profiles (2, 4) and (3, 3) are then θ and $1 - \theta$. In contrast, choosing the constant map $x \mapsto J$ with $J \sim \pi$ also gives the same P , but all six starting states meet at time one. Thus even this particularly simple transition matrix supports markedly different joint behavior.

3. A general construction from bipartite graphs

Let $G = (S, E)$ be a finite simple bipartite graph with at least one edge and no isolated vertices. Write $m = |E|$. A subset $A \subset S$ will be called a color class if every edge has exactly one endpoint in A . Let ρ be any probability law on such subsets. Choose $A \sim \rho$ independently of a uniform oriented edge (U, V) and define $f_{A,U,V}$ by (7), with A in place of A_b .

Proposition 3.1. *The resulting independent random map coupling has*

$$P_{ij} = \pi_j = \frac{\deg_G(j)}{2m} > 0. \quad (11)$$

For every allowed sequence of maps, its eventual partition is $\{A^{(1)}, S \setminus A^{(1)}\}$, where $A^{(1)}$ is the first chosen color class. Each of these two blocks has stationary mass $1/2$.

Proof. Both coordinates of a uniform oriented edge have distribution $\deg_G(j)/(2m)$, proving (11). Every allowed map sends edges to edges, so the proof of (10) applies without change. Finally, each edge has precisely one endpoint in A , whence

$$\sum_{j \in A} \deg_G(j) = m, \quad \pi(A) = \frac{m}{2m} = \frac{1}{2}.$$

The complement has the remaining mass $1/2$. $\square$

The auxiliary graph may be disconnected even though P is strictly positive since the maps may send a color class to an endpoint in any connected component.

3.1 Exactly when this graph construction has varying sizes

Suppose G has connected components $G_1, \dots, G_k$. Let their bipartition sizes be (a_i, b_i) , relabeled so that $a_i \geq b_i$, and put

$$n = |S|, \quad \delta_i = a_i - b_i \geq 0.$$

Since each component is connected and has an edge, its proper two-coloring is unique up to exchange of colors. Thus the possible cardinalities of a color class satisfy

$$|A| = \frac{1}{2} \left(n + \sum_{i=1}^k \varepsilon_i \delta_i \right), \quad \varepsilon_i \in \{-1, 1\}. \quad (12)$$

Its unordered profile depends only on the absolute value of the signed sum:

$$\text{prof}\{A, S \setminus A\} = \left(\frac{n - |\sum_i \varepsilon_i \delta_i|}{2}, \frac{n + |\sum_i \varepsilon_i \delta_i|}{2} \right).$$

Proposition 3.2. *Assume ρ gives positive probability to every proper two-coloring. The profile in Proposition 3.1 is nonconstant if and only if at least two connected components have unequal bipartition sizes. Consequently, six vertices are necessary and sufficient for a counterexample within this graph construction.*

Proof. If at most one δ_i is positive, then $|\sum_i \varepsilon_i \delta_i|$ is independent of the signs. If at least two are positive, let $D = \sum_i \delta_i$ and choose j with $0 < \delta_j < D$. All positive signs give absolute sum D . Reversing just the j th sign gives $|D - 2\delta_j| < D$. Both choices have positive probability and yield different profiles.

A connected bipartite graph with an edge and unequal bipartition sizes has at least three vertices. Two such components require at least six vertices. The graph in (5) attains that bound. Proposition 4.4 below proves the same lower bound for all finite irreducible independent random map couplings, by a different argument. $\square$

For an explicit infinite family, let G be the disjoint union of two stars $K_{1,d}$, with $d \geq 2$. Choose with equal probabilities between putting both centers in one color class and putting one center together with the leaves of the other star. Proposition 3.1 gives profiles

$$(2, 2d) \quad \text{and} \quad (d+1, d+1), \tag{13}$$

each with probability $1/2$. The two centers each have stationary mass $1/4$; each of the $2d$ leaves has mass $1/(4d)$. In either profile, each block again has mass $1/2$.

4. Why the stationary masses are equal

The equality of masses in Proposition 3.1 reflects a classical fact about finite random mappings. Equal-weight partitions occur in Friedman's road coloring analysis [2, Theorem 2.1]. The minimum rank fiber statement appears in Yano [5, Theorem 4.1] and in Budzban and Feinsilver [1, Theorem 4.8]. We give a self-contained proof, including the passage from minimum rank maps to eventual classes. The theorem numbers in these two references refer to the linked preprints. Only irreducibility is needed for the version below.

Let $\mathcal{A} = \text{supp } \mu$ and let $\mathcal{T}$ be the finite semigroup of nonempty compositions of maps from $\mathcal{A}$. For $g \in \mathcal{T}$, define its rank to be $|g(S)|$, and define

$$r = \min_{g \in \mathcal{T}} |g(S)|. \tag{14}$$

A *fiber* of h is a nonempty set $h^{-1}\{z\}$, with $z \in h(S)$. A subset is a *transversal* of a partition if it contains exactly one point from every block.

Lemma 4.1 (Transversals of minimum rank fibers). *If $g, h \in \mathcal{T}$ both have rank r , then $g(S)$ is a transversal of the fiber partition of h . Moreover, every $f \in \mathcal{T}$ acts injectively on $g(S)$.*

Proof. The composition $h \circ g$ has rank at least r by minimality and at most r because $g(S)$ has r elements. Therefore h is injective on $g(S)$. Its r distinct values must be all of $h(S)$, proving the transversal assertion. The same argument applied to $f \circ g$ proves the final assertion. $\square$

Theorem 4.2 (Equal stationary masses). *Let P in (2) be irreducible, with stationary distribution π , and let r be as in (14). Every fiber C of every rank- r map in $\mathcal{T}$ satisfies*

$$\pi(C) = \frac{1}{r}. \tag{15}$$

Almost surely, $\mathcal{C}_\infty$ is the fiber partition of a rank- r map in $\mathcal{T}$. Consequently it has r blocks, each of stationary mass $1/r$.

Proof. Fix a rank- r map h and one of its fibers C . Also fix any rank- r image $R = g(S)$. For each $t \geq 0$, let $R_t = H_t(R)$ and assign mass $1/r$ to each of its points. By Lemma 4.1, H_t is injective on R and R_t is another rank- r image. Every R_t therefore meets C in exactly one point. Pathwise,

$$\frac{1}{r} \sum_{x \in R} \mathbf{1}_{\{H_t(x) \in C\}} = \frac{1}{r}. \quad (16)$$

Let ν_R be the uniform probability measure on R . Taking expectations in (16) and using the marginal Markov chains gives

$$(\nu_R P^t)(C) = \frac{1}{r} \quad (t \geq 0). \quad (17)$$

For completeness, the Cesàro averages $\bar{\nu}_N = N^{-1} \sum_{t=0}^{N-1} \nu_R P^t$ converge to π . Indeed, the probability simplex is compact, and

$$\bar{\nu}_N P - \bar{\nu}_N = \frac{\nu_R P^N - \nu_R}{N} \longrightarrow 0.$$

Every subsequential limit is thus stationary, and irreducibility gives the unique stationary law π . Applying this to (17) proves (15). This argument also covers periodic P .

It remains to identify the eventual partition. Choose a word $w = f_\ell \circ \dots \circ f_1$ of rank r , with each $f_j \in \mathcal{A}$. In each disjoint block of ℓ independent draws, the ordered word $(f_1, \dots, f_\ell)$ has probability

$$p = \prod_{j=1}^{\ell} \mu(f_j) > 0.$$

The probability that it fails to occur in the first N such blocks is $(1-p)^N$, which tends to zero. At the end of its first occurrence, the full composition is $w \circ H$ for the preceding composition H . Its image has at most r points and by (14), exactly r points.

Thereafter the rank cannot decrease below r . Fiber partitions can only merge under further left composition; with their number fixed at r , no merger is possible. Thus the partition at this finite time is exactly $\mathcal{C}_\infty$. The already proved equality applies to all its blocks. $\square$

Corollary 4.3. *If π is uniform on n states, then every grand coupling has deterministic eventual profile*

$$\text{prof}(\mathcal{C}_\infty) = (n/r, \dots, n/r)$$

with r entries, where r is its minimum rank. In particular r divides n . A random eventual profile for an irreducible chain requires a nonuniform stationary distribution.

Proof. For every eventual block C , Theorem 4.2 gives $|C|/n = \pi(C) = 1/r$. $\square$

Proposition 4.4 (Six states are minimal). *An independent random map grand coupling of an irreducible chain on at most five states has an almost surely deterministic eventual cardinality profile. Consequently, Theorem 1.1 has the smallest possible state space size under these assumptions.*

Proof. Fix a coupling, and let r be its minimum rank. By Theorem 4.2, every eventual block has mass $1/r$. Since irreducibility gives $\pi_x > 0$ for every state, a state x is a singleton block if and only if $\pi_x = 1/r$. The set of singleton blocks is therefore deterministic.

Let k be the number of these singleton states and let $q = r - k$ be the number of remaining blocks. Both numbers are fixed. Each remaining block has size at least two. If $q = 0$, all blocks are singletons. If $q = 1$, the sole remaining block has the fixed size $n - k$. If $n \leq 5$, the only other possibility is $q = 2$, and its two sizes sum to $n - k \leq 5$. Two integers each at least two with fixed sum at most five have the unique unordered pair $(2, 2)$ or $(2, 3)$. Thus the profile cannot vary. The six-state construction supplies the matching upper bound. $\square$

In the six-state example the masses can also be checked immediately:

$$\begin{aligned}\pi(A_0) = \pi(\{1, 3, 4, 6\}) &= 4 \cdot \frac{1}{8} = \frac{1}{2}, & \pi(S \setminus A_0) = \pi(\{2, 5\}) &= 2 \cdot \frac{1}{4} = \frac{1}{2}, \\ \pi(A_1) = \pi(\{1, 3, 5\}) &= \frac{1}{8} + \frac{1}{8} + \frac{1}{4} = \frac{1}{2}, & \pi(S \setminus A_1) = \pi(\{2, 4, 6\}) &= \frac{1}{4} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.\end{aligned}$$

Thus the random cardinalities are compatible with a rigid deterministic mass constraint. Equal stationary mass does not imply equal counting measure when the states have different stationary weights. The number of classes is two in either case, and each class has stationary mass $1/2$. Their cardinalities nevertheless have profiles $(2, 4)$ and $(3, 3)$: four states of mass $1/8$ balance two states of mass $1/4$, while a different grouping gives three states on each side, thus illustrating how unequal stationary weights permit variation in cardinalities. Corollary 4.3 shows that such variation is impossible when the stationary distribution is uniform.

5. Transformation semigroups and the preserved graph

The sixteen maps provide a small concrete instance of the transformation semigroup viewpoint developed in, for example, Budzban and Feinsilver [1]. The following facts can be verified directly:

Proposition 5.1. *The sixteen maps in (8) are closed under composition. Every such map has rank two. For any two of them, f and g ,*

$$\text{fib}(g \circ f) = \text{fib}(f), \quad (g \circ f)(S) = g(S). \quad (18)$$

Here $\text{fib}(f)$ denotes the partition into nonempty fibers of f .

Proof. Write $f = f_{b,u,v}$. Since $\{u, v\}$ is an edge and g preserves edges, the pair $(g(u), g(v))$ is an oriented edge. Hence

$$g \circ f_{b,u,v} = f_{b,g(u),g(v)}$$

is another map in the list. Because $g(u) \neq g(v)$, the two fibers of f remain distinct. Finally, g takes its two values on the endpoints of every edge, so $g(\{u, v\}) = g(S)$. $\square$

The semigroup therefore has exactly two fiber partitions, four image sets, and two bijections between the blocks of any chosen partition and the vertices of any chosen image, hence $2 \cdot 4 \cdot 2 = 16$ elements. Equation (18) explains why the rightmost, earliest map fixes the eventual classes while later maps change their locations.

There is also an exact graph interpretation. Given a transformation semigroup $\mathcal{T}$ on S , its *kernel graph* has an edge $\{x, y\}$ precisely when no $g \in \mathcal{T}$ identifies x and y ; see Schaefer [4]. In the present example the kernel graph is exactly G , not just a graph preserved by the maps. To see this, record the membership of each state in the two color classes:

x	1	2	3	4	5	6
$(\mathbf{1}_{A_0}(x), \mathbf{1}_{A_1}(x))$	(1, 1)	(0, 0)	(1, 1)	(1, 0)	(0, 1)	(1, 0).

Two states are separated by both partitions precisely when their bit pairs are complementary, which pairs are exactly the four edges in (5). Every other pair lies together in at least one of the two partitions and is identified by the corresponding maps.

Moreover, the sixteen maps are exactly the rank-two endomorphisms of G . If an edge-preserving map has rank two, its image must be an edge, and its two fibers give a proper two-coloring. Each of the two path components admits just two choices of color, so G has exactly two unordered proper bipartitions. Choosing the bipartition, an image edge, and the assignment of the two blocks to its endpoints gives exactly the sixteen maps above.

6. A direct connection to a previous coupling by state splitting

The preceding construction is not the only way to obtain varying cardinalities. A general state-splitting operation turns randomness of membership in a partition into randomness of ordinary cardinalities. We give its details to make the relationship to the earlier example explicit.

Let S be a finite state space with a random map coupling F_t . Replace each $x \in S$ by a nonempty finite set T_x , with the sets T_x pairwise disjoint, and let

$$T = \bigsqcup_{x \in S} T_x, \quad q : T \longrightarrow S, \quad q(y) = x \text{ for } y \in T_x.$$

Choose a strictly positive probability distribution η_x on each T_x . Independently at each time, choose a section $s_t : S \rightarrow T$ by choosing $s_t(x)$ according to η_x , independently over x and independently of F_t . The identity $q \circ s_t = \text{id}_S$ holds, and each section is injective. Define

$$\tilde{F}_t = s_t \circ F_t \circ q. \tag{19}$$

Proposition 6.1 (State splitting). *The lifted maps are independent and identically distributed. Their marginal transition matrix and stationary distribution are*

$$\tilde{P}_{yz} = P_{q(y), q(z)} \eta_{q(z)}(z), \quad \tilde{\pi}_z = \pi_{q(z)} \eta_{q(z)}(z). \tag{20}$$

If P is irreducible, so is $\tilde{P}$; if P is also aperiodic, so is $\tilde{P}$. The eventual lifted partition is

$$\tilde{\mathcal{C}}_\infty = \{q^{-1}(C) : C \in \mathcal{C}_\infty\}. \tag{21}$$

In particular, with $m_x = |T_x|$, its block cardinalities are the sums $\sum_{x \in C} m_x$ over the original eventual blocks.

Proof. The pairs (F_t, s_t) are independent and identically distributed. Conditioning on $F_t(q(y))$ proves the first formula in (20). Since $q \circ s_j$ is the identity, all intermediate sections cancel under composition:

$$\tilde{F}_t \circ \cdots \circ \tilde{F}_1 = s_t \circ (F_t \circ \cdots \circ F_1) \circ q \quad (t \geq 1). \tag{22}$$

Injectivity of s_t implies that the images of $y, z \in T$ on the left are equal exactly when $H_t(q(y)) = H_t(q(z))$. This proves (21); points in the same initial fiber of q meet at time one.

Summing the first formula in (20) over an intermediate fiber yields, by induction,

$$(\tilde{P}^t)_{yz} = (P^t)_{q(y), q(z)} \eta_{q(z)}(z), \quad t \geq 1. \tag{23}$$

Positive probabilities in the base chain therefore lift to positive probabilities between any chosen representatives. This proves irreducibility. The positive return times to y and to $q(y)$ also agree, proving the assertion about aperiodicity. Finally,

$$(\tilde{\pi}\tilde{P})_z = \sum_{x \in S} \pi_x P_{x, q(z)} \eta_{q(z)}(z) = \pi_{q(z)} \eta_{q(z)}(z) = \tilde{\pi}_z,$$

where the probabilities within every fiber sum to one. $\square$

This operation also preserves the stationary mass of each block:

$$\tilde{\pi}(q^{-1}(C)) = \pi(C). \quad (24)$$

The cardinality of the lifted block is $|q^{-1}(C)| = \sum_{x \in C} m_x$. Thus the choice of multiplicities determines how membership in the original blocks is reflected in their lifted sizes. The multiplicities are fixed before any maps are sampled, so variation in those sizes occurs under one fixed lifted coupling law. If distinct subsets of S have distinct sums of multiplicities, each lifted block size identifies its original block. The entire lifted profile then determines the original partition, as used in the following corollary.

Corollary 6.2. *Any independent random map coupling of a finite irreducible chain whose eventual partition is not almost surely constant admits a finite state-splitting lift whose eventual cardinality profile is not almost surely constant.*

Proof. Enumerate $S = \{1, \dots, n\}$ and take $m_x = 2^{x-1}$. Distinct subsets of S have distinct sums of the m_x , by uniqueness of binary expansion. Thus the multiset of block sums uniquely determines the original unordered partition. Two distinct partitions of positive probability give two distinct lifted profiles of positive probability. Proposition 6.1 supplies the required irreducible coupling. $\square$

6.1 Splitting the four-state example

Grimmett and Holmes [3, Example 3.10] use the four maps

$$(1, 2, 1, 2), \quad (1, 2, 2, 1), \quad (3, 4, 3, 4), \quad (3, 4, 4, 3). \quad (25)$$

Their two possible eventual partitions are

$$\mathcal{D}_0 = \{\{1, 3\}, \{2, 4\}\}, \quad \mathcal{D}_1 = \{\{1, 4\}, \{2, 3\}\}. \quad (26)$$

Both have profile $(2, 2)$. Take equal probabilities on the four maps. Directly from (25), the transition matrix is

$$P_0 = \begin{pmatrix} 1/2 & 0 & 1/2 & 0 \\ 0 & 1/2 & 0 & 1/2 \\ 1/4 & 1/4 & 1/4 & 1/4 \\ 1/4 & 1/4 & 1/4 & 1/4 \end{pmatrix}. \quad (27)$$

Note that P is irreducible, aperiodic and doubly stochastic. Each map has image either $\{1, 2\}$ or $\{3, 4\}$, and every map separates both of these pairs. Thus the first map fixes the eventual partition, and the two possibilities in (26) each have probability $1/2$.

Now choose multiplicities

$$(m_1, m_2, m_3, m_4) = (2, 1, 2, 1), \quad (28)$$

with uniform section choices in each fiber. There are six lifted states. The block sums are

base partition	lifted block sizes	sorted profile
$\mathcal{D}_0$	$(m_1 + m_3, m_2 + m_4) = (4, 2)$	$(2, 4)$
$\mathcal{D}_1$	$(m_1 + m_4, m_2 + m_3) = (3, 3)$	$(3, 3)$

Proposition 6.1 gives a second six-state counterexample, again with probability $1/2$ for each profile. Its stationary masses are $1/8$ at the four copies of states 1 and 3 and $1/4$ at the copies of states 2 and 4.

This lift has eight distinct maps with equal probabilities: For each base map, its image contains one doubled state and one single state, and only the section choice at that doubled

image state affects the lifted map. It is a different coupling from the sixteen-map construction. In particular, its transition matrix retains zeros from (27); it is irreducible and aperiodic but not strictly positive. The lift proves that an elementary modification of the earlier random partition example suffices to produce random cardinalities.

Appendix. The complete map list and exact verification

For each oriented edge (u, v) , the two maps in the corresponding row below are $f_{0,u,v}$ and $f_{1,u,v}$. Every entry is the ordered tuple $(f(1), \dots, f(6))$. Each of the sixteen maps has probability $1/16$.

(u, v)	$f_{0,u,v}$	$f_{1,u,v}$
(1, 2)	(1, 2, 1, 1, 2, 1)	(1, 2, 1, 2, 1, 2)
(2, 1)	(2, 1, 2, 2, 1, 2)	(2, 1, 2, 1, 2, 1)
(2, 3)	(2, 3, 2, 2, 3, 2)	(2, 3, 2, 3, 2, 3)
(3, 2)	(3, 2, 3, 3, 2, 3)	(3, 2, 3, 2, 3, 2)
(4, 5)	(4, 5, 4, 4, 5, 4)	(4, 5, 4, 5, 4, 5)
(5, 4)	(5, 4, 5, 5, 4, 5)	(5, 4, 5, 4, 5, 4)
(5, 6)	(5, 6, 5, 5, 6, 5)	(5, 6, 5, 6, 5, 6)
(6, 5)	(6, 5, 6, 6, 5, 6)	(6, 5, 6, 5, 6, 5)

The accompanying source package contains a short script, `verify.py` which checks the marginal transition matrix, edge preservation, closure under all 16^2 compositions, and the eight-map state-splitting construction.

References

- [1] G. Budzban and Ph. Feinsilver, *A hierarchical structure of transformation semigroups with applications to probability limit measures*, Journal of Difference Equations and Applications **18** (2012), no. 8, 1405–1434. doi:10.1080/10236198.2011.639365. Preprint: arXiv:1103.0235.
- [2] J. Friedman, *On the road coloring problem*, Proceedings of the American Mathematical Society **110** (1990), no. 4, 1133–1135. doi:10.1090/S0002-9939-1990-0953004-8.
- [3] G. R. Grimmett and M. Holmes, *Coalescence in Markov chains*, Journal of Theoretical Probability **39** (2026), article 49. doi:10.1007/s10959-026-01500-w. Preprint: arXiv:2510.13572.
- [4] A. Schaefer, *Generating sets of the kernel graph and the inverse problem in synchronization theory*, preprint (2016), arXiv:1601.04295.
- [5] K. Yano, *Random walk in a finite directed graph subject to a road coloring*, Journal of Theoretical Probability **26** (2013), no. 1, 259–283. Preprint: arXiv:1005.0079.