



UNIVERSITÀ DI PISA

Dipartimento di Fisica “E. Fermi”

Corso di Laurea Magistrale in Fisica

Mott localisation, antiferromagnetism and nematic superconductivity in the extended Hubbard model with non-local pairing

Relatore:

Prof. Giacomo Mazza

Candidato:

Alessandro Gori

*Anno accademico 2025/2026
Draft: May 26, 2026*

*Ai miei meravigliosi nonni:
Aladina, Ferdinando, Giuliano e Matilde.*

Abstract

High temperature superconductivity in copper oxides (cuprates) represents one of the most striking phenomena in condensed matter physics. These materials host superconducting states with the highest known critical temperatures, emerging upon doping an antiferromagnetic insulator compound. The antiferromagnetic (AF) insulating and superconducting (SC) phases defy standard descriptions in terms of band picture and conventional phonon-mediated superconductivity. In contrast, these phenomena are genuine signatures of strong electronic correlations which hinder effective single-particle descriptions of the many-electron system.

After 40 years since their discovery, it is widely accepted that the Hubbard model on the square lattice provides a minimal description of the low-energy electronic properties of cuprates. The model describes the competition between the electronic kinetic energy, modelled by nearest-neighbours hopping t , and the Coulomb repulsion, modelled by a purely local interaction potential U . At half-filling, this competition results in the so-called “Mott localisation” phenomenon, which distinguishes the metallic phase for $U/t \ll 1$ from an insulating phase for $U/t \gg 1$. The latter is characterised by the almost complete localisation of electrons due to the high energetic cost of hopping towards occupied sites. Interestingly, this basic competition is intimately related to AF ordering and, upon doping, to the formation of the SC state with characteristic d -wave symmetry.

The fact that a purely repulsive interaction U is responsible for the formation of a SC state is remarkable, being superconductivity typically associated with an effective attractive interaction. In fact, a d -wave SC state can also be obtained, in general, by considering an attractive potential between electrons of neighbouring sites. Motivated by this observation, in this thesis, we studied a modified version of the Hubbard model, dubbed as the “extended Hubbard model”, in which we include a nearest-neighbours attractive potential $-V$. The general goal of this work is to explore the interplay between the purely local repulsion and the non-local attraction. It is worth mentioning that this is not only motivated by the fact that both interactions are, in principle, responsible for d -wave superconductivity, but also by recent experimental observations which strongly suggest that the term $-V$ needs to be included in order to accurately describe the magnetic spectra of some specific compounds.

In this thesis, we study the extended Hubbard model using the Hartree-Fock method. Namely, we approximate the ground state by implementing a variational search in the subspace of the many-body Hilbert space of gaussian states. Despite its simplicity, the Hartree-Fock method provides a powerful framework to study competition among different phases, originating from different symmetry breaking mechanisms. In particular, we study the competition between three different phases: the normal phase, characterised by no symmetry breaking, the AF phase, characterised by a lowering of $SU(2)$ symmetry, and the SC phase, characterised by the breaking of $U(1)$ charge conservation.

This thesis work led to a number of original results, which can be summarised by the general observation that the non-local attraction $-V$ acts cooperatively with the local repulsion U for the stabilisation of the various phases hosted by the Hubbard model.

For the normal phase, we find that the non-local attraction acts as a source of Mott localisation, leading to an effective reduction of the electronic bandwidth. In particular, for values of V larger than a critical value and close to half-filling, we find a strong Mott localisation characterised by the complete flattening of the electronic bands. Interestingly, this result shows an example of Mott localisation that can be described at the level of the Hartree-Fock approximation, whereas Mott localisation in the conventional Hubbard model usually requires more sophisticated approximations.

For the AF phase, we restrict our analysis to the half-filled case. We find that the pairing-induced Mott localisation leads to a significant enhancement of the magnetisation, which primarily originates from the local repulsion U .

Finally, we focus on the superconducting phase and derive self-consistency equations in both singlet and triplet Cooper pairing channels, showing how Cooper pairing can be mediated simultaneously by both the local repulsion and the non-local attraction. Therefore, we numerically investigate the singlet channel and find evidence of “symmetry mixing”, namely, the formation of a superconducting gap that is neither symmetric nor anti-symmetric under rotations of $\pi/2$ in the plane. We show that symmetry mixing is connected to the emergence of a finite nematic order parameter, that signals the breaking of planar rotational symmetry. This nematic symmetry breaking results in the fact that the renormalisation of bands becomes anisotropic, producing a difference in electronic kinetics in the x and y directions.

To summarise, the original results obtained in this thesis show that the non-local pairing provides a different route towards the formation of key strongly correlated phenomena, ranging from Mott localisation to nematic superconductivity, and call for future investigation using more sophisticated methods such as, for example, Dynamical Mean Field Theory.

Contents

Bibliography	3
------------------------------	---

List of symbols and abbreviations

AF	Antiferromagnetic
BCS	Bardeen-Cooper-Schrieffer (theory)
BZ	Brillouin zone
β	Inverse temperature
DoF	Degree of freedom
DoS	Density of states
EHM	Extended Hubbard model
EV(s)	Expectation value(s)
FS	Fermi surface
HF	Hartree-Fock
HFP(s)	Hartree-Fock parameter(s)
HM	Hubbard model
L_ℓ	Number of lattice sites in direction ℓ
iHF	Iterative Hartree-Fock
MBZ	Magnetic Brillouin zone
MFT	Mean-Field Theory
NBZ	Nematic Brillouin zone
NN(s)	Nearest neighbor(s)
PHS	Particle-hole symmetry
RMP(s)	Renormalized model parameter(s)
SC	Superconducting
SDW	Spin-density wave
SSB	Spontaneous symmetry breaking
T_c	Critical temperature

Acknowledgements

Questa tesi, così come tutti gli anni di Pisa, sono tutt'altro che un lavoro individuale. Se non avessi avuto attorno a me una rete di persone splendide a sostenermi, incoraggiarmi e aiutarmi in quei numerosi momenti di difficoltà, sarebbe andato tutto molto diversamente. Quindi, mi prendo un po' di spazio per i dovuti ringraziamenti.

Ringrazio per primo il Prof. Mazza, che mi ha fatto da relatore, mi ha proposto l'argomento di tesi e mi ha accompagnato durante tutto il percorso di preparazione di questo testo. Lo ringrazio specialmente per la pazienza e la dedizione mostrata nell'aiutarmi a rendere una valanga di calcoli qualcosa di umanamente intellegibile.

Ringrazio le amiche e gli amici di Fisica per gli anni passati assieme a maledire gli esami, a distribuire appunti, a scroccare buffet in Edificio E e seguire scuole estive *appassionanti* in giro per l'Europa, insomma, a viverci un po' meglio questa cosa pesante e complicata che è studiare Fisica. Lo stesso ringraziamento lo estendo a tutte le persone fuori da Fisica con cui ho condiviso questi anni (e non dite che i Fisici stanno solo tra loro, mica siamo lucchesi), nello specifico a chi con me ha abitato casa sgarrupata e con mille problemi, ma nella quale non si può dire che non ci siamo divertiti.

Ringrazio tutte le persone con cui ho condiviso la rappresentanza studentesca in CGS, compreso il gruppo che ci ha preceduto e quello – neoeletto, congratulazioni! – che verrà dopo di noi. Abbiamo provato a fare qualcosa per rendere un po' meglio il Dipartimento, e mi sembra che sia andata bene, tutto sommato. E se no, pazienza: almeno ci siamo divertiti. Ringrazio e abbraccio le compagne e i compagni di Exploit, che tutti i giorni mi mostrano che un'altra via è possibile, e che si può camminare insieme in quella direzione.

Ringrazio tutta la mia famiglia, che chi mi conosce sa essere numerosa, quindi mi permetto di esplicitare solo alcuni dei ringraziamenti personali. Ai miei genitori, che hanno sempre creduto in me e che mi hanno supportato in tutto, e non scordiamocelo, mi hanno insegnato a stare al mondo. A mio fratello, che per me è sempre stato un faro e un'ispirazione meravigliosa, con il quale condivido *cose che solo tra fratelli*, e quindi neanche ci provo a scriverle, non sarei in grado.

...

... e grazie per tutto il pesce! [D. A.]

Bibliography

- [1] J. G. Bednorz and K. A. Muller. “Possible high-Tc superconductivity in the Ba-La-Cu-O system”. en. In: *Zeitschrift für Physik B Condensed Matter* 64.2 (June 1986), pp. 189–193. ISSN: 1431-584X. DOI: [10.1007/BF01303701](https://doi.org/10.1007/BF01303701). URL: <https://doi.org/10.1007/BF01303701> (visited on 04/03/2026).
- [2] Liangzi Deng et al. “Ambient-pressure 151 K superconductivity in HgBa₂Ca₂Cu₃O₈ via pressure quench”. In: *Proceedings of the National Academy of Sciences* 123.11 (Mar. 2026), e2536178123. DOI: [10.1073/pnas.2536178123](https://www.pnas.org/doi/10.1073/pnas.2536178123). URL: <https://www.pnas.org/doi/10.1073/pnas.2536178123> (visited on 04/03/2026).
- [3] B. Keimer et al. “From quantum matter to high-temperature superconductivity in copper oxides”. en. In: *Nature* 518.7538 (Feb. 2015), pp. 179–186. ISSN: 0028-0836, 1476-4687. DOI: [10.1038/nature14165](https://www.nature.com/articles/nature14165). URL: <https://www.nature.com/articles/nature14165> (visited on 03/15/2025).
- [4] Giuseppe Grosso and Giuseppe Pastori Parravicini. *Solid State Physics*. Second Edition. Academic Press, 2014.
- [5] Patrick A. Lee, Naoto Nagaosa, and Xiao-Gang Wen. “Doping a Mott insulator: Physics of high-temperature superconductivity”. en. In: *Reviews of Modern Physics* 78.1 (Jan. 2006), pp. 17–85. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.78.17](https://link.aps.org/doi/10.1103/RevModPhys.78.17). URL: <https://link.aps.org/doi/10.1103/RevModPhys.78.17> (visited on 05/07/2026).
- [6] J. Hubbard. “Electron Correlations in Narrow Energy Bands”. en. In: *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* 276.1365 (1963), pp. 238–257. URL: <http://www.jstor.org/stable/2414761>.
- [7] F. C. Zhang et al. “A renormalised Hamiltonian approach to a resonant valence bond wave-function”. en. In: *Superconductor Science and Technology* 1.1 (June 1988), p. 36. ISSN: 0953-2048. DOI: [10.1088/0953-2048/1/1/009](https://iopscience.iop.org/article/10.1088/0953-2048/1/1/009). URL: <https://iopscience.iop.org/article/10.1088/0953-2048/1/1/009> (visited on 05/11/2026).
- [8] D. J. Scalapino. “A common thread: The pairing interaction for unconventional superconductors”. en. In: *Reviews of Modern Physics* 84.4 (Oct. 2012), pp. 1383–1417. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.84.1383](https://link.aps.org/doi/10.1103/RevModPhys.84.1383). URL: <https://link.aps.org/doi/10.1103/RevModPhys.84.1383> (visited on 04/03/2026).
- [9] Daniel P. Arovas et al. “The Hubbard Model”. In: *Annual Review of Condensed Matter Physics* 13. Volume 13, 2022 (2022), pp. 239–274. ISSN: 1947-5462. DOI: <https://doi.org/10.1146/annurev-conmatphys-031620-102024>. URL: <https://www.annualreviews.org/content/journals/10.1146/annurev-conmatphys-031620-102024>.
- [10] Jan Zaanen. “Watching Rush Hour in the World of Electrons”. In: *Science* 315.5817 (Mar. 2007), pp. 1372–1373. DOI: [10.1126/science.1139805](https://www.science.org/doi/10.1126/science.1139805). URL: <https://www.science.org/doi/10.1126/science.1139805> (visited on 04/03/2026).
- [11] Antoine Georges et al. “Dynamical mean-field theory of strongly correlated fermion systems and the limit of infinite dimensions”. en. In: *Reviews of Modern Physics* 68.1 (Jan. 1996), pp. 13–125. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.68.13](https://link.aps.org/doi/10.1103/RevModPhys.68.13). URL: <https://link.aps.org/doi/10.1103/RevModPhys.68.13> (visited on 03/15/2025).

- [12] Gabriel Kotliar and Dieter Vollhardt. “Strongly Correlated Materials: Insights From Dynamical Mean-Field Theory”. In: *Physics Today* 57 (Mar. 2004). ADS Bibcode: 2004PhT....57c..53K, pp. 53–59. ISSN: 0031-9228. DOI: [10.1063/1.1712502](https://doi.org/10.1063/1.1712502). URL: <https://ui.adsabs.harvard.edu/abs/2004PhT....57c..53K> (visited on 05/11/2026).
- [13] J. Brad Marston and Ian Affleck. “Large- n limit of the Hubbard-Heisenberg model”. en. In: *Physical Review B* 39.16 (June 1989), pp. 11538–11558. ISSN: 0163-1829. DOI: [10.1103/PhysRevB.39.11538](https://doi.org/10.1103/PhysRevB.39.11538). URL: <https://link.aps.org/doi/10.1103/PhysRevB.39.11538> (visited on 05/07/2026).
- [14] Piers Coleman. *Introduction to Many-Body Physics*. Cambridge University Press, 2015.
- [15] P. W. Anderson. “Resonating valence bonds: A new kind of insulator?”. In: *Materials Research Bulletin* 8.2 (Feb. 1973), pp. 153–160. ISSN: 0025-5408. DOI: [10.1016/0025-5408\(73\)90167-0](https://doi.org/10.1016/0025-5408(73)90167-0). URL: <https://www.sciencedirect.com/science/article/pii/0025540873901670> (visited on 05/11/2026).
- [16] P. Fazekas and P. W. Anderson. “On the ground state properties of the anisotropic triangular antiferromagnet”. en. In: *Philosophical Magazine* 30.2 (Aug. 1974), pp. 423–440. ISSN: 0031-8086. DOI: [10.1080/14786439808206568](https://doi.org/10.1080/14786439808206568). URL: <http://www.tandfonline.com/doi/abs/10.1080/14786439808206568> (visited on 05/11/2026).
- [17] M. T. Béal-Monod, C. Bourbonnais, and V. J. Emery. “Possible superconductivity in nearly antiferromagnetic itinerant fermion systems”. en. In: *Physical Review B* 34.11 (Dec. 1986), pp. 7716–7720. ISSN: 0163-1829. DOI: [10.1103/PhysRevB.34.7716](https://doi.org/10.1103/PhysRevB.34.7716). URL: <https://link.aps.org/doi/10.1103/PhysRevB.34.7716> (visited on 05/11/2026).
- [18] K. Miyake, S. Schmitt-Rink, and C. M. Varma. “Spin-fluctuation-mediated even-parity pairing in heavy-fermion superconductors”. en. In: *Physical Review B* 34.9 (Nov. 1986), pp. 6554–6556. ISSN: 0163-1829. DOI: [10.1103/PhysRevB.34.6554](https://doi.org/10.1103/PhysRevB.34.6554). URL: <https://link.aps.org/doi/10.1103/PhysRevB.34.6554> (visited on 05/11/2026).
- [19] D. J. Scalapino, E. Loh, and J. E. Hirsch. “ d -wave pairing near a spin-density-wave instability”. In: *Physical Review B* 34.11 (Dec. 1986), pp. 8190–8192. DOI: [10.1103/PhysRevB.34.8190](https://doi.org/10.1103/PhysRevB.34.8190). URL: <https://link.aps.org/doi/10.1103/PhysRevB.34.8190> (visited on 05/11/2026).
- [20] C. Walsh et al. “Superconductivity in the two-dimensional Hubbard model with cellular dynamical mean-field theory: A quantum impurity model analysis”. en. In: *Physical Review B* 108.7 (Aug. 2023), p. 075163. ISSN: 2469-9950, 2469-9969. DOI: [10.1103/PhysRevB.108.075163](https://doi.org/10.1103/PhysRevB.108.075163). URL: <https://link.aps.org/doi/10.1103/PhysRevB.108.075163> (visited on 05/23/2026).
- [21] Mi Jiang. “Enhancing d -wave superconductivity with nearest-neighbor attraction in the extended Hubbard model”. en. In: *Physical Review B* 105.2 (Jan. 2022), p. 024510. ISSN: 2469-9950, 2469-9969. DOI: [10.1103/PhysRevB.105.024510](https://doi.org/10.1103/PhysRevB.105.024510). URL: <https://link.aps.org/doi/10.1103/PhysRevB.105.024510> (visited on 05/08/2026).
- [22] Hari Padma et al. “Beyond-Hubbard pairing in a cuprate ladder”. en. In: *Physical Review X* 15.2 (May 2025), p. 021049. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.15.021049](https://doi.org/10.1103/PhysRevX.15.021049). URL: <https://link.aps.org/doi/10.1103/PhysRevX.15.021049> (visited on 04/03/2026).
- [23] R. Micnas, J. Ranninger, and S. Robaszkiewicz. “Superconductivity in narrow-band systems with local nonretarded attractive interactions”. In: *Reviews of Modern Physics* 62.1 (Jan. 1990). Publisher: American Physical Society, pp. 113–171. DOI: [10.1103/RevModPhys.62.113](https://doi.org/10.1103/RevModPhys.62.113). URL: <https://link.aps.org/doi/10.1103/RevModPhys.62.113> (visited on 01/06/2026).
- [24] J. E. Hirsch. “Two-dimensional Hubbard model: Numerical simulation study”. In: *Phys. Rev. B* 31 (7 Apr. 1985), pp. 4403–4419. DOI: [10.1103/PhysRevB.31.4403](https://doi.org/10.1103/PhysRevB.31.4403). URL: <https://link.aps.org/doi/10.1103/PhysRevB.31.4403>.
- [25] Robin Scholle et al. “Comprehensive mean-field analysis of magnetic and charge orders in the two-dimensional Hubbard model”. In: *Phys. Rev. B* 108 (3 July 2023), p. 035139. DOI: [10.1103/PhysRevB.108.035139](https://doi.org/10.1103/PhysRevB.108.035139). URL: <https://link.aps.org/doi/10.1103/PhysRevB.108.035139>.

- [26] Zhangkai Cao et al. “Dominant p -wave pairing induced by nearest-neighbor attraction in the square-lattice extended Hubbard model”. In: *Phys. Rev. B* 111 (2 Jan. 2025), p. 024509. DOI: [10.1103/PhysRevB.111.024509](https://doi.org/10.1103/PhysRevB.111.024509). URL: <https://link.aps.org/doi/10.1103/PhysRevB.111.024509>.
- [27] Avinash Singh and Zlatko Tešanović. “Collective excitations in a doped antiferromagnet”. en. In: *Physical Review B* 41.1 (Jan. 1990), pp. 614–631. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.41.614](https://doi.org/10.1103/PhysRevB.41.614). URL: <https://link.aps.org/doi/10.1103/PhysRevB.41.614> (visited on 12/12/2025).
- [28] Avinash Singh, Zlatko Tešanović, and Ju H. Kim. “Doped Hubbard antiferromagnet: Instability and effective interactions”. en. In: *Physical Review B* 44.14 (Oct. 1991), pp. 7757–7759. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.44.7757](https://doi.org/10.1103/PhysRevB.44.7757). URL: <https://link.aps.org/doi/10.1103/PhysRevB.44.7757> (visited on 12/12/2025).
- [29] Alexander Wietek et al. “Stripes, Antiferromagnetism, and the Pseudogap in the Doped Hubbard Model at Finite Temperature”. en. In: *Physical Review X* 11.3 (July 2021), p. 031007. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.11.031007](https://doi.org/10.1103/PhysRevX.11.031007). URL: <https://link.aps.org/doi/10.1103/PhysRevX.11.031007> (visited on 12/12/2025).
- [30] Kazuyoshi Yoshimi, Takeo Kato, and Hideaki Maebashi. “Enhanced Spin Susceptibility toward the Charge-Ordering Transition in a Two-Dimensional Extended Hubbard Model”. In: *Journal of the Physical Society of Japan* 78.10 (Oct. 2009), p. 104002. ISSN: 0031-9015. DOI: [10.1143/JPSJ.78.104002](https://doi.org/10.1143/JPSJ.78.104002). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.78.104002> (visited on 04/17/2026).
- [31] Ashvin Vishwanath. “Dirac Nodes and Quantized Thermal Hall Effect in the Mixed State of d-wave Superconductors”. In: *Physical Review B* 66.6 (Aug. 2002). arXiv:cond-mat/0104213, p. 064504. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.66.064504](https://doi.org/10.1103/PhysRevB.66.064504). URL: <http://arxiv.org/abs/cond-mat/0104213> (visited on 01/05/2026).
- [32] P. Senarath Yapa et al. “Mixed-symmetry superconductivity and the energy gap”. en. In: *Physica C: Superconductivity and its Applications* 633 (June 2025), p. 1354719. ISSN: 09214534. DOI: [10.1016/j.physc.2025.1354719](https://doi.org/10.1016/j.physc.2025.1354719). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0921453425000723> (visited on 01/06/2026).
- [33] Joel Hutchinson and Frank Marsiglio. “Mixed temperature-dependent order parameters in the extended Hubbard model”. en. In: *Journal of Physics: Condensed Matter* 33.6 (Feb. 2021), p. 065603. ISSN: 0953-8984, 1361-648X. DOI: [10.1088/1361-648X/abc801](https://doi.org/10.1088/1361-648X/abc801). URL: <https://iopscience.iop.org/article/10.1088/1361-648X/abc801> (visited on 01/06/2026).
- [34] C. Zhou and H. J. Schulz. “Density of states and tunneling spectra in two-dimensional d-wave superconductors”. In: *Physical Review B* 45.13 (Apr. 1992). Publisher: American Physical Society, pp. 7397–7401. DOI: [10.1103/PhysRevB.45.7397](https://doi.org/10.1103/PhysRevB.45.7397). URL: <https://link.aps.org/doi/10.1103/PhysRevB.45.7397> (visited on 01/06/2026).
- [35] James F. Annett. “Symmetry of the order parameter for high-temperature superconductivity”. In: *Advances in Physics* 39.2 (Apr. 1990). Publisher: Taylor & Francis .eprint: <https://doi.org/10.1080/00018739000101481>, pp. 83–126. ISSN: 0001-8732. DOI: [10.1080/00018739000101481](https://doi.org/10.1080/00018739000101481). URL: <https://doi.org/10.1080/00018739000101481> (visited on 01/06/2026).
- [36] Manfred Sgrist and Kazuo Ueda. “Phenomenological theory of unconventional superconductivity”. In: *Reviews of Modern Physics* 63.2 (Apr. 1991). Publisher: American Physical Society, pp. 239–311. DOI: [10.1103/RevModPhys.63.239](https://doi.org/10.1103/RevModPhys.63.239). URL: <https://link.aps.org/doi/10.1103/RevModPhys.63.239> (visited on 01/07/2026).
- [37] Anna E. Böhrer et al. “Nematicity and nematic fluctuations in iron-based superconductors”. en. In: *Nature Physics* 18.12 (Dec. 2022), pp. 1412–1419. ISSN: 1745-2481. DOI: [10.1038/s41567-022-01833-3](https://doi.org/10.1038/s41567-022-01833-3). URL: <https://www.nature.com/articles/s41567-022-01833-3>.

- [38] Mingpu Qin et al. “Absence of Superconductivity in the Pure Two-Dimensional Hubbard Model”. In: *Physical Review X* 10.3 (July 2020), p. 031016. DOI: [10.1103/PhysRevX.10.031016](https://doi.org/10.1103/PhysRevX.10.031016). URL: <https://link.aps.org/doi/10.1103/PhysRevX.10.031016> (visited on 03/21/2026).
- [39] Jan Zaanen and Olle Gunnarsson. “Charged magnetic domain lines and the magnetism of high- T_c oxides”. In: *Physical Review B* 40.10 (Oct. 1989), pp. 7391–7394. DOI: [10.1103/PhysRevB.40.7391](https://doi.org/10.1103/PhysRevB.40.7391). URL: <https://link.aps.org/doi/10.1103/PhysRevB.40.7391> (visited on 03/21/2026).
- [40] Masaru Kato et al. “Soliton Lattice Modulation of Incommensurate Spin Density Wave in Two Dimensional Hubbard Model -A Mean Field Study-”. In: *Journal of the Physical Society of Japan* 59.3 (Mar. 1990), pp. 1047–1058. ISSN: 0031-9015. DOI: [10.1143/JPSJ.59.1047](https://doi.org/10.1143/JPSJ.59.1047). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.59.1047> (visited on 03/21/2026).
- [41] Eugene Kogan and Godfrey Gumbs. “Green’s Functions and DOS for Some 2D Lattices”. en. In: *Graphene* 10.01 (2021), pp. 1–12. ISSN: 2169-3439, 2169-3471. DOI: [10.4236/graphene.2021.101001](https://doi.org/10.4236/graphene.2021.101001). URL: <https://www.scirp.org/journal/paperinformation?paperid=104591> (visited on 03/26/2026).
- [42] Charles L. Cleveland and Rodrigo Medina A. “Obtaining a Heisenberg Hamiltonian from the Hubbard model”. en. In: *American Journal of Physics* 44.1 (Jan. 1976), pp. 44–46. ISSN: 0002-9505, 1943-2909. DOI: [10.1119/1.10537](https://doi.org/10.1119/1.10537). URL: <https://pubs.aip.org/ajp/article/44/1/44/1050180/Obtaining-a-Heisenberg-Hamiltonian-from-the> (visited on 05/08/2026).
- [43] Eva Pavarini et al., eds. *Many-body physics: from Kondo to Hubbard: lecture notes of the Autumn School on Correlated Electrons 2015: at Forschungszentrum Jülich, 21-25 September 2015*. en. Band 5. Meeting Name: Autumn School on Correlated Electrons. Forschungszentrum Jülich, 2015. ISBN: 978-3-95806-074-6.
- [44] Gabriele Giuliani and Giovanni Vignale. *Quantum Theory of the Electron Liquid*. Cambridge University Press, 2005.
- [45] Michele Fabrizio. *A Course in Quantum Many-Body Theory*. Springer, 2022.
- [46] Eva Pavarini et al., eds. *Correlated electrons: from models to materials: lecture notes of the Autumn School Correlated Electrons 2012: at Forschungszentrum Jülich, 3-7 September 2012*. eng. Schriften des Forschungszentrums Jülich. Reihe Modeling and Simulation Band 2. Meeting Name: Autumn School Correlated Electrons. Jülich: Forschungszentrum Jülich, Zentralbibliothek, Verl, 2012. ISBN: 978-3-89336-796-2.