

Numerical analysis of superconducting phases in the extended Hubbard model with non-local pairing

University of Pisa, a.y. 2025-2026

Alessandro Gori*

Thesis for the Master's degree in Physics

Abstract

[To be continued...]

- Appendix on s -wave SC ghost simulations on HM
- Appendix on triplet pairing in MFT self-consistent scheme
- Set ComputerModern font in all CairoMakie figures

Contents

1	Theoretical introduction	3
1.1	Antiferromagnetic ordering in the Hubbard model	3
1.2	The Extended Fermi-Hubbard model	3
1.2.1	Experimental insight on NN attraction	4
2	Mean-field theory discussion of the EHM	5
2.1	Introduction to Hartree-Fock methods	5
2.1.1	Slater determinants and generalities	5
2.1.2	Wick's theorem applied to Slater determinants	6
2.1.3	Gaussian states and Slater determinants	7
2.1.4	Sketch of HF variational method	7
2.2	Extended Hubbard model symmetries and Wick's theorem	8
2.2.1	Spatial and topological symmetries	8
2.2.2	Spin and charge symmetries	8
2.2.3	Particle-hole symmetry at half-filling	9
2.3	MFT on Hubbard lattices	11
2.3.1	MFT general procedure, self-consistency and Wick's theorem	11
2.3.2	MFT treatment of the pure Hubbard model	12
2.4	The non-local attraction $\hat{H}_V$ as a source of symmetry-breaking interactions	13
2.4.1	Real space description	14
2.4.2	Fourier transform of the model	15
2.5	Iterative computational approach to MFT	17
2.5.1	Sketch of the Hartree-Fock iterative algorithm	17
2.5.2	Self-consistency and RMPs	18
2.5.3	Fixed density simulations and the role of μ	19
2.5.4	The reject-mix-iterate convergence scheme	20
2.6	Square lattice spatial symmetry channels	20

*a.gori23@studenti.unipi.it / nepero27178@github.com

Bibliography

25

List of symbols and abbreviations

AF	Antiferromagnetic
BCS	Bardeen-Cooper-Schrieffer (theory)
BZ	Brillouin zone
β	Inverse temperature
$\Delta^{(\gamma)}$	γ -wave component of the gap function
DoF	Degree of freedom
EHM	Extended Hubbard model
EV(s)	Expectation value(s)
FS	Fermi surface
HF	Hartree-Fock
HFP(s)	Hartree-Fock parameter(s)
HM	Hubbard model
L_ℓ	Number of lattice sites in direction ℓ
LRT	Linear Response Theory
MBZ	Magnetic Brillouin zone
MFT	Mean-Field Theory
NBZ	Nematic Brillouin zone
NN(s)	Nearest neighbor(s)
RWC(s)	Relevant Wick's contraction(s)
RMP(s)	Renormalized model parameter(s)
SC	Superconductor
SDW	Spin-density wave
SSB	Spontaneous symmetry breaking
T_c	Critical temperature

Introduction

This thesis project is about my favorite ice cream flavor. [To be continued...]

Chapter 1

Theoretical introduction

[To be continued...]

1.1 Antiferromagnetic ordering in the Hubbard model

Consider the ordinary Hubbard model:

$$\hat{H} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \quad t, U > 0 \quad (1.1)$$

The two competing mechanisms are site-hopping of amplitude t and local repulsion of amplitude U . For this model defined on a bipartite lattice at half filling and fixed electron number, it is well known [hirsch1985hubbard] that, below a certain critical temperature T_c and above some (small) critical repulsion U_c/t , the ground-state acquires antiferromagnetic (AF) long-range ordering, schematically depicted in Fig. 1.1a. The mechanism for the formation of the AF phase takes advantage of virtual hopping, as described in App. ??; the Mean-Field Theory (MFT) treatment of ferromagnetic-antiferromagnetic orderings in 2D Hubbard lattices is discussed in App. ??.

In this chapter the discussion is limited to the two-dimensional square lattice Hubbard model. The lattice considered has L_{ℓ} sites on side $\ell = x, y$, thus a total of $L_x L_y$ sites. The total number of single electron states is given by $D = 2L_x L_y$. All theoretical discussion neglects border effects, thus considering $D \rightarrow +\infty$.

1.2 The Extended Fermi-Hubbard model

The Extended Fermi-Hubbard model is defined by:

$$\hat{H} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'} \quad (1.2)$$




Figure 1.1 | Schematic representation of the AF phase. Fig. 1.1a shows a portion of the square lattice with explicit representation of the spin for each site. Fig. 1.1b divides the square lattice $\mathcal{S}$ in two polarized sublattices $\mathcal{S}_a, \mathcal{S}_b$. The AF phase results from the interaction of two inversely polarized “ferromagnetic” square lattices.

Note that, on a square lattice, we can perform the summation over NN just as

$$\sum_{\langle ij \rangle} \equiv \sum_{i \in \mathcal{S}_a} [\delta_{j=i+\delta_x} + \delta_{j=i-\delta_x} + \delta_{j=i+\delta_y} + \delta_{j=i-\delta_y}]$$

where notation of Fig. 1.1b has been used. The last term represents an effective attraction between neighboring electrons, of amplitude V . Such an interaction is believed [cao2025p-wave] necessary to describe the insurgence of high- T_c superconductivity in cuprate SCs. [To be continued...]

1.2.1 Experimental insight on NN attraction

[Add:

- High T_c SC in cuprates;
- Experimental evidence of topological SC;
- Insertion of the non-local attraction;

]

Chapter 2

Mean-field theory discussion of the EHM

[Introduction: this is a technical chapter about Hartree Fock applied to EHM.]

2.1 Introduction to Hartree-Fock methods

The Hartree-Fock (HF) method is a variational approach to determine the ground-state wavefunction of a many-body hamiltonian. The wavefunction is searched in a variational scheme as the one minimising energy, limiting search to the subspace of gaussian states. Let us introduce these concepts. The sections that follow are based on the discussions of [6, 8].

2.1.1 Slater determinants and generalities

For a electronic system, it follows from fermionic algebra that the many-body wavefunction needs to be antisymmetric under interchange of coordinates and spins $(\mathbf{r}_i, \sigma_i)$, $(\mathbf{r}_j, \sigma_j)$ of any couple of electrons. The simplest way to build a many-body wavefunction satisfying this rule is “to antisymmetrize” a set of orthonormal N one-body spin-orbitals, being N the total number of electrons. Let $\{\psi_i\}$ be said set,

$$\frac{1}{2\mathcal{V}} \sum_{\sigma} \int d\mathbf{r} \psi_i(\mathbf{r}, \sigma) \psi_j(\mathbf{r}, \sigma) = \delta_{ij}$$

being $\mathcal{V}$ the total volume. Let then the antisymmetrization operator

$$\mathcal{A} \equiv \frac{1}{\sqrt{N!}} \sum_{P \in S_N} (-1)^{\sigma(P)} P$$

being S_N the $N!$ elements symmetric group of N objects permutations, and $\sigma(P)$ the “parity factor” of any permutation $P \in S_N$. $\sigma(P)$ counts the number of exchanges needed to achieve a given configuration. This way, an even number of exchanges gives $+1$, an odd number gives -1 . The permutation operator exchanges spin-coordinates couples,

$$P : (\mathbf{r}, \sigma) \leftrightarrow (\mathbf{r}', \sigma')$$

thus any given pair is exchanged with another.

Equipped with $\mathcal{A}$, we define the many-body wavefunction

$$\Psi(\mathbf{r}_1, \sigma_1; \mathbf{r}_2, \sigma_2; \dots; \mathbf{r}_N, \sigma_N) = \mathcal{A} \left\{ \prod_{i=1}^N \psi_i(\mathbf{r}_i, \sigma_i) \right\}$$

which by construction is antisymmetric under exchange of coordinates and spins pairs. This class of wavefunctions, obtained as antisymmetric combinations of product states, are representable as so-called Slater determinants,

$$\Psi(\mathbf{r}_1, \sigma_1; \mathbf{r}_2, \sigma_2; \dots; \mathbf{r}_N, \sigma_N) = \frac{1}{\sqrt{N!}} \det \begin{bmatrix} \psi_1(\mathbf{r}_1, \sigma_1) & \psi_1(\mathbf{r}_2, \sigma_2) & \cdots & \psi_1(\mathbf{r}_N, \sigma_N) \\ \psi_2(\mathbf{r}_1, \sigma_1) & \psi_2(\mathbf{r}_2, \sigma_2) & \cdots & \psi_2(\mathbf{r}_N, \sigma_N) \\ \vdots & \vdots & & \vdots \\ \psi_N(\mathbf{r}_1, \sigma_1) & \psi_N(\mathbf{r}_2, \sigma_2) & \cdots & \psi_N(\mathbf{r}_N, \sigma_N) \end{bmatrix}$$

Determinants of matrices with two equal rows or two equal columns are null. Thus, all spin-orbitals must differ, as well as no single spin-orbital ψ_i can be doubly occupied (which is, all spin-coordinate couples must be different). Pauli exclusion principle is included in this framework.

Any normalised combinations of Slater determinants gives an antisymmetric wavefunction, although not necessarily representable as a single Slater determinant. In HF theory, we search for the best possible choice of orthonormal one-body spin orbitals $\{\psi_1, \psi_2, \dots, \psi_N\}$ to approximate the true ground-state of the model. The space of possible “Slater-like” many-body wavefunctions is the parametric space where we perform the variational constrained search.

The EHM is formulated in second-quantisation. It is then useful to represent Slater-like states in this language. Let us map all the spin-orbitals from the discussion above to a couple of annihilation-creation fermionic operators,

$$\psi_i \rightarrow \hat{c}_i, \hat{c}_i^\dagger$$

Any Slater state can be expressed in second-quantisation as:

$$|\Psi\rangle = \prod_{i=1}^N \hat{c}_i^\dagger |\Omega\rangle \quad (2.1)$$

being $|\Omega\rangle$ the vacuum.

Note that up to now we never required explicitly $\hat{c}_i$ to be “standard” electronic operators. In fact, let M be any unitary matrix, and

$$\hat{a}_i \equiv \sum_j M_{ij} \hat{c}_j \quad \hat{a}_i^\dagger \equiv \sum_j M_{ji}^* \hat{c}_j^\dagger$$

be a combination of annihilation operators. The new operators satisfy Fermi anticommutation rules,

$$\begin{aligned} \{\hat{a}_i, \hat{a}_j^\dagger\} &= \sum_{i'} M_{ii'} \sum_{j'} M_{jj'} \{\hat{c}_{i'}, \hat{c}_{j'}^\dagger\} \\ &= \sum_{i'} M_{ii'} \sum_{j'} M_{j'i}^* \delta_{i'j'} \\ &= [MM^\dagger]_{ij} \\ &= \delta_{ij} \end{aligned}$$

being M unitary. Express Eq. (2.1) in the new basis:

$$\begin{aligned} |\Psi\rangle &= \prod_{i=1}^N \hat{c}_i^\dagger |\Omega\rangle \\ &= \prod_{i=1}^N \sum_j M_{ij} \hat{a}_j^\dagger |\Omega\rangle \end{aligned}$$

The concise expression of Eq. (2.1) may appear more complicated in a different basis. The core concept is that Slater determinants represent, for a specific choice of fermionic operators, a non-interacting Fermi gas. HF methods assume the ground state to be of the form of Eq. (2.1) for an unknown set of fermionic operators, and then look for the best approximation within all these possible Slater states.

2.1.2 Wick’s theorem applied to Slater determinants

In this picture, second-quantised fermionic $\hat{c}$ operators span, in general, a subset of all possible spin-orbitals (states in Hilbert space). Let $\hat{c}_i$ with $i > N$ identify all spin-orbitals not already included in the definition of $|\Psi\rangle$ of Eq. (2.1). Then it holds:

$$\begin{aligned} 0 \leq i \leq N & : \quad \hat{c}_i^\dagger |\Psi\rangle = 0 \\ i > N & : \quad \hat{c}_i |\Psi\rangle = 0 \end{aligned}$$

Let:

$$\hat{a}_i \equiv \begin{cases} \hat{c}_i^\dagger & \text{for } 0 \leq i \leq N \\ \hat{c}_i & \text{for } i > N \end{cases}$$

These new operators satisfy a Fermi algebra $\{\hat{a}_i, \hat{a}_j^\dagger\} = \delta_{ij}$. They are annihilation and creation operators above the “vacuum” state $|\Psi\rangle$. In other words, any Slater state can be seen as a vacuum state for a suitably chosen set of fermionic operators.

This property is particularly handy for evaluation of expectation values (EV). By applying the transformation described above, any $\mathcal{O}(c)$ operator $\hat{C}_\ell$ (either $\hat{c}_i$ or $\hat{c}_i^\dagger$ for some i) can be mapped to its $\mathcal{O}(a)$ counterpart $\hat{A}_\ell$. Now, Wick’s theorem states that

$$\begin{aligned} \hat{C}_1 \hat{C}_2 \cdots \hat{C}_{2k} &= N \left\{ \hat{C}_1 \hat{C}_2 \cdots \hat{C}_{2k} \right\} \\ &+ \sum_{(\alpha\beta)} (-1)^{\zeta_{\alpha\beta}} D_{\alpha\beta} N \left\{ \prod_{i \neq \alpha, \beta} \hat{C}_i \right\} \\ &+ \sum_{(\alpha\beta)(\gamma\delta)} (-1)^{\zeta_{\alpha\beta}} D_{\alpha\beta} (-1)^{\zeta_{\gamma\delta}} D_{\gamma\delta} N \left\{ \prod_{i \neq \alpha, \beta, \gamma, \delta} \hat{C}_i \right\} \\ &+ \cdots \end{aligned}$$

where $N\{\cdots\}$ indicates normal ordering, $D_{\alpha\beta}$ is the contraction of the pair $\hat{C}_\alpha \hat{C}_\beta$ and $\zeta_{\alpha\beta}$ is a symmetry factor included to take care of the number of fermionic algebra. Sums are performed over pairs of indices, implicitly taken to be different from each other. Now, to take averages of this complicated operator over $|\Psi\rangle$ is in general non trivial. If we change basis and express all operators in terms of $\hat{a}, \hat{a}^\dagger$,

$$\hat{C}_i \rightarrow \hat{A}_i$$

the above decomposition holds identically, with slight change of notation. Now, taking a zero temperature average computed on $|\Psi\rangle$, all normal-ordered terms disappear,

$$\langle \Psi | N\{\hat{A}_a \hat{A}_b \cdots\} | \Psi \rangle = 0$$

thus in the decomposition above only fully contracted terms survive. Going back to $\hat{c}$ operators, we can re-express fully contracted terms in the original basis. In this procedure, the only key property of Slater states we used was the fact that they are vacuum to some fermionic basis. Finite-temperature extension of Wick’s theorem as well as more complicated change of basis are possible, but the concept remains the same: on a Slater state, which is essentially a non-interacting state for some set of fermions, Wick’s theorem is particularly well-behaved.

From the point above it follows that

$$\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta \rangle = \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\delta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\gamma \rangle}_{\text{Hartree}} - \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \langle \hat{c}_\gamma^\dagger \hat{c}_\delta \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \langle \hat{c}_\gamma \hat{c}_\delta \rangle}_{\text{Cooper}}$$

being the average intended on a Slater state. The subscript indicate the name we will give to these terms throughout the text.

2.1.3 Gaussian states and Slater determinants

In this text we make use of an extended version of HF method, often referred to as Hartree-Fock-Bogoliubov (HFB). Minimisation is performed not only over Slater determinants, but a larger set: gaussian states. Introducing the concept of Slater determinants was useful to describe intuitively how HF methods work, but now we extend to a broader set.

[To be continued...]

2.1.4 Sketch of HF variational method

[To be continued...]

2.2 Extended Hubbard model symmetries and Wick's theorem

The general aim of this project is to study the phase diagram of the EHM by comparing ground-state energies of different phases. The phases we consider in next chapters for the EHM are the normal state (N), the antiferromagnetic ordering (AF), given by a non-uniform distribution of charge in each spin sector, and the anisotropic superconducting (SC) phase, described by a uniformly distributed charge allowing for Cooper pairing instabilities.

At the end of this section, in Tab. 2.2 we summarize all symmetries with the relative operations. Throughout the entire thesis we will refer to specific SSB for given groups to establish a specific phase.

2.2.1 Spatial and topological symmetries

Let us start from the trivial topology of the lattice we are working on. The EHM of Eq. (1.2) is evidently translational invariant on both the x and y directions. We refer to these groups as $T_1^{(x)}$, $T_1^{(y)}$. The “1” subscript indicates the one-site translations under which the model is symmetric.

Evidently, the following statement holds:

$$T_n^{(\ell)} \supset T_{kn}^{(\ell)} \quad k \in \mathbb{N}$$

A model which is invariant for n -sites translations in the ℓ direction, also is for kn -sites translations. Then we can break translation symmetry in a given direction reducing it to a smaller group. For example, the commensurate AF phase we analyze halves translational invariance,

$$\bigotimes_{\ell=x,y} T_1^{(\ell)} \xrightarrow{\text{SSB}} \bigotimes_{\ell=x,y} T_2^{(\ell)}$$

since it deals with two-site periodic spin-density waves (SDWs); instead the SC phase preserves translational invariance.

Another natural symmetry of the EHM is the four-fold point group C_4 of $\pi/2$ rotations, expressing the square lattice property being mapped onto itself under said rotations. A similar property as for the translations holds for point groups,

$$C_n \supset C_{n/k} \quad k, n/k \in \mathbb{N}$$

which expresses the fact that, if a system is topologically invariant under n -fold rotations and $k \in \mathbb{N}$ exists such that $n/k \in \mathbb{N}$, then the smaller point group is automatically included. As for translations, we can break the point group reducing it to a smaller subgroup, say

$$C_4 \xrightarrow{\text{SSB}} C_2$$

This is what happens in general with the effective arising of anisotropic corrections to the kinetic hamiltonian due to non-local interactions, as will be discussed in Sec. ???. For our purposes, we will leave C_4 unbroken both in the AF and SC phases.

Finally, the system also is inversion-symmetric: let I_2 be the inversion group on the two-dimensional square lattice, containing trivially just the inversion operator and the identity. In general, rotations and mirroring are different operations from a group-theoretical point of view. However, when it comes to the square lattice, operatively it holds

$$I_2 \equiv C_2$$

To rotate by π is the same as to mirror the lattice. The inversion symmetry is then already included in C_4 , and we leave it unspecified.

2.2.2 Spin and charge symmetries

When it comes to Fermi algebra and operators, a large symmetry of the EHM is a global $U(2)$ symmetry [2] given by the transformation

$$\hat{c}_{i\sigma} \rightarrow \sum_{\sigma'} \mathcal{U}_{\sigma\sigma'} \hat{c}_{i\sigma'} \quad \text{with} \quad \mathcal{U}_{\sigma\sigma'} \in U(2) \quad (2.2)$$




Figure 2.1 | Graphic representation of the concept of $SU(2) \rightarrow U^z(1)$ SSB. The solid color identifies, in both cases, the rotations domain for the current symmetry. When SSB takes place in a magnetic phase, the magnetization order parameter acquires a precise direction which was previously invariant over the full spherical domain. After SSB, the hamiltonian is rotationally invariant only for rotations *around* the selected magnetization vector.

Phase	SSBs
AF	$T_1^{(\ell)} \rightarrow T_2^{(\ell)}$ $SU(2) \rightarrow U^z(1)$
SC	$U^c(1) \rightarrow 0$

Table 2.1 | SSB interpretation of the AF and SC phases. In the last line, the 0 on the SSB right-hand side indicates a completely disrupted symmetry.

The proof is trivial and follows from the fact that $\mathcal{U}^\dagger \mathcal{U} = \mathcal{U} \mathcal{U}^\dagger = \mathbb{I}$ [add proof?]. Note that the the following decomposition holds,

$$U(2) = SU(2) \otimes U^c(1) \quad (2.3)$$

which expresses the separate charge conservation and spin rotational symmetries. In detail:

- $U^c(1)$ charge conservation, coming from Eq. (2.3) decomposition. By breaking this symmetry, we allow for the total charge to fluctuate in the ground state. These fluctuations are only allowed in the SC phase in the present text;
- $SU(2)$ spin rotation symmetry, also , coming from Eq. (2.3) decomposition. Note that:

$$SU(2) \supset U^z(1)$$

The z subscript we included to distinguish from the charge group. Hence $SU(2)$ symmetry can be broken, selecting a particular vectorial direction for the order parameter (we call z), and reduced to a smaller $U^z(1)$ symmetry which is essentially expressed by the conservation of the magnitude of the order parameter.

The last point is important: for the AF phase, for example, there is no need for the magnetization vector to be directed along the physical z -axis, which is the one perpendicular to the lattice plane. $SU(2)$ symmetry breaks when (staggered) magnetization is established along a particular direction, but spin rotations around said direction still are ground state symmetries. This is shown schematically in Fig. 2.1.

2.2.3 Particle-hole symmetry at half-filling

The half-filled EHM is of particular interest due to its particle-hole symmetry (PHS), which is immediately disrupted at infinitesimal doping. Let us prove that. Consider the generic unitary

operation

$$P : \hat{c}_{i\sigma} \rightarrow e^{i\eta_{i\sigma}} \hat{h}_{i\sigma}^\dagger$$

being in general η_i a phase dependent on the fermionic quantum state ($i\sigma$). Inserting the above transformation in the EHM, we get

$$P\{\hat{H}\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma})} \hat{h}_{i\sigma} \hat{h}_{j\sigma}^\dagger + U \sum_i \hat{h}_{i\uparrow} \hat{h}_{i\downarrow} \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\uparrow}^\dagger - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger$$

For the two interactions $\hat{H}_U$, $\hat{H}_V$ the phase factors cancelled out. We aim to find the conditions under which the above hamiltonian is mapped onto the EHM of (1.2),

$$P\{\hat{H}\} \stackrel{?}{=} -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma} + U \sum_i \hat{h}_{i\uparrow}^\dagger \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}$$

Before starting the derivation, which is an extension of the argument of Arovas et al. [2] about the conventional HM, let us discuss some properties of the hole operators. Due to the presence of the phase factors in the PHS operative definition, the Fermi anticommutation rules hold regardlessly of the phase factor:

$$\begin{aligned} \{\hat{h}_{i\sigma}, \hat{h}_{j\sigma'}^\dagger\} &= e^{i(\eta_{i\sigma} - \eta_{j\sigma'})} \{\hat{c}_{i\sigma}^\dagger, \hat{c}_{j\sigma'}\} \\ &= e^{i(\eta_{i\sigma} - \eta_{j\sigma'})} \delta_{ij} \delta_{\sigma\sigma'} \\ &= \delta_{ij} \delta_{\sigma\sigma'} \end{aligned}$$

and evidently:

$$\hat{h}_{i\sigma} \hat{h}_{i\sigma}^\dagger = \hat{n}_{i\sigma}$$

being $\hat{n}_{i\sigma}$ the fermionic number operator on site i with spin σ . It follows, for quartic interactions

$$\begin{aligned} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger &= -\hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \\ &= \hat{h}_{i\sigma} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= -\hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= -\hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'} - \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \left(\hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \right) \delta_{ij} \delta_{\sigma\sigma'} \\ &= \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} \hat{h}_{i\sigma} + (\hat{n}_{i\sigma} + \hat{n}_{j\sigma'}) - 1 - \left(\hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \right) \delta_{ij} \delta_{\sigma\sigma'} \end{aligned}$$

The first term of the last passage is mirrored with respect to the starting term. Now, let us deal with the kinetic, local and non-local interactions separately.

Kinetic term. Consider the kinetic term,

$$P\{\hat{H}_t\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma})} \hat{h}_{i\sigma} \hat{h}_{j\sigma}^\dagger$$

It suffices to anticommute the two fermionic operators, and exchange site labels $i \leftrightarrow j$, to get

$$P\{\hat{H}_t\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma} + \pi)} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma}$$

A necessary condition on the unitary operation P is the following,

$$\eta_{i\sigma} + \eta_{j\sigma} = (2k - 1)\pi \quad \text{with} \quad k \in \mathbb{Z}$$

valid for NN sites.

Local repulsion. Substitute in $\hat{H}_U$ the above derivation,

$$\begin{aligned} P\{\hat{H}_U\} &= U \sum_i \hat{h}_{i\uparrow} \hat{h}_{i\downarrow} \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\uparrow}^\dagger \\ &= U \sum_i \hat{h}_{i\uparrow}^\dagger \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} + U \sum_i (\hat{n}_{i\uparrow} + \hat{n}_{i\downarrow}) - U \sum_i (1) \\ &= U \sum_i \hat{h}_{i\uparrow}^\dagger \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} + U(\hat{N} - L_x L_y) \end{aligned}$$

Thus, being for the half-filled state

$$\langle \hat{N} \rangle = L_x L_y$$

preserves PHS in the local sector.

Non-local attraction. Proceed identically for $\hat{H}_V$,

$$\begin{aligned} P\{\hat{H}_V\} &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger \\ &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} \hat{h}_{i\sigma} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} (\hat{n}_{i\sigma} + \hat{n}_{j\sigma'}) + V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} (1) \\ &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} \hat{h}_{i\sigma} - 2zV(\hat{N} - L_x L_y) \end{aligned}$$

with $z = 4$ the lattice coordination number. In the last passage, we used that on the square lattice

$$\begin{aligned} \sum_{\langle ij \rangle} (1) &= \sum_{i \in \mathcal{S}/2} \sum_{j \in \text{NN}(i)} (1) \\ &= \frac{L_x L_y}{2} \times z \end{aligned} \tag{2.4}$$

which will turn out quite often. As before, an half-filled phase preserves PHS in the non-local sector. In last passage we used the fact that summing over NNs means overcounting each site $z/2$ times, and spin degeneracy.

No other condition on phases is derived, thus we conclude the half-filled EHM preserves PHS if the two sublattices $\mathcal{S}_a, \mathcal{S}_b$ the square lattice can be divided into, being a (trivial) bipartite lattice, differ by a π phase difference under PHS transform. In other words, if P is defined by the unitary operation

$$P : \begin{cases} \hat{c}_{i\sigma} \rightarrow \hat{h}_{i\sigma}^\dagger & \text{for } i \in \mathcal{S}_a \\ \hat{c}_{i\sigma} \rightarrow -\hat{h}_{i\sigma}^\dagger & \text{for } i \in \mathcal{S}_b \end{cases}$$

and the system is half-filled, PHS is respected. Any other phase choice, as long as it alternates by π on the two sublattices, is good enough. To summarize, we report in Tab. 2.2 all identified symmetries.

2.3 MFT on Hubbard lattices

This section serves as groundwork for everything that follows. We here define the MFT scheme as well as the iterative HF computational scheme we use to estimate parametrically the relevant physical quantities of the various phases. We start by recalling the general features of MFT and then move to the specificity of the EHM, apply Wick's theorem and finally move to algorithmic aspects.

2.3.1 MFT general procedure, self-consistency and Wick's theorem

The Mean Field Theory approach to any fermionic quartic operator is given by the following procedure: given the hamiltonian made of the quadratic operators $\hat{A}$ and $\hat{B}$

$$\hat{H} \equiv \hat{A}\hat{B}$$

Symmetry	Operations
$T_1^{(x)} \otimes T_1^{(y)}$	One-site discrete translations separately in both directions
C_4	Four-fold point group (real $\pi/2$ angle rotations)
$U^c(1)$	Charge-conserving global phase shifts
$SU(2)$	Three-dimensional rotations in spin space
PHS	Particle-hole exchange (only at half filling)
$U^z(1)$	Reduced two-dimensional rotations in spin space

Table 2.2 | Model symmetries and relative group operations. Note that $U^c(1)$ represents the charge conserving group, while $U^z(1)$ represents the subgroup obtained by breaking the $SU(2)$ symmetry with a vector order parameter and preserving its amplitude. Since the latter is “derived” from $SU(2)$, we separate it from the four main symmetries of the model.

we define the fluctuations

$$\delta\hat{A} \equiv \hat{A} - \langle\hat{A}\rangle \quad \delta\hat{B} \equiv \hat{B} - \langle\hat{B}\rangle$$

thus the following chain holds:

$$\begin{aligned} \hat{A}\hat{B} &= \langle\hat{A}\rangle\langle\hat{B}\rangle + \delta\hat{A}\langle\hat{B}\rangle + \langle\hat{A}\rangle\delta\hat{B} + \delta\hat{A}\delta\hat{B} \\ &= -\langle\hat{A}\rangle\langle\hat{B}\rangle + \hat{A}\langle\hat{B}\rangle + \langle\hat{A}\rangle\hat{B} + \underbrace{\delta\hat{A}\delta\hat{B}}_{\mathcal{O}(\delta)^2} \end{aligned}$$

which implies

$$\langle\hat{H}\rangle = \langle\hat{A}\rangle\langle\hat{B}\rangle + \langle\delta\hat{A}\delta\hat{B}\rangle$$

Thus, if $|\langle\hat{A}\rangle\langle\hat{B}\rangle| \gg |\langle\delta\hat{A}\delta\hat{B}\rangle|$, in a statistical sense we can neglect the $\mathcal{O}(\delta)^2$ part of the hamiltonian and approximate

$$\hat{H} \simeq \hat{A}\langle\hat{B}\rangle + \langle\hat{A}\rangle\hat{B} - \langle\hat{A}\rangle\langle\hat{B}\rangle \implies \langle\hat{H}\rangle \simeq \langle\hat{A}\rangle\langle\hat{B}\rangle \quad (2.5)$$

Self consistency of MFT procedure is thus embodied by the condition

$$\left| \frac{\langle\delta\hat{A}\delta\hat{B}\rangle}{\langle\hat{A}\rangle\langle\hat{B}\rangle} \right| \ll 1 \implies \left| \frac{\langle\hat{A}\hat{B}\rangle}{\langle\hat{A}\rangle\langle\hat{B}\rangle} - 1 \right| \ll 1 \quad (2.6)$$

which essentially states that the least the fluctuations, the better the approximation. This rule of thumb is necessary in order to interpret the physical relevance of numerical MFT results. We will come back to this in the end, for now assuming the validity of MFT approximation and deriving the mathematical results.

2.3.2 MFT treatment of the pure Hubbard model

The pure Hubbard model, $\hat{H}_t + \hat{H}_U$, offers a simple framework for MFT analysis of the AF phase: this is described in detail in App. ???. The key passage is there given by the MFT approximation

$$\hat{n}_{i\uparrow}\hat{n}_{i\downarrow} \simeq \hat{n}_{i\uparrow}\langle\hat{n}_{i\downarrow}\rangle + \langle\hat{n}_{i\uparrow}\rangle\hat{n}_{i\downarrow} + (\text{constants}) \quad (2.7)$$

from which the AF structure is simply recovered. In terms of the discussion of the above paragraph, we have $\hat{A} \equiv \hat{n}_{i\uparrow}$ and $\hat{B} \equiv \hat{n}_{i\downarrow}$. However the “full” operator is given by

$$\hat{n}_{i\uparrow}\hat{n}_{i\downarrow} = \hat{c}_{i\uparrow}^\dagger \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \hat{c}_{i\uparrow}$$

whose terms we can contract in three different ways following Wick’s Theorem:

$$\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \hat{c}_{i\uparrow}\rangle \simeq \underbrace{\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger\rangle \langle\hat{c}_{i\downarrow} \hat{c}_{i\uparrow}\rangle}_{\text{Cooper}} - \underbrace{\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}\rangle \langle\hat{c}_{i\downarrow}^\dagger \hat{c}_{i\uparrow}\rangle}_{\text{Fock}} + \underbrace{\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow}\rangle \langle\hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow}\rangle}_{\text{Hartree}} \quad (2.8)$$

where the $-$ sign in the Fock term comes from fermionic anticommutation rules.

Term	Normal phase	Antiferromagnet	Superconductor
Hartree			
Fock	Prohibited	Prohibited	Prohibited
Cooper	Prohibited	Prohibited	

Table 2.3 | Summary of Wick's terms selection rules and relevant contributions to the effective hamiltonian for each phase limitedly to $\hat{H}_U$.




Figure 2.2 | Schematic representation of the four NNs of a given site i for a planar square lattice.

As a first approximation, the theorem is assumed to hold (which, in a BCS-like fashion, is equivalent to assuming for the ground-state to be a coherent state). The AF ground state reduces translational invariance and rotational symmetry, as explained in Tab. 2.1. Thus, of the three terms above, when considering the pure Hubbard model in AF phase:

- Cooper fluctuations are suppressed, because they break charge conservation;
- Similarly the Fock term is null as well because if $i = j$ and $\sigma' = \bar{\sigma}$ (as is for the local interaction, which contains the operator $\hat{n}_{i\uparrow}\hat{n}_{i\downarrow}$) the expectation values involved are describing a process breaking the survivor $U^z(1)$ spin symmetry.

Thus, correctly, the Wick's decomposition of pure HM only involves Hartree-terms of Eq. (2.8), getting us back to (2.7). The distinction between the normal phase and the AF phase is given by whether the Hartree fluctuations break or not translational symmetry: AF is indeed a specific SDW ordering which weakens the latter. In general the three parts of the decomposition (Hartree, Fock, Cooper) need to be considered altogether: this is what is done in the next chapters. What said here extends naturally to the EHM: the relevant contributions for each phase given by the local repulsion $\hat{U}$ in the EHM are listed in Tab. 2.3.

2.4 The non-local attraction $\hat{H}_V$ as a source of symmetry-breaking interactions

Let us now extend the discussion to the EHM and consider the NN non-local attraction,

$$\hat{H}_V \equiv -V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma\sigma'} \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}'\sigma'} \quad (2.9)$$

where we changed notation, and replaced the site index i with its 2D position on the lattice

$$i \rightarrow \mathbf{r} = (x, y)$$

Let also the lattice versors,

$$\underline{\mathbf{x}} \equiv (1, 0) \quad \underline{\mathbf{y}} \equiv (0, 1)$$

as in Fig. 2.2.

Evidently the hamiltonian can be decomposed in various spin terms,

$$\begin{aligned} \hat{H}_V &= \sum_{\sigma\sigma'} \hat{H}_V^{\sigma\sigma'} \\ &= \underbrace{\hat{H}_V^{\uparrow\uparrow} + \hat{H}_V^{\downarrow\downarrow}}_{\text{Same-spin}} + \underbrace{\hat{H}_V^{\uparrow\downarrow} + \hat{H}_V^{\downarrow\uparrow}}_{\text{Opposite-spin}} \end{aligned} \quad (2.10)$$

The role of said terms is crucial in establishing the pairing channel of the dominant physical phase. As an example, the s.s. terms can be seen as a possible source of triplet pairing superconductivity. Next section are devoted to analyze said terms and derive a simple analytical result in reciprocal space.

2.4.1 Real space description

Let us start from the real space form of Eq. (2.10). To carry out a summation over nearest neighbors $\langle \mathbf{r}\mathbf{r}' \rangle$ of a square lattice means precisely to sum over all links of the lattice. Then we can identify the generic opposite-spin (o.s.) term $\hat{H}_V^{\sigma\bar{\sigma}}$ as the one collecting the σ operators of sublattice $\mathcal{S}_a$ and $\bar{\sigma}$ operators of sublattice $\mathcal{S}_b$. The o.s. non-local interactions can be written as a sum of terms over just one of the two sublattices $\mathcal{S}_a$ and $\mathcal{S}_b$. Define the site hamiltonian

$$\hat{h}_V^{(\mathbf{r})} \equiv -V \sum_{\boldsymbol{\delta}} (\hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}+\boldsymbol{\delta}\downarrow} + \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}-\boldsymbol{\delta}\downarrow}) \quad \text{with} \quad \boldsymbol{\delta} \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\} \quad (2.11)$$

Then, it follows,

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} &= \overbrace{\sum_{\mathbf{r} \in \mathcal{S}_a} \hat{h}_V^{(\mathbf{r})}}^{\hat{H}_V^{\uparrow\downarrow}} + \overbrace{\sum_{\mathbf{r} \in \mathcal{S}_b} \hat{h}_V^{(\mathbf{r})}}^{\hat{H}_V^{\downarrow\uparrow}} \\ &= \sum_{\mathbf{r} \in \mathcal{S}} \hat{h}_V^{(\mathbf{r})} \end{aligned} \quad (2.12)$$

Here the notation of Fig. 1.1b is used. For each site $\mathbf{r}$ in a given sublattice, the nearest neighbors sites are four – all in the other sublattice. Similarly, the same-spin (s.s.) hamiltonian decomposes as

$$\hat{H}_V^{(\text{s.s.})} = -V \sum_{\mathbf{r} \in \mathcal{S}_a} \sum_{\boldsymbol{\delta}, \sigma} (\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\boldsymbol{\delta}\sigma} + \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}-\boldsymbol{\delta}\sigma}) \quad \text{with} \quad \boldsymbol{\delta} \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}$$

Opposite-spin terms. Consider first the o.s. terms of Eq. (2.11): take e.g. the NNs term $\hat{n}_{i\uparrow} \hat{n}_{\mathbf{r}+\boldsymbol{\delta}\downarrow}$. As in Eq. (2.8), taking Wick's contractions,

$$\begin{aligned} \langle \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}+\boldsymbol{\delta}\downarrow} \rangle &= \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow}^\dagger \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle \\ &\simeq \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Cooper}} - \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow} \rangle \langle \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow}^\dagger \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\downarrow} \rangle}_{\text{Hartree}} \end{aligned}$$

Identical decompositions are given for all others NNs. Of the three terms above:

- The Cooper term breaks $U^c(1)$ charge symmetry, allowing for superconducting instabilities;
- The Fock term breaks $U^z(1)$ symmetry, because it accounts for a site hop *plus* spin flip process;
- The Hartree term can reduce both translational invariance and spin rotational symmetry.

As a general rule of thumb, throughout this text we will discard the o.s. Fock term everywhere, preserving a robust $U^z(1)$ symmetry. For the normal state as well as for the AF instability only the Hartree term is to be accounted; for the SC phase also the Cooper term contributes. Note that for superconducting instabilities, due to superexchange mechanism (as explained in App. ??) the o.s. term accounts for singlet pairing as well as zero-projection triplet pairing. Which channel is preferred, is a matter of thermodynamic advantage.

Sector	Term	Normal phase	Antiferromagnet	Superconductor
o.s.	Hartree			
	Fock	Prohibited	Prohibited	Prohibited
	Cooper	Prohibited	Prohibited	
s.s.	Hartree			
	Fock			
	Cooper	Prohibited	Prohibited	

Table 2.4 Summary of Wick's terms selection rules and relevant contributions to the effective hamiltonian for each phase limitedly to $\hat{H}_V$, assuming preservation of $U^z(1)$ symmetry.

Same-spin terms. Consider then the same-spin terms: take e.g. the term $\hat{n}_{\mathbf{r}\uparrow}\hat{n}_{\mathbf{r}+\delta\uparrow}$. As above,

$$\begin{aligned}
\langle \hat{n}_{\mathbf{r}\uparrow}\hat{n}_{\mathbf{r}+\delta\uparrow} \rangle &= \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow} \hat{c}_{\mathbf{r}\uparrow} \rangle \\
&= \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}+\delta\uparrow} \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Cooper}} - \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow} \rangle \langle \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow} \rangle}_{\text{Hartree}}
\end{aligned}$$

Identical consideration as in the above paragraph hold for each term. The only difference with the o.s. terms is given by the Fock term: since the spin-flip process is absent, now the Fock fluctuations actually contribute as an effective NN hopping term. In other words, this term does not break $U^z(1)$ symmetry and thus is perfectly legitimate in each phase. Tab. 2.4 summarizes the selection rules hereby presented.

2.4.2 Fourier transform of the model

It is useful to derive analytically the reciprocal-space form of the three hamiltonian parts. By the sign $\stackrel{\mathcal{F}}{=}$, we indicate a passage where a Fourier-transform is performed. $\mathcal{F}$ -transform is obtained according to the convention Fourier transform it according to the convention

$$\hat{c}_{\mathbf{r}\sigma} \stackrel{\mathcal{F}}{=} \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{-i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma} \quad (2.13)$$

Kinetic part. The kinetic part is the easiest to transform. Apply the definition,

$$\begin{aligned}
-t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}'\sigma} + \text{h.c.}] &\stackrel{\mathcal{F}}{=} -t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma}^\dagger \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k}' \in \text{BZ}} e^{-i\mathbf{k}' \cdot \mathbf{r}'} \hat{c}_{\mathbf{k}'\sigma} + \text{h.c.} \\
&= -2t \sum_{\mathbf{k}, \mathbf{k}'} \sum_{\sigma} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma} \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{S}/2} e^{i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}} \left[e^{ik'_x} + e^{-ik'_x} + e^{ik'_y} + e^{-ik'_y} \right] \\
&= \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \epsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \quad (2.14)
\end{aligned}$$

where the bare bands were defined,

$$\epsilon_{\mathbf{k}} \equiv -2t (\cos k_x + \cos k_y) \quad (2.15)$$

Non-local attraction. Consider a generic bond, say, the one connecting NNs $\mathbf{r}$ and $\mathbf{r} \pm \delta$. Then:

$$\begin{aligned}
\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r} \pm \delta \sigma'} &= \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r} \pm \delta \sigma'}^\dagger \hat{c}_{\mathbf{r} \pm \delta \sigma'} \hat{c}_{\mathbf{r}\sigma} \\
&\stackrel{\mathcal{F}}{=} \frac{1}{(L_x L_y)^2} \sum_{\mathbf{k}_1, \dots, \mathbf{k}_4} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{r}} e^{\pm i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta} \hat{c}_{\mathbf{k}_1\sigma}^\dagger \hat{c}_{\mathbf{k}_2\sigma'}^\dagger \hat{c}_{\mathbf{k}_3\sigma'} \hat{c}_{\mathbf{k}_4\sigma}
\end{aligned}$$

Then, the interaction at site $\mathbf{r}$, spin σ with its NNs at spin σ' – indicated as $(\mathbf{r}\sigma\sigma')$ – is given by

$$\begin{aligned}
(\mathbf{r}\sigma\sigma') &= -V \sum_{\delta \in \{\underline{x}, \underline{y}\}} [\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\delta\sigma'} + \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}-\delta\sigma'}] \\
&\stackrel{\mathcal{F}}{=} -\frac{V}{(L_x L_y)^2} \sum_{\delta} \sum_{\mathbf{k}_1, \dots, \mathbf{k}_4} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} \\
&\quad \times \left(e^{i(\mathbf{k}_2-\mathbf{k}_3) \cdot \delta} + e^{-i(\mathbf{k}_2-\mathbf{k}_3) \cdot \delta} \right) \hat{c}_{\mathbf{k}_1\sigma}^\dagger \hat{c}_{\mathbf{k}_2\sigma'}^\dagger \hat{c}_{\mathbf{k}_3\sigma'} \hat{c}_{\mathbf{k}_4\sigma} \\
&= -\frac{2V}{(L_x L_y)^2} \sum_{\delta} \sum_{\mathbf{k}_1, \dots, \mathbf{k}_4} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} \cos[(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta] \hat{c}_{\mathbf{k}_1\sigma}^\dagger \hat{c}_{\mathbf{k}_2\sigma'}^\dagger \hat{c}_{\mathbf{k}_3\sigma'} \hat{c}_{\mathbf{k}_4\sigma}
\end{aligned}$$

The full non-local interaction is given by summing over all sites of $\mathcal{S}/2$ (which is, one sublattice of $\mathcal{S}$). This gives back momentum conservation,

$$\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{S}/2} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} = \frac{1}{2} \delta_{\mathbf{k}_1+\mathbf{k}_2=\mathbf{k}_3+\mathbf{k}_4}$$

Let $\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4 = \mathbf{K}$, and define $\mathbf{k}, \mathbf{k}'$ such that

$$\mathbf{k}_1 \equiv \mathbf{K} + \mathbf{k} \quad \mathbf{k}_2 \equiv \mathbf{K} - \mathbf{k} \quad \mathbf{k}_3 \equiv \mathbf{K} - \mathbf{k}' \quad \mathbf{k}_4 \equiv \mathbf{K} + \mathbf{k}' \quad \delta \mathbf{k} \equiv \mathbf{k} - \mathbf{k}'$$

Sums over these variables must be intended as over the Brillouin Zone (BZ). Hence:

$$\begin{aligned}
\hat{H}_V &= \sum_{\mathbf{r} \in \mathcal{S}_a} \sum_{\sigma\sigma'} (\mathbf{r}\sigma\sigma') \\
&\stackrel{\mathcal{F}}{=} -\frac{V}{L_x L_y} \sum_{\sigma\sigma'} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \cos(\delta \mathbf{k} \cdot \delta) \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma'} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \\
&= -\frac{V}{L_x L_y} \sum_{\sigma\sigma'} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma'} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \quad (2.16)
\end{aligned}$$

which decomposes in s.s. and o.s. channels as:

$$\begin{aligned}
\hat{H}_V &\stackrel{\mathcal{F}}{=} \overbrace{-\frac{V}{L_x L_y} \sum_{\sigma} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{s.s.}} \\
&\quad \underbrace{-\frac{V}{L_x L_y} \sum_{\sigma} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\bar{\sigma}}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\bar{\sigma}} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma}}_{\text{o.s.}}
\end{aligned}$$

The o.s. part can be simplified: since

$$\hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\bar{\sigma}}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\bar{\sigma}} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} = \hat{c}_{\mathbf{K}-\mathbf{k}\bar{\sigma}}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \hat{c}_{\mathbf{K}-\mathbf{k}'\bar{\sigma}}$$

due to fermionic anti-commutation rules, and since the composite transform

$$\mathbf{k} \rightarrow -\mathbf{k} \quad \mathbf{k}' \rightarrow -\mathbf{k}'$$

leaves the form factor $\cos(\delta k_x) + \cos(\delta k_y)$ invariant, we reduce the spin sum to two times the $\sigma = \uparrow$ part,

$$\hat{H}_V^{(\text{o.s.})} = -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \quad (2.17)$$

whiel the explicit s.s. sector is given immediately by:

$$\begin{aligned}
\hat{H}_V^{(\text{s.s.})} &= -\frac{V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \\
&\quad \times \left[\hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\uparrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} + \hat{c}_{\mathbf{K}+\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\downarrow} \right] \quad (2.18)
\end{aligned}$$

The result here obtained will be used various times in next chapters. Note that different Wick contraction schemes will be referred to as:

$$\overbrace{c_{\mathbf{K}+\mathbf{k}\sigma}^\dagger c_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger c_{\mathbf{K}-\mathbf{k}'\sigma'} c_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{Cooper contraction}} \quad (2.19)$$

$$\overbrace{c_{\mathbf{K}+\mathbf{k}\sigma}^\dagger c_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger c_{\mathbf{K}-\mathbf{k}'\sigma'} c_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{Fock contraction}} \quad (2.20)$$

$$\overbrace{c_{\mathbf{K}+\mathbf{k}\sigma}^\dagger c_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger c_{\mathbf{K}-\mathbf{k}'\sigma'} c_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{Hartree contraction}} \quad (2.21)$$

When contracting in the Fock channel, a minus sign must be included.

Local repulsion. A somewhat identical argument can be carried out for the local interaction,

$$\hat{H}_U = U \sum_{\mathbf{r} \in \mathcal{S}} \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}\downarrow}$$

The only difference here is made by the fact that the interaction is repulsive on-site, thus the final structure factor in the Fourier transform disappears as well as the minus sign,

$$\hat{H}_U \stackrel{\mathcal{F}}{=} \frac{U}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \quad (2.22)$$

When contracting these fermionic operators, identical considerations as for the non-local counterpart (2.16) hold. Note the similarity of this Fourier transform with the o.s. sector of the non-local attraction, given in Eq. (2.16). With this, we conclude the general introduction to the model and pass to the setup of the Hartree-Fock (HF) algorithm.

2.5 Iterative computational approach to MFT

This section is devoted to the explanation of the general iterative strategy we use to simulate the EHM through iterative Mean-Field Theory (iMFT). The simplicity of the algorithm is only possible thanks to the violent reduction of complexity MFT is able to reach. We use an iterative self-consistent scheme, thus solving a finite set of equation searching for a fixed point in parameters space for each phase.

2.5.1 Sketch of the Hartree-Fock iterative algorithm

The general sketch of a self-consistent Hartree-Fock algorithm in our context is simple enough. Start from a fermionic quartic hamiltonian,

$$\hat{H} = \sum_{\alpha\beta} A_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta + \sum_{\alpha\beta\gamma\delta} B_{\alpha\beta\gamma\delta} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta + \text{h.c.} \quad (2.23)$$

where $A_{\alpha\beta}$, $B_{\alpha\beta\gamma\delta}$ are model parameters. We limit ourselves to quartic hamiltonians, but the idea can be extended. As discussed before, in this kind of context MFT is employed by applying Wick's theorem. This way, the quartic hamiltonian is effectively reduced to a quadratic hamiltonian,

$$\hat{H} \simeq \hat{H}^{(\text{eff})} = \sum_{\alpha\beta} C_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta + \sum_{\alpha\beta} D_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger + \text{h.c.} \quad (2.24)$$

where now the new parameters, which we will refer to as Hartree-Fock parameters (HFP), depend on the model itself via the equations

$$C_{\alpha\beta} = A_{\alpha\beta} + \sum_{\gamma\delta} [B_{\alpha\gamma\delta\beta} - B_{\alpha\gamma\beta\delta}] \langle \hat{c}_\gamma^\dagger \hat{c}_\delta \rangle$$

$$D_{\alpha\beta} = \sum_{\gamma\delta} B_{\alpha\beta\gamma\delta} \langle \hat{c}_\gamma \hat{c}_\delta \rangle$$

These equations are determined by many-body expectation values, which in general are computed by the means of thermal averages. A self consistent Hartree-Fock scheme goes as follows:




Figure 2.3 | Schematization of the self-consistent Hartree Fock method described in Sec. 2.5. The effective model is solved, the solution is used to estimate self-consistently the Hartree Fock vector, which is then used to update iteratively the model itself. HFP_i indicates the initializer HFPs vector, HFP_c the converged HFPs vector.

1. Apply MFT to the quartic hamiltonian, recovering its general theoretic effective form;
2. Impose selection rules on the parameters $C_{\alpha\beta}$, $D_{\alpha\beta}$, eventually eliminating those not respecting the symmetry sector we are working in. As an example: if we choose not to break charge conservation, set $\forall \alpha, \beta : D_{\alpha\beta} = 0$;
3. Choose a starting value and a coherent symmetry structure for the remaining parameters;
4. Enter the iterative scheme of Fig. 2.3:
 - a) Based on the current HFPs, determine the value $\mu[C, D]$ giving the correct density;
 - b) Diagonalize the effective hamiltonian $\hat{H}^{(\text{eff})}[C, D] - \mu[C, D]\hat{N}$ and recover the ground-state $|\Psi[C, D]\rangle$;
 - c) Compute the new HFPs $C'_{\alpha\beta}$, $D'_{\alpha\beta}$ via the self consistent equations;
 - d) Initialize the new effective hamiltonian $\hat{H}^{(\text{eff})}[C', D']$ and repeat the cycle until the HFPs have converged to stable values.

The stable values we are referring to in the last point are our solutions. The resulting vector is a fixed point in the parametric recursive map defined by the set of MFT self-consistent equations.

Of course, to work with any of these algorithms we need to know how to diagonalize the hamiltonian. In the context of the EHM at thermal equilibrium, being the phases we are simulating simple enough, we know analytically the diagonalized hamiltonian, and making use of this knowledge we are able to reduce the self consistent scheme to a simple iterative algorithm. An explicit example of HF algorithm used in present text, somewhat simplified due to the presence of a single HFP, is given in App. ??.

2.5.2 Self-consistency and RMPs

As will be thoroughly discussed in the following derivations for each physical phase we treat, treating the EHM via MFT keeping under strict control the effects of Wick's theorem produces ground states whose parametric properties (band structures, gap functions, chemical potential) are controlled self-consistently by the model two-body interactions $\hat{H}_U$ and $\hat{H}_V$. Let us define the concept of Renormalized model parameter (RMP), indicating any of the model parameters that is effectively renormalized via interactions.

Let us propose a relevant example of RMP. Suppose that under MFT, for a given phase, the decomposition $\hat{H}_V$ has a net effect of renormalizing the hopping amplitude

$$t \xrightarrow{\hat{H}_V} \tilde{t}$$

In other words, the decomposition of $\hat{H}_V$ produces a free Fermionic hamiltonian whose diffusive parameter is $\tilde{t}$. We define $\tilde{t}$ as an RMP, being the result of the original model parameter t plus the effect of interactions.

2.5.3 Fixed density simulations and the role of μ

In the context of the present text, all simulations are performed at fixed density, determining self-consistently the value of the chemical potential. This means that, for a given step i and the relative HFPs vector $\mathbf{v}_i$, and for a fixed density $n \in [0, 1]$, μ_i (the chemical potential at current step) is determined as

$$\mu_i : \min_{\mu} \left\{ \frac{1}{2L_x L_y} \sum_{\mathbf{k}} \sum_{\sigma} \langle \hat{n}_{\mathbf{k}\sigma} \rangle \Big|_{\mathbf{v}=\mathbf{v}_i} - n \right\} \quad (2.25)$$

being $\mathbf{v} = \mathbf{v}_i$ the HFPs vector at current step. Averaging is performed over the ground state determined parametrically by the current value of the HFPs.

Hence, if we are able to perform averages and know both $\mathbf{v}_i$ and n , at step i we infer the correct chemical potential. Now, as we will discuss $\tilde{\mu}$ will turn out to be an RMP itself, meaning that theoretically

$$\mu \rightarrow \mu + F[\mathbf{v}] \equiv \tilde{\mu}[\mathbf{v}]$$

being F some functional of the HFPs. In other words, a common effect of interactions under MFT will be precisely to shift the chemical potential. At step i , keeping the density fixed, the chemical potential we are determining numerically and using in our equations is then $\tilde{\mu}_i$. In simpler words, the designed algorithm “does not know” about chemical potential renormalization and thus just determines the number that solves Eq. (2.25). But we know that this number is made of two contributions: the “physical” μ_i and the effective shift produced by the model itself, $F[\mathbf{v}_i]$. So, when it comes to μ_i , this consideration becomes important.

It is the case of the ground state energy estimation. Let us abstract from the EHM and MFT, and consider a generic hamiltonian $\hat{H}$. To find the ground state $|\Psi\rangle$ means precisely to solve

$$\frac{\delta}{\delta \langle \Psi |} \langle \Psi | \hat{H} | \Psi \rangle = 0$$

Now, working with a fixed particles number N means we are performing a constrained minimization (which is, taking the functional derivative) on the composite operator

$$\hat{H} - \mu(\hat{N} - N) \equiv \hat{K} + \mu N$$

being μ the Lagrange multiplier, $\hat{N}$ the number operator and $\hat{K} \equiv \hat{H} - \mu\hat{N}$. Finding the constrained ground state parametrized by a vector $\mathbf{v}$ now means to solve

$$\mathbf{v} \Rightarrow |\Psi[\mathbf{v}]\rangle : \begin{cases} \frac{\delta}{\delta \langle \Psi |} \langle \Psi | \hat{K} + \mu N | \Psi \rangle \Big|_{|\Psi\rangle=|\Psi[\mathbf{v}]\rangle} = 0 \\ \langle \Psi[\mathbf{v}] | \hat{N} | \Psi[\mathbf{v}] \rangle = N \end{cases}$$

In the first line μN is derived to zero. The constraint is imposed in the second line. This means that, when it comes to determining the parametric ground state, $\hat{K}$ is functionally minimized. In this context μ (the Lagrange multiplier) sums with other effective terms arisen from algebraic manipulations of $\hat{H}$ giving a grand total of $\tilde{\mu}$. The number operator $\hat{N}$ average is performed over the many-body ground state $|\Psi[\mathbf{v}]\rangle$ with the total effective chemical potential being $\tilde{\mu}$, not μ . However, when it comes to computing energy,

$$\begin{aligned} \langle \hat{H} \rangle &= \langle \Psi[\mathbf{v}] | \hat{K} | \Psi[\mathbf{v}] \rangle + \mu N \\ &= \Omega[\mathbf{v}] + (\tilde{\mu}[\mathbf{v}] - F[\mathbf{v}])N \end{aligned}$$

being Ω the expectation value (EV) of $\hat{K}$. In the last passage we wrote everything in terms of functionals of the HFPs vector $\mathbf{v}$. Using this relation, when the algorithm has converged, we can estimate the physical energy.

2.5.4 The reject-mix-iterate convergence scheme

Up to now, we did not specify when we consider the algorithm to “have converged”. From a purely theoretical point of view, considering the set of self-consistency equations provided by MFT as a nonlinear map in HFPs space, we consider the run finished when our algorithm, following a non-trivial path in HFPs space, comes close enough to a fixed point of the map; which is, a point which gets mapped onto itself. Explicitly, we consider the run over and converged whenever the following logic condition is satisfied for each component of $\mathbf{v}$:

$$\forall v_j \quad : \quad |v_j - A_j(\mathbf{v})| \leq \Delta v_j$$

being $A(\mathbf{v})$ the algorithmic image of the HFP vector $\mathbf{v}$, and A_j its j -th component. The specification of Δv_j is clearly arbitrary. We are dealing in general with quantities of order 10^{-1} , so convergence values of order $10^{-3} \div 10^{-4}$ will be generally sufficient.

When the acceptance condition is not satisfied, which is, when at least one component of $\mathbf{v}$ is mapped by the HF algorithm to a new image value for a total displacement larger than the respective component of $\Delta \mathbf{v}$, the step is rejected and iteration continues. This lets us define the mixing parameter $g \in [0, 1]$, which represents the deterministic fractional combination of the current step with its image value to obtain a new step,

$$\mathbf{v}_{i+1} \equiv gA(\mathbf{v}_i) + (1 - g)\mathbf{v}_i \quad (2.26)$$

and becomes a simple but useful knob to tune algorithmic evolution. Its usefulness comes from the core concept that, even if physically there must exist a unique solution to the model (thus just one self-consistent HFP vector, eventually of all zeros) such a fixed point could be hard to reach following a pure iterative numeric scheme on a given map (which would be the $g = 1$ case of the above equation) for a variety of reasons. By moving “slower” (low g) with an accurate optimization of the mixing, as will be seen concretely in Sec. ??, it is possible to reach convergence in a controlled way.

When it comes to HF iterative schemes of this kind, it can happen with some frequency that the algorithm remains stuck in an infinite loop between a set of points mathematically linked by the self-consistent map but not representing a physical solution (because of their reciprocal absolute distance in HFP space). In some other situations, evolutions could turn out to be rather slow. Because of this it is reasonable to define a parameter p , such that the number of iterations is kept $i \leq p$. If the algorithm failed to converge in p steps, the point is saved as “NaN” (Not a number) and the algorithm moves to another simulation. Introducing this simple rule, runtime is kept under control and we gain an easy method to identify the regions of parameter space requiring more care – just read from the raw data where NaNs have been saved.

2.6 Square lattice spatial symmetry channels

In next sections we introduce spatial symmetry structures and discuss their relation with point groups. We will always refer to various spatial structures as “symmetry channels”, throughout the system is able to establish a given order parameter defined in space.

Looking at $\hat{H}$, the main difference between its two-body interactions $\hat{H}_U$ and $\hat{H}_V$ of course relies in locality (first) and non-locality (second). Both operators in principle are able to break C_4 symmetry, as will later be discussed extensively.

For the 2D square lattice, various spatial symmetries are possible. Let i be a specific site, and consider a generic complex function g_{ij} whose domain is given by the site itself and its four NNs (the domain concides with the points of Fig. 2.2). The function can be written in the C_4 basis

$$g_{ij} = \sum_{\gamma} g^{(\gamma)} \varphi_{ij}^{(\gamma)}$$

where $g^{(\gamma)} \in \mathbb{C}$ are the g_{ij} symmetry channels coefficients, while $\varphi_{ij}^{(\gamma)}$ are the form factors listed in Tab. 2.5 and plotted in Fig. 2.5, a simple orthonormal rearrangement of the harmonics basis.




Figure 2.4 | Form factors at different topologies, as listed in Tab. 2.5. In figures five sites are represented: the hub site i and its four NN. Solid lines represent non-zero values for φ_{δ} , while dashed lines represent vanishing factors.

Structure	Form factor	Graph
s-wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(s)} = \delta_{\mathbf{r}\mathbf{r}'}$	Fig. 2.4a
Extended s-wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(s^*)} = \frac{1}{2} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 2.4b
p_x -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(p_x)} = \frac{1}{\sqrt{2}} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}}]$	Fig. 2.4c
p_y -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(p_y)} = \frac{1}{\sqrt{2}} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 2.4d
$d_{x^2-y^2}$ -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(d)} = \frac{1}{2} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}} - \delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 2.4e

Table 2.5 | Form factors for a function defined on a square lattice, with a prefactor included for normalization. We use lattice vector notation here. The last column indicates the graph representation of each contribution given in Fig. 2.4. Subscript $x^2 - y^2$ is omitted for notational clarity.

When dealing with phases which respect translational symmetry, or at most reduce it to a weaker symmetry by shrinking the Brillouin Zone, it's useful to work in reciprocal space, as discussed in Sec. 2.4. As said, a particular phase can show specific spatial symmetries, and so will the Hartree-Fock parameters. The key point to notice is that self-consistency equations involve sums (or integrals) to be computed over fermionic states, which can be heavily simplified by employing said symmetries. Consider the form factors of Tab. 2.5, and their Fourier transforms of Tab. 2.6. They satisfy the relations:

$$\frac{1}{L_x L_y} \sum_{ij} \varphi_{ij}^{(\gamma)} \varphi_{ij}^{(\gamma)*} = \delta_{\gamma\gamma'} \quad \frac{1}{L_x L_y} \sum_{\mathbf{k}} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} = \delta_{\gamma\gamma'}$$

If these symmetries are developed by the HFPs, it's of no use to integrate over the entire BZ, since

Structure	Structure factor	Graph
s -wave	$\varphi_{\mathbf{k}}^{(s)} = 1$	Fig. 2.4a
Extended s -wave	$\varphi_{\mathbf{k}}^{(s^*)} = \cos k_x + \cos k_y$	Fig. 2.4b
p_x -wave	$\varphi_{\mathbf{k}}^{(p_x)} = i\sqrt{2} \sin k_x$	Fig. 2.4c
p_y -wave	$\varphi_{\mathbf{k}}^{(p_y)} = i\sqrt{2} \sin k_y$	Fig. 2.4d
$d_{x^2-y^2}$ -wave	$\varphi_{\mathbf{k}}^{(d)} = \cos k_x - \cos k_y$	Fig. 2.4e

Table 2.6 | Structure factors derived from the correlation structures of Tab. 2.5. The functions hereby defined are orthonormal, and are plotted in Fig- 2.5.




Figure 2.5 | Representation of the various structure factors listed in Tab. 2.6 on the BZ. For the s^* and $d_{x^2-y^2}$ -wave factors, the real part is plotted; for the p_ℓ factors, the imaginary part is plotted.

different regions linked by periodicity of the form factors will lead to redundant calculations.

Finally, a very useful detail is given by the following argument. Consider the equation

$$\cos(\delta k_x) + \cos(\delta k_y) = \cos k_x \cos k'_x + \sin k_x \sin k'_x + \cos k_y \cos k'_y + \sin k_y \sin k'_y$$

For the sake of readability, the notations

$$c_\ell \equiv \cos k_\ell \quad s_\ell \equiv \sin k_\ell \quad c'_\ell \equiv \cos k'_\ell \quad s'_\ell \equiv \sin k'_\ell$$

are used. Group the four terms above,

$$\underbrace{(c_x c'_x + c_y c'_y)}_{\text{Symmetric}} + \underbrace{(s_x s'_x + s_y s'_y)}_{\text{Anti-symmetric}} \quad (2.27)$$

The first two exhibit inversion symmetry for both arguments $\mathbf{k}, \mathbf{k}'$; the second two exhibit anti-symmetry. Decoupling the symmetric part, we obtain

$$c_x c'_x + c_y c'_y = \frac{1}{2}(c_x + c_y)(c'_x + c'_y) + \frac{1}{2}(c_x - c_y)(c'_x - c'_y)$$

which finally gives:

$$\begin{aligned} \cos(\delta k_x) + \cos(\delta k_y) &= \frac{1}{2}(c_x + c_y)(c'_x + c'_y) && (s^* \text{-wave}) \\ &+ s_x s'_x && (p_x \text{-wave}) \\ &+ s_y s'_y && (p_y \text{-wave}) \\ &+ \frac{1}{2}(c_x - c_y)(c'_x - c'_y) && (d_{x^2-y^2} \text{-wave}) \end{aligned}$$

or, in a more compact form

$$\cos(\delta k_x) + \cos(\delta k_y) = \frac{1}{2} \sum_{\gamma} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} \quad \text{for } \gamma \in \{s^*, p_x, p_y, d_{x^2-y^2}\} \quad (2.28)$$

$$= \frac{1}{2} \sum_{\gamma} \varphi_{\mathbf{k}}^{(\gamma)*} \varphi_{\mathbf{k}'}^{(\gamma)} \quad (2.29)$$

This decomposition will become particularly handy in later calculations.

Bibliography

- [1] James F. Annett. “Symmetry of the order parameter for high-temperature superconductivity”. In: *Advances in Physics* 39.2 (Apr. 1990). Publisher: Taylor & Francis .eprint: <https://doi.org/10.1080/00018739000101481>, pp. 83–126. ISSN: 0001-8732. DOI: [10.1080/00018739000101481](https://doi.org/10.1080/00018739000101481). URL: <https://doi.org/10.1080/00018739000101481> (visited on 01/06/2026).
- [2] Daniel P. Arovas et al. “The Hubbard Model”. In: *Annual Review of Condensed Matter Physics* 13. Volume 13, 2022 (2022), pp. 239–274. ISSN: 1947-5462. DOI: <https://doi.org/10.1146/annurev-conmatphys-031620-102024>. URL: <https://www.annualreviews.org/content/journals/10.1146/annurev-conmatphys-031620-102024>.
- [3] Anna E. Böhrer et al. “Nematicity and nematic fluctuations in iron-based superconductors”. en. In: *Nature Physics* 18.12 (Dec. 2022), pp. 1412–1419. ISSN: 1745-2481. DOI: [10.1038/s41567-022-01833-3](https://doi.org/10.1038/s41567-022-01833-3). URL: <https://www.nature.com/articles/s41567-022-01833-3>.
- [4] Zhangkai Cao et al. “Dominant p -wave pairing induced by nearest-neighbor attraction in the square-lattice extended Hubbard model”. In: *Phys. Rev. B* 111 (2 Jan. 2025), p. 024509. DOI: [10.1103/PhysRevB.111.024509](https://doi.org/10.1103/PhysRevB.111.024509). URL: <https://link.aps.org/doi/10.1103/PhysRevB.111.024509>.
- [5] Piers Coleman. *Introduction to Many-Body Physics*. Cambridge University Press, 2015.
- [6] Michele Fabrizio. *A Course in Quantum Many-Body Theory*. Springer, 2022.
- [7] Gabriele Giuliani and Giovanni Vignale. *Quantum Theory of the Electron Liquid*. Cambridge University Press, 2005.
- [8] Giuseppe Grosso and Giuseppe Pastori Parravicini. *Solid State Physics*. Second Edition. Academic Press, 2014.
- [9] J. E. Hirsch. “Two-dimensional Hubbard model: Numerical simulation study”. In: *Phys. Rev. B* 31 (7 Apr. 1985), pp. 4403–4419. DOI: [10.1103/PhysRevB.31.4403](https://doi.org/10.1103/PhysRevB.31.4403). URL: <https://link.aps.org/doi/10.1103/PhysRevB.31.4403>.
- [10] Joel Hutchinson and Frank Marsiglio. “Mixed temperature-dependent order parameters in the extended Hubbard model”. en. In: *Journal of Physics: Condensed Matter* 33.6 (Feb. 2021), p. 065603. ISSN: 0953-8984, 1361-648X. DOI: [10.1088/1361-648X/abc801](https://doi.org/10.1088/1361-648X/abc801). URL: <https://iopscience.iop.org/article/10.1088/1361-648X/abc801> (visited on 01/06/2026).
- [11] Masaru Kato et al. “Soliton Lattice Modulation of Incommensurate Spin Density Wave in Two Dimensional Hubbard Model -A Mean Field Study-”. In: *Journal of the Physical Society of Japan* 59.3 (Mar. 1990), pp. 1047–1058. ISSN: 0031-9015. DOI: [10.1143/JPSJ.59.1047](https://doi.org/10.1143/JPSJ.59.1047). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.59.1047> (visited on 03/21/2026).
- [12] R. Micnas, J. Ranninger, and S. Robaszkiewicz. “Superconductivity in narrow-band systems with local nonretarded attractive interactions”. In: *Reviews of Modern Physics* 62.1 (Jan. 1990). Publisher: American Physical Society, pp. 113–171. DOI: [10.1103/RevModPhys.62.113](https://doi.org/10.1103/RevModPhys.62.113). URL: <https://link.aps.org/doi/10.1103/RevModPhys.62.113> (visited on 01/06/2026).

- [13] Robin Scholle et al. “Comprehensive mean-field analysis of magnetic and charge orders in the two-dimensional Hubbard model”. In: *Phys. Rev. B* 108 (3 July 2023), p. 035139. DOI: [10.1103/PhysRevB.108.035139](https://doi.org/10.1103/PhysRevB.108.035139). URL: <https://link.aps.org/doi/10.1103/PhysRevB.108.035139>.
- [14] P. Senarath Yapa et al. “Mixed-symmetry superconductivity and the energy gap”. en. In: *Physica C: Superconductivity and its Applications* 633 (June 2025), p. 1354719. ISSN: 09214534. DOI: [10.1016/j.physc.2025.1354719](https://doi.org/10.1016/j.physc.2025.1354719). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0921453425000723> (visited on 01/06/2026).
- [15] Manfred Sgrist and Kazuo Ueda. “Phenomenological theory of unconventional superconductivity”. In: *Reviews of Modern Physics* 63.2 (Apr. 1991). Publisher: American Physical Society, pp. 239–311. DOI: [10.1103/RevModPhys.63.239](https://doi.org/10.1103/RevModPhys.63.239). URL: <https://link.aps.org/doi/10.1103/RevModPhys.63.239> (visited on 01/07/2026).
- [16] Simons Collaboration on the Many-Electron Problem et al. “Absence of Superconductivity in the Pure Two-Dimensional Hubbard Model”. In: *Physical Review X* 10.3 (July 2020), p. 031016. DOI: [10.1103/PhysRevX.10.031016](https://doi.org/10.1103/PhysRevX.10.031016). URL: <https://link.aps.org/doi/10.1103/PhysRevX.10.031016> (visited on 03/21/2026).
- [17] Avinash Singh and Zlatko Tešanović. “Collective excitations in a doped antiferromagnet”. en. In: *Physical Review B* 41.1 (Jan. 1990), pp. 614–631. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.41.614](https://doi.org/10.1103/PhysRevB.41.614). URL: <https://link.aps.org/doi/10.1103/PhysRevB.41.614> (visited on 12/12/2025).
- [18] Ashvin Vishwanath. “Dirac Nodes and Quantized Thermal Hall Effect in the Mixed State of d-wave Superconductors”. In: *Physical Review B* 66.6 (Aug. 2002). arXiv:cond-mat/0104213, p. 064504. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.66.064504](https://doi.org/10.1103/PhysRevB.66.064504). URL: <http://arxiv.org/abs/cond-mat/0104213> (visited on 01/05/2026).
- [19] Alexander Wietek et al. “Stripes, Antiferromagnetism, and the Pseudogap in the Doped Hubbard Model at Finite Temperature”. en. In: *Physical Review X* 11.3 (July 2021), p. 031007. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.11.031007](https://doi.org/10.1103/PhysRevX.11.031007). URL: <https://link.aps.org/doi/10.1103/PhysRevX.11.031007> (visited on 12/12/2025).
- [20] Jan Zaanen and Olle Gunnarsson. “Charged magnetic domain lines and the magnetism of high- T_c oxides”. In: *Physical Review B* 40.10 (Oct. 1989), pp. 7391–7394. DOI: [10.1103/PhysRevB.40.7391](https://doi.org/10.1103/PhysRevB.40.7391). URL: <https://link.aps.org/doi/10.1103/PhysRevB.40.7391> (visited on 03/21/2026).
- [21] C. Zhou and H. J. Schulz. “Density of states and tunneling spectra in two-dimensional d-wave superconductors”. In: *Physical Review B* 45.13 (Apr. 1992). Publisher: American Physical Society, pp. 7397–7401. DOI: [10.1103/PhysRevB.45.7397](https://doi.org/10.1103/PhysRevB.45.7397). URL: <https://link.aps.org/doi/10.1103/PhysRevB.45.7397> (visited on 01/06/2026).