

# Numerical analysis of superconducting phases in the extended Hubbard model with non-local pairing

University of Pisa, a.y. 2025-2026

Alessandro Gori\*

Thesis for the Master's degree in Physics

## Abstract

[To be continued...]

- Appendix on  $s$ -wave SC ghost simulations on HM
- Set ComputerModern font in all CairoMakie figures

## Contents

|          |                                                                    |          |
|----------|--------------------------------------------------------------------|----------|
| <b>1</b> | <b>Superconducting instability</b>                                 | <b>3</b> |
| 1.1      | The SC phase and its symmetries                                    | 3        |
| 1.2      | Local effects: (impossibility of) superconductivity in the HM      | 4        |
| 1.2.1    | Local Hartree channel in the SC phase: analytic derivation         | 5        |
| 1.2.2    | Local Hartree contraction energy                                   | 5        |
| 1.2.3    | Local Cooper channel in the SC phase: analytic derivation          | 5        |
| 1.2.4    | Local Cooper contraction energy                                    | 6        |
| 1.2.5    | Superconductivity in the HM                                        | 6        |
| 1.2.6    | Features of the SC solution                                        | 9        |
| 1.3      | Extension to the EHM                                               | 10       |
| 1.4      | Non-local Hartree effects: chemical potential renormalisation      | 12       |
| 1.4.1    | Non-local Hartree channel in the SC phase: analytic derivation     | 12       |
| 1.4.2    | Non-local Hartree contraction energy                               | 12       |
| 1.5      | Fock effects: bare bands renormalization                           | 12       |
| 1.5.1    | Fock channel in the SC phase: analytic derivation                  | 12       |
| 1.5.2    | Fock contraction energy                                            | 14       |
| 1.6      | Non-local Cooper effects: anisotropic pairing                      | 14       |
| 1.6.1    | Non-local Cooper o.s. channel in the SC phase: analytic derivation | 15       |
| 1.7      | Full MFT hamiltonian and anisotropic pairing                       | 16       |
| 1.8      | The zero-temperature ground-state at half-filling                  | 17       |
| 1.9      | SC free energy density                                             | 17       |
| 1.9.1    | The gap energy                                                     | 19       |
| 1.10     | Theoretical constraints on the singlet SC phase HFPs               | 19       |
| 1.11     | Discussion of numeric results (singlet SC)                         | 19       |
| 1.11.1   | Multi-symmetry singlet superconductivity                           | 20       |
| 1.11.2   | Rotational SSB and nematic superconductivity                       | 20       |
| 1.12     | Theoretical constraints on the triplet SC phase HFPs               | 20       |
| 1.13     | Discussion of numeric results (triplet SC)                         | 22       |

---

\*a.gori23@studenti.unipi.it / nepero27178@github.com

Bibliography

23

---

### List of symbols and abbreviations

---

|                     |                                              |
|---------------------|----------------------------------------------|
| AF                  | Antiferromagnetic                            |
| BCS                 | Bardeen-Cooper-Schrieffer (theory)           |
| BZ                  | Brillouin zone                               |
| $\beta$             | Inverse temperature                          |
| $\Delta^{(\gamma)}$ | $\gamma$ -wave component of the gap function |
| DoF                 | Degree of freedom                            |
| EHM                 | Extended Hubbard model                       |
| EV(s)               | Expectation value(s)                         |
| FS                  | Fermi surface                                |
| HF                  | Hartree-Fock                                 |
| HFP(s)              | Hartree-Fock parameter(s)                    |
| HM                  | Hubbard model                                |
| $L_\ell$            | Number of lattice sites in direction $\ell$  |
| iHF                 | Iterative Hartree-Fock                       |
| LRT                 | Linear Response Theory                       |
| MBZ                 | Magnetic Brillouin zone                      |
| MFT                 | Mean-Field Theory                            |
| NBZ                 | Nematic Brillouin zone                       |
| NN(s)               | Nearest neighbor(s)                          |
| PH                  | Particle-hole                                |
| PHS                 | Particle-hole symmetry                       |
| RWC(s)              | Relevant Wick's contraction(s)               |
| RMP(s)              | Renormalized model parameter(s)              |
| SC                  | Superconductor                               |
| SDW                 | Spin-density wave                            |
| SSB                 | Spontaneous symmetry breaking                |
| $T_c$               | Critical temperature                         |

---



# Introduction

This thesis project is about my favorite ice cream flavor. [To be continued...]



# Chapter 1

## Superconducting instability

In this chapter the HF methods introduced in Sec. ?? are applied to the SC phase. We study a translationally invariant superconductor, which breaks the conservation of total charge. Analytic calculations will lead us to various effects: bare bands renormalisation, anisotropic Cooper pairing and nematicity. Our description of the SC phase is closely related to the BCS solution for pure  $s$ -wave superconductivity.

As for previous phases, this chapter is divided in three parts. First, analysis of the SC phase symmetries based on the discussion of Chap. ??, and implementation of HF methods. Second, search for theoretical constraints on the self-consistent solution. Third, numeric results.

### 1.1 The SC phase and its symmetries

As for antiferromagnetism, superconductivity can establish in a variety of ways. We deal with the simplest realisation. In Sec. ?? we anticipated that the SC phase is described by the two SSBs

$$\begin{aligned} \mathrm{U}^z(1) &\xrightarrow{\text{SSB}} 0 \\ \mathrm{C}_4 &\xrightarrow{\text{SSB}} \mathrm{C}_2 \end{aligned}$$

Translational invariance is unbroken. Then electronic density needs to be uniform across all sites, similarly to the normal phase discussed in Chap. ?. Non-uniform density distributions are inconsistent with translational invariance in both directions. We have, then

$$\langle \hat{n}_{\mathbf{r}\sigma} \rangle = n \quad (1.1)$$

which is identical to Eq. (??) for the normal phase.

Recall Tabs. ?? and ??, collecting the selection rules for the local repulsion  $\hat{H}_U$  and the non-local attraction  $\hat{H}_V$ . From Eqns. (1.2), (??) for the SC phase we need to account for the following contractions:

|                           |                                 |
|---------------------------|---------------------------------|
| $\hat{H}_U$               | Hartree and Cooper contractions |
| $\hat{H}_V$ (o.s. sector) | Hartree and Cooper contractions |
| $\hat{H}_V$ (s.s. sector) | Hartree and Fock contractions   |

We anticipate their roles in Tab. 1.1.

The SSB of  $\mathrm{U}^z(1)$  allows for EVs like  $\langle \hat{c}^\dagger \hat{c}^\dagger \rangle$  to be non-zero. This leads to the formation of Cooper pairs. These are composite quantum objects with a specific spin-spatial structure. Consider the generic Cooper fluctuation

$$\langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\bar{\sigma}}^\dagger \rangle \quad \text{with } i, j \text{ nearest neighbours}$$

We work in o.s. sector because we know from Tab. ?? that the s.s. sector is prohibited. From basic Quantum Mechanics we know the summation rules of the spin algebra  $\mathfrak{su}(2)$ ,

$$\frac{1}{2} \otimes \frac{1}{2} = 0 \oplus 1$$

| Operator    | Sector | RWCs    | Net effect                                          |
|-------------|--------|---------|-----------------------------------------------------|
| $\hat{H}_U$ |        | Hartree | $\mu$ shift by $-nU$                                |
|             |        | Cooper  | Isotropic pairing in singlet channel                |
| $\hat{H}_V$ | o.s.   | Hartree | $\mu$ shift by $nzV$                                |
|             |        | Cooper  | Anisotropic pairing in singlet and triplet channels |
|             | s.s.   | Hartree | $\mu$ shift by $nzV$                                |
|             |        | Fock    | Band renormalisation and resymmetrisation           |

**Table 1.1** | Relevant Wick's contractions and net effect in the SC phase. This RWCs table is more structured than Tabs. ?? and ??.

| Spatial structure            | Pairing channel |
|------------------------------|-----------------|
| Symmetric wave function      | Singlet pairing |
| Anti-symmetric wave function | Triplet pairing |

**Table 1.2** | Relation of the Cooper pairing channel with the wavefunction spatial symmetry (intended as the inversion  $(x, y) \rightarrow (-x, -y)$ ).

The two pairing channels are, at this level, the singlet channel associated to total spin 0 and the triplet channel associated to total spin 1. Now, when applying HF method to the quartic operators, we get the decomposition of Eq. (??). We can impose to the approximated HF ground-state a particular spatial structure – say, inversion symmetry or antisymmetry. This reflects in the requirement that Cooper pairs show the same spatial structure.

This gives a selection rule over Cooper pairings: if we work with inversion-symmetric structures – say,  $s^*$ -wave or  $d$ -wave – the pairing will take place in singlet channel, allowing us to eliminate the triplet pairing. Working with antisymmetric structures, inversely, we can eliminate the singlet channel. This concept is summarised in Tab. 1.2.

We proceed as we did for the AF phase: in order to highlight the physical effects of  $\hat{H}_V$ , we first analyse the SC phase limitedly to the conventional HM. In next section we prove that BCS-like superconductivity is impossible at HF level on the pure HM, with just the local repulsion  $\hat{H}_U$ . It is the presence of the non-local attraction that enables Cooper pairing mechanism. Before starting, let

$$\hat{n}_{\mathbf{k}\sigma} \equiv \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \quad (1.2)$$

$$\hat{\phi}_{\mathbf{k}} \equiv \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \quad (1.3)$$

where  $\hat{n}_{\mathbf{k}\sigma}$  was defined as for the AF phase, in Eq. (??). The HF approximation in reciprocal space to the EHM will be written in terms of these operators.

## 1.2 Local effects: (impossibility of) superconductivity in the HM

Let us start from the local effects. Following Tab. ?? and Eqns. (1.2), (??), we know the local repulsion decomposes in Hartree and Cooper channels,

$$\begin{aligned}
\hat{H}_U &= U \sum_{\mathbf{r} \in \mathcal{L}} \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}\downarrow} \\
&\simeq U \sum_{\mathbf{r} \in \mathcal{L}} \left( \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \langle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \rangle + \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} - \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \rangle \right) \quad (\text{Hartree}) \\
&\quad + \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \langle \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle + \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \rangle \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} - \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle \quad (\text{Cooper})
\end{aligned}$$

In next sections we deal with each channel separately.



### 1.2.1 Local Hartree channel in the SC phase: analytic derivation

For the Hartree channel, being the density uniform across sites, we can re-use the entirety of Sec. ?? for the normal phase. Substituting Eq. (1.1), we get the approximation of Eq. (??),

$$\hat{H}_U \simeq nU \times \hat{N} - E_{H/U}^{(SC)} + (\text{Cooper channel}) \quad (1.4)$$

thus chemical potential renormalization  $\tilde{\mu} \rightarrow \mu - nU$  is present also in the SC phase.

### 1.2.2 Local Hartree contraction energy

As for the section above, we use the result of Sec. ?? for the normal phase. From Eq. (??)

$$E_{H/U}^{(SC)} = L_x L_y \times n^2 U \quad (1.5)$$

For a locally repulsive model, the Hartree channel contributes negatively to ground state energy. Indeed, contraction energy is subtracted from the hamiltonian. We now move to the Cooper channel.

### 1.2.3 Local Cooper channel in the SC phase: analytic derivation

We move to reciprocal space, through the transformation of Eq. (??). Consider the Cooper channel of Eq. (1.2). In reciprocal space we get:

$$\begin{aligned} \hat{H}_U \simeq (\text{Hartree channel}) + \frac{U}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} & \left[ \langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \right. \\ & \left. + \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \langle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \rangle - \langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \rangle \right] \end{aligned} \quad (1.6)$$

We are not breaking translational invariance, thus only Cooper fluctuations with net zero total momentum are allowed. This means only  $\mathbf{K} = \mathbf{0}$  contributes. Let us prove this: consider the EV of the Cooper pair

$$\langle \hat{c}_{\mathbf{k}_1\uparrow}^\dagger \hat{c}_{\mathbf{k}_2\downarrow}^\dagger \rangle$$

for two momenta  $\mathbf{k}_1, \mathbf{k}_2$ . Moving to real space via an  $\mathcal{F}$ -transform, we get

$$\langle \hat{c}_{\mathbf{k}_1\uparrow}^\dagger \hat{c}_{\mathbf{k}_2\downarrow}^\dagger \rangle \stackrel{\mathcal{F}}{=} \frac{1}{L_x L_y} \sum_{\mathbf{r}_1 \in \mathcal{L}} \sum_{\mathbf{r}_2 \in \mathcal{L}} e^{-i(\mathbf{k}_1 \cdot \mathbf{r}_1 + \mathbf{k}_2 \cdot \mathbf{r}_2)} \langle \hat{c}_{\mathbf{r}_1\uparrow}^\dagger \hat{c}_{\mathbf{r}_2\downarrow}^\dagger \rangle$$

where we used the convention of Eq. (??). Let now  $\mathbf{r}_1 \rightarrow \mathbf{r}$  and  $\mathbf{r}_2 \rightarrow \Delta\mathbf{r}$ . Translational invariance implies that

$$g(\Delta\mathbf{r}) = \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\Delta\mathbf{r}\downarrow}^\dagger \rangle$$

is independent of  $\mathbf{r}$ . Thus we have:

$$\begin{aligned} \langle \hat{c}_{\mathbf{k}_1\uparrow}^\dagger \hat{c}_{\mathbf{k}_2\downarrow}^\dagger \rangle & \stackrel{\mathcal{F}}{=} \sum_{\Delta\mathbf{r} \in \mathcal{L}} e^{-i\mathbf{k}_2 \cdot \Delta\mathbf{r}} g(\Delta\mathbf{r}) \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} e^{-i(\mathbf{k}_1 + \mathbf{k}_2) \cdot \mathbf{r}} \\ & = \sum_{\Delta\mathbf{r} \in \mathcal{L}} e^{-i\mathbf{k}_2 \cdot \Delta\mathbf{r}} g(\Delta\mathbf{r}) \delta_{\mathbf{k}_1 = \mathbf{k}_2} \end{aligned}$$

where we used the identity of Eq. (??). Then the above EV vanishes for  $\mathbf{k}_1 \neq \mathbf{k}_2$ . Getting back to Eq. (1.6), this proves that only pairs at null total momentum can have nonzero EVs.

Let now  $w_{\mathbf{k}}$  be the EV of  $\phi_{\mathbf{k}}^\dagger$ , and  $w^{(s)}$  its  $s$ -wave projection:

$$w_{\mathbf{k}} \equiv \langle \phi_{\mathbf{k}}^\dagger \rangle \quad (1.7)$$

$$w^{(s)} \equiv \frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} w_{\mathbf{q}} \quad (1.8)$$

Eq. (1.6) reduces to

$$\hat{H}_U \simeq (\text{Hartree channel}) + \frac{U}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} \left[ \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \hat{\phi}_{\mathbf{k}'} + \hat{\phi}_{\mathbf{k}}^\dagger \langle \hat{\phi}_{\mathbf{k}'} \rangle \right] - E_{C/U}^{(\text{SC})} \quad (1.9)$$

being  $E_{C/U}^{(\text{SC})}$  the HM Cooper contraction energy, written explicitly in Sec. 1.2.4. Let now

$$\Delta^{(s)} \equiv -U w^{(s)} \quad (1.10)$$

This is the  $s$ -wave component of the gap function. Note that, in general,  $\Delta^{(s)} \in \mathbb{C}$ . However, we can always choose  $\Delta^{(s)}$  as purely real. This is due to the fact that the hamiltonian is gauge-invariant, then it suffices to apply the gauge map of Eq. (??) to the fermionic operators with

$$\eta \equiv \frac{\arg \Delta^{(s)}}{2}$$

to obtain  $\Delta^{(s)} \in \mathbb{R}$ . For the sake of generality, we proceed with  $\Delta^{(s)} \in \mathbb{C}$ . Although this choice gives no advantage now, we implement it anyway in order to derive the most general form of the equations that follow. This is useful for the generalisation to the EHM.

In the end, we reduce  $\hat{H}_U$  from Eq. (1.9) to

$$\hat{H}_U \simeq (\text{Hartree channel}) - \sum_{\mathbf{k} \in \text{BZ}} \left[ \Delta^{(s)} \hat{\phi}_{\mathbf{k}} + \Delta^{(s)*} \hat{\phi}_{\mathbf{k}}^\dagger \right] - E_{C/U}^{(\text{SC})} \quad (1.11)$$

This last decomposition resembles Eq. (??), obtained for the AF phase in the Hartree channel.

#### 1.2.4 Local Cooper contraction energy

The Cooper contraction energy in the Cooper channel is given by:

$$\begin{aligned} E_{C/U}^{(\text{SC})} &\equiv \frac{U}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \langle \hat{\phi}_{\mathbf{k}'} \rangle \\ &= L_x L_y \times U |w^{(s)}|^2 \end{aligned} \quad (1.12)$$

As for the Hartree channel, also the Cooper channel contributes negatively to ground-state energy.

#### 1.2.5 Superconductivity in the HM

We proceed here as we did for the AF phase in Sec. ???. We start by giving a matrix representation of the approximated hamiltonian, then map it to a pseudospin problem and finally specify how to diagonalise it to retrieve its ground state.

**Matrix representation.** Collecting the results of the above paragraphs, we obtain the quadratic approximation to the Hubbard hamiltonian in the SC phase,

$$\begin{aligned} \hat{H}_t + \hat{H}_U &\simeq \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \epsilon_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \\ &\quad - \sum_{\mathbf{k} \in \text{BZ}} \left[ \Delta^{(s)} \hat{\phi}_{\mathbf{k}} + \Delta^{(s)*} \hat{\phi}_{\mathbf{k}}^\dagger \right] + nU \times \hat{N} - \left[ E_{H/U}^{(\text{SC})} + E_{C/U}^{(\text{SC})} \right] \end{aligned} \quad (1.13)$$

We define the shifted bare bands:

$$\xi_{\mathbf{k}} \equiv \epsilon_{\mathbf{k}} - \tilde{\mu}$$

Notice we are using the renormalised chemical potential. Due to inversion symmetry, it holds  $\xi_{\mathbf{k}} = \xi_{-\mathbf{k}}$ . Using Fermi anticommutation rules, the kinetic operator can be written as:

$$\begin{aligned}
\hat{H}_t - \tilde{\mu}\hat{N} &= \sum_{\sigma} \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}\sigma} \hat{n}_{\mathbf{k}\sigma} \\
&= \sum_{\mathbf{k} \in \text{BZ}} (\xi_{\mathbf{k}} \hat{n}_{\mathbf{k}\uparrow} + \xi_{-\mathbf{k}} \hat{n}_{-\mathbf{k}\downarrow}) \\
&= \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}} \left( \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} + 1 \right) \\
&= \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}} \left( \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \right) + E_a^{(\text{SC})}
\end{aligned}$$

being  $E_a^{(\text{SC})}$  the “anticommutation energy”

$$E_a^{(\text{SC})} \equiv \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}} \quad (1.14)$$

In the end, the hamiltonian is approximated by:

$$\begin{aligned}
\hat{H}_t + \hat{H}_U &\simeq \sum_{\mathbf{k} \in \text{BZ}} \epsilon_{\mathbf{k}} \left[ \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\downarrow}^{\dagger} \right] + E_a^{(\text{SC})} \\
&\quad - \sum_{\mathbf{q} \in \text{BZ}} \left[ \Delta^{(s)} \hat{\phi}_{\mathbf{q}} + \Delta^{(s)*} \hat{\phi}_{\mathbf{q}}^{\dagger} \right] + nU \times \hat{N} - \left[ E_{\text{H}/U}^{(\text{SC})} + E_{\text{C}/U}^{(\text{SC})} \right]
\end{aligned}$$

with  $E_{\text{H}/U}^{(\text{SC})}$ ,  $E_{\text{C}/U}^{(\text{SC})}$  the contraction energies arising from Hartree and Cooper channels, and  $E_a^{(\text{SC})}$  the energy shift resulting from anticommutation rules, given respectively in Eqns. (1.5), (1.12), and (1.14). We recognised the  $s$ -wave gap  $\Delta^{(s)}$  from Eq. (1.10).

Let the Nambu spinor

$$\hat{\Phi}_{\mathbf{k}} \equiv \begin{bmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \end{bmatrix}$$

The approximated hamiltonian can be written in matrix form as:

$$\hat{H}_t + \hat{H}_U - \mu\hat{N} \simeq \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \hat{\Phi}_{\mathbf{k}}^{\dagger} h_{\mathbf{k}} \hat{\Phi}_{\mathbf{k}} + E_a^{(\text{SC})} - \left[ E_{\text{H}/U}^{(\text{SC})} + E_{\text{C}/U}^{(\text{SC})} \right]$$

being

$$h_{\mathbf{k}} \equiv \begin{bmatrix} \xi_{\mathbf{k}} & -\Delta_{\mathbf{k}}^* \\ -\Delta_{\mathbf{k}} & -\xi_{\mathbf{k}} \end{bmatrix} \quad \text{with} \quad \xi_{\mathbf{k}} \equiv \epsilon_{\mathbf{k}} - \tilde{\mu}$$

and  $\Delta_{\mathbf{k}} = \Delta^{(s)}$ .

**Pseudospin representation.** The hamiltonian can be written equivalently as

$$h_{\mathbf{k}} = \xi_{\mathbf{k}} \tau^z - \text{Re}\{\Delta_{\mathbf{k}}\} \tau^x - \text{Im}\{\Delta_{\mathbf{k}}\} \tau^y$$

Let now the pseudospin operator  $\hat{\mathbf{s}}_{\mathbf{k}}$

$$\hat{s}_{\mathbf{k}}^{\alpha} \equiv \hat{\Phi}_{\mathbf{k}}^{\dagger} \tau^{\alpha} \hat{\Phi}_{\mathbf{k}} \quad \text{with} \quad \hat{\mathbf{s}}_{\mathbf{k}\sigma} = \begin{bmatrix} \hat{s}_{\mathbf{k}\sigma}^x \\ \hat{s}_{\mathbf{k}\sigma}^y \\ \hat{s}_{\mathbf{k}\sigma}^z \end{bmatrix}$$

and the pseudofield  $\mathbf{b}_{\mathbf{k}}$ :

$$\mathbf{b}_{\mathbf{k}} \equiv \begin{bmatrix} -\text{Re}\{\Delta_{\mathbf{k}}\} \\ -\text{Im}\{\Delta_{\mathbf{k}}\} \\ \xi_{\mathbf{k}} \end{bmatrix}$$

| EV                                       | Meaning                                                                              |
|------------------------------------------|--------------------------------------------------------------------------------------|
| $\langle \hat{s}_{\mathbf{k}}^x \rangle$ | $\langle \hat{\phi}_{\mathbf{k}} + \hat{\phi}_{\mathbf{k}}^\dagger \rangle$          |
| $\langle \hat{s}_{\mathbf{k}}^y \rangle$ | $i \langle \hat{\phi}_{\mathbf{k}} - \hat{\phi}_{\mathbf{k}}^\dagger \rangle$        |
| $\langle \hat{s}_{\mathbf{k}}^z \rangle$ | $\langle \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{-\mathbf{k}\downarrow} \rangle - 1$ |

**Table 1.3** | List of the various expectation values (EVs) of each pseudospin component and their meaning in terms of fermionic operators.

For a purely real gap, the pseudofield has no  $y$  component. In general, different gauges rotate the pseudofield around the  $z$  axis.

As was for the AF phase, these operators satisfy a  $\mathfrak{su}(2)$  algebra up to an constant, irrelevant in the context of this discussion. They satisfy the identities:

$$\hat{s}_{\mathbf{k}}^x = \hat{\phi}_{\mathbf{k}} + \hat{\phi}_{\mathbf{k}}^\dagger \quad (1.15)$$

$$\hat{s}_{\mathbf{k}}^y = i \left( \hat{\phi}_{\mathbf{k}} - \hat{\phi}_{\mathbf{k}}^\dagger \right) \quad (1.16)$$

$$\hat{s}_{\mathbf{k}}^z = \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{-\mathbf{k}\downarrow} - 1 \quad (1.17)$$

We report the EVs related to these identities Tab. 1.3.

We express the approximated hamiltonian in pseudospin representation as:

$$\hat{H}_t + \hat{H}_U - \mu \hat{N} \simeq \sum_{\mathbf{k} \in \text{BZ}} \mathbf{b}_{\mathbf{k}} \cdot \hat{\mathbf{s}}_{\mathbf{k}} + nU\hat{N} + E_{\text{a}}^{(\text{SC})} - \left[ E_{\text{H}/U}^{(\text{SC})} + E_{\text{C}/U}^{(\text{SC})} \right] \quad (1.18)$$

This operator is similar enough to the one found for the AF phase, in Eq. (??). It describes a set of pseudospin subject to a uniform field. As for the AF phase, we define the Bogoliubov bands  $E_{\mathbf{k}}$

$$E_{\mathbf{k}} \equiv \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$

The key difference with the AF Bogoliubov bands of Eq. (??) is that  $E_{\mathbf{k}}$  now are functions of  $\tilde{\mu}$ , since  $\xi_{\mathbf{k}}$  depends on the chemical potential. In other words, now the Bogoliubov bands depend on the shifted bands  $\xi_{\mathbf{k}}$  instead of the bare bands  $\epsilon_{\mathbf{k}}$ .

Moreover, let us define the angles  $\theta_{\mathbf{k}}$ ,  $\zeta_{\mathbf{k}}$  as

$$\cos 2\theta_{\mathbf{k}} = \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \quad \sin 2\theta_{\mathbf{k}} \cos 2\zeta_{\mathbf{k}} = \frac{\text{Re}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}} \quad \sin 2\theta_{\mathbf{k}} \sin 2\zeta_{\mathbf{k}} = \frac{\text{Im}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}}$$

Note that choosing the gauge  $\eta = \arg \Delta^{(s)}/2$  the imaginary part of the gap is zero, and this reflects in  $2\zeta_{\mathbf{k}} = k \times 2\pi$  with  $k \in \mathbb{Z}$ .

**Diagonalisation.** For the SC phase, diagonalisation of the HF hamiltonian follows the same procedure as was for the AF phase. Indeed, the diagonalising matrix is given by

$$\begin{aligned} W_{\mathbf{k}} &\equiv e^{-i\theta_{\mathbf{k}}\tau^y} e^{-i\zeta_{\mathbf{k}}\tau^z} \\ &= \begin{bmatrix} \cos \theta_{\mathbf{k}} & -\sin \theta_{\mathbf{k}} \\ \sin \theta_{\mathbf{k}} & \cos \theta_{\mathbf{k}} \end{bmatrix} \begin{bmatrix} e^{-i\zeta_{\mathbf{k}}} & \\ & e^{i\zeta_{\mathbf{k}}} \end{bmatrix} \\ &= \begin{bmatrix} \cos \theta_{\mathbf{k}} e^{-i\zeta_{\mathbf{k}}} & -\sin \theta_{\mathbf{k}} e^{i\zeta_{\mathbf{k}}} \\ \sin \theta_{\mathbf{k}} e^{-i\zeta_{\mathbf{k}}} & \cos \theta_{\mathbf{k}} e^{i\zeta_{\mathbf{k}}} \end{bmatrix} \end{aligned} \quad (1.19)$$

and the graphic procedure is the same of Fig. ??, with  $\epsilon \rightarrow \xi$ . We get:

$$d_{\mathbf{k}} = W_{\mathbf{k}} h_{\mathbf{k}} W_{\mathbf{k}}^\dagger \quad \text{with} \quad d_{\mathbf{k}} \equiv \begin{bmatrix} E_{\mathbf{k}} & \\ & -E_{\mathbf{k}} \end{bmatrix} \quad (1.20)$$

In the tilted frame we have the expression for the Nambu spinors:

$$\hat{\Gamma}_{\mathbf{k}} = W_{\mathbf{k}} \hat{\Psi}_{\mathbf{k}} \quad (1.21)$$

Explicitly:

$$\begin{aligned}\hat{\Gamma}_{\mathbf{k}} &= \begin{bmatrix} \cos \theta_{\mathbf{k}} e^{-i\zeta_{\mathbf{k}}} & -\sin \theta_{\mathbf{k}} e^{i\zeta_{\mathbf{k}}} \\ \sin \theta_{\mathbf{k}} e^{-i\zeta_{\mathbf{k}}} & \cos \theta_{\mathbf{k}} e^{i\zeta_{\mathbf{k}}} \end{bmatrix} \begin{bmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{bmatrix} \\ &= \begin{bmatrix} \cos \theta_{\mathbf{k}} e^{-i\zeta_{\mathbf{k}}} \hat{c}_{\mathbf{k}\uparrow} - \sin \theta_{\mathbf{k}} e^{i\zeta_{\mathbf{k}}} \hat{c}_{-\mathbf{k}\downarrow}^\dagger \\ \sin \theta_{\mathbf{k}} e^{-i\zeta_{\mathbf{k}}} \hat{c}_{\mathbf{k}\uparrow} + \cos \theta_{\mathbf{k}} e^{i\zeta_{\mathbf{k}}} \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{bmatrix} \equiv \begin{bmatrix} \hat{\gamma}_{\mathbf{k}}^{(+)} \\ \hat{\gamma}_{\mathbf{k}}^{(-)} \end{bmatrix}\end{aligned}\quad (1.22)$$

The new Fermionic operators describe a Fermi gas on distributed on two mirrored bands,

$$\hat{H}_t + \hat{H}_U - \tilde{\mu} \hat{N} \simeq \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} E_{\mathbf{k}} \left( \hat{\gamma}_{\mathbf{k}}^{(+)\dagger} \hat{\gamma}_{\mathbf{k}}^{(+)} - \hat{\gamma}_{\mathbf{k}}^{(-)\dagger} \hat{\gamma}_{\mathbf{k}}^{(-)} \right) + (\dots)$$

with  $(\dots)$  the various energy shifts.

### 1.2.6 Features of the SC solution

In this section we give some details about the HF solution for the translationally-invariant SC phase. We prove that in the conventional HM, with local repulsion  $\hat{H}_U$ , isotropic BCS-like superconductivity is forbidden. This serves as ground to motivate the passage to the EHM.

**Pseudospin EVs.** The EVs of pseudospin operators in the various directions are given by:

$$\begin{aligned}\langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^x \hat{\Psi}_{\mathbf{k}} \rangle &= -\sin 2\theta_{\mathbf{k}} \cos 2\zeta_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle \\ \langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^y \hat{\Psi}_{\mathbf{k}} \rangle &= -\sin 2\theta_{\mathbf{k}} \sin 2\zeta_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle \\ \langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^z \hat{\Psi}_{\mathbf{k}} \rangle &= \cos 2\theta_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle\end{aligned}$$

In the tilted frame, the averaged  $z$  projection of pseudospin is:

$$\langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle = f(E_{\mathbf{k}}; \beta, 0) - f(-E_{\mathbf{k}}; \beta, 0) \quad (1.23)$$

Compared with the respective relation for the AF phase, Eq. (??), the main difference here is that the chemical potential inside the Fermi-Dirac distribution is set to zero. This is due to the fact that  $\tilde{\mu}$  is already included in the renormalised bands  $E_{\mathbf{k}}$ . The approximated quadratic hamiltonian describes a gas of  $\gamma$  fermions at null chemical potential.

Let us define the thermal factors:

$$\text{Th}_{\mathbf{k}}^{(\pm)}[\beta] \equiv f(-E_{\mathbf{k}}; \beta, 0) \pm f(E_{\mathbf{k}}; \beta, 0) \quad (1.24)$$

In the end we have the EVs:

$$\langle \hat{s}_{\mathbf{k}}^x \rangle = \frac{\text{Re}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}} \text{Th}_{\mathbf{k}}^{(-)}[\beta] \quad (1.25)$$

$$\langle \hat{s}_{\mathbf{k}}^y \rangle = \frac{\text{Im}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}} \text{Th}_{\mathbf{k}}^{(-)}[\beta] \quad (1.26)$$

$$\langle \hat{s}_{\mathbf{k}}^z \rangle = -\frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \text{Th}_{\mathbf{k}}^{(-)}[\beta] \quad (1.27)$$

Note that, as is expressed in Tab. 1.3, the  $z$  component of the pseudospin is related to density, being  $1 + \langle \hat{s}_{\mathbf{k}}^z \rangle = \langle \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{-\mathbf{k}\downarrow} \rangle$ .

**$s$ -wave gap self-consistency equation.** Applying the HF method to the conventional HM in SC phase, we obtained just one HFP:  $w^{(s)}$ . Let us derive its self-consistency equation. First, let

$$\tau^\pm \equiv \frac{\tau^x \pm i\tau^y}{2}$$

Recall Eq. (1.8) and expand it, to obtain

$$\begin{aligned}
w^{(s)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \\
&= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^+ \hat{\Phi}_{\mathbf{k}} \rangle \\
&= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\langle \hat{s}_{\mathbf{k}}^x \rangle + i \langle \hat{s}_{\mathbf{k}}^y \rangle}{2} \\
&= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\Delta_{\mathbf{k}}}{E_{\mathbf{k}}} \text{Th}_{\mathbf{k}}^{(-)}[\beta]
\end{aligned} \tag{1.28}$$

Last line is the explicit form of the self-consistency equation for  $w^{(s)}$ .

**Impossibility of  $s$ -wave superconductivity.** Let us define the critical  $U$ ,

$$U_c[\beta] \equiv \left[ \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\text{Th}_{\mathbf{k}}^{(-)}[\beta]}{E_{\mathbf{k}}} \right]^{-1} \tag{1.29}$$

We call it “critical” because its value gives a sort of boundary for convergence. Indeed, as we show in App. ??, for the SC phase in conventional HM, we are able to optimise parametrically the mixing parameter  $g$  in order to ensure convergence, based on the ratio  $U/U_c[\beta]$ .

Let us show that superconductivity is not possible in the conventional HM with  $U > 0$ . Eq. (1.28) can be rewritten as:

$$w^{(s)} = -\frac{U}{U_c[\beta]} w^{(s)}$$

By construction  $U_c[\beta] > 0$  (the thermal factor is negative-defined). Hence the unique possible self-consistent solution for the equation above is  $w^{(s)} = 0$ . This implication terminates the argument: superconductivity cannot establish self-consistently at HF level in the repulsive HM. If instead  $U < 0$ , which is, if the model is attractive, the aforementioned implication is not true and a self-consistent non-zero solution is possible.

**Density.** In order to estimate density, we can use the  $z$  projection of the pseudospin operators. From Eq. (1.17) we know

$$\hat{s}_{\mathbf{k}}^z + 1 = \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{\mathbf{k}\downarrow}$$

Then, density can be estimated as:

$$\begin{aligned}
n &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{-\mathbf{k}\downarrow} \rangle \\
&= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\langle \hat{s}_{\mathbf{k}}^z \rangle + 1) \\
&= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left( 1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \text{Th}_{\mathbf{k}}^{(-)}[\beta] \right)
\end{aligned} \tag{1.30}$$

We can now move to the EHM.

### 1.3 Extension to the EHM

Add now the non-local interaction  $\hat{H}_V$ . As we did for the AF phase, we look for a solution with an identical mathematical structure as the one here described for the conventional HM. We define the renormalised quantities

$$\xi_{\mathbf{k}} \rightarrow \tilde{\xi}_{\mathbf{k}} \quad E_{\mathbf{k}} \rightarrow \tilde{E}_{\mathbf{k}} \quad \Delta_{\mathbf{k}} \rightarrow \tilde{\Delta}_{\mathbf{k}} \quad \mu \rightarrow \tilde{\mu}$$

To be completely correct, the chemical potential was already renormalized in the HM derivation in Sec. ?? . It here acquires yet another renormalization. The band energies renormalization is simply

$$\tilde{E}_{\mathbf{k}} \equiv \sqrt{\tilde{\xi}_{\mathbf{k}}^2 + |\tilde{\Delta}_{\mathbf{k}}|^2}$$

Let the renormalised angle:

$$\sin 2\tilde{\theta}_{\mathbf{k}} = \frac{|\tilde{\Delta}_{\mathbf{k}}|}{\tilde{E}_{\mathbf{k}}} \quad \cos 2\tilde{\theta}_{\mathbf{k}} = \frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \quad (1.31)$$

and similarly the angle  $\tilde{\zeta}_{\mathbf{k}}$ ,

$$\sin 2\tilde{\zeta}_{\mathbf{k}} = \frac{\text{Re}\{\tilde{\Delta}_{\mathbf{k}}\}}{|\tilde{\Delta}_{\mathbf{k}}|} \quad \cos 2\tilde{\zeta}_{\mathbf{k}} = \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{|\tilde{\Delta}_{\mathbf{k}}|} \quad (1.32)$$

The pseudospin EVs are renormalised too,

$$\langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^x \hat{\Phi}_{\mathbf{k}} \rangle = -\sin 2\tilde{\theta}_{\mathbf{k}} \sin 2\tilde{\zeta}_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle \quad (1.33)$$

$$\langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^y \hat{\Phi}_{\mathbf{k}} \rangle = -\sin 2\tilde{\theta}_{\mathbf{k}} \cos 2\tilde{\zeta}_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle \quad (1.34)$$

$$\langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^z \hat{\Phi}_{\mathbf{k}} \rangle = \cos 2\tilde{\theta}_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle \quad (1.35)$$

Let also:

$$\langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle = f(\tilde{E}_{\mathbf{k}}; \beta, 0) - f(-\tilde{E}_{\mathbf{k}}; \beta, 0) \quad (1.36)$$

$$\tilde{\text{Th}}_{\mathbf{k}}^{(\pm)}[\beta, \tilde{\mu}] \equiv f(-\tilde{E}_{\mathbf{k}}; \beta, 0) \pm f(\tilde{E}_{\mathbf{k}}; \beta, 0) \quad (1.37)$$

For the SC phase, the thermal factors reduce to:

$$\begin{aligned} \tilde{\text{Th}}_{\mathbf{k}}^{(+)} &= 1 \\ \tilde{\text{Th}}_{\mathbf{k}}^{(-)} &= -\tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \end{aligned}$$

Hence, pseudospin expectation values are:

$$\langle \hat{s}_{\mathbf{k}}^x \rangle = -\frac{\text{Re}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \quad (1.38)$$

$$\langle \hat{s}_{\mathbf{k}}^y \rangle = -\frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \quad (1.39)$$

$$\langle \hat{s}_{\mathbf{k}}^z \rangle = \frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \quad (1.40)$$

These equations are an extension of Eqns. (1.38), (1.39), (1.40) for the EHM.

We can rewrite the equations for  $w^{(s)}$  and  $n$  accordingly,

$$w^{(s)} = -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \quad (1.41)$$

$$n = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[ 1 + \cos 2\tilde{\theta}_{\mathbf{k}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right] \quad (1.42)$$

where we used Eqns. (1.28), (1.30).

From Tabs. ?? and ?? we know none of the three channels (Hartree, Fock, Cooper) is subject to selection rules. Thus all terms of the decomposition of  $\hat{H}_V$  given in Eq. (??) contribute. As always, we break down each.

## 1.4 Non-local Hartree effects: chemical potential renormalisation

We start by analysing the non-local Hartree channel,

$$\hat{H}_V \simeq -V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \left( \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle \right) + (\text{Fock channel}) + (\text{Cooper channel})$$

where we used Eq. (??). This discussion is based on that for the normal phase in Sec. ??.

### 1.4.1 Non-local Hartree channel in the SC phase: analytic derivation

Without repeating the calculations, the non-local attraction  $\hat{H}_V$  is reduced in the Hartree channel to

$$\hat{H}_V \simeq -2nzV \times \hat{N} - E_{\text{H}/V}^{(\text{SC})} + (\text{all the rest})$$

This, together with the  $\hat{H}_U$  Hartree channel decomposition

$$\hat{H}_U \simeq nU \times \hat{N} - E_{\text{H}/U}^{(\text{SC})}$$

gives the chemical potential full renormalization, identical to Eq. (??),

$$\tilde{\mu} \equiv \mu + n(2zV - U) \quad (1.43)$$

### 1.4.2 Non-local Hartree contraction energy

The contraction energy for the non-local sector was derived in Eq. (??),

$$E_{\text{H}/V}^{(\text{SC})} = -L_x L_y \times 2zn^2V$$

and together with the local counterpart of Eq. (??)

$$E_{\text{H}/U}^{(\text{SC})} = L_x L_y \times n^2U$$

Now we move to the more interesting Fock sector.

## 1.5 Fock effects: bare bands renormalization

As done for the other phases, we here discuss the bands renormalization effect derived from the Fock sector of the non-local attraction. From Eq. (??), we have

$$\hat{H}_V \simeq (\text{Hartree channel}) + V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \left( \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rangle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} \rangle \right) + (\text{Cooper channel})$$

This discussion is essentially analogous the one done for the normal phase in Sec. ??. As for previous phases, following the selection rules of Tab. ??, only the s.s. channel contributes. Therefore we use from now on, in this section,  $\sigma = \sigma'$ .

### 1.5.1 Fock channel in the SC phase: analytic derivation

Let us define, as done for the normal phase in Eq. (??) and for the AF phase in Eq. (??),

$$u_{\mathbf{k}\sigma} \equiv \langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \rangle \quad (1.44)$$



Eq. (??) stands perfectly valid, translated to the SC phase

$$V \sum_{\langle ij \rangle} \sum_{\sigma} \left[ \langle \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} \rangle \hat{c}_{j\sigma}^{\dagger} \hat{c}_{i\sigma} + \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} \langle \hat{c}_{j\sigma}^{\dagger} \hat{c}_{i\sigma} \rangle - \langle \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} \rangle \langle \hat{c}_{j\sigma}^{\dagger} \hat{c}_{i\sigma} \rangle \right] \\ \stackrel{\mathcal{F}}{=} \sum_{\gamma, \sigma} \left[ \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} u_{\mathbf{k}\sigma} \right] \times V \sum_{\mathbf{k}' \in \text{BZ}} \varphi_{\mathbf{k}'}^{(\gamma)*} \hat{c}_{\mathbf{k}'\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} - E_{\text{F}/V}^{(\text{SC})} \quad (1.45)$$

being

$$E_{\text{F}/V}^{(\text{SC})} \equiv \frac{V}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} \sum_{\sigma} [\cos(\delta k_x) + \cos(\delta k_y)] \langle \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} \rangle \langle \hat{c}_{\mathbf{k}'\sigma}^{\dagger} \hat{c}_{\mathbf{k}'\sigma} \rangle \quad (1.46)$$

To get to this point, we employed the assumption

$$\langle \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}'\sigma'} \rangle = \delta_{\mathbf{k}=\mathbf{k}'} \delta_{\sigma=\sigma'} \langle \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} \rangle$$

which essentially states translational invariance and spin degeneracy. This assumption is verified by the BCS ground state of Eq. ???. Let us define each  $\gamma$ -wave component as

$$u_{\sigma}^{(\gamma)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} \rangle$$

These are our HFPs in the Fock channel. The SC phase does not break inversion symmetry: then, as we concluded for the other phases,  $p_{\ell}$  components are null. The renormalised bands are, as in Eqns. (??), (??),

$$\tilde{\epsilon}_{\mathbf{k}} \equiv \epsilon_{\mathbf{k}} + u^{(s^*)} V (\cos k_x + \cos k_y) + u^{(d)} V (\cos k_x - \cos k_y)$$

Let us compute the EVs of  $u_{\mathbf{k}\sigma}$  in the two spin sectors,

$$u_{\mathbf{k}\uparrow} = \langle \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} \rangle \\ = \left\langle \hat{\Phi}_{\mathbf{k}}^{\dagger} \frac{\mathbb{I} + \tau^z}{2} \hat{\Phi}_{\mathbf{k}} \right\rangle \\ = \frac{1}{2} \left[ \langle \hat{\Gamma}_{\mathbf{k}}^{\dagger} \hat{\Gamma}_{\mathbf{k}} \rangle + \langle \hat{s}_{\mathbf{k}}^z \rangle \right] \\ = \frac{1}{2} \left[ \tilde{\text{Th}}_{\mathbf{k}}^{(+)}[\beta] - \cos(2\tilde{\theta}_{\mathbf{k}}) \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta] \right] \\ = \frac{1}{2} \left[ 1 - \cos(2\tilde{\theta}_{\mathbf{k}}) \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right] \quad (1.47)$$

having we used Eq. (??) and following. For the spin  $\downarrow$  EV,

$$u_{-\mathbf{k}\downarrow} = \langle \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow} \rangle \\ = 1 - \langle \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \rangle \\ = 1 - \left\langle \hat{\Phi}_{\mathbf{k}}^{\dagger} \frac{\mathbb{I} - \tau^z}{2} \hat{\Phi}_{\mathbf{k}} \right\rangle \\ = 1 - \frac{1}{2} \left[ \langle \hat{\Gamma}_{\mathbf{k}}^{\dagger} \hat{\Gamma}_{\mathbf{k}} \rangle - \langle \hat{s}_{\mathbf{k}}^z \rangle \right] \\ = 1 - \frac{1}{2} \left[ \tilde{\text{Th}}_{\mathbf{k}}^{(+)}[\beta] + \cos 2\tilde{\theta}_{\mathbf{k}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta] \right] \\ = 1 - \frac{1}{2} \left[ 1 + \cos(2\tilde{\theta}_{\mathbf{k}}) \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right] \quad (1.48)$$

From the above results, note first that

$$u_{\mathbf{k}\uparrow} = u_{-\mathbf{k}\downarrow}$$

| Symmetry            | Self-consistency equation                                                                                                                                                                                                  | Graph   |
|---------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| $s^*$ -wave         | $u^{(s^*)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) \left[ 1 - \frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right]$ | Fig. ?? |
| $d_{x^2-y^2}$ -wave | $u^{(d)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) \left[ 1 - \frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right]$   | Fig. ?? |

Table 1.4 | Symmetry resolved self-consistency equations for the harmonics of  $u_{\mathbf{k}}$ .

which is coherent with our requirement of a uniform-density, spin uniform and inversion symmetric phase. Note also that Eqns. (1.47), (1.48) combine to give the expression for density of Eq. (1.42).

If we take the  $\tilde{\theta}_{\mathbf{k}} \rightarrow 0$  limit, we must retrieve the Fermi-Dirac distribution. Indeed, this limit coincides with  $U \rightarrow 0$ . Explicitly:

$$\begin{aligned}
\lim_{\tilde{\theta}_{\mathbf{k}} \rightarrow 0} u_{\mathbf{k}\sigma} &= \frac{1}{2} \left[ 1 + \tanh\left(\frac{\beta \tilde{\xi}_{\mathbf{k}}}{2}\right) \right] \\
&= \frac{1}{2} \left[ \frac{e^{\beta \tilde{\xi}_{\mathbf{k}}/2} + e^{-\beta \tilde{\xi}_{\mathbf{k}}/2}}{e^{\beta \tilde{\xi}_{\mathbf{k}}/2} + e^{-\beta \tilde{\xi}_{\mathbf{k}}/2}} - \frac{e^{\beta \tilde{\xi}_{\mathbf{k}}/2} - e^{-\beta \tilde{\xi}_{\mathbf{k}}/2}}{e^{\beta \tilde{\xi}_{\mathbf{k}}/2} + e^{-\beta \tilde{\xi}_{\mathbf{k}}/2}} \right] \\
&= f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})
\end{aligned}$$

Correctly, the null-gap limit (for which the angle reduces to zero) gives back a perfectly degenerate Fermi gas.

We can then derive the  $u_{\mathbf{k}}$ -related HFPs self-consistency equations. In next passages, we use  $\sigma = \uparrow$  but could have used the opposite due to spin degeneracy discussed above. Then:

$$\begin{aligned}
u^{(\gamma)} &\equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{n}_{\mathbf{k}\uparrow} \rangle \\
&= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \left[ 1 - \cos(2\tilde{\theta}_{\mathbf{k}}) \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right]
\end{aligned} \tag{1.49}$$

reported explicitly for the only two inversion symmetric structures in Tab. 1.4.

### 1.5.2 Fock contraction energy

In terms of the energy shift due to Wick's contraction, Eq. (??) holds identically

$$E_{\text{F}/V}^{(\text{SC})} = L_x L_y \times V \left[ |u^{(s^*)}|^2 + |u^{(d)}|^2 \right]$$

with the only obvious difference that the  $u^{(\gamma)}$  are computed on the SC ground state by the means of Eqns. (1.47), (1.48).

## 1.6 Non-local Cooper effects: anisotropic pairing

In this section we deal with the Cooper part of Eq. (??),

$$\begin{aligned}
\hat{H}_V &= \simeq (\text{Hartree channel}) + (\text{Fock channel}) \\
&- V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \left( \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \langle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \rangle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \rangle \langle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle \right)
\end{aligned}$$

We know from Eq. (??) that the non-local attraction can be separated in o.s. and s.s. channels. As we reported in Tab. ?? and stated in Sec. 1.1, the only non-zero contribution comes from the o.s. sector.

### 1.6.1 Non-local Cooper o.s. channel in the SC phase: analytic derivation

The o.s. Cooper sector contributes both to singlet and triplet pairing. Note that in singlet pairing channel the spatial wavefunction must be inversion-symmetric. On the contrary, in triplet pairing channel it needs to be inversion-antisymmetric.

We move to reciprocal space. From Eq. (??), we get

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} \simeq (\text{Hartree channel}) \\ - \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \times \left[ \langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \right. \\ \left. + \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \langle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \rangle - \langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \rangle \right] \end{aligned}$$

Identical considerations as for the local repulsion hold, and uniquely the  $\mathbf{K} = \mathbf{0}$  term gives a non-zero contribution, in general. Therefore:

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} \simeq (\text{Hartree channel}) \\ - \frac{2V}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \times \left[ \langle \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \rangle \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} \right. \\ \left. + \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \langle \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} \rangle - \langle \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \rangle \langle \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} \rangle \right] \end{aligned}$$

Then, using the harmonics decomposition of Eq. (??), we end up with:

$$\hat{H}_V^{(\text{o.s.})} \simeq (\text{Hartree channel}) - \frac{V}{L_x L_y} \sum_{\gamma} \sum_{\mathbf{k}, \mathbf{k}'} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} \left[ w_{\mathbf{k}} \hat{\phi}_{\mathbf{k}'} + w_{\mathbf{k}}^* \hat{\phi}_{\mathbf{k}'}^\dagger \right] - E_{C/V/\text{o.s.}}^{(\text{SC})} \quad (1.50)$$

In the equation above the contraction energy is given by

$$E_{C/V/\text{o.s.}}^{(\text{SC})} \equiv - \frac{V}{L_x L_y} \sum_{\gamma} \sum_{\mathbf{k}, \mathbf{k}'} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} w_{\mathbf{k}} w_{\mathbf{k}'}^* \quad (1.51)$$

Then, let the harmonics  $w^{(\gamma)}$  be

$$w^{(\gamma)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} w_{\mathbf{k}}$$

Eq. (1.50) becomes

$$\hat{H}_V^{(\text{o.s.})} \simeq (\text{Hartree channel}) - V \sum_{\gamma} \sum_{\mathbf{k} \in \text{BZ}} \left[ w^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} \hat{\phi}_{\mathbf{k}} + \text{h.c.} \right] - E_{C/V/\text{o.s.}}^{(\text{SC})}$$

extending the symmetry sum to  $\gamma \in \{s^*, p_x, p_y, d_{x^2-y^2}\}$ .

For the  $\gamma$ -wave component of the superconducting order parameter, the following chain, similar to Eq. (1.28), holds:

$$\begin{aligned} w^{(\gamma)} &\equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^+ \hat{\Phi}_{\mathbf{k}} \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\langle \hat{s}_{\mathbf{k}}^x \rangle + i \langle \hat{s}_{\mathbf{k}}^y \rangle}{2} \\ &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{T}}\text{h}_{\mathbf{k}}^{(-)}[\beta] \end{aligned} \quad (1.52)$$

| Symmetry            | Self-consistency equation                                                                                                                                                                                  | Graph   |
|---------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| $s$ -wave           | $w^{(s)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$                         | Fig. ?? |
| $s^*$ -wave         | $w^{(s^*)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$ | Fig. ?? |
| $p_x$ -wave         | $w^{(p_x)} = \frac{i\sqrt{2}}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \sin k_x \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$      | Fig. ?? |
| $p_y$ -wave         | $w^{(p_y)} = \frac{i\sqrt{2}}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \sin k_y \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$      | Fig. ?? |
| $d_{x^2-y^2}$ -wave | $w^{(d)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$   | Fig. ?? |

Table 1.5 | Symmetry resolved self-consistency equations for the harmonics of  $w_{\mathbf{k}}$ .

where  $\tau^+ = (\tau^x + i\tau^y)/2$ . We used Eqns. (1.33), (1.34) and (1.36) as well as the thermal factor defined in Eq. (1.24). The derivation above holds for all symmetries, (including  $s$ -wave). Explicitly, five self-consistency equations are then derived and listed in Tab. 1.5.

Of these, the singlet channel admits for non-zero fluctuations of the superconducting order parameter uniquely in the  $s^*$  and  $d$  channels. This is because the gap function is inversion-symmetric and  $p_\ell$  channels equations integrate it with an antisymmetric function.

## 1.7 Full MFT hamiltonian and anisotropic pairing

Let now the total contraction energy

$$E_c^{(\text{SC})} \equiv - \left[ E_{\text{H}/U}^{(\text{SC})} + E_{\text{C}/U}^{(\text{SC})} + E_{\text{H}/V}^{(\text{SC})} + E_{\text{F}/V}^{(\text{SC})} + E_{\text{C}/V/\text{o.s.}}^{(\text{SC})} \right]$$

By composing all three channels we are able to derive the final form of the MFT gran-canonical hamiltonian,

$$\hat{H} - \mu \hat{N} \simeq \sum_{\mathbf{k} \in \text{BZ}} \tilde{\xi}_{\mathbf{k}} \left[ \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} + \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow} \right] - \sum_{\mathbf{k} \in \text{BZ}} \left[ \tilde{\Delta}_{\mathbf{k}} \hat{\phi}_{\mathbf{k}} + \tilde{\Delta}_{\mathbf{k}}^* \hat{\phi}_{\mathbf{k}}^\dagger \right] + E_a^{(\text{SC})} + E_c^{(\text{SC})}$$

having we included the “anticommutation energy” also defined for the conventional HM in Eq. (??),

$$E_a^{(\text{SC})} \equiv \sum_{\mathbf{k} \in \text{BZ}} \tilde{\xi}_{\mathbf{k}}$$

coming from fermionic anticommutation relations that are employed in order to get the kinetic term. The renormalized bands are

$$\tilde{\xi}_{\mathbf{k}} = \epsilon_{\mathbf{k}} + V \sum_{\gamma \in \{s^*, d\}} u^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} - \tilde{\mu} \quad (1.53)$$

and the full gap function by

$$\tilde{\Delta}_{\mathbf{k}} = -U w^{(s)} + V \sum_{\gamma} w^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} \quad (1.54)$$

We can identify the gap harmonics by

$$\tilde{\Delta}^{(s)} = -U w^{(s)} \quad \tilde{\Delta}^{(\gamma)} = V w^{(\gamma)} \quad \text{being} \quad \gamma \in \{s^*, p_x, p_y, d\}$$

As anticipated, the singlet channel admits three components only,

$$\text{Singlet channel} \quad : \quad \tilde{\Delta}_{\mathbf{k}} = \tilde{\Delta}^{(s)} + \tilde{\Delta}^{(s^*)}(\cos k_x + \cos k_y) + \tilde{\Delta}^{(d)}(\cos k_x - \cos k_y)$$

making the gap function purely real. An inverse statement holds for the triplet pairing, for which

$$\text{Triplet channel} \quad : \quad \tilde{\Delta}_{\mathbf{k}} = i\sqrt{2} \left[ \tilde{\Delta}^{(p_x)} \sin k_x + \tilde{\Delta}^{(p_y)} \sin k_y \right]$$

In this case, the gap is purely imaginary. When working in a specific channel, the gap function is either purely real or purely imaginary. This lets us discuss the shape of the renormalised ground-state.

## 1.8 The zero-temperature ground-state at half-filling

The HF approximation to the true ground-state in the SC phase is obtained by aligning the each pseudospin with the respective pseudo-field. Let the up and down states,

$$|\uparrow_{\mathbf{k}}\rangle \quad |\downarrow_{\mathbf{k}}\rangle$$

Using now Eq. (1.23), we have

$$\langle \downarrow_{\mathbf{k}} | \hat{s}_{\mathbf{k}}^z + \hat{\mathbb{I}}_{\mathbf{k}} | \downarrow_{\mathbf{k}} \rangle = 0$$

$$\langle \uparrow_{\mathbf{k}} | \hat{s}_{\mathbf{k}}^z + \hat{\mathbb{I}}_{\mathbf{k}} | \uparrow_{\mathbf{k}} \rangle = 2$$

so up to a phase

$$|\downarrow_{\mathbf{k}}\rangle \sim |\Omega_{\mathbf{k}}\rangle \quad |\uparrow_{\mathbf{k}}\rangle \sim \hat{\phi}_{\mathbf{k}}^\dagger |\Omega_{\mathbf{k}}\rangle$$

The derivations carried out for the AF phase in Sec. ?? can be re-used. We end up with the state:

$$|\Psi\rangle = \bigotimes_{\mathbf{k} \in \text{BZ}} \left[ \cos \tilde{\theta}_{\mathbf{k}} + e^{2i\tilde{\zeta}_{\mathbf{k}}} \sin \tilde{\theta}_{\mathbf{k}} \hat{\phi}_{\mathbf{k}}^\dagger \right] |\Omega_{\mathbf{k}}\rangle$$

which is an extension of the conventional BCS state. The phase factor angle is the gap “tilt” in the complex plane. This is proven by:

$$\begin{aligned} \langle \Psi | \hat{s}_{\mathbf{k}}^x | \Psi \rangle &= \langle \Psi | \hat{\phi}_{\mathbf{k}} + \hat{\phi}_{\mathbf{k}}^\dagger | \Psi \rangle \\ &= \sin \tilde{\theta}_{\mathbf{k}} \cos \tilde{\theta}_{\mathbf{k}} \left( e^{2i\tilde{\zeta}_{\mathbf{k}}} + e^{-2i\tilde{\zeta}_{\mathbf{k}}} \right) \\ &= \sin(2\tilde{\theta}_{\mathbf{k}}) \cos(2\tilde{\zeta}_{\mathbf{k}}) \end{aligned}$$

$$\begin{aligned} \langle \Psi | \hat{s}_{\mathbf{k}}^y | \Psi \rangle &= i \langle \Psi | \hat{\phi}_{\mathbf{k}} - \hat{\phi}_{\mathbf{k}}^\dagger | \Psi \rangle \\ &= i \sin \tilde{\theta}_{\mathbf{k}} \cos \tilde{\theta}_{\mathbf{k}} \left( e^{-2i\tilde{\zeta}_{\mathbf{k}}} - e^{2i\tilde{\zeta}_{\mathbf{k}}} \right) \\ &= \sin(2\tilde{\theta}_{\mathbf{k}}) \sin(2\tilde{\zeta}_{\mathbf{k}}) \end{aligned}$$

These results are coherent with Eqns. (1.33) as well as (1.34) and allow us to identify the phase with angle of the pseudofield in the  $xy$  plane.

## 1.9 SC free energy density

Free energy density of the SC phase is derived following the procedure used for the other two phases. As for the AF phase of Sec. ??, two fermionic bands are present; however, recalling the antiferromagnet entropy density of Eq. (??), the key differences are here given by the fact that the chemical potential inside the Fermi-Dirac factors is null and the sum is extended on the full BZ (but not later on spin index),

$$\begin{aligned} s^{(\pm)} \equiv -\frac{k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} & \left[ \left( 1 - f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \right) \ln \left( 1 - f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \right) \right. \\ & \left. + f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \ln f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \right] \quad (1.55) \end{aligned}$$

with identical meaning of the notation. Using the relation

$$\frac{1}{e^x + 1} + \frac{1}{e^{-x} + 1} = 1$$

we conclude that

$$s^{(-)} = s^{(+)} \implies s \equiv s^{(-)} + s^{(+)} = 2s^{(+)}$$

being  $s$  the total entropy density. The five contributions to free energy are:

1. Quasiparticles energy:

$$\begin{aligned} e_g^{(\text{SC})} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta] \\ &= -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \end{aligned}$$

2. Anticommutation energy:

$$\begin{aligned} e_a^{(\text{SC})} &\equiv \frac{E_a^{(\text{SC})}}{L_x L_y} \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{\xi}_{\mathbf{k}} \end{aligned}$$

3. Contraction energy:

$$e_c^{(\text{SC})} = -U \left[ n^2 + |w^{(s)}|^2 \right] + V \left[ 2zn^2 - \sum_{\gamma} |u^{(\gamma)}|^2 + \sum_{\gamma} |w^{(\gamma)}|^2 \right]$$

4. Entropic energy

$$e_s^{(\text{SC})} = \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln \left( 1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) - \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} f(\tilde{E}_{\mathbf{k}}; \beta, 0)$$

5. Chemical potential correction

$$e_n^{(\text{SC})} = \mu n$$

Compose everything

$$f_{\text{MFT}}^{(\text{SC})} \equiv e_g^{(\text{SC})} + e_a^{(\text{SC})} + e_c^{(\text{SC})} + e_s^{(\text{SC})} + e_n^{(\text{SC})}$$

and use that:

$$\tanh\left(\frac{x}{2}\right) + \frac{2}{e^x + 1} = 1$$

In the end we get:

$$\begin{aligned} f_{\text{MFT}}^{(\text{SC})} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[ \tilde{\xi}_{\mathbf{k}} - \tilde{E}_{\mathbf{k}} \right] + \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln \left( 1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \\ &\quad - U \left[ n^2 + |w^{(s)}|^2 \right] + V \left[ 2zn^2 - \sum_{\gamma} |u^{(\gamma)}|^2 + \sum_{\gamma} |w^{(\gamma)}|^2 \right] + \mu \end{aligned}$$

As was for the normal and AF phases, we recognize the chemical potential shift of Eq. (1.43),

$$\begin{aligned} f_{\text{MFT}}^{(\text{SC})} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[ \tilde{\xi}_{\mathbf{k}} - \tilde{E}_{\mathbf{k}} \right] + \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln \left( 1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \\ &\quad - U |w^{(s)}|^2 - V \left[ \sum_{\gamma} |u^{(\gamma)}|^2 - \sum_{\gamma} |w^{(\gamma)}|^2 \right] + \tilde{\mu} n \quad (1.56) \end{aligned}$$

### 1.9.1 The gap energy

Similarly to Sec. ??, let us prove the following equation about the gap energy:

$$-U|w^{(s)}|^2 + V \sum_{\gamma} |w^{(\gamma)}|^2 = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{|\tilde{\Delta}_{\mathbf{k}}|^2}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \quad (1.57)$$

Expand the right-hand side through Eq. (1.54),

$$\begin{aligned} \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{|\tilde{\Delta}_{\mathbf{k}}|^2}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[ -U w^{(s)*} + V \sum_{\gamma} w^{(\gamma)*} \varphi_{\mathbf{k}}^{(\gamma)} \right] \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \\ &= -U w^{(s)*} \times \overbrace{\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)}^{w^{(s)}} \\ &\quad + V \sum_{\gamma} w^{(\gamma)*} \times \underbrace{\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)}_{w^{(\gamma)}} \end{aligned}$$

having we used the self-consistency equations of Tab. 1.5. This proves Eq.(1.57), which is similar to Eq. (??) with one exception. Since the right-hand side of Eq.(1.57) is positive, for  $V > 0$  we have the constraint

$$\sum_{\gamma} |w^{(\gamma)}|^2 \geq M^{(\text{SC})} \quad \text{being} \quad M^{(\text{SC})} \equiv |w^{(s)}|^2 \frac{U}{V}$$

Now, as we state in Sec. 1.2.6, for the pure HM we have identically  $w^{(s)} = 0$ . The pure HM cannot exhibit BCS-like superconductivity. For the EHM instead we see that:

- In the strong coupling limit  $U \ll V$ , even small  $s$ -wave gaps impose larger  $\gamma \neq s$  components;
- The  $s$ -wave gap  $w^{(s)}$  cannot converge to a finite value if for any  $\gamma \neq s$  we have  $w^{(\gamma)} = 0$ .

Of course, Eq. (1.57) is also solved by a null gap on both sides: this indicates that when the constraint is impossible to fulfil self-consistently, the gap closes.

## 1.10 Theoretical constraints on the singlet SC phase HFPs

[To be continued...]


## 1.11 Discussion of numeric results (singlet SC)

In this section we discuss numeric results for the singlet pairing SC phase. The HFPs vector has five components:

$$\mathbf{v} = \begin{bmatrix} u^{(s^*)} \\ u^{(d)} \\ w^{(s)} \\ w^{(s^*)} \\ w^{(d)} \end{bmatrix} \quad \mathbf{v} \in \mathbb{R}^5$$

This phase is described by three RMPs:

$$\begin{aligned} \tilde{t}_{ij}^{(s^*)} &\equiv \left( t - \frac{u^{(s^*)} V}{2} \right) \varphi_{ij}^{(s^*)} \\ \tilde{t}_{ij}^{(d)} &\equiv -\frac{u^{(s^*)} V}{2} \varphi_{ij}^{(d)} \\ \tilde{\mu} &\equiv \mu + n(2zV - U) \end{aligned}$$



**Figure 1.1** | Dependency of the  $s$ -wave gap varying the doping  $\delta$  and the local repulsion  $U$ . In the top panel we compare three doping levels  $\delta = 0.0, 0.2, 0.4$  varying  $0 \leq U \leq 8$ . In the bottom panel we compare three repulsion values  $U = 0.0, 5.0, 10.0$  varying  $0 \leq \delta \leq 0.45$ .

We ran simulations at  $\beta = 100.0$  within the boundaries:

$$0 \leq U \leq 20 \quad 0 \leq V \leq 8 \quad 0 \leq \delta \leq 0.45$$

For the normal phase we did not vary  $U$ ; for the AF phase we did not vary  $\delta$ . Here, we need to vary both in order to explore the entire parametric space.

### 1.11.1 Multi-symmetry singlet superconductivity

In singlet channel, the most general gap function is a combination of  $s$ ,  $s^*$  and  $d$  components. However, whether the actual gap function  $\tilde{\Delta}_{\mathbf{k}}$  exhibits all these symmetries is not an immediate question to answer. **Superconductivity in cuprates is observed in  $d$ -wave channel.** In Fig. 1.1 we report our findings for the HFP  $w^{(d)}$ , recalling that the  $d$ -wave gap is given by:

$$\tilde{\Delta}^{(d)} = V w^{(d)}$$


This gap component is approximately independent of  $U$ , as is visible in the top panel (Fig. 1.1a), and increases with  $V$ . Its dependence on doping  $\delta$  is much different, as is visible in the bottom panel (Fig. 1.1b). At strong coupling ( $V \gtrsim 5$ )  $w^{(d)}$  decreases with doping, maintaining non-zero values up to almost unitary filling. Note that the data show an hole at  $\delta = 0.4$ . We were unable to understand why the algorithm failed to converge at that point, and fine-tuning of  $g$  did not help in this case.

### 1.11.2 Rotational SSB and nematic superconductivity


## 1.12 Theoretical constraints on the triplet SC phase HFPs

[To be continued...]






a |  $U$  dependency of  $w^{(s)}$  at various  $\delta$ s.




b |  $\delta$  dependency of  $w^{(s)}$  at various  $U$ s.

**Figure 1.2** | Dependency of the  $s$ -wave gap varying the doping  $\delta$  and the local repulsion  $U$ . In the top panel we compare three doping levels  $\delta = 0.0, 0.2, 0.4$  varying  $0 \leq U \leq 8$ . In the bottom panel we compare three repulsion values  $U = 0.0, 5.0, 10.0$  varying  $0 \leq \delta \leq 0.45$ .



a |  $U$  dependency of  $w^{(s)}$  at various  $\delta$ s.



b |  $\delta$  dependency of  $w^{(s)}$  at various  $U$ s.

**Figure 1.3** | Dependency of the  $s$ -wave gap varying the doping  $\delta$  and the local repulsion  $U$ . In the top panel we compare three doping levels  $\delta = 0.0, 0.2, 0.4$  varying  $0 \leq U \leq 8$ . In the bottom panel we compare three repulsion values  $U = 0.0, 5.0, 10.0$  varying  $0 \leq \delta \leq 0.45$ .

### 1.13 Discussion of numeric results (triplet SC)

[To be continued...]

# Bibliography

- [1] Avinash Singh and Zlatko Tešanović. “Collective excitations in a doped antiferromagnet”. en. In: *Physical Review B* 41.1 (Jan. 1990), pp. 614–631. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.41.614](https://doi.org/10.1103/PhysRevB.41.614). URL: <https://link.aps.org/doi/10.1103/PhysRevB.41.614> (visited on 12/12/2025).
- [2] Avinash Singh, Zlatko Tešanović, and Ju H. Kim. “Doped Hubbard antiferromagnet: Instability and effective interactions”. en. In: *Physical Review B* 44.14 (Oct. 1991), pp. 7757–7759. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.44.7757](https://doi.org/10.1103/PhysRevB.44.7757). URL: <https://link.aps.org/doi/10.1103/PhysRevB.44.7757> (visited on 12/12/2025).
- [3] Alexander Wietek et al. “Stripes, Antiferromagnetism, and the Pseudogap in the Doped Hubbard Model at Finite Temperature”. en. In: *Physical Review X* 11.3 (July 2021), p. 031007. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.11.031007](https://doi.org/10.1103/PhysRevX.11.031007). URL: <https://link.aps.org/doi/10.1103/PhysRevX.11.031007> (visited on 12/12/2025).
- [4] Eugene Kogan and Godfrey Gumbs. “Green’s Functions and DOS for Some 2D Lattices”. en. In: *Graphene* 10.01 (2021), pp. 1–12. ISSN: 2169-3439, 2169-3471. DOI: [10.4236/graphene.2021.101001](https://doi.org/10.4236/graphene.2021.101001). URL: <https://www.scirp.org/journal/paperinformation?paperid=104591> (visited on 03/26/2026).
- [5] J. G. Bednorz and K. A. Muller. “Possible high-T<sub>c</sub> superconductivity in the Ba-La-Cu-O system”. en. In: *Zeitschrift für Physik B Condensed Matter* 64.2 (June 1986), pp. 189–193. ISSN: 1431-584X. DOI: [10.1007/BF01303701](https://doi.org/10.1007/BF01303701). URL: <https://doi.org/10.1007/BF01303701> (visited on 04/03/2026).
- [6] Liangzi Deng et al. “Ambient-pressure 151 K superconductivity in HgBa<sub>2</sub>Ca<sub>2</sub>Cu<sub>3</sub>O<sub>8</sub> via pressure quench”. In: *Proceedings of the National Academy of Sciences* 123.11 (Mar. 2026), e2536178123. DOI: [10.1073/pnas.2536178123](https://doi.org/10.1073/pnas.2536178123). URL: <https://www.pnas.org/doi/10.1073/pnas.2536178123> (visited on 04/03/2026).
- [7] B. Keimer et al. “From quantum matter to high-temperature superconductivity in copper oxides”. en. In: *Nature* 518.7538 (Feb. 2015), pp. 179–186. ISSN: 0028-0836, 1476-4687. DOI: [10.1038/nature14165](https://doi.org/10.1038/nature14165). URL: <https://www.nature.com/articles/nature14165> (visited on 03/15/2025).
- [8] D. J. Scalapino. “A common thread: The pairing interaction for unconventional superconductors”. en. In: *Reviews of Modern Physics* 84.4 (Oct. 2012), pp. 1383–1417. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.84.1383](https://doi.org/10.1103/RevModPhys.84.1383). URL: <https://link.aps.org/doi/10.1103/RevModPhys.84.1383> (visited on 04/03/2026).
- [9] Hari Padma et al. “Beyond-Hubbard pairing in a cuprate ladder”. en. In: *Physical Review X* 15.2 (May 2025), p. 021049. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.15.021049](https://doi.org/10.1103/PhysRevX.15.021049). URL: <https://link.aps.org/doi/10.1103/PhysRevX.15.021049> (visited on 04/03/2026).
- [10] Antoine Georges et al. “Dynamical mean-field theory of strongly correlated fermion systems and the limit of infinite dimensions”. en. In: *Reviews of Modern Physics* 68.1 (Jan. 1996), pp. 13–125. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.68.13](https://doi.org/10.1103/RevModPhys.68.13). URL: <https://link.aps.org/doi/10.1103/RevModPhys.68.13> (visited on 03/15/2025).
- [11] J. E. Hirsch. “Two-dimensional Hubbard model: Numerical simulation study”. In: *Phys. Rev. B* 31 (7 Apr. 1985), pp. 4403–4419. DOI: [10.1103/PhysRevB.31.4403](https://doi.org/10.1103/PhysRevB.31.4403). URL: <https://link.aps.org/doi/10.1103/PhysRevB.31.4403>.

- [12] Robin Scholle et al. “Comprehensive mean-field analysis of magnetic and charge orders in the two-dimensional Hubbard model”. In: *Phys. Rev. B* 108 (3 July 2023), p. 035139. DOI: [10.1103/PhysRevB.108.035139](https://doi.org/10.1103/PhysRevB.108.035139). URL: <https://link.aps.org/doi/10.1103/PhysRevB.108.035139>.
- [13] Zhangkai Cao et al. “Dominant  $p$ -wave pairing induced by nearest-neighbor attraction in the square-lattice extended Hubbard model”. In: *Phys. Rev. B* 111 (2 Jan. 2025), p. 024509. DOI: [10.1103/PhysRevB.111.024509](https://doi.org/10.1103/PhysRevB.111.024509). URL: <https://link.aps.org/doi/10.1103/PhysRevB.111.024509>.
- [14] Daniel P. Arovas et al. “The Hubbard Model”. In: *Annual Review of Condensed Matter Physics* 13. Volume 13, 2022 (2022), pp. 239–274. ISSN: 1947-5462. DOI: [10.1146/annurev-conmatphys-031620-102024](https://doi.org/10.1146/annurev-conmatphys-031620-102024). URL: <https://www.annualreviews.org/content/journals/10.1146/annurev-conmatphys-031620-102024>.
- [15] Kazuyoshi Yoshimi, Takeo Kato, and Hideaki Maebashi. “Enhanced Spin Susceptibility toward the Charge-Ordering Transition in a Two-Dimensional Extended Hubbard Model”. In: *Journal of the Physical Society of Japan* 78.10 (Oct. 2009), p. 104002. ISSN: 0031-9015. DOI: [10.1143/JPSJ.78.104002](https://doi.org/10.1143/JPSJ.78.104002). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.78.104002> (visited on 04/17/2026).
- [16] Ashvin Vishwanath. “Dirac Nodes and Quantized Thermal Hall Effect in the Mixed State of d-wave Superconductors”. In: *Physical Review B* 66.6 (Aug. 2002). arXiv:cond-mat/0104213, p. 064504. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.66.064504](https://doi.org/10.1103/PhysRevB.66.064504). URL: <http://arxiv.org/abs/cond-mat/0104213> (visited on 01/05/2026).
- [17] P. Senarath Yapa et al. “Mixed-symmetry superconductivity and the energy gap”. en. In: *Physica C: Superconductivity and its Applications* 633 (June 2025), p. 1354719. ISSN: 09214534. DOI: [10.1016/j.physc.2025.1354719](https://doi.org/10.1016/j.physc.2025.1354719). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0921453425000723> (visited on 01/06/2026).
- [18] Joel Hutchinson and Frank Marsiglio. “Mixed temperature-dependent order parameters in the extended Hubbard model”. en. In: *Journal of Physics: Condensed Matter* 33.6 (Feb. 2021), p. 065603. ISSN: 0953-8984, 1361-648X. DOI: [10.1088/1361-648X/abc801](https://doi.org/10.1088/1361-648X/abc801). URL: <https://iopscience.iop.org/article/10.1088/1361-648X/abc801> (visited on 01/06/2026).
- [19] C. Zhou and H. J. Schulz. “Density of states and tunneling spectra in two-dimensional d-wave superconductors”. In: *Physical Review B* 45.13 (Apr. 1992). Publisher: American Physical Society, pp. 7397–7401. DOI: [10.1103/PhysRevB.45.7397](https://doi.org/10.1103/PhysRevB.45.7397). URL: <https://link.aps.org/doi/10.1103/PhysRevB.45.7397> (visited on 01/06/2026).
- [20] James F. Annett. “Symmetry of the order parameter for high-temperature superconductivity”. In: *Advances in Physics* 39.2 (Apr. 1990). Publisher: Taylor & Francis eprint: <https://doi.org/10.1080/00018739000101481>, pp. 83–126. ISSN: 0001-8732. DOI: [10.1080/00018739000101481](https://doi.org/10.1080/00018739000101481). URL: <https://doi.org/10.1080/00018739000101481> (visited on 01/06/2026).
- [21] R. Micnas, J. Ranninger, and S. Robaszkiewicz. “Superconductivity in narrow-band systems with local nonretarded attractive interactions”. In: *Reviews of Modern Physics* 62.1 (Jan. 1990). Publisher: American Physical Society, pp. 113–171. DOI: [10.1103/RevModPhys.62.113](https://doi.org/10.1103/RevModPhys.62.113). URL: <https://link.aps.org/doi/10.1103/RevModPhys.62.113> (visited on 01/06/2026).
- [22] Manfred Sgrist and Kazuo Ueda. “Phenomenological theory of unconventional superconductivity”. In: *Reviews of Modern Physics* 63.2 (Apr. 1991). Publisher: American Physical Society, pp. 239–311. DOI: [10.1103/RevModPhys.63.239](https://doi.org/10.1103/RevModPhys.63.239). URL: <https://link.aps.org/doi/10.1103/RevModPhys.63.239> (visited on 01/07/2026).
- [23] Anna E. Böhrer et al. “Nematicity and nematic fluctuations in iron-based superconductors”. en. In: *Nature Physics* 18.12 (Dec. 2022), pp. 1412–1419. ISSN: 1745-2481. DOI: [10.1038/s41567-022-01833-3](https://doi.org/10.1038/s41567-022-01833-3). URL: <https://www.nature.com/articles/s41567-022-01833-3>.

- [24] Simons Collaboration on the Many-Electron Problem et al. “Absence of Superconductivity in the Pure Two-Dimensional Hubbard Model”. In: *Physical Review X* 10.3 (July 2020), p. 031016. DOI: [10.1103/PhysRevX.10.031016](https://doi.org/10.1103/PhysRevX.10.031016). URL: <https://link.aps.org/doi/10.1103/PhysRevX.10.031016> (visited on 03/21/2026).
- [25] Jan Zaanen and Olle Gunnarsson. “Charged magnetic domain lines and the magnetism of high- $T_c$  oxides”. In: *Physical Review B* 40.10 (Oct. 1989), pp. 7391–7394. DOI: [10.1103/PhysRevB.40.7391](https://doi.org/10.1103/PhysRevB.40.7391). URL: <https://link.aps.org/doi/10.1103/PhysRevB.40.7391> (visited on 03/21/2026).
- [26] Masaru Kato et al. “Soliton Lattice Modulation of Incommensurate Spin Density Wave in Two Dimensional Hubbard Model -A Mean Field Study-”. In: *Journal of the Physical Society of Japan* 59.3 (Mar. 1990), pp. 1047–1058. ISSN: 0031-9015. DOI: [10.1143/JPSJ.59.1047](https://doi.org/10.1143/JPSJ.59.1047). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.59.1047> (visited on 03/21/2026).
- [27] Eva Pavarini et al., eds. *Many-body physics: from Kondo to Hubbard: lecture notes of the Autumn School on Correlated Electrons 2015: at Forschungszentrum Jülich, 21-25 September 2015*. en. Band 5. Meeting Name: Autumn School on Correlated Electrons. Forschungszentrum Jülich, 2015. ISBN: 978-3-95806-074-6.
- [28] Piers Coleman. *Introduction to Many-Body Physics*. Cambridge University Press, 2015.
- [29] Giuseppe Grosso and Giuseppe Pastori Parravicini. *Solid State Physics*. Second Edition. Academic Press, 2014.
- [30] Gabriele Giuliani and Giovanni Vignale. *Quantum Theory of the Electron Liquid*. Cambridge University Press, 2005.
- [31] Michele Fabrizio. *A Course in Quantum Many-Body Theory*. Springer, 2022.