



UNIVERSITÀ DI PISA

Dipartimento di Fisica “E. Fermi”

Corso di Laurea Magistrale in Fisica

Mott localisation, antiferromagnetism and nematic superconductivity in the extended Hubbard model with non-local pairing

Relatore:

Prof. Giacomo Mazza

Candidato:

Alessandro Gori

*Anno accademico 2025/2026
Draft: May 20, 2026*

Contents

1	Introduction	3
1.1	Cuprates and the square Hubbard model	4
1.1.1	Mott insulation in cuprates	4
1.1.2	The insurgence of antiferromagnetism	5
1.1.3	Unconventional superconductivity	6
1.2	The Extended Hubbard model	6
1.3	Outline of this thesis	7
2	Introduction to Hartree-Fock methods	9
2.1	Variational principle and gaussian states	9
2.2	The HF variational method	10
2.2.1	Properties of the target quadratic hamiltonian	11
2.2.2	Self-consistent variational solution	12
2.2.3	Operative hamiltonian approximation	15
2.3	The iHF algorithm	15
2.3.1	Fixed density simulations and the role of μ	17
2.3.2	The convergence scheme	18
3	Phase transitions and spontaneous symmetry breaking in the extended Hubbard model	21
3.1	Symmetries of the extended Hubbard model on the square lattice	21
3.1.1	Lattice symmetries	21
3.1.2	Spin and charge symmetries	22
3.1.3	Particle-hole symmetry at half-filling	26
3.2	The HF method and symmetry breaking	28
3.2.1	Local repulsion	29
3.2.2	Non-local attraction	31
3.3	The EHM in reciprocal space	33
3.4	Square lattice spatial symmetry channels	36
4	Mott localisation in the normal phase	41
4.1	The normal phase and its symmetries	41
4.2	Application of Hartree-Fock methods to the normal phase	42
4.2.1	Effects of the Hartree terms: chemical potential renormalisation	43
4.2.2	Effects of the Fock terms: bare bands renormalisation	44
4.3	The HF hamiltonian and its properties	47
4.3.1	Nematicity: d -wave bands and anisotropic hopping	48
4.3.2	The entropy density of the normal ground-state	50
4.3.3	The free energy density of the normal ground-state	50
4.4	Constraints on the self-consistent solution	52
4.5	Discussion of numeric results	55
4.5.1	Mott localisation	56
4.5.2	Solution free energy and entropy densities	57
4.5.3	Convergence problems and mixing fine-tuning	58

CONTENTS

5	Antiferromagnetic phase	61
5.1	The AF phase and its symmetries	61
5.1.1	Symmetries of the commensurate AF Ansatz	62
5.1.2	The AF Ansatz in reciprocal space	64
5.2	The AF phase in the conventional HM	65
5.2.1	Derivation of the HF hamiltonian	66
5.2.2	Diagonalisation of the HF hamiltonian	67
5.3	Generalisation to the EHM	73
5.3.1	Effects of the non-local Hartree terms: chemical potential renormalisation	76
5.3.2	Effects of the Fock terms: gap and bare bands renormalisation	78
5.4	The HF hamiltonian and its properties	84
5.4.1	Gap complexification and AF flavours	84
5.4.2	The entropy density of the AF ground-state	85
5.4.3	The free energy density of the AF ground-state	86
5.5	Analysis of the solution	87
5.6	Discussion of numeric results	89
5.6.1	Mott localisation in the AF phase	89
5.6.2	Mott-induced enhancement of magnetisation	92
5.6.3	Solution free energy density	93
5.6.4	Fixed-point interpretation of HF solution	94
6	Superconducting instability	97
6.1	The SC phase and its symmetries	97
6.2	The SC phase in the conventional HM	98
6.2.1	Derivation of the HF hamiltonian	98
6.2.2	Diagonalisation of the HF hamiltonian	100
6.3	Generalisation to the EHM	104
6.3.1	Effects of non-local Hartree terms: chemical potential renormalisation	105
6.3.2	Effects of Fock terms: bare bands renormalisation	106
6.3.3	Effects of non-local Cooper terms: generalised pairing	107
6.4	The HF hamiltonian and its properties	110
6.4.1	Gap symmetries and anisotropic pairing	110
6.4.2	The entropy density of the SC ground-state	111
6.4.3	The free energy density of the SC ground-state	112
6.5	Analysis of the solution	113
6.6	Discussion of numeric results for the singlet channel	115
6.6.1	Multi-symmetry singlet superconductivity	115
6.6.2	Rotational SSB and nematic superconductivity	120
6.6.3	Isotropic bands shrinking	122
6.6.4	Solution free energy and entropy densities	124
6.7	A final note	126
A	Superexchange and virtual hopping in Hubbard lattices	127
A.1	Singlet pairing in Hubbard lattices	127
A.2	Direct exchange and virtual hopping	128
A.3	Superexchange in cuprates	130
B	Gaussian states	131
B.1	Slater determinants	131
B.2	Wick's theorem applied to Slater determinants	132
B.3	Generalisation to Bogoliubov-Valatin states	133
C	Connection with mean-field theories	137
C.1	Mean-field theory	137
C.2	Hartree-Fock methods	137
D	Antiferromagnetism in the HM	141

D.1	The AF ground-state for the conventional HM	141
D.2	Explicit AF ground state at half-filling and $\beta \rightarrow \infty$	142
D.3	Instability of the commensurate AF phase	143
D.4	iHF results	144
D.5	Analytic expression of self-consistent magnetisation at half-filling	145
E	Theoretical constraints on the HFPs	149
E.1	Normal phase	149
E.1.1	Mixed $s^* \oplus d$ bands	149
E.1.2	Bands shrinking in C_4	151
E.1.3	Bands shrinking in $C_2/\mathbb{Z}_2$	152
E.2	AF phase	156
E.2.1	Bands shrinking in C_4	157
E.2.2	Selection rules on the magnetic HFPs in sAF phase	157
E.2.3	Selection rules on the magnetic HFPs in aAF phase	162
E.2.4	Bands shrinking in $C_2/\mathbb{Z}_2$	163
F	Mixing optimization in HM model	167
F.1	Map analysis for s -wave superconductivity in HM	167
F.1.1	Constraints for algorithmic evolution	167
	Bibliography	169

Chapter 1

Introduction

A major breakthrough in Condensed Matter Physics was the discovery of superconductivity in copper oxide compounds (often referred to as “cuprates”). Until 1986, when Bednorz and Muller found evidence of a superconducting transition in Ba-La-Cu-O compounds around 30 K [9], superconductivity had been observed up to critical temperatures (T_c) of few Kelvins. The new class of copper oxides rapidly took over in the field of high- T_c superconductivity, increasing critical temperatures to around 130 K at ambient pressure. Today, the highest observed value is $T_c \approx 151$ K at ambient pressure on the cuprate $\text{HgBa}_2\text{Ca}_2\text{Cu}_3\text{O}_{8+\delta}$ [10]. It is worth mentioning that this result was obtained in recent months after almost 30 years from the previous record.

Cuprates are very rich materials, with a peculiar phase diagram that we sketch in Fig. 1.1. Undoped copper oxides are Mott insulators and show an antiferromagnetic ground state, which is disrupted both by electronic and hole doping. As charge carriers are inserted, various phases emerge: strange metal and pseudogap regimes, and most notably superconductivity [5]. The latter, when discovered, was particularly unexpected: indeed, copper oxides based compounds are at ordinary conditions very poor conductors. The understanding of such a complex phase diagram and the interplay of all physical phenomena active inside these materials is, to date, subject of great scientific effort and still an open problem.

In this thesis we investigate both theoretically and numerically the interplay of Mott insulation, antiferromagnetism and superconductivity in the context of a simplified model of two-dimensional cuprates. This introduction is organised as follows: in Sec. 1.1 we describe the general features of cuprates and how these are described through the Hubbard model. In Sec. 1.2 we discuss how the Hubbard model fails to address some observed phenomena, and present an extended version with an added non-local attraction. Finally, in Sec. 1.3 we give an outline of this thesis.




Figure 1.1 | Sketch of the phase diagram of cuprates. These curves result from graphic elaboration and do not refer to a specific physical dataset.




Figure 1.2 | Schematics of a typical layered copper oxide compound. The inter layers are represented as solid planes, while the electrically active layers as a square grid hosting copper on its nodes and oxygen on its edges.

1.1 Cuprates and the square Hubbard model

The crystal structure of copper oxide superconductors is given by a stacking of 2D CuO_2 sheets separated by inert layers of insulating materials. We give a sketch of the material in Fig. 1.2. The intermediate layers act as reservoirs of electrons (or holes) and dope the active CuO_2 sheets [8]. Due to the presence of these intermediate inert layers, the CuO_2 sheets are weakly coupled to each other, and can be studied independently. In these sheets, copper and oxygens are arranged as a checkerboard lattice: oxygen atoms form a square lattice, and in the center of each cell is placed a copper atom. At null doping, due to the high electronegativity of oxygens, copper atoms are ionised and left with one hole in their d shell, while oxygen atoms reach a closed shell configuration. Therefore, at low energy the kinetics of electrons is given by the hopping among the d orbitals of Cu^{2+} ions, arranged in a planar square lattice [30].

Electrons repel each other through Coulomb interaction. The simplest model capable of describing this situation is the single-band Hubbard model [34],

$$\hat{H}_{\text{HM}} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \quad (t, U > 0), \quad (1.1)$$

being t the hopping energy and U the Coulomb repulsion. Sum is intended on the spin index $\sigma \in \{\uparrow, \downarrow\}$ and the sites i, j of a planar square lattice $\mathcal{L}$. The notation $\langle ij \rangle$ indicates nearest neighbours (NNs). Note that the Coulomb repulsion is limited to be a on-site interaction.

The Hubbard model captures the low-energy properties of copper oxides [30, 33] and is capable of showing a rich variety of physical phenomena [11, 17]. Indeed, despite its simplicity, it can describe Mott insulation, antiferromagnetism and superconductivity. We give a brief description of these three phenomena in next subsections.

1.1.1 Mott insulation in cuprates

Conventional band theory describes the electronic properties of materials on the assumption that electron-electron interaction is negligible. Within this framework, a material forms a band insulator whenever the conduction band fills up. Due to Pauli principle, electrons fill all states of the band and have no room left for low-energy scattering, hence the filled band is completely ineffective for conduction. On the other hand, if the conduction band is partially filled, the material is a metal and conducts electricity [8]. In general, band theory predicts a metallic behaviour if the number of electrons per unit cell is odd; an insulating behaviour if is even.

In undoped copper oxides, the conduction band is not completely filled. Indeed, copper is ionised in the crystal and each Cu^{2+} ion accommodates one hole in its most external orbital. Band theory would predict a metallic nature for these compounds, but the observed state is insulating. The reason for the failure of band theory lies in the fact that electron-electron interaction is not negligible. Cuprates are Mott insulators, not band insulators: their insulating properties do not derive from a single-particle band being filled up and separated by a gap from another band at higher energy; instead, from an amplification of Coulomb repulsion that “freezes” electrons on the lattice sites [5]. One can picture this situation as a sort of traffic jam, where electrons do not move because the energetic cost of having two electrons on the same site would be too high [29], as we show in Fig. [Missing]. This produces an insulating behaviour: electrons are completely localised and conductivity is low.

The Hubbard model describes naturally Mott physics [11]. Consider for instance the two-sites Hubbard model,

$$\hat{H} = -t \left(\hat{c}_{1\uparrow}^\dagger \hat{c}_{2\uparrow} + \hat{c}_{2\downarrow}^\dagger \hat{c}_{1\downarrow} \right) + U (\hat{n}_{1\uparrow} \hat{n}_{1\downarrow} + \hat{n}_{2\uparrow} \hat{n}_{2\downarrow}), \quad (1.2)$$

where the subscript indicates the site index. In the strong coupling limit $U \gg t$, we can treat the kinetic terms perturbatively. Indeed, let E_i be the energy given by having i electrons on one site. At order zero in perturbation theory, the difference in energy between the state with one site doubly occupied and the other empty, and the state with two singly occupied sites is

$$E_2 + E_0 - 2E_1 = U.$$

The zeroth order ground-state, then, is a combination of states where both sites are singly occupied. Insertion of the kinetic terms as a perturbative correction does not change the scenario (see App. A for details).

In the Hubbard model, the transition from a metallic state to a Mott insulator is controlled by the strength of the Coulomb repulsion U , and impacts visibly the density of states (DoS), which undergoes significant changes [40]. Let us sketch the general behaviour: to start, let $U = 0$ and consider a lattice with one electron per site (half-filling). Electrons are free and do not interact, therefore move freely across the lattice. Due to Pauli exclusion principle, they form a Fermi sea that occupies the lower half a metallic band with a width W . As we increase the value of U , it becomes increasingly harder for electrons to move. When U becomes sufficiently large (with respect to W), the electronic density of states splits in two bands. The lower band, filled, accommodates all completely localised electrons; in the upper band, empty, are all the states where a site gets doubly occupied. This splitting is of order $\sim U$. At intermediate values of U , the DoS presents a peculiar three-peaks shape, which is the result of a mixture of the two Mott bands, and a residual of the original free band around the Fermi energy. This is shown in Fig. [Missing].

[Figure: Hubbard bands]

One of the signatures of a Mott transition, thus, is the flattening of the bare single-particle bands.

1.1.2 The insurgence of antiferromagnetism

It is widely known that the ground-state of undoped copper oxides exhibits antiferromagnetic ordering [5, 11, 14]. The insurgence of this phase can be connected to Mott physics in the Hubbard model. Indeed, in the strong coupling regime $U \gg t$, the Hubbard hamiltonian is effectively approximated by the Heisenberg hamiltonian with antiferromagnetic coupling [31],

$$\hat{H}_{\text{HM}} \simeq \frac{4t^2}{U} \sum_{\langle ij \rangle} \hat{\mathbf{S}}_i \cdot \hat{\mathbf{S}}_j,$$

where $\hat{\mathbf{S}}_i$ is the spin operator on site i .

Antiferromagnetism is established through the “super-exchange mechanism”, which takes advantage of the fact that electrons are localised on-site. This is best explained on the two-sites Hubbard model toy model of Eq. (1.2): as we explained in Sec. 1.1.1, the kinetic term can be

treated as a perturbation to the fully-localised ground-state, which favours the formation of a spin singlet between the two sites (see App. A for details). A spin singlet

The idea that singlet pairing across nearest-neighbouring sites could explain the insurgence of antiferromagnetism has been explored. A many-body state resulting from a superposition of spin singlets across different sites has been proposed [35, 36] to capture the relation between antiferromagnetism and Mott physics.

1.1.3 Unconventional superconductivity

Injection of charge (electrons or holes) in copper oxide compounds disrupts antiferromagnetic ordering and establishes a variety of peculiar phases [5]. Among these, the high temperature superconducting phase is significantly different from that described by Bardeen-Cooper-Schrieffer (BCS) theory. Because of this, is denominated “unconventional”.

In BCS theory, couples of electrons are “glued” together by an effective attraction that is mediated by lattice phonons. This mechanism is referred to as “Cooper pairing”. The resulting state is a condensate of Cooper pairs, with an energy spectrum that is fully gapped and thus no low-energy excitations. The gap $\Delta_{\mathbf{k}}$, as a function of momentum, is isotropic (s -wave) and represents the energy cost of breaking a Cooper pair glued by phonons.

In copper oxides, superconductivity is established with significantly different features. Pairing is still active, but is not mediated by phonons. Most notably, the superconducting gap is not isotropic but changes sign under rotations of $\pi/2$ on the plane [11]. Because of this, we talk about $d_{x^2-y^2}$ -wave superconductivity. A consequence is the appearance of gapless quasiparticle excitations in the energy spectrum. Indeed, a $d_{x^2-y^2}$ -wave gap has nodal lines (continuous sets of points, in reciprocal space, where it is null) which, in a square lattice geometry, result in a set of nodes around which dispersion is linearised (Dirac cones) [42]. Moreover, Mott physics and $d_{x^2-y^2}$ -wave superconductivity are related. Indeed, such a spatial structure of the gap function is understood to be a manifestation of the fact that electrons are spatially correlated in such a way to avoid occupying the same site [11].

The source of pairing in cuprates is still object of debate. In the phase diagram of copper oxides, the $d_{x^2-y^2}$ -wave superconducting region “emerges” from the antiferromagnetic region, as doping is increased at low temperature. Hence, it has been proposed that the same Mott-induced spin fluctuations that lead to the formation of the antiferromagnetic state could act as an effective Cooper pairing [42] [37–39]. Electron-phonon coupling has been studied too, but recent developments suggest that it could act as a source of additional interactions – namely, a nearest neighbours attraction mediated by phonons [32]. This idea leads to the next section, where we present and motivate the extension to the Hubbard model that we studied.

1.2 The Extended Hubbard model

The Hubbard model of Eq. (1.1) has been long considered a fair description of the main features of copper oxides [11]. There are evidences, however, that the model could be missing some interactions that must be accounted to describe d -wave superconductivity in cuprates, for instance a nearest-neighbours attraction [32]. Moreover, recent developments have shown numerically that inclusion of a non-local attraction between neighbouring sites is necessary to reproduce some features of the excitation spectrum of doped cuprates [12].

Moreover, it has been known since early Hartree-Fock calculations that adding a non-local attraction contributes to $d_{x^2-y^2}$ -wave superconductivity on the square lattice [24]. With these motivation, we propose the following line of reasoning:

1. We insert “artificially” a non-local attraction in the Hubbard model, knowing that it leads to $d_{x^2-y^2}$ -wave superconductivity;
2. We study the resulting model, searching for manifestations of Mott physics and understanding how the newly added interaction relates to it;
3. We analyse how antiferromagnetism and superconductivity are impacted by the findings of the step above.

To summarise: we know that Mott insulation originates $d_{x^2-y^2}$ -wave superconductivity. It is worth asking if the opposite implication holds.

We define the Extended Hubbard model (EHM) by adding a non-local attraction to the Hubbard model of Eq. (1.1),

$$\hat{H}_{\text{EHM}} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'} \quad (t, U, V > 0). \quad (1.3)$$

The newly added interaction is independent of spin index and acts on nearest neighbours. We will study the ground state of this model by the means of theoretical and numerical Hartree-Fock methods, varying the strength of the interactions U , V and electronic density. In particular, we will explore the competition of three phases (namely: normal, antiferromagnetic and superconducting) as we vary the model parameters and assess which one is the most stable. We will limit to finite square lattices and low temperatures.

1.3 Outline of this thesis

In this section we give an outline of next chapters.

- Chap. 2 We introduce the Hartree-Fock method for a generic fermionic quartic hamiltonian, and discuss how it is used to find a variational approximation to its ground state. Then, we explain how this problem can be re-stated in terms of a finite set of self-consistency equations. In the end, we propose an iterative algorithmic scheme to solve these equations.
- Chap. 3 We discuss the symmetries of the extended Hubbard model of Eq. (1.3). Then we give a definition of the three phases (normal, antiferromagnetic, superconducting) and describe each based on which of the original symmetries is spontaneously broken.
- Chap. 4 We apply the Hartree-Fock method to the normal phase and derive the self-consistency equations. Then we present numeric results that compose evidence of V -induced Mott localisation.
- Chap. 5 We apply the Hartree-Fock method to the antiferromagnetic phase and derive the self-consistency equations. First, we discuss antiferromagnetism in the conventional Hubbard model of Eq. (1.1), hence setting $V = 0$. Then we let $V > 0$ and discuss how the Hartree-Fock solution changes. In the end, we present numeric results that show how antiferromagnetic ordering is enhanced (at half-filling) by V -induced Mott localisation.
- Chap. 6 We apply the Hartree-Fock method to the antiferromagnetic phase and derive the self-consistency equations. We discuss the impossibility (at Hartree-Fock level) of s -wave superconductivity in the conventional Hubbard model of Eq. (1.1), setting $V = 0$. Then we let $V > 0$ and discuss how the presence of a non-local attraction gives rise to a stable superconducting solution. In the end, we discuss numeric results that show:
 - a) The competition among (extended) s -wave and $d_{x^2-y^2}$ -wave superconductivity, and how the two symmetries can mix in a region of parameters space;
 - b) The insurgence of nematic effects (namely, breaking of planar rotational symmetry) related to the mixing of gap symmetries;
 - c) The presence of Mott localisation in superconducting phase.

Chap. ?? We summarise our findings and *thank for all the fish*.

Appendices are organised as follows:

App. A We discuss the superexchange mechanism already introduced in Sec. 1.1.2.

- App. [B](#) We define the set of gaussian states, which are used in the Hartree-Fock method discussed in Chap. [2](#).
- App. [C](#) We explain how Hartree-Fock methods and mean-field theory are related.
- App. [D](#) We give more details about antiferromagnetism in the conventional Hubbard model.
- App. [E](#) We derive a set of theoretical constraints on the Hartree-Fock solution for the normal and AF phase, derived respectively in Chaps. [4](#) and [5](#).
- App. [F](#) We discuss one concrete example of algorithmic optimisation that we used extensively for all phases. The example is discussed in the context of (absence of) s -wave superconductivity in the conventional Hubbard model.

Chapter 2

Introduction to Hartree-Fock methods

In this chapter we give a general introduction to the Hartree-Fock (HF) variational method and its numerical implementation. In Quantum Mechanics, variational methods allow to approximate the ground state of a many-body hamiltonian by minimising the energy computed for a specific subspace of physical states. In particular, in the HF method we approximate the real ground-state with the best possible many-body state that is an eigenstate of a non-interacting fermionic hamiltonian.

Despite its simplicity, the method is very powerful for describing various types of quantum many-body phenomena and, in particular, spontaneous symmetry breaking (SSB). The description of HF method in this chapter is largely based on [8, 44].

2.1 Variational principle and gaussian states

Let $\hat{H}$ be a generic hamiltonian acting on the Hilbert space $\mathcal{H}$. For any many-body state $|\Psi\rangle \in \mathcal{H}$, the energy is given by the expectation value (EV)

$$E[\Psi] = \langle \Psi | \hat{H} | \Psi \rangle.$$

Let $|\Psi_0\rangle$ be the ground state of $\hat{H}$, and $E_0 \equiv E[\Psi_0]$. Then

$$\forall |\Psi\rangle \in \mathcal{H} \quad : \quad E[\Psi] \geq E_0,$$

since the ground-state minimises energy, by definition. Let $\mathcal{S} \subset \mathcal{H}$ be a subspace of states, ad let $|\Psi\rangle$ be a state inside that locally minimises energy,

$$|\Psi\rangle \in \mathcal{S} \quad \frac{\delta E[\Psi]}{\delta |\Psi\rangle} = 0.$$

If the true ground state belongs to $\mathcal{S}$, the found local minimum coincides with $|\Psi_0\rangle$; otherwise, approximates it. In Fig. 2.1 we give a pictorial representation of this idea: the state $|\Psi\rangle \in \mathcal{S}$ is an approximation, found by constrained minimisation, of the true ground-state.

In HF methods, we identify $\mathcal{S}$ with the set of so-called “gaussian states”. These are all the ground-states of quadratic hamiltonians, namely the hamiltonians that are bilinear in the fermionic creations operators (see App. B for details). The restriction to gaussian states significantly simplifies the computation of expectation values. Indeed, thanks to Wick’s theorem, the expectation value of quartic operators exactly decomposes as

$$\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta \rangle = \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\delta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\gamma \rangle}_{\text{Hartree}} - \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\gamma \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\delta \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \langle \hat{c}_\gamma \hat{c}_\delta \rangle}_{\text{Cooper}}. \quad (2.1)$$

The subscripts indicate the standard names to the different types of contractions, that we adopt throughout the text.

Operatively, one can think of the HF method as an approximation of the the hamiltonian with a non-interacting hamiltonian

$$\hat{H} \stackrel{\text{HF}}{\simeq} \hat{H}_{\text{HF}} = \sum_{\alpha} E_{\alpha} \hat{\gamma}_{\alpha}^{\dagger} \hat{\gamma}_{\alpha},$$




Figure 2.1 | Pictorial representation of the search for the best approximation $|\Psi\rangle$ to the true ground-state inside the subspace $\mathcal{S} \subset \mathcal{H}$.

where $\hat{\gamma}_\alpha$ are fermionic operators determined by the minimisation procedure. These new operators satisfy Fermi anticommutation rules

$$\{\hat{\gamma}_\alpha, \hat{\gamma}_\beta^\dagger\} = \delta_{\alpha\beta},$$

and are connected to the conventional electronic operators $\hat{c}_\alpha$ through an unitary map. See App. B for details.

Let us stress that the operatorial approximation $\overset{\text{HF}}{\simeq}$ is a shorthand notation that must be intended as follows: $\hat{H}_{\text{HF}}$ is that non-interacting operator whose ground-state is the best approximation to the ground-state of $\hat{H}$. This does not mean that in general the two operators are related.

2.2 The HF variational method

We now discuss the implementation of the variational HF method for the generic quartic hamiltonian,

$$\hat{H} = \sum_{\alpha,\beta} A_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} B_{\alpha\beta\gamma\delta} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta, \quad (2.2)$$

where

$$A_{\alpha\beta} = A_{\beta\alpha}^* \quad \text{and} \quad B_{\alpha\beta\gamma\delta} = B_{\delta\gamma\beta\alpha}^*, \quad (2.3)$$

to ensure hermiticity. Moreover, the following equation must hold:

$$B_{\alpha\beta\gamma\delta} = B_{\beta\alpha\delta\gamma}, \quad (2.4)$$

due to the property $\hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta = \hat{c}_\beta^\dagger \hat{c}_\alpha^\dagger \hat{c}_\delta \hat{c}_\gamma$, which follows from Fermi anticommutation rules.

Our aim is to approximate the above hamiltonian with a non-interacting hamiltonian. The most general quadratic hamiltonian is given by

$$\hat{H} \overset{\text{HF}}{\simeq} \hat{H}_{\text{HF}} \equiv E_0 + \sum_{\alpha,\beta} k_{\alpha\beta} \hat{\gamma}_\alpha^\dagger \hat{\gamma}_\beta. \quad (2.5)$$

As is specified in Sec. B.3, γ and c operators must be linked by an unitary map. To find the best approximation to the original hamiltonian means to identify:

- the offset E_0 ;
- the fermionic operators $\hat{\gamma}_\alpha, \hat{\gamma}_\alpha^\dagger$;

- the $k_{\alpha\beta}$ coefficients.

Note that, in terms of the algebraic solution, E_0 is irrelevant. To determine it correctly is only relevant in order to get the correct approximation of the actual ground-state energy. Let us discuss briefly the features of $\hat{H}_{\text{HF}}$ and then propose a solution.

2.2.1 Properties of the target quadratic hamiltonian

In this section we enumerate some useful properties of $\hat{H}_{\text{HF}}$. We will use also the following notation:

$$\hat{a}_\alpha^- \equiv \hat{c}_\alpha \quad \text{and} \quad \hat{a}_\alpha^+ \equiv \hat{c}_\alpha^\dagger.$$

Due to linearity of the unitary mapping that connects $\hat{c}$ and $\hat{\gamma}$ operators, the operator $\hat{H}_{\text{HF}}$ of Eq. (2.5) can also be expressed as the most general quadratic hamiltonians for electronic operators:

$$\hat{H}_{\text{HF}} = E_0 + \sum_{\alpha,\beta} \sum_{p,q} h_{\alpha\beta}^{pq} \hat{a}_\alpha^p \hat{a}_\beta^q. \quad (2.6)$$

By changing basis from $\hat{\gamma}$ to $\hat{c}$ operators, we moved all our ignorance into the new $h_{\alpha\beta}^{pq}$ matrix elements. These satisfy the relations:

$$h_{\alpha\beta}^{++} = \left(h_{\beta\alpha}^{--} \right)^*, \quad (2.7)$$

$$h_{\beta\alpha}^{+-} = -h_{\alpha\beta}^{-+}. \quad (2.8)$$

Because of these properties, we actually need to determine just half of the fields, since the other half will be uniquely determined.

Proof 2.1 — Symmetry properties of the tensor $h_{\alpha\beta}^{pq}$.

Let us start by proving Eq. (2.7). The HF hamiltonian can be represented through Nambu spinors as

$$\begin{aligned} \hat{H}_{\text{HF}} &= E_0 + \hat{\Phi}^\dagger \mathbf{h} \hat{\Phi} \\ &= E_0 + \begin{bmatrix} \hat{\mathbf{c}} \\ \hat{\mathbf{c}}^\dagger \end{bmatrix}^\dagger \begin{bmatrix} h^{+-} & h^{++} \\ h^{--} & h^{-+} \end{bmatrix} \begin{bmatrix} \hat{\mathbf{c}} \\ \hat{\mathbf{c}}^\dagger \end{bmatrix}. \end{aligned}$$

Our hamiltonian must be hermitian. We conclude that

$$\hat{H}_{\text{HF}}^\dagger = \hat{H}_{\text{HF}} \quad \implies \quad h^{++} = \left[h^{--} \right]^\dagger,$$

which reads, element-wise:

$$h_{\alpha\beta}^{++} = \left(h_{\beta\alpha}^{--} \right)^*.$$

This proves Eq. (2.7).

Eq. (2.8) is proven as follows. We can set, without loss of generality,

$$\text{Tr } h^{+-} + \text{Tr } h^{-+} = 0.$$

This passage is guaranteed by the fact that if $\mathbf{h}$ is not traceless, we can redefine it by subtracting its trace and including the latter in the energy shift E_0 . Then

$$\begin{aligned} \hat{H}_{\text{HF}} &= E_0 + \sum_{\alpha,\beta} \left[h_{\alpha\beta}^{+-} \hat{c}_\alpha^\dagger \hat{c}_\beta + h_{\alpha\beta}^{-+} \hat{c}_\alpha \hat{c}_\beta^\dagger \right] + (\text{same sign terms}) \\ &= E_0 + \sum_{\alpha,\beta} \left[h_{\alpha\beta}^{+-} \left(\delta_{\alpha\beta} - \hat{c}_\beta \hat{c}_\alpha^\dagger \right) + h_{\alpha\beta}^{-+} \left(\delta_{\alpha\beta} - \hat{c}_\beta^\dagger \hat{c}_\alpha \right) \right] + (\text{same sign terms}) \\ &= E_0 - \sum_{\alpha,\beta} \left[h_{\beta\alpha}^{+-} \hat{c}_\alpha \hat{c}_\beta^\dagger + h_{\beta\alpha}^{-+} \hat{c}_\alpha^\dagger \hat{c}_\beta \right] + (\text{same sign terms}). \end{aligned}$$

In the second line we got rid of the traces sum and exchanged indices $\alpha \leftrightarrow \beta$. Confronting last and first row we conclude:

$$h_{\beta\alpha}^{+-} = -h_{\alpha\beta}^{-+},$$

which proves Eq. (2.8).

2.2.2 Self-consistent variational solution

This section gives the operational method to obtain the variational solution. We follow and extend the argument proposed by Fabrizio [44]. Let

$$\Delta_{\alpha\beta}^{pq} \equiv \langle \hat{a}_{\alpha}^p \hat{a}_{\beta}^q \rangle,$$

thus, explicitly

$$\Delta_{\alpha\beta}^{+-} = \langle \hat{c}_{\alpha}^{\dagger} \hat{c}_{\beta} \rangle, \quad \Delta_{\alpha\beta}^{-+} = \langle \hat{c}_{\alpha} \hat{c}_{\beta}^{\dagger} \rangle, \quad \Delta_{\alpha\beta}^{++} = \langle \hat{c}_{\alpha}^{\dagger} \hat{c}_{\beta}^{\dagger} \rangle, \quad \Delta_{\alpha\beta}^{--} = \langle \hat{c}_{\alpha} \hat{c}_{\beta} \rangle. \quad (2.9)$$

The first two are “normal densities”; the second two are “anomalous densities”. The reason for such names is the fact that these expectation values are non-zero if total charge conservation is broken. Conjugation of the last two gives a symmetry relation similar to Eq. (2.7),

$$\left(\Delta_{\alpha\beta}^{++} \right)^{\dagger} = \Delta_{\beta\alpha}^{--}. \quad (2.10)$$

Here $\Delta_{\alpha\beta}^{pq}$ are functions of the fields $h_{\alpha\beta}^{pq}$. These quantities satisfy the equation:

$$\sum_{\alpha,\beta} \sum_{p,q} h_{\alpha\beta}^{pq} \frac{\partial \Delta_{\alpha\beta}^{pq}}{\partial h_{\mu\nu}^{rs}} = 0. \quad (2.11)$$

Let us prove this.

Proof 2.2 — Proof of Eq. (2.11).

The HF ground-state energy is given by

$$\begin{aligned} E_{\text{HF}} &= \langle \hat{H}_{\text{HF}} \rangle \\ &= E_0 + \sum_{\alpha,\beta} \sum_{p,q} h_{\alpha\beta}^{pq} \Delta_{\alpha\beta}^{pq}, \end{aligned}$$

where the expectation value is taken over the (up to now unknown) HF ground-state. Deriving it, we obtain

$$\begin{aligned} \frac{\partial E_{\text{HF}}}{\partial h_{\mu\nu}^{rs}} &= \frac{\partial}{\partial h_{\mu\nu}^{rs}} \sum_{\alpha,\beta} \sum_{p,q} h_{\alpha\beta}^{pq} \Delta_{\alpha\beta}^{pq} \\ &= \Delta_{\mu\nu}^{rs} + \sum_{\alpha,\beta} \sum_{p,q} h_{\alpha\beta}^{pq} \frac{\partial \Delta_{\alpha\beta}^{pq}}{\partial h_{\mu\nu}^{rs}}. \end{aligned} \quad (2.12)$$

From Hellman-Feynman theorem [8], we know that for a parametric hamiltonian it holds:

$$\begin{aligned} \frac{\partial E_{\text{HF}}}{\partial h_{\mu\nu}^{rs}} &= \langle \Psi | \frac{\partial \hat{H}_{\text{HF}}}{\partial h_{\mu\nu}^{rs}} | \Psi \rangle \\ &= \langle \Psi | \frac{\partial}{\partial h_{\mu\nu}^{rs}} \sum_{\alpha,\beta} \sum_{p,q} h_{\alpha\beta}^{pq} \hat{a}_{\alpha}^p \hat{a}_{\beta}^q | \Psi \rangle \\ &= \langle \hat{a}_{\mu}^r \hat{a}_{\nu}^s \rangle \\ &= \Delta_{\mu\nu}^{rs}. \end{aligned} \quad (2.13)$$

Eqs. (2.12) and (2.13), combined, imply

$$\sum_{\alpha,\beta} \sum_{p,q} h_{\alpha\beta}^{pq} \frac{\partial \Delta_{\alpha\beta}^{pq}}{\partial h_{\mu\nu}^{rs}} = 0,$$

which ends the proof.

We now use Eq. (2.11) to minimise the variational energy $E = \langle \hat{H} \rangle$. Performing Wick's contractions, the variational energy becomes

$$\begin{aligned} E &= \langle \hat{H} \rangle \\ &= \sum_{\alpha,\beta} A_{\alpha\beta} \langle \hat{c}_{\alpha}^{\dagger} \hat{c}_{\beta} \rangle + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} B_{\alpha\beta\gamma\delta} \langle \hat{c}_{\alpha}^{\dagger} \hat{c}_{\beta}^{\dagger} \hat{c}_{\gamma} \hat{c}_{\delta} \rangle \\ &= \sum_{\alpha,\beta} A_{\alpha\beta} \langle \hat{a}_{\alpha}^{+} \hat{a}_{\beta}^{-} \rangle + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} B_{\alpha\beta\gamma\delta} \left(\langle \hat{a}_{\alpha}^{+} \hat{a}_{\beta}^{+} \rangle \langle \hat{a}_{\gamma}^{-} \hat{a}_{\delta}^{-} \rangle - \langle \hat{a}_{\alpha}^{+} \hat{a}_{\gamma}^{-} \rangle \langle \hat{a}_{\beta}^{+} \hat{a}_{\delta}^{-} \rangle + \langle \hat{a}_{\alpha}^{+} \hat{a}_{\delta}^{-} \rangle \langle \hat{a}_{\beta}^{+} \hat{a}_{\gamma}^{-} \rangle \right) \\ &= \sum_{\alpha,\beta} A_{\alpha\beta} \Delta_{\alpha\beta}^{+-} + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} B_{\alpha\beta\gamma\delta} \left(\Delta_{\alpha\beta}^{++} \Delta_{\gamma\delta}^{--} - \Delta_{\alpha\gamma}^{+-} \Delta_{\beta\delta}^{+-} + \Delta_{\alpha\delta}^{+-} \Delta_{\beta\gamma}^{+-} \right) \\ &= \sum_{\alpha,\beta} A_{\alpha\beta} \Delta_{\alpha\beta}^{+-} + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} (B_{\alpha\gamma\delta\beta} - B_{\alpha\gamma\beta\delta}) \Delta_{\alpha\beta}^{+-} \Delta_{\gamma\delta}^{+-} + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} B_{\alpha\beta\gamma\delta} \Delta_{\alpha\beta}^{++} \Delta_{\gamma\delta}^{--}, \end{aligned} \quad (2.14)$$

where in the second line, we inserted the relations (2.9). The HF approximation to the real ground state it obtained as the ground-state of a quadratic hamiltonian whose fields $h_{\alpha\beta}^{pq}$ satisfy the following relations:

$$h_{\alpha\beta}^{+-} = \frac{1}{2} \left[A_{\alpha\beta} + \sum_{\gamma,\delta} (B_{\alpha\gamma\delta\beta} - B_{\alpha\gamma\beta\delta}) \Delta_{\gamma\delta}^{+-} \right], \quad (2.15)$$

$$h_{\alpha\beta}^{++} = \sum_{\gamma,\delta} \frac{B_{\alpha\beta\gamma\delta} \Delta_{\gamma\delta}^{--}}{2}. \quad (2.16)$$

These equations, operationally, give the instructions to build the best possible fermionic non-interacting hamiltonian to reproduce our original hamiltonian. These equations are proven as follows:

Proof 2.3 — Proof of Eqs. (2.15), (2.16).

Consider E as expressed in Eq. (2.14). Upon taking the derivative, we obtain

$$\begin{aligned} \frac{\partial E}{\partial h_{\mu\nu}^{rs}} &= \sum_{\alpha,\beta} A_{\alpha\beta} \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} (B_{\alpha\gamma\delta\beta} - B_{\alpha\gamma\beta\delta}) \left(\frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} \Delta_{\gamma\delta}^{+-} + \Delta_{\alpha\beta}^{+-} \frac{\partial \Delta_{\gamma\delta}^{+-}}{\partial h_{\mu\nu}^{rs}} \right) \\ &\quad + \frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} B_{\alpha\beta\gamma\delta} \left(\frac{\partial \Delta_{\alpha\beta}^{++}}{\partial h_{\mu\nu}^{rs}} \Delta_{\gamma\delta}^{--} + \Delta_{\alpha\beta}^{++} \frac{\partial \Delta_{\gamma\delta}^{--}}{\partial h_{\mu\nu}^{rs}} \right), \end{aligned}$$

which, by using Eq. (2.4), becomes:

$$\begin{aligned} \frac{\partial E}{\partial h_{\mu\nu}^{rs}} &= \sum_{\alpha,\beta} \left[A_{\alpha\beta} + \sum_{\gamma,\delta} (B_{\alpha\gamma\delta\beta} - B_{\alpha\gamma\beta\delta}) \Delta_{\gamma\delta}^{+-} \right] \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} \\ &\quad + \sum_{\alpha,\beta,\gamma,\delta} \frac{B_{\alpha\beta\gamma\delta} \Delta_{\gamma\delta}^{--}}{2} \frac{\partial \Delta_{\alpha\beta}^{++}}{\partial h_{\mu\nu}^{rs}} + \sum_{\alpha,\beta,\gamma,\delta} \frac{B_{\alpha\beta\gamma\delta} \Delta_{\alpha\beta}^{++}}{2} \frac{\partial \Delta_{\gamma\delta}^{--}}{\partial h_{\mu\nu}^{rs}}. \end{aligned} \quad (2.17)$$

Let us define

$$C_{\alpha\beta} \equiv \frac{1}{2} \left[A_{\alpha\beta} + \sum_{\gamma,\delta} (B_{\alpha\gamma\delta\beta} - B_{\alpha\gamma\beta\delta}) \Delta_{\gamma\delta}^{+-} \right] \quad \text{and} \quad D_{\alpha\beta} \equiv \sum_{\gamma,\delta} \frac{B_{\alpha\beta\gamma\delta} \Delta_{\gamma\delta}^{--}}{2}. \quad (2.18)$$

From Fermi anticommutation rules, we have

$$\Delta_{\alpha\beta}^{+-} = \delta_{\alpha\beta} - \Delta_{\beta\alpha}^{-+} \quad \Rightarrow \quad \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} = - \frac{\partial \Delta_{\beta\alpha}^{-+}}{\partial h_{\mu\nu}^{rs}},$$

which implies

$$\begin{aligned} \sum_{\alpha,\beta} 2C_{\alpha\beta} \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} &= \sum_{\alpha,\beta} \left(C_{\alpha\beta} \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} + C_{\alpha\beta} \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} \right) \\ &= \sum_{\alpha,\beta} \left(C_{\alpha\beta} \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} - C_{\beta\alpha} \frac{\partial \Delta_{\alpha\beta}^{-+}}{\partial h_{\mu\nu}^{rs}} \right). \end{aligned}$$

Moreover, using Eqs. (2.3), (2.10),

$$\sum_{\alpha,\beta} \frac{B_{\alpha\beta\gamma\delta} \Delta_{\alpha\beta}^{++}}{2} = \sum_{\alpha,\beta} \left(\frac{B_{\delta\gamma\beta\alpha} \Delta_{\beta\alpha}^{--}}{2} \right)^*,$$

and, recognising $D_{\gamma\delta}$ from Eq. (2.18), we conclude

$$\sum_{\alpha,\beta,\gamma,\delta} \frac{B_{\alpha\beta\gamma\delta} \Delta_{\alpha\beta}^{++}}{2} \frac{\partial \Delta_{\gamma\delta}^{--}}{\partial h_{\mu\nu}^{rs}} = \sum_{\gamma,\delta} D_{\delta\gamma}^* \frac{\partial \Delta_{\gamma\delta}^{--}}{\partial h_{\mu\nu}^{rs}}.$$

We recollect everything and impose the solution to be a stationary point of energy,

$$\frac{\partial E}{\partial h_{\mu\nu}^{rs}} \stackrel{!}{=} 0.$$

The notation “ $\stackrel{!}{=}$ ” indicates, throughout the thesis, that we are imposing some condition. Eq. (2.17) becomes

$$\frac{\partial E}{\partial h_{\mu\nu}^{rs}} = \sum_{\alpha,\beta} \left(C_{\alpha\beta} \frac{\partial \Delta_{\alpha\beta}^{+-}}{\partial h_{\mu\nu}^{rs}} - C_{\beta\alpha} \frac{\partial \Delta_{\alpha\beta}^{-+}}{\partial h_{\mu\nu}^{rs}} + D_{\alpha\beta} \frac{\partial \Delta_{\alpha\beta}^{++}}{\partial h_{\mu\nu}^{rs}} + D_{\beta\alpha}^* \frac{\partial \Delta_{\alpha\beta}^{--}}{\partial h_{\mu\nu}^{rs}} \right) \stackrel{!}{=} 0. \quad (2.19)$$

Comparing this equation with Eq. (2.11), we conclude that it must hold

$$h_{\alpha\beta}^{+-} = C_{\alpha\beta} \quad \text{and} \quad h_{\alpha\beta}^{++} = D_{\alpha\beta}.$$

Recalling the explicit form of $C_{\alpha\beta}$ and $D_{\alpha\beta}$ from Eq. (2.18), we prove Eqs. (2.15), (2.16).

We notice that the right-hand side of both Eqs. (2.15), (2.16) depends on the fields themselves, as $\Delta_{\alpha\beta}^{pq}$ are defined as expectation values of two-points fermionic operators over the ground-state $|\Psi\rangle$. For a given set of fields $h_{\alpha\beta}^{pq}$, the latter is determined (up to some unitary transformation) as the ground-state of a fermionic non-interacting model. In other words, being $\mathbf{h}$ the tensor containing all fields,

$$\Delta_{\alpha\beta}^{pq}[\mathbf{h}] = \langle \Psi[\mathbf{h}] | \hat{a}_{\alpha}^p \hat{a}_{\beta}^q | \Psi[\mathbf{h}] \rangle. \quad (2.20)$$

The three Eqs. (2.15), (2.16) and (2.20) define the solution of the HF variational method. Specifically, we need to determine $\Delta_{\alpha\beta}^{pq}$ such that, for given fields, Eq. (2.20) is respected.

The list of parameters $\{\Delta_{\alpha\beta}^{pq}\}$ to be determined self-consistently may be not that short: by looking at Eq. (2.20), the indices α, β, p, q are spanning a large Hilbert space. However, this equation can be manipulated and transformed, reducing the set of parameters to be determined

self-consistently to a very manageable number. For each phase we define these last as “Hartree-Fock parameters” (HFPs): a set of objects self-consistently determined by a set of equivalent, but transformed, version of Eq. (2.20).

The great advantage of this scheme is that, with an appropriate choice of the Hilbert space basis (for example, moving to reciprocal space for a translationally-invariant model), we can implement easily model symmetries.

2.2.3 Operative hamiltonian approximation

In this section we give an alternative expression for Eqs. (2.15), (2.16), by using an equivalent (but more straightforward) operatorial approximation. For a generic two-body quartic operator, the HF approximation is given by the decomposition

$$\begin{aligned} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\beta \hat{c}_\alpha &\stackrel{\text{HF}}{\simeq} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \langle \hat{c}_\beta \hat{c}_\alpha \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \hat{c}_\beta \hat{c}_\alpha - \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \langle \hat{c}_\beta \hat{c}_\alpha \rangle & (\text{Cooper}) \\ &- \hat{c}_\alpha^\dagger \hat{c}_\beta \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle - \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \hat{c}_\beta^\dagger \hat{c}_\alpha + \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle & (\text{Fock}) \\ &+ \hat{c}_\alpha^\dagger \hat{c}_\alpha \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \hat{c}_\beta^\dagger \hat{c}_\beta - \langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle. & (\text{Hartree}) \end{aligned}$$

The proof of this statement is given in App. C, where we also argue the interpretation of HF method as a mean-field theory (MFT). Let us summarise this section through a practical rule to decompose the quartic part of the local repulsion $\hat{H}_U$ and the non-local attraction $\hat{H}_V$ in the EHM, by applying Eq. (C.9).

Local repulsion. Applying the HF method, the local repulsion $\hat{H}_U$ is approximated by the quadratic operator:

$$\begin{aligned} \hat{H}_U &= U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} & (2.21) \\ &\stackrel{\text{HF}}{\simeq} U \sum_i \left(\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \langle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \rangle + \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} - \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle \langle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \rangle \right. & (\text{Hartree}) \\ &- \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow} \langle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\uparrow} \rangle - \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow} \rangle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\uparrow} + \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow} \rangle \langle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\uparrow} \rangle & (\text{Fock}) \\ &\left. + \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \langle \hat{c}_{i\downarrow} \hat{c}_{i\uparrow} \rangle + \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \rangle \hat{c}_{i\downarrow} \hat{c}_{i\uparrow} - \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \rangle \langle \hat{c}_{i\downarrow} \hat{c}_{i\uparrow} \rangle \right). & (\text{Cooper}) \end{aligned}$$

Non-local attraction. Applying the HF method, the local repulsion $\hat{H}_V$ is approximated by the quadratic operator:

$$\begin{aligned} \hat{H}_V &= -V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'} & (2.22) \\ &\stackrel{\text{HF}}{\simeq} -V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle \right. & (\text{Hartree}) \\ &- \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} \rangle - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rangle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} \rangle & (\text{Fock}) \\ &\left. + \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \langle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \rangle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \rangle \langle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle \right). & (\text{Cooper}) \end{aligned}$$

We will use these decomposition referring to each sector in dedicated chapters.

2.3 The iHF algorithm

In this section we describe the algorithm we used to compute Eq. (2.20). We refer to this technique as “iterative Hartree Fock” (iHF). Let $\mathbf{v}$ be the vector containing all HFPs. Suppose we have N




Figure 2.2 | Schematisation of the self-consistent Hartree Fock method described in Sec. 2.3. The effective model is solved, the solution is used to estimate self-consistently the Hartree Fock vector $\mathbf{v}$, which is then used to update iteratively the model itself. The notation $\mathbf{v}_0$ indicates the initialiser HFPs vector, $\mathbf{v}_c$ the converged HFPs vector.

HFPs v_i

$$\mathbf{v} \in \mathbb{C}^N \quad \text{with} \quad \mathbf{v} \equiv \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_N \end{bmatrix}.$$

In order to get a physical solution, each HFP must satisfy a self-consistency equation similar to Eq. (2.20). Each equation has the structure:

$$v_i = (\text{Some non-linear function of } \mathbf{v}).$$

Let A be the non-linear operator collecting all these functions. The problem of finding a self-consistent solution can be reformulated as follows: $\mathbf{v}$ represents a physical solution if it is a fixed point of the map A ,

$$A: \mathbb{C}^N \rightarrow \mathbb{C}^N \quad : \quad A(\mathbf{v}) = \mathbf{v} \quad \implies \quad \mathbf{v} \text{ is a solution.} \quad (2.23)$$

We implement the search for a self-consistent solution as follows: first, we obtain an analytic expression for A . Then we choose an initialiser $\mathbf{v}_0$ and apply A to it, obtaining a new HFPs vector. Iterating this procedure, we “follow” the non-linear map A until we converge. Specifically:

1. We start with a generic quartic hamiltonian:

$$\hat{H} = \sum_{\alpha\beta} A_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta + \sum_{\alpha\beta\gamma\delta} B_{\alpha\beta\gamma\delta} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta.$$

We choose the model parameters and set the electronic density target;

2. Following the discussion of Sec. 2.2, we approximate analytically $\hat{H}$ with a quadratic hamiltonian

$$\hat{H} \simeq \hat{H}_{\text{HF}} \equiv \sum_{\alpha\beta} \left[h_{\alpha\beta}^{+-} \hat{c}_\alpha^\dagger \hat{c}_\beta - h_{\beta\alpha}^{+-} \hat{c}_\alpha \hat{c}_\beta^\dagger + h_{\alpha\beta}^{++} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger + \left(h_{\beta\alpha}^{++} \right)^* \hat{c}_\alpha \hat{c}_\beta \right],$$

with the fields given by Eqs. (2.15), (2.16);

3. We impose selection rules on the fields $h_{\alpha\beta}^{+-}$, $h_{\alpha\beta}^{++}$, eventually eliminating those not respecting the symmetry sector we are targeting. For example, if we choose not to break total charge conservation, we set $\forall \alpha, \beta: h_{\alpha\beta}^{++} = 0$;
4. We choose a starting value for the HFPs vector $\mathbf{v}_0$;

5. We set $\mathbf{v} = \mathbf{v}_0$ and enter the iterative scheme of Fig. 2.2:
 - a) Get a numeric approximation of the hamiltonian for the current value of $\mathbf{v}$, $\hat{H}_{\text{HF}}[\mathbf{v}]$;
 - b) Determine the value $\mu[\mathbf{v}]$ giving the desired density;
 - c) Diagonalise the effective hamiltonian $\hat{H}_{\text{HF}}[\mathbf{v}] - \mu[\mathbf{v}]\hat{N}$, and obtain the approximated ground-state $|\Psi[\mathbf{v}]\rangle$;
 - d) Compute $A(\mathbf{v})$ using self-consistency equations;
 - e) If the algorithm converged, exit; otherwise, update the value of $\mathbf{v}$ repeat from point a).

In the following subsections we describe technical details concerning the chemical potential fixing and the convergence criteria used for the practical implementation of the above iterative scheme.

2.3.1 Fixed density simulations and the role of μ

In this thesis all derivations and simulations are carried out at fixed target density. This is done by fixing the chemical potential μ during iterations of the algorithm. This must be done iteratively: changing at each step the HFPs vector, we are actively modifying the approximate hamiltonian. Chemical potential must be determined accordingly to maintain the required density.

For a given choice of HFPs vector and chemical potential, the ground state $|\Psi[\mathbf{v}, \mu]\rangle$ is uniquely determined; then we compute density as

$$n(\mathbf{v}, \mu) \equiv \frac{1}{\mathcal{V}} \sum_{\alpha} \langle \Psi[\mathbf{v}, \mu] | \hat{c}_{\alpha}^{\dagger} \hat{c}_{\alpha} | \Psi[\mathbf{v}, \mu] \rangle ,$$

being $\mathcal{V}$ the volume. Suppose we fix density to a value $n_0 \in [0, 1]$. At step i , naming the current HFPs vector $\mathbf{v}_i$, μ_i (the chemical potential at current step) is determined as

$$\mu_i \equiv \min_{\mu} |n(\mathbf{v}_i, \mu) - n_0| . \quad (2.24)$$

The density n must be a monotonically increasing function of μ : then we can translate the above equation, which would inspire a minimisation procedure, to a much simpler node-search rule:

$$\mu_i = \mu \mid_{\delta n(\mathbf{v}_i, \mu)=0} \quad \text{being} \quad \delta n(\mathbf{v}_i, \mu) \equiv n(\mathbf{v}_i, \mu) - n_0 .$$

In order to find where the function changes sign (which is, the position of μ_i) we use a basic bisection algorithm:

1. Set an arbitrary tolerance $\Delta n > 0$ and a shifting parameter $\Delta \mu > 0$;
2. Choose starting lower and upper bounds $\mu_L < \mu_U$;
3. Compute $\delta n(\mathbf{v}_i, \mu_L)$. Evaluate:
 - if $|\delta n(\mathbf{v}_i, \mu_L)| \leq \Delta n$, set $\mu_i = \mu_L$ and halt;
 - if $\delta n(\mathbf{v}_i, \mu_L) < -\Delta n$ do nothing;
 - if $\delta n(\mathbf{v}_i, \mu_L) > \Delta n$, shift the lower bound down, $\mu_L \rightarrow \mu_L - \Delta \mu$ and repeat from step 3;
4. Compute $\delta n(\mathbf{v}_i, \mu_U)$. Evaluate:
 - if $|\delta n(\mathbf{v}_i, \mu_U)| \leq \Delta n$, set $\mu_i = \mu_U$ and halt;
 - if $\delta n(\mathbf{v}_i, \mu_U) > \Delta n$ do nothing;
 - if $\delta n(\mathbf{v}_i, \mu_U) < -\Delta n$, shift the upper bound up, $\mu_U \rightarrow \mu_U + \Delta \mu$ and repeat from step 4;
5. At this point, the the lower bound underestimates density while the upper bound overestimates. Start iterations:
 - a) Compute the midpoint,

$$\mu_M = \frac{\mu_L + \mu_U}{2} .$$

b) Evaluate:

- If $|\delta n(\mathbf{v}_i, \mu_M)| \leq \Delta n$, set $\mu_i = \mu_M$ and halt;
- If $\delta n(\mathbf{v}_i, \mu_M) > \Delta n$, set $\mu_U = \mu_M$ and repeat from point a);
- If $\delta n(\mathbf{v}_i, \mu_M) < -\Delta n$, set $\mu_L = \mu_M$ and repeat from point a);

The bottleneck of the procedure is our ability to compute expectation values. This depends on how much we are able to simplify analytically the computations for each phase.

As will turn out in the derivation, chemical potential will be renormalised by interaction. This means that applying self-consistent HF methods to the EHM will produce an effective hamiltonian containing a shift to the chemical potential,

$$\mu \rightarrow \mu + F[\mathbf{v}] \equiv \tilde{\mu}[\mathbf{v}],$$

being F some function of the HFPs. The chemical potential is an example of Renormalised Model Parameter (RMP).

Note that at step i of the iHF algorithm, our bisection procedure determines $\tilde{\mu}_i$, not μ_i . This consideration is relevant when dealing with ground state energy estimation. Let us abstract from the EHM and iHF, and consider a generic hamiltonian $\hat{H}$. To find the ground state $|\Psi\rangle$ means precisely to solve

$$\frac{\delta}{\delta \langle \Psi |} \langle \Psi | \hat{H} | \Psi \rangle = 0.$$

Now, working with a fixed particles number N means we are performing a constrained minimisation (which is, taking the functional derivative) on the composite operator

$$\hat{H} - \mu(\hat{N} - N) \equiv \hat{K} + \mu N,$$

being μ the Lagrange multiplier, $\hat{N}$ the number operator and $\hat{K} \equiv \hat{H} - \mu\hat{N}$. Finding the constrained ground state parametrised by a vector $\mathbf{v}$ now means to solve

$$\mathbf{v} \longrightarrow |\Psi[\mathbf{v}]\rangle \quad : \quad \begin{cases} \frac{\delta}{\delta \langle \Psi |} \langle \Psi | \hat{K} + \mu N | \Psi \rangle \Big|_{|\Psi\rangle=|\Psi[\mathbf{v}]\rangle} = 0, \\ \langle \Psi[\mathbf{v}] | \hat{N} | \Psi[\mathbf{v}] \rangle = N. \end{cases}$$

In the first line μN is derived to zero; the constraint is imposed in the second line. Energy is given by:

$$\begin{aligned} \langle \hat{H} \rangle &= \langle \Psi[\mathbf{v}] | \hat{K} | \Psi[\mathbf{v}] \rangle + \mu N \\ &= \Omega[\mathbf{v}] + (\tilde{\mu}[\mathbf{v}] - F[\mathbf{v}])N, \end{aligned}$$

being Ω the expectation value (EV) of $\hat{K}$. To correctly compute energy, we need to add $+\mu N$; since we determine $\tilde{\mu}[\mathbf{v}] \neq \mu$, everything needs to be expressed in terms of it. Using this relation, when the algorithm has converged, we can estimate ground-state energy.

2.3.2 The convergence scheme

Let us discuss briefly the convergence criterion of our algorithm and the iteration mechanism. The iHF scheme we design is a sequential loop that runs up to a logical halt condition. In order to specify such condition, we need to define our notion of “convergence”. Recall the definition of $A(\mathbf{v})$, the non-linear map in HFPs space whose components are given by a set of self-consistency equations. We claim to have reached convergence when $\mathbf{v}$ comes “close enough” to a fixed point of the map A , as in Eq. (2.23). Explicitly, we stop whenever the following logic condition is satisfied for each component of $\mathbf{v} = (v_1, v_2, \dots)$:

$$\forall v_j \quad : \quad |v_j - A_j(\mathbf{v})| \leq \Delta v_j,$$

being $A(\mathbf{v})$ the map image of the HFP vector $\mathbf{v}$, and A_j its j -th component. The tolerance Δv_j is arbitrary. If just one component of $\mathbf{v}$ is mapped “too far” (with respect to Δv_j) the step is rejected

and iterations continue. In this text we deal in with quantities of order 10^{-1} , so convergence values of order $10^{-3} \div 10^{-4}$ will be in general sufficient.

At step i , we check the distance of the two vectors $A(\mathbf{v}_i)$ and $\mathbf{v}_i$ and if possible, halt. If instead convergence is not reached, we need repeat the cycle with a new initialiser $\mathbf{v}_{i+1}$. A very simple choice would be to set $\mathbf{v}_{i+1} = \mathbf{v}_i$. We do something slightly different:

$$\mathbf{v}_{i+1} \equiv gA(\mathbf{v}_i) + (1 - g)\mathbf{v}_i. \quad (2.25)$$

The “mixing parameter” g is a simple but useful knob to tune algorithmic evolution. Its usefulness comes from this observation: even if a fixed point exists, we don’t know how easy it is to actually get there following A . Setting $g = 1$, the new initialiser is just the image of the previous initialiser. By moving “slower” ($g < 1$) with an accurate optimisation of the mixing, it is possible to reach convergence in a controlled way.

Let us give some concrete justification about the relevance of controlling g . When it comes to HF iterative schemes of this kind, it can sometimes happen for the algorithm to remain stuck in infinite loops. An example of this behaviour is given in Sec. 4.5.3. A typical situation is the following: during algorithm evolution we reach a couple of points $\mathbf{v}^{(a)}, \mathbf{v}^{(b)}$ such that

$$\dots \xrightarrow{A, g} \mathbf{v}^{(a)} \xrightarrow{A, g} \mathbf{v}^{(b)} \xrightarrow{A, g} \mathbf{v}^{(a)} \xrightarrow{A, g} \mathbf{v}^{(b)} \xrightarrow{A, g} \dots$$

The notation “ $\xrightarrow{A, g}$ ” indicates one algorithmic step at mixing parameter g . We are stuck forever in this loop. Now, to predict if couples like $(\mathbf{v}^{(a)}, \mathbf{v}^{(b)})$ exist, how they depend on g , where they are located in HFPs space... is impossible at this level. Thus there is no immediate way to avoid these loops *a priori*. One workaround we found to work in most cases, however, was to just make g smaller. We gained empirical evidence that to “slow down” in HFPs space makes numeric instabilities less frequent. The trade-off, however, is that if we move too slow, runtime could get very large. Note that inaccurate setup of the algorithm could lead to resources-intensive runs also in “ordinary” situations, when we are not stuck in a loop.

To keep the number of iterations under control, we also defined an upper bound p to the total number of steps. If after p steps we algorithm failed to converge, it halts. In that case, $\mathbf{v}$ is saved as a “NaN” (Not a number) vector and we move to another simulation. Introducing this simple rule, runtime is controlled and we gain an easy method to identify the regions of parameter space requiring more care – just read from the raw data where NaNs have been saved and repeat simulations fine-tuning the parameters. This basically means to lower g and increase p . These considerations end this technical introductory chapter on iHF methods.

Chapter 3

Phase transitions and spontaneous symmetry breaking in the extended Hubbard model

In this chapter we discuss the symmetries of the extended Hubbard model, and classify the various phases discussed in this thesis based on the spontaneous symmetry breaking (SSB) mechanism. Therefore, we describe the explicit implementation of the HF method to investigate phase transitions hosted by the model.

3.1 Symmetries of the extended Hubbard model on the square lattice

In this section, we present the symmetries of the extended Hubbard model (EHM) introduced in Eq. (1.3)

$$\hat{H}_{\text{EHM}} = \underbrace{-t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma}}_{\hat{H}_t} + \underbrace{U \sum_{i \in \mathcal{L}} \hat{n}_{i\uparrow} \hat{n}_{i\downarrow}}_{\hat{H}_U} - \underbrace{V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'}}_{\hat{H}_V},$$

where $t, U, V > 0$ and $\langle ij \rangle$ is a shorthand notation to indicate nearest neighbours (NNs) on the lattice $\mathcal{L}$. The symmetries of the model can be divided in two groups: the ones related to the geometry of the square lattice, and those related to the algebraic properties of $\hat{H}_{\text{EHM}}$.

3.1.1 Lattice symmetries

We study the EHM on the planar square lattice, with one orbital per site. We indicate with L_x the number of sites in direction x , and L_y in direction y . The hamiltonian of Eq. (1.3) respects three fundamental lattice symmetries:

1. Discrete translations symmetry in both the x and y directions. We refer to the groups of one-site translations as $T_1^{(x)}, T_1^{(y)}$. It holds:

$$T_n^{(\ell)} \supset T_{kn}^{(\ell)} \quad \text{where } k \in \mathbb{N}. \quad (3.1)$$

Namely, a shift in direction ℓ by kn sites can be decomposed in k shift by n sites. Invariance by n -sites shifts, thus, implies invariance also for kn -sites shifts.

2. Discrete $\pi/2$ rotations symmetry, described by the four-fold cyclic point group C_4 , containing rotations by angle $\pi/2, \pi, 3\pi/2$ and the identity. A similar property as for the translations holds for cyclic point groups,

$$C_n \supset C_{n/k} \quad \text{where } k, n/k \in \mathbb{N}. \quad (3.2)$$

Invariance for rotations of angle $2\pi/n$ implies invariance for rotations of angle $k \times 2\pi/n$.




Figure 3.1 | Square lattice $\mathcal{L}$ bipartition in two sublattices $\mathcal{S}_1$ (red), $\mathcal{S}_2$ (blue).

3. Inversion symmetry. We define I_2 as the inversion group in two-dimensional space, containing the reflection operator and identity. Notice that

$$I_2 \equiv C_2, \quad (3.3)$$

implying that, in 2D space, a rotation by π around the origin and the reflection of the position of a point are the same operation.

The planar square lattice is bipartite: it can be divided in two square lattices, as represented in Fig. 3.1. Two sites are “connected” if linked by a non-zero hopping operator. Which is, if they are NNs. Note that odd-site translations leave the hamiltonian invariant but exchange the two sublattices, while even-sites translations don’t. Specifically, let us consider a graph $\mathcal{L}$ and the set of its vertices $V(\mathcal{L}) = \{v_i\}$. Let $E(\mathcal{L})$ be the set of graph edges,

$$E(\mathcal{L}) = \{(v, w) \mid v, w \in V(\mathcal{L}) \wedge v, w \text{ are connected}\}.$$

The graph is said to be bipartite if two sub-graphs $\mathcal{S}_1, \mathcal{S}_2$ exist such that:

- The vertices of $\mathcal{L}$ split in the two sub-graphs,

$$V(\mathcal{L}) = V(\mathcal{S}_1) \cup V(\mathcal{S}_2).$$

- Vertices of one sub-graph are linked only to vertices of the other sub-graph. In other words

$$v, w \in \mathcal{S}_i \implies (v, w) \notin E(\mathcal{L}).$$

The two sublattices of the square lattice are coloured differently in Fig. 3.1.

3.1.2 Spin and charge symmetries

The conventional HM shows a global $U(2)$ symmetry, extended to $SU(2) \times SU(2)/\mathbb{Z}_2 = SO(4)$ symmetry at half-filling on a bipartite lattice [17], such as the square lattice. For the EHM, we limit to prove that we can identify a $U(2) \subset SO(4)$ global symmetry.

$U^z(1)$ symmetry and charge conservation. The EHM is symmetric under the global gauge transformation

$$\hat{c}_{i\sigma} \rightarrow e^{-i\eta} \hat{c}_{i\sigma}, \quad \hat{c}_{i\sigma}^\dagger \rightarrow e^{i\eta} \hat{c}_{i\sigma}^\dagger \quad \text{where } \eta \in \mathbb{R}. \quad (3.4)$$

To prove this, it suffices to note that any product of fermionic operators with an equal number of annihilations and creations is gauge-symmetric,

$$\hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rightarrow e^{i\eta} \hat{c}_{i\sigma}^\dagger e^{-i\eta} \hat{c}_{j\sigma'} = \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}. \quad (3.5)$$

Hence the kinetic part $\hat{H}_t$ and the interactions $\hat{H}_U, \hat{H}_V$ are invariant. We will refer to this symmetry also as “gauge symmetry”.

We observe that the on-site density is gauge invariant,

$$\hat{n}_{i\sigma} \rightarrow \hat{n}_{i\sigma}. \quad (3.6)$$

Since $[\hat{H}, \hat{N}] = 0$, it follows that the total number of particles (charge)

$$\hat{N} = \sum_{i\sigma} \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma}$$

is a good quantum number.

Gauge symmetry is described by the transformation of Eq. (3.4), which is a pure phase shift by η . The gauge group is $U^c(1)$, with a superscript “c” indicating “charge”.

Global spin symmetry Let the local spin operator

$$\hat{S}_i^\alpha \equiv \frac{1}{2} \hat{\Psi}_i^\dagger \tau^\alpha \hat{\Psi}_i, \quad (3.7)$$

where we defined the spinors $\hat{\Psi}_i$

$$\hat{\Psi}_i \equiv \begin{bmatrix} \hat{c}_{i\uparrow} \\ \hat{c}_{i\downarrow} \end{bmatrix}, \quad (3.8)$$

and the Pauli matrices

$$\tau^x \equiv \begin{bmatrix} & 1 \\ 1 & \end{bmatrix} \quad \tau^y \equiv \begin{bmatrix} & -i \\ i & \end{bmatrix} \quad \tau^z \equiv \begin{bmatrix} 1 & \\ & -1 \end{bmatrix}.$$

The spin operators of Eq. (3.7) obey an $\mathfrak{su}(2)$ algebra,

$$[\hat{S}_j^\alpha, \hat{S}_j^\beta] = i \sum_\gamma \epsilon_{\alpha\beta\gamma} \hat{S}_j^\gamma, \quad (3.9)$$

with $\epsilon_{\alpha\beta\gamma}$ the Levi-Civita tensor.

Proof 3.1 — Proof of Eq. (3.9): spin operators of Eq. (3.7) respect the $\mathfrak{su}(2)$ algebra.

To prove that $\mathfrak{su}(2)$ algebra is respected, we must compute all possible commutators. This involves many very similar calculations. For the sake of conciseness, we sketch the proof for $\alpha = x, \beta = y$:

$$\begin{aligned} [\hat{S}_j^x, \hat{S}_j^y] &= \frac{1}{4} \left[\hat{\Psi}_j^\dagger \begin{bmatrix} & 1 \\ 1 & \end{bmatrix} \hat{\Psi}_j, \hat{\Psi}_j^\dagger \begin{bmatrix} & -i \\ i & \end{bmatrix} \hat{\Psi}_j \right] \\ &= \frac{1}{4} \left[\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} + \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow}, -i \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} + i \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \right] \\ &= \frac{i}{4} \left[\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow}, \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \right] - \frac{i}{4} \left[\hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow}, \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} \right] \\ &= \frac{i}{2} \left[\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow}, \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \right] \\ &= \frac{i}{2} \left(\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} - \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} \right) \\ &= \frac{i}{2} \left(-\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \hat{c}_{j\uparrow} + \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\uparrow} + \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\uparrow} \hat{c}_{j\downarrow} - \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \right) \\ &= \frac{i}{2} \left(\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\uparrow} - \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \right) \\ &= i \hat{S}_j^z. \end{aligned}$$

We used Fermi anticommutation rules, $\{\hat{c}_{i\sigma}, \hat{c}_{j\sigma'}^\dagger\} = \delta_{ij} \delta_{\sigma\sigma'}$. Identical derivations can be carried out for all other commutators, giving in the end the $\mathfrak{su}(2)$ spin algebra.

We define the on-site raising and lowering operators:

$$\hat{S}_j^+ \equiv \frac{\hat{S}_j^x + i\hat{S}_j^y}{2}, \quad \hat{S}_j^- \equiv \frac{\hat{S}_j^x - i\hat{S}_j^y}{2}, \quad (3.10)$$

and the global spin operators

$$\hat{S}^\alpha \equiv \sum_{i \in \mathcal{L}} \hat{S}_i^\alpha, \quad [\hat{S}^\alpha, \hat{S}^\beta] = i \sum_{\gamma} \epsilon_{\alpha\beta\gamma} \hat{S}^\gamma, \quad (3.11)$$

being $\mathcal{L}$ the lattice. Explicitly,

$$\hat{S}^z \equiv \frac{1}{2} \sum_{i \in \mathcal{L}} (\hat{n}_{i\uparrow} - \hat{n}_{i\downarrow}), \quad \hat{S}^+ \equiv \frac{1}{2} \sum_{i \in \mathcal{L}} \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}, \quad \hat{S}^- \equiv (\hat{S}^+)^\dagger.$$

The EHM hamiltonian and the number operator commute with the global spin operator. For the kinetic part $\hat{H}_t$ and the local repulsion $\hat{H}_U$ this is a well known result [17, 41],

$$[\hat{H}_t + \hat{H}_U, \hat{\mathbf{S}}] = 0.$$

Also $\hat{H}_V$ and $\hat{N}$ commute with each component of the global spin given in Eq. (3.11),

$$[\hat{H}_V, \hat{\mathbf{S}}] = 0, \quad [\hat{N}, \hat{\mathbf{S}}] = 0. \quad (3.12)$$

Proof 3.2 — Proof of Eq. (3.12): $\hat{H}_V$ and $\hat{N}$ are SU(2) symmetric..

The proof is organised through a sequence of results:

1. The non-local attraction can be written as:

$$\begin{aligned} \hat{H}_V &= V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'} \\ &= V \sum_{\langle ij \rangle} \hat{\Psi}_i^\dagger \hat{\Psi}_i \hat{\Psi}_j^\dagger \hat{\Psi}_j, \end{aligned} \quad (3.13)$$

having we used $\hat{\Psi}_i^\dagger \hat{\Psi}_i = \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} + \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow}$.

2. Similarly, the number operator is given by:

$$\begin{aligned} \hat{N} &= \sum_{i \in \mathcal{L}} \sum_{\sigma} \hat{n}_{i\sigma} \\ &= \sum_{i \in \mathcal{L}} \hat{\Psi}_i^\dagger \hat{\Psi}_i. \end{aligned} \quad (3.14)$$

3. Let now $\hat{R}(\boldsymbol{\theta})$ be any global spin rotation,

$$\hat{R}(\boldsymbol{\theta}) \equiv \exp(-i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}) \quad \text{where} \quad \boldsymbol{\theta} = (\theta_x, \theta_y, \theta_z),$$

with $\hat{\mathbf{S}}$ the global spin given in Eq. (3.11). Spinors transform as:

$$\hat{R}(\boldsymbol{\theta}) \hat{\Psi}_i \hat{R}(\boldsymbol{\theta})^\dagger = \mathcal{R}(\boldsymbol{\theta}) \hat{\Psi}_i,$$

with $\mathcal{R}(\boldsymbol{\theta}) \in \text{SU}(2)$ the unitary matrix associated to said rotation,

$$\mathcal{R}(\boldsymbol{\theta}) = \exp\left(-\frac{i}{2} \boldsymbol{\theta} \cdot \boldsymbol{\tau}\right) \quad \text{where} \quad \boldsymbol{\tau} = (\tau_x, \tau_y, \tau_z). \quad (3.15)$$

Let us prove this. Let $\hat{A} = i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}$ e $\hat{B} = \hat{\Psi}_i$. The following identity, a consequence of Baker-Campbell-Housdorff formula, holds:

$$e^{\hat{A}}\hat{B}e^{-\hat{A}} = \hat{B} + [\hat{A}, \hat{B}] + \frac{1}{2!}[\hat{A}, [\hat{A}, \hat{B}]] + \frac{1}{3!}[\hat{A}, [\hat{A}, [\hat{A}, \hat{B}]]] + \dots$$

The first order of the expansion gives:

$$\begin{aligned} [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, \hat{\Psi}_i] &= i \sum_{\alpha} \theta_{\alpha} [\hat{S}^{\alpha}, \hat{\Psi}_i] \\ &= -\frac{i}{2} \sum_{\alpha} \theta_{\alpha} \tau^{\alpha} \hat{\Psi}_i \\ &= -\frac{i}{2} \boldsymbol{\theta} \cdot \boldsymbol{\tau} \hat{\Psi}_i. \end{aligned}$$

We omitted some purely algebraic passages. The second order is given by

$$\begin{aligned} [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, \hat{\Psi}_i]] &= [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, -\frac{i}{2}(\boldsymbol{\theta} \cdot \boldsymbol{\tau}) \hat{\Psi}_i] \\ &= -\frac{i}{2} \boldsymbol{\theta} \cdot \boldsymbol{\tau} [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, \hat{\Psi}_i] \\ &= \frac{1}{4} (\boldsymbol{\theta} \cdot \boldsymbol{\tau})^2 \hat{\Psi}_i, \end{aligned}$$

and so on. Collecting all terms, we obtain

$$\begin{aligned} \hat{R}(\boldsymbol{\theta}) \hat{\Psi}_i \hat{R}(\boldsymbol{\theta})^{\dagger} &= \hat{\Psi}_i - \frac{i}{2} \boldsymbol{\theta} \cdot \boldsymbol{\tau} \hat{\Psi}_i + \frac{1}{2! \times 4} (\boldsymbol{\theta} \cdot \boldsymbol{\tau})^2 \hat{\Psi}_i + \dots \\ &= \sum_{k=1}^{\infty} \frac{1}{k!} \left(-\frac{i}{2} \boldsymbol{\theta} \cdot \boldsymbol{\tau} \right)^k \hat{\Psi}_i \\ &= \mathcal{R}(\boldsymbol{\theta}) \hat{\Psi}_i. \end{aligned}$$

In the last line, we identified the series as the Taylor expansion of the exponential, recognising the expression for $\mathcal{R}(\boldsymbol{\theta})$ given in Eq. (3.15).

4. Then, the group acts on the product $\hat{\Psi}_i^{\dagger} \hat{\Psi}_i$ as

$$\begin{aligned} \hat{R} \hat{\Psi}_i^{\dagger} \hat{\Psi}_i \hat{R}^{\dagger} &= \hat{R} \hat{\Psi}_i^{\dagger} \hat{R}^{\dagger} \hat{R} \hat{\Psi}_i \hat{R}^{\dagger} \\ &= \hat{\Psi}_i^{\dagger} \mathcal{R}^{\dagger} \mathcal{R} \hat{\Psi}_i \\ &= \hat{\Psi}_i^{\dagger} \hat{\Psi}_i \end{aligned}$$

being $\mathcal{R} \in \text{SU}(2)$. Recalling Eqs. (3.13), (3.14), we conclude that $\hat{H}_V$ and $\hat{N}$ are $\text{SU}(2)$ symmetric – which means, it does not change under global spin rotations. This fact implies Eq. (3.12).

We observe that the above proof can be used to show that the on-site number operator is spin-invariant,

$$\hat{n}_{i\sigma} = \hat{\Psi}_i^{\dagger} \hat{\Psi}_i \implies [\hat{n}_{i\sigma}, \hat{\mathbf{S}}] = 0. \quad (3.16)$$

The EHM is symmetric under global spin rotations. Specifically, both $\hat{H}_U$ and $\hat{H}_V$ are symmetric also under local spin rotations, whereas the kinetic operator is symmetric only under global spin rotations.

The $\text{U}^c(1)$ gauge symmetry and the $\text{SU}(2)$ spin rotations symmetry can be combined,

$$\text{U}^c(1) \otimes \text{SU}(2) = \text{U}(2).$$




Figure 3.2 | Graphic representation of the domain of SU(2) rotations (left figure) and U^z(1) rotations (right figure). For any given direction **m**, the group of rotations around said direction is contained in the larger group SU(2).

We will refer to U(2) symmetry to indicate full charge-spin conservation. Finally, note the group of 3D rotations contains the subgroup of 2D rotations around a given axis z ,

$$U^z(1) \subset SU(2),$$

where the z superscript serves to distinguish from the gauge group. The z axis does not need to be identified with the “real” z axis, perpendicular with the lattice plane. Any given direction works. In Fig. 3.2 we show schematically how U^z(1) is a subgroup of SU(2) for a given direction m .

3.1.3 Particle-hole symmetry at half-filling

At half-filling (equal number of electrons and lattice sites), the EHM on the square lattice exhibits particle-hole symmetry (PHS). The presence of this additional symmetry is a consequence of the fact that the square lattice is bipartite. The generic unitary particle-hole (PH) transformation is given by

$$P : \hat{c}_{i\sigma} \rightarrow e^{i\eta_{i\sigma}} \hat{h}_{i\sigma}^\dagger, \quad (3.17)$$

being in general η_i a phase dependent on the fermionic quantum state ($i\sigma$). Inserting the above transformation in the EHM, we get

$$P\{\hat{H}\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma})} \hat{h}_{i\sigma} \hat{h}_{j\sigma}^\dagger + U \sum_i \hat{h}_{i\uparrow} \hat{h}_{i\downarrow} \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\uparrow}^\dagger - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger.$$

For the two interactions $\hat{H}_U$, $\hat{H}_V$ the phase factors cancelled out. For a specific choice of the η_i phases,

$$P : \begin{cases} \hat{c}_{i\sigma} \rightarrow \hat{h}_{i\sigma}^\dagger & \text{for } i \in \mathcal{S}_1, \\ \hat{c}_{i\sigma} \rightarrow -\hat{h}_{i\sigma}^\dagger & \text{for } i \in \mathcal{S}_2, \end{cases} \quad (3.18)$$

the above hamiltonian is mapped onto the EHM of (1.3),

$$P\{\hat{H}\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma} + U \sum_{i \in \mathcal{L}} \hat{h}_{i\uparrow}^\dagger \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}.$$

The equality $P\{\hat{H}\} = \hat{H}$ expresses PHS, which holds as long as we work at half-filling.

Proof 3.3 — PHS at half filling, with η_i given by Eq. (3.18).

This proof is an extension of the argument carried out for the conventional HM [17]. For the

hole operators, Fermi anticommutation rules hold regardlessly of the phase factor:

$$\begin{aligned}\{\hat{h}_{i\sigma}, \hat{h}_{j\sigma'}^\dagger\} &= e^{i(\eta_{i\sigma} - \eta_{j\sigma'})} \{\hat{c}_{i\sigma}^\dagger, \hat{c}_{j\sigma'}\} \\ &= e^{i(\eta_{i\sigma} - \eta_{j\sigma'})} \delta_{ij} \delta_{\sigma\sigma'} \\ &= \delta_{ij} \delta_{\sigma\sigma'}.\end{aligned}$$

The on-site density operator is given by:

$$\hat{h}_{i\sigma} \hat{h}_{i\sigma}^\dagger = \hat{n}_{i\sigma},$$

being $\hat{n}_{i\sigma}$ the fermionic number operator on site i with spin σ . It follows, for quartic interactions

$$\begin{aligned}\hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger &= -\hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \\ &= \hat{h}_{i\sigma} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= -\hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} - \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= -\hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'} - \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \left(\hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \right) \delta_{ij} \delta_{\sigma\sigma'} \\ &= \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} \hat{h}_{i\sigma} + (\hat{n}_{i\sigma} + \hat{n}_{j\sigma'}) - 1 - \left(\hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \right) \delta_{ij} \delta_{\sigma\sigma'}.\end{aligned}\quad (3.19)$$

The first term of the last passage is mirrored with respect to the starting term. In next paragraphs we derive the constraint of Eq. (3.18) and the requirement to work at half-filling.

Kinetic term. Consider the kinetic term,

$$\hat{H}_t = -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma}.$$

We apply the PH transform of Eq. (3.17)

$$P\{\hat{H}_t\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma})} \hat{h}_{i\sigma} \hat{h}_{j\sigma}^\dagger.$$

Anti-commutating the two fermionic operators, and exchanging site labels $i \leftrightarrow j$, we get:

$$P\{\hat{H}_t\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma} + \pi)} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma}.$$

To obtain $P\{\hat{H}_t\} \stackrel{!}{=} \hat{H}_t$, a necessary condition on the unitary operation P is

$$\eta_{i\sigma} + \eta_{j\sigma} \stackrel{!}{=} (2k - 1)\pi \quad \text{with} \quad k \in \mathbb{Z}, \quad (3.20)$$

valid for NN sites. This requirement on the phases η_i gives Eq. (3.18).

Local repulsion. Substitute Eq. (3.19) in $\hat{H}_U$,

$$\begin{aligned}P\{\hat{H}_U\} &= U \sum_{i \in \mathcal{L}} \hat{h}_{i\uparrow} \hat{h}_{i\downarrow} \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\uparrow}^\dagger \\ &= U \sum_{i \in \mathcal{L}} \hat{h}_{i\uparrow}^\dagger \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} + U \sum_{i \in \mathcal{L}} (\hat{n}_{i\uparrow} + \hat{n}_{i\downarrow}) - U \sum_{i \in \mathcal{L}} (1) \\ &= U \sum_i \hat{h}_{i\uparrow}^\dagger \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} + U(\hat{N} - L_x L_y).\end{aligned}$$

Symmetry	Operations
$T_1^{(x)} \otimes T_1^{(y)}$	One-site discrete translations separately in both directions
C_4	Four-fold point group (real $\pi/2$ angle rotations)
$U^c(1)$	Charge-conserving global phase shifts
$SU(2)$	Three-dimensional rotations in global spin space

Table 3.1 | Model symmetries and relative group operations. Note that $U^c(1)$ represents the gauge symmetry linked to charge conservation.

At half filling, $\langle \hat{N} \rangle = L_x L_y$. The unitary map of Eq. (3.17), independently of the condition of Eq. (3.20), leaves the hamiltonian identical at half-filling.

Non-local attraction. We proceed identically for $\hat{H}_V$,

$$\begin{aligned}
 P\{\hat{H}_V\} &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger \\
 &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} \hat{h}_{i\sigma} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} (\hat{n}_{i\sigma} + \hat{n}_{j\sigma'}) + V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} (1) \\
 &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} \hat{h}_{i\sigma} - 2zV(\hat{N} - L_x L_y),
 \end{aligned}$$

with $z = 4$ the lattice coordination number. In the last line, we used that on the square lattice

$$\begin{aligned}
 \sum_{\langle ij \rangle} (1) &= \sum_{i \in \mathcal{L}/2} z \\
 &= \frac{L_x L_y}{2} \times z.
 \end{aligned} \tag{3.21}$$

As before, the non-local attraction is particle-hole symmetric at half-filling. We conclude that the entire EHM exhibits PHS at half filling, if P is defined as in Eq. (3.18).

As for the conventional HM [17], this result can be generalised to any bipartite lattice. In turn, if a lattice is not bipartite, PHS is not present.

We report in Tab. 3.1 all identified symmetries.

3.2 The HF method and symmetry breaking

In this thesis, we use the HF method to investigate the different many-body phases hosted by the EHM. We classify these phases in terms of the symmetry broken by spontaneous symmetry breaking (SSB) mechanism:

- *Normal phase*: no broken symmetry.
- *Commensurate antiferromagnetic phase*: simultaneous $SU(2) \xrightarrow{\text{SSB}} U^z(1)$ and $T_1^{(x)} \otimes T_1^{(y)} \xrightarrow{\text{SSB}} T_2^{(x)} \otimes T_2^{(y)}$.
- *Nematic superconducting phase*: simultaneous $U^c(1) \xrightarrow{\text{SSB}} 0$ and $C_4 \xrightarrow{\text{SSB}} C_2$, namely superconductivity and breaking of planar rotational symmetry, observed for example in iron-based superconductors [26].

In the subsections that follow we discuss how to implement these SSBs in the context of the HF method. Specifically, as is described in Chap. 2, we investigate each of these phases by approximating the EHM hamiltonian

$$\hat{H} \stackrel{\text{HF}}{\simeq} \hat{H}_{\text{HF}},$$




Figure 3.3 | Schematic representation of the four NNs of a given site i for a planar square lattice.

with an operator $\hat{H}_{\text{HF}}$ that respects the symmetries of the given phase. We start with the general decomposition of the two interactions $\hat{H}_U$ and $\hat{H}_V$, given respectively in Eqs. (2.21) and (2.22), and discuss the symmetry properties of each sector.

In the following, we switch from the site index $i \in \mathcal{L}$ to the vector notation $\mathbf{r} \in \mathbb{N}^2$, indicating the site 2D position on the lattice as

$$i \rightarrow \mathbf{r} = (x, y),$$

with $x, y \in \mathbb{N}$. The lattice versors are:

$$\underline{\mathbf{x}} \equiv (1, 0) \quad \text{and} \quad \underline{\mathbf{y}} \equiv (0, 1),$$

as is shown in Fig. 3.3.

3.2.1 Local repulsion

The local repulsion $\hat{H}_U$ is given by:

$$\begin{aligned} \hat{H}_U &= U \sum_{\mathbf{r} \in \mathcal{L}} \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}\downarrow} \\ &= U \sum_{\mathbf{r} \in \mathcal{L}} \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow}. \end{aligned}$$

Applying Wick's theorem and using the denominations of Eq. (2.1) we have

$$\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle = \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \rangle}_{\text{Hartree}} - \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \rangle \langle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Cooper}}, \quad (3.22)$$

where the equality holds exactly as long as we perform the averaging operation on the subspace of gaussian states. We now discuss the above expectation values (EVs) separately, obtaining for each phase a set of “symmetry selection rules”. We define a symmetry selection rule as a theoretical argument that allows us to assess if a given EV is zero, based on symmetry considerations.

Hartree term. The Hartree term of the above decomposition is a product of density EVs,

$$\langle \hat{n}_{\mathbf{r}\uparrow} \rangle \langle \hat{n}_{\mathbf{r}\downarrow} \rangle.$$

From Eq. (3.6), (3.16) we know that single-site number operators are invariant under the entire group $U(2)$. Moreover, shifting or rotating the lattice leaves density operators unchanged. Thus no symmetry selection rule applies, and the above EVs are in principle non-zero.

Fock terms. For the Fock term, we notice that

$$\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} = \hat{S}_{\mathbf{r}}^+,$$

being $\hat{S}_{\mathbf{r}}^+$ the local raising operator. From $\mathfrak{su}(2)$ algebra we have:

$$[\hat{S}_{\mathbf{r}}^+, \hat{S}_{\mathbf{r}}^\alpha] \neq 0 \quad \text{where} \quad \alpha \in \{x, y, z\}.$$

If the ground-state is invariant under global spin rotations, $\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \rangle$ must be zero. instead, gauge symmetry gives no symmetry selection rules, due to Eq. (3.5).

Term from $\hat{H}_U$	Normal phase	Antiferromagnet	Superconductor
Hartree	Allowed	Allowed	Allowed
Fock	Prohibited	Prohibited	Prohibited
Cooper	Prohibited	Prohibited	Allowed

Table 3.2 | Summary of symmetry selection rules for the Wick's decomposition of Eq. (3.22) of the local repulsion $\hat{H}_U$.

Cooper terms. The Cooper term can be non-zero only if charge is not conserved. In terms of the gauge symmetry, it suffices to apply the map of Eq. (3.4) to $\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger$,

$$\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \rightarrow e^{2i\eta} \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger,$$

to conclude that this operator is in general not gauge invariant. It is instead SU(2)-symmetric.

Proof 3.4 — Local Cooper terms are SU(2)-symmetric.

We must compute the commutators of $\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger$ with $\hat{S}^z$, $\hat{S}^+$:

$$\begin{aligned} [\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{S}^z] &= \left[\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger, \frac{1}{2} \sum_{\mathbf{r}' \in \mathcal{L}} (\hat{n}_{\mathbf{r}'\uparrow} - \hat{n}_{\mathbf{r}'\downarrow}) \right] \\ &= \frac{1}{2} [\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}] \\ &= \frac{1}{2} [\hat{c}_{\mathbf{r}\uparrow}^\dagger, \hat{n}_{\mathbf{r}\uparrow}] \hat{c}_{\mathbf{r}\downarrow}^\dagger - \frac{1}{2} \hat{c}_{\mathbf{r}\uparrow}^\dagger [\hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{n}_{\mathbf{r}\downarrow}]. \end{aligned}$$

Since:

$$\begin{aligned} [\hat{c}_{\mathbf{r}\sigma}^\dagger, \hat{n}_{\mathbf{r}\sigma}] &= \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma} - \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma} \hat{c}_{\mathbf{r}\sigma}^\dagger \\ &= -\hat{c}_{\mathbf{r}\sigma}^\dagger (1 - \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma}) \\ &= -\hat{c}_{\mathbf{r}\sigma}^\dagger, \end{aligned} \tag{3.23}$$

we have:

$$[\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{S}^z] = -\frac{1}{2} \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger + \frac{1}{2} \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger = 0.$$

Hence, $\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger$ is U^z(1)-symmetric.

Moreover, considering the raising operator:

$$\begin{aligned} [\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{S}^+] &= \left[\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger, \frac{1}{2} \sum_{\mathbf{r}' \in \mathcal{L}} \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \right] \\ &= [\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}] \\ &= [\hat{c}_{\mathbf{r}\uparrow}^\dagger, \hat{c}_{\mathbf{r}\uparrow}^\dagger] \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} - (\hat{c}_{\mathbf{r}\uparrow}^\dagger)^2 [\hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{c}_{\mathbf{r}\downarrow}] \\ &= 0, \end{aligned}$$

where we used the facts that every operator commutes with itself, and a squared creation operator is null. These commutation relations, together, prove that Cooper EVs are legitimately non-zero for a SU(2)-preserving phase.

In Tab. 3.2 we collect the symmetry selection rules for the Hartree, Fock and Cooper terms of Eq. (3.22), based on the symmetry specifications given for each phase in above paragraphs. Note that Fock-like EVs are prohibited for all phases.

3.2.2 Non-local attraction

The non-local attraction $\hat{H}_V$ is given by:

$$\begin{aligned}\hat{H}_V &= -V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma, \sigma'} \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}'\sigma'} \\ &= V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma, \sigma'} \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}'\sigma'}^\dagger \hat{c}_{\mathbf{r}'\sigma'} \hat{c}_{\mathbf{r}\sigma}.\end{aligned}\quad (3.24)$$

Let

$$\hat{H}_V^{\sigma\sigma'} \equiv -V \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}'\sigma'}.$$

The interaction decomposes in two “spin sectors”

$$\begin{aligned}\hat{H}_V &= \underbrace{\hat{H}_V^{\uparrow\uparrow} + \hat{H}_V^{\downarrow\downarrow}}_{\text{Same spin}} + \underbrace{\hat{H}_V^{\uparrow\downarrow} + \hat{H}_V^{\downarrow\uparrow}}_{\text{Opposite spin}} \\ &\equiv \hat{H}^{(\text{s.s.})} + \hat{H}^{(\text{o.s.})},\end{aligned}\quad (3.25)$$

where we further separated in s.s. and o.s. operators.

The operator $\hat{H}^{(\text{s.s.})}$ can be written as follows:

$$\begin{aligned}\hat{H}_V^{(\text{s.s.})} &\equiv \hat{H}_V^{\uparrow\uparrow} + \hat{H}_V^{\downarrow\downarrow} \\ &= -V \sum_{\mathbf{r} \in \mathcal{L}/2} \sum_{\delta \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}} \sum_{\sigma \in \{\uparrow, \downarrow\}} (\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\delta\sigma} + \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}-\delta\sigma}),\end{aligned}\quad (3.26)$$

where the sum over $\mathcal{L}/2$, in last line, means that we are summing over just one sublattice (it does not matter which one). Applying Wick’s theorem in the subspace of gaussian states, we get:

$$\begin{aligned}\langle \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\delta\sigma} \rangle &= \langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma} \hat{c}_{\mathbf{r}\sigma} \rangle \\ &= \underbrace{\langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma} \rangle \langle \hat{c}_{\mathbf{r}+\delta\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma} \rangle}_{\text{Hartree}} - \underbrace{\langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma} \rangle \langle \hat{c}_{\mathbf{r}+\delta\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma}^\dagger \rangle \langle \hat{c}_{\mathbf{r}+\delta\sigma} \hat{c}_{\mathbf{r}\sigma} \rangle}_{\text{Cooper}}.\end{aligned}\quad (3.27)$$

Proof 3.5 — Proof of Eq. (3.26).

Let $f(\mathbf{r}, \mathbf{r}')$ be a function of two variables on the square lattice. Summing over NNs, we get the identity

$$\sum_{\langle \mathbf{r}\mathbf{r}' \rangle} f(\mathbf{r}, \mathbf{r}') = \sum_{\mathbf{r}\mathbf{r}'} f(\mathbf{r}, \mathbf{r}') \left(\delta_{\mathbf{r}'=\mathbf{r}+\underline{\mathbf{x}}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{\mathbf{x}}} + \delta_{\mathbf{r}'=\mathbf{r}+\underline{\mathbf{y}}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{\mathbf{y}}} \right),$$

where we used the notation of Fig. 3.3. The above sum is identically written as:

$$\sum_{\langle \mathbf{r}\mathbf{r}' \rangle} f(\mathbf{r}, \mathbf{r}') = \sum_{\mathbf{r} \in \mathcal{L}/2} \sum_{\delta \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}} [f(\mathbf{r}, \mathbf{r} + \delta) + f(\mathbf{r}, \mathbf{r} - \delta)]. \quad (3.28)$$

The symbol “ $\mathcal{L}/2$ ” indicates one of the two sublattices $\mathcal{S}_1, \mathcal{S}_2$ represented in Fig. 3.1. Inserting Eq. (3.28) in Eq. (3.24), we obtain:

$$\hat{H}_V = -V \sum_{\mathbf{r} \in \mathcal{L}/2} \sum_{\delta \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}} \sum_{\sigma, \sigma'} (\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\delta\sigma'} + \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}-\delta\sigma'}). \quad (3.29)$$

By taking the s.s. sector of the above equation, namely $\sigma = \sigma'$, and performing Wick’s contractions, we obtain Eq. (3.26).

The operator $\hat{H}^{(\text{o.s.})}$ can be written as follows:

$$\begin{aligned}\hat{H}_V^{(\text{o.s.})} &\equiv \hat{H}_V^{\uparrow\downarrow} + \hat{H}_V^{\downarrow\uparrow} \\ &= -V \sum_{\mathbf{r} \in \mathcal{L}} \sum_{\delta \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}} (\hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}+\delta\downarrow} + \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}-\delta\downarrow}).\end{aligned}\quad (3.30)$$

From Wick's theorem:

$$\begin{aligned} \langle \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}\pm\delta\downarrow} \rangle &= \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\pm\delta\downarrow}^\dagger \hat{c}_{\mathbf{r}\pm\delta\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle \\ &= \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}\pm\delta\downarrow}^\dagger \hat{c}_{\mathbf{r}\pm\delta\downarrow} \rangle}_{\text{Hartree}} - \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\pm\delta\downarrow} \rangle \langle \hat{c}_{\mathbf{r}\pm\delta\downarrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\pm\delta\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}\pm\delta\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Cooper}}. \end{aligned} \quad (3.31)$$

Again, equality holds in the subspace of gaussian states.

Proof 3.6 — Proof of Eq. (3.30).

Consider Eq. (3.29), restricting to $\sigma = \uparrow$, $\sigma' = \downarrow$ and setting $\mathcal{L}/2 = \mathcal{S}_1$. We obtain

$$\hat{H}_V^{\uparrow\downarrow} = -V \sum_{\mathbf{r} \in \mathcal{S}_1} \sum_{\delta \in \{\underline{x}, \underline{y}\}} (\hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}+\delta\downarrow} + \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}-\delta\downarrow}).$$

Note that spin $\uparrow$ operators act on the sublattice $\mathcal{S}_1$; spin $\downarrow$ operators act on the sublattice $\mathcal{S}_2$. If instead, from Eq. (3.29), we restrict to $\sigma = \downarrow$, $\sigma' = \uparrow$ and set $\mathcal{L}/2 = \mathcal{S}_2$, we get:

$$\hat{H}_V^{\downarrow\uparrow} = -V \sum_{\mathbf{r} \in \mathcal{S}_2} \sum_{\delta \in \{\underline{x}, \underline{y}\}} (\hat{n}_{\mathbf{r}\downarrow} \hat{n}_{\mathbf{r}+\delta\uparrow} + \hat{n}_{\mathbf{r}\downarrow} \hat{n}_{\mathbf{r}-\delta\uparrow}).$$

We observe that if $\mathbf{r} \in \mathcal{S}_2$, all its NNs belong to $\mathcal{S}_1$. Again, spin $\uparrow$ operators act on the sublattice $\mathcal{S}_1$; spin $\downarrow$ operators act on the sublattice $\mathcal{S}_2$.

Therefore, summing $\hat{H}_V^{\uparrow\downarrow} + \hat{H}_V^{\downarrow\uparrow}$, we obtain

$$\hat{H}_V^{\uparrow\downarrow} + \hat{H}_V^{\downarrow\uparrow} = -V \sum_{\mathbf{r} \in \mathcal{L}} \sum_{\delta \in \{\underline{x}, \underline{y}\}} (\hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}+\delta\downarrow} + \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}-\delta\downarrow}),$$

which coincides with Eq. (3.30).

The separation of $\hat{H}_V$ in o.s. and s.s. sector is necessary in order to derive a set of symmetry selection rules for the Wick's contractions of Eqs. (3.27), (3.31). This is done in next paragraphs.

Hartree terms. As was for the local repulsion $\hat{H}_U$, the non-local Hartree terms of Eqs. (3.30), (3.26) are not subject to symmetry selection rules for any symmetry of Tab. 3.1. Thus they are allowed for all sectors.

Fock terms. The Fock EVs from the two sectors behave differently. Let us compute the following generic commutation relations:

$$\begin{aligned} [\hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma'}, \hat{S}^z] &= \left[\hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma'}, \frac{1}{2} \sum_{\mathbf{r}' \in \mathcal{L}} (\hat{n}_{\mathbf{r}'\uparrow} - \hat{n}_{\mathbf{r}'\downarrow}) \right] \\ &= \frac{(-1)^{\delta_{\sigma=\downarrow}}}{2} [\hat{c}_{\mathbf{r}\sigma}^\dagger, \hat{n}_{\mathbf{r}\sigma}] \hat{c}_{\mathbf{r}+\delta\sigma'} + \frac{(-1)^{\delta_{\sigma'=\downarrow}}}{2} \hat{c}_{\mathbf{r}\sigma}^\dagger [\hat{c}_{\mathbf{r}+\delta\sigma'}, \hat{n}_{\mathbf{r}+\delta\sigma'}] \\ &= \left(-\frac{(-1)^{\delta_{\sigma=\downarrow}}}{2} + \frac{(-1)^{\delta_{\sigma'=\downarrow}}}{2} \right) \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma'}, \end{aligned} \quad (3.32)$$

where in last passage we used Eq. (3.23) and its conjugate version,

$$[\hat{c}_{\mathbf{r}+\delta\sigma'}, \hat{n}_{\mathbf{r}+\delta\sigma}] = \hat{c}_{\mathbf{r}+\delta\sigma'}.$$

Now, if $\sigma = \sigma'$ the last passage of Eq. (3.32) is zero; otherwise is not. This proves that the o.s. Fock EVs are non-zero only for a phase that completely breaks SU(2); instead, in the s.s. sector we can have non-zero EVs for a phase that reduces SU(2) to U^z(1).

Sector	Term from $\hat{H}_V$	Normal phase	Antiferromagnet	Superconductor
o.s.	Hartree	Allowed	Allowed	Allowed
	Fock	Prohibited	Prohibited	Prohibited
	Cooper	Prohibited	Prohibited	Allowed
s.s.	Hartree	Allowed	Allowed	Allowed
	Fock	Allowed	Allowed	Allowed
	Cooper	Prohibited	Prohibited	Prohibited

Table 3.3 | Summary of symmetry selection rules for the Wick's decomposition of Eqs. (3.31), (3.27) of the non-local attraction $\hat{H}_V$.

Cooper terms. Cooper terms are not gauge invariant. The proof is analogous to the one given for $\hat{H}_U$. Instead, the o.s. and s.s. spin sectors behave differently under global spin rotations. With an identical procedure as for Eq. (3.32), we get:

$$\begin{aligned}
\left[\hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma'}^\dagger, \hat{S}^z \right] &= \left[\hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma'}^\dagger, \frac{1}{2} \sum_{\mathbf{r}' \in \mathcal{L}} (\hat{n}_{\mathbf{r}'\uparrow} - \hat{n}_{\mathbf{r}'\downarrow}) \right] \\
&= \frac{(-1)^{\delta_{\sigma=\downarrow}}}{2} [\hat{c}_{\mathbf{r}\sigma}^\dagger, \hat{n}_{\mathbf{r}\sigma}] \hat{c}_{\mathbf{r}+\delta\sigma'}^\dagger + \frac{(-1)^{\delta_{\sigma'=\downarrow}}}{2} \hat{c}_{\mathbf{r}\sigma}^\dagger [\hat{c}_{\mathbf{r}+\delta\sigma'}^\dagger, \hat{n}_{\mathbf{r}+\delta\sigma'}] \\
&= \left(-\frac{(-1)^{\delta_{\sigma=\downarrow}}}{2} - \frac{(-1)^{\delta_{\sigma'=\downarrow}}}{2} \right) \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\delta\sigma'}^\dagger.
\end{aligned} \tag{3.33}$$

We have a “mirrored” situation with respect to the Fock terms. For the Cooper channel, the s.s. EVs are zero if $U^z(1)$ is unbroken, while the o.s. sector are not.

In Tab. 3.3 we summarise these considerations, listing a set of symmetry selection rules about the EVs of Eqs. (3.31), (3.27), dividing in the two sectors “o.s.” and “s.s.”. By identifying these symmetry selection rules, we set grounds for the detailed derivation of each chapter and heavily simplify application of HF model to the EHM.

3.3 The EHM in reciprocal space

The EHM is symmetric under discrete lattice translations. We will investigate phases that do not (completely) break translational invariance. Therefore, It is convenient to move to reciprocal space, by Fourier-transforming the hamiltonian. We will use the following convention:

$$\hat{c}_{\mathbf{r}\sigma} \stackrel{\mathcal{F}}{=} \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{-i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma}. \tag{3.34}$$

Throughout the text, $\stackrel{\mathcal{F}}{=}$ will indicate a passage where a Fourier transform is used. “BZ” is shorthand notation for the Brillouin Zone.

It is useful to recall that the following equation for Kronecker's delta holds:

$$\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} e^{i\mathbf{k} \cdot \mathbf{r}} = \delta_{\mathbf{k}=\mathbf{0}}. \tag{3.35}$$

On one sublattice:

$$\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}/2} e^{i\mathbf{k} \cdot \mathbf{r}} = \frac{1}{2} \delta_{\mathbf{k}=\mathbf{0}}. \tag{3.36}$$

The notation $\mathcal{L}/2$ is shorthand to indicate either $\mathcal{S}_1$ or $\mathcal{S}_2$.

Kinetic part. The kinetic operators transforms as follows:

$$\hat{H}_t \stackrel{\mathcal{F}}{=} \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \epsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}, \tag{3.37}$$

where we defined the bare bands:

$$\epsilon_{\mathbf{k}} \equiv -2t (\cos k_x + \cos k_y) . \quad (3.38)$$

Proof 3.7 — Proof of Eq. (3.37): $\mathcal{F}$ -transform of $\hat{H}_t$.

Consider the kinetic operator $\hat{H}_t$:

$$\hat{H}_t = -t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}'\sigma} + \text{h.c.}] .$$

Applying the convention of Eq. (3.34), we get:

$$\begin{aligned} \hat{H}_t &\stackrel{\mathcal{F}}{=} -t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \left(\frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \right) \left(\frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k}' \in \text{BZ}} e^{-i\mathbf{k}' \cdot \mathbf{r}'} \hat{c}_{\mathbf{k}'\sigma} \right) + \text{h.c.} \\ &= -2t \sum_{\mathbf{k}, \mathbf{k}'} \sum_{\sigma} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}'\sigma} \times \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}/2} e^{i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}} \times \sum_{\delta \in \{\underline{x}, \underline{y}\}} e^{i\mathbf{k}' \cdot \delta} . \end{aligned}$$

We used Eq. (3.28) and recognised that the h.c. term gives an identical contribution. We recognise in the central term the relation of Eq. (3.36). Hence,

$$\begin{aligned} \hat{H}_t &\stackrel{\mathcal{F}}{=} -2t \sum_{\mathbf{k}, \mathbf{k}'} \sum_{\sigma} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} \left[\frac{e^{ik_x} + e^{-ik_x}}{2} + \frac{e^{ik_y} + e^{-ik_y}}{2} \right] \\ &= \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \epsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} , \end{aligned}$$

which proves Eq. (3.37).

Local repulsion. The local repulsion transforms as follows:

$$\hat{H}_U \stackrel{\mathcal{F}}{=} \frac{U}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} . \quad (3.39)$$

Sums over the variables $\mathbf{K}, \mathbf{k}, \mathbf{k}'$ must be intended as over the BZ.

Proof 3.8 — Proof of Eq. (3.39): $\mathcal{F}$ -transform of $\hat{H}_U$.

Using the convention of Eq. (3.34), we get for $\hat{H}_U$:

$$\begin{aligned} \hat{H}_U &= U \sum_{\mathbf{r} \in \mathcal{L}} \hat{c}_{\mathbf{r}\uparrow}^{\dagger} \hat{c}_{\mathbf{r}\downarrow}^{\dagger} \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \\ &\stackrel{\mathcal{F}}{=} U \sum_{\mathbf{r} \in \mathcal{L}} \frac{1}{(L_x L_y)^2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} e^{i\mathbf{k}_1 \cdot \mathbf{r}} \hat{c}_{\mathbf{k}_1\uparrow}^{\dagger} e^{i\mathbf{k}_2 \cdot \mathbf{r}} \hat{c}_{\mathbf{k}_2\downarrow}^{\dagger} e^{-i\mathbf{k}_3 \cdot \mathbf{r}} \hat{c}_{\mathbf{k}_3\downarrow} e^{-i\mathbf{k}_4 \cdot \mathbf{r}} \hat{c}_{\mathbf{k}_4\uparrow} \\ &= \frac{U}{L_x L_y} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \hat{c}_{\mathbf{k}_1\uparrow}^{\dagger} \hat{c}_{\mathbf{k}_2\downarrow}^{\dagger} \hat{c}_{\mathbf{k}_3\downarrow} \hat{c}_{\mathbf{k}_4\uparrow} \times \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} \\ &= \frac{U}{L_x L_y} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \hat{c}_{\mathbf{k}_1\uparrow}^{\dagger} \hat{c}_{\mathbf{k}_2\downarrow}^{\dagger} \hat{c}_{\mathbf{k}_3\downarrow} \hat{c}_{\mathbf{k}_4\uparrow} \delta_{\mathbf{k}_1+\mathbf{k}_2=\mathbf{k}_3+\mathbf{k}_4} , \end{aligned}$$

where we used Eq. (3.35). Let $\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4 = \mathbf{K}$, and define $\mathbf{k}, \mathbf{k}'$ such that

$$\mathbf{k}_1 \equiv \mathbf{K} + \mathbf{k}, \quad \mathbf{k}_2 \equiv \mathbf{K} - \mathbf{k}, \quad \mathbf{k}_3 \equiv \mathbf{K} - \mathbf{k}', \quad \mathbf{k}_4 \equiv \mathbf{K} + \mathbf{k}' .$$

Substituting, we get the expression of Eq. (3.39).

Non-local attraction. In o.s. sector, the non-local attraction transforms as:

$$\hat{H}_V^{(o.s.)} \stackrel{\mathcal{F}}{=} -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow}. \quad (3.40)$$

In the s.s. sector:

$$\begin{aligned} \hat{H}_V^{(s.s.)} \stackrel{\mathcal{F}}{=} & -\frac{V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \\ & \times \left[\hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\uparrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} + \hat{c}_{\mathbf{K}+\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\downarrow} \right]. \end{aligned} \quad (3.41)$$

Sums over the variables $\mathbf{K}, \mathbf{k}, \mathbf{k}'$ must be intended as over the BZ.

Proof 3.9 — Proof of Eqs. (3.40), (3.41): $\mathcal{F}$ -transform of $\hat{H}_V$.

To prove Eqs. (3.40), (3.41), it is useful to start by defining:

$$h_{\mathbf{r}\sigma\sigma'} \equiv -V \sum_{\boldsymbol{\delta} \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}} (\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\boldsymbol{\delta}\sigma'} + \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}-\boldsymbol{\delta}\sigma'}) ,$$

with:

$$\hat{H}_V = \sum_{\mathbf{r} \in \mathcal{L}/2} \sum_{\sigma, \sigma'} h_{\mathbf{r}\sigma\sigma'} .$$

The product of densities on neighbouring sites transforms as:

$$\begin{aligned} \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\boldsymbol{\delta}\sigma'} &= \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\sigma'}^\dagger \hat{c}_{\mathbf{r}+\boldsymbol{\delta}\sigma'} \hat{c}_{\mathbf{r}\sigma} \\ &\stackrel{\mathcal{F}}{=} \frac{1}{(L_x L_y)^2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} e^{\pm i(\mathbf{k}_2-\mathbf{k}_3) \cdot \boldsymbol{\delta}} \hat{c}_{\mathbf{k}_1\sigma}^\dagger \hat{c}_{\mathbf{k}_2\sigma'}^\dagger \hat{c}_{\mathbf{k}_3\sigma'} \hat{c}_{\mathbf{k}_4\sigma} . \end{aligned}$$

Then

$$\begin{aligned} h_{\mathbf{r}\sigma\sigma'} &\stackrel{\mathcal{F}}{=} -\frac{V}{(L_x L_y)^2} \sum_{\boldsymbol{\delta} \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} \\ &\quad \times \left(e^{i(\mathbf{k}_2-\mathbf{k}_3) \cdot \boldsymbol{\delta}} + e^{-i(\mathbf{k}_2-\mathbf{k}_3) \cdot \boldsymbol{\delta}} \right) \hat{c}_{\mathbf{k}_1\sigma}^\dagger \hat{c}_{\mathbf{k}_2\sigma'}^\dagger \hat{c}_{\mathbf{k}_3\sigma'} \hat{c}_{\mathbf{k}_4\sigma} \\ &= -\frac{2V}{(L_x L_y)^2} \sum_{\boldsymbol{\delta} \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} \cos[(\mathbf{k}_2 - \mathbf{k}_3) \cdot \boldsymbol{\delta}] \hat{c}_{\mathbf{k}_1\sigma}^\dagger \hat{c}_{\mathbf{k}_2\sigma'}^\dagger \hat{c}_{\mathbf{k}_3\sigma'} \hat{c}_{\mathbf{k}_4\sigma} . \end{aligned}$$

The full non-local interaction is given by summing over all sites of $\mathcal{L}/2$ (which is, one sublattice). Applying Eq. (3.36) gives back momentum conservation

$$\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}/2} e^{i[(\mathbf{k}_1+\mathbf{k}_2)-(\mathbf{k}_3+\mathbf{k}_4)] \cdot \mathbf{r}} = \frac{1}{2} \delta_{\mathbf{k}_1+\mathbf{k}_2=\mathbf{k}_3+\mathbf{k}_4} ,$$

where we used Eq. (3.36). Define $\mathbf{k}, \mathbf{k}'$ and $\mathbf{K}$ as we did for the local repulsion, and

$$\delta \mathbf{k} \equiv \mathbf{k} - \mathbf{k}' .$$

Hence:

$$\begin{aligned}\hat{H}_V &= \sum_{\mathbf{r} \in \mathcal{L}/2} \sum_{\sigma, \sigma'} h_{\mathbf{r}\sigma\sigma'} \\ &\stackrel{\mathcal{F}}{=} -\frac{V}{L_x L_y} \sum_{\sigma, \sigma'} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \cos(\delta \mathbf{k} \cdot \boldsymbol{\delta}) \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma'} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \\ &= -\frac{V}{L_x L_y} \sum_{\sigma, \sigma'} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma'} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma}. \quad (3.42)\end{aligned}$$

Finally, we can separate in same spin and opposite spin channels. This gives:

$$\begin{aligned}\hat{H}_V &\stackrel{\mathcal{F}}{=} \overbrace{-\frac{V}{L_x L_y} \sum_{\sigma} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{s.s.}} \\ &\quad - \underbrace{\frac{V}{L_x L_y} \sum_{\sigma} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\bar{\sigma}}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\bar{\sigma}} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma}}_{\text{o.s.}}.\end{aligned}$$

The s.s. part confirms Eq. (3.41). The o.s. part can be simplified: since

$$\hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\bar{\sigma}}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\bar{\sigma}} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} = \hat{c}_{\mathbf{K}-\mathbf{k}\bar{\sigma}}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \hat{c}_{\mathbf{K}-\mathbf{k}'\bar{\sigma}}$$

due to fermionic anti-commutation rules, and since the composite transform

$$\mathbf{k} \rightarrow -\mathbf{k} \quad \text{and} \quad \mathbf{k}' \rightarrow -\mathbf{k}'$$

leaves the form factor $\cos(\delta k_x) + \cos(\delta k_y)$ invariant, we reduce the spin sum to two times the $\sigma = \uparrow$ part. This confirms Eq. (3.40)

In reciprocal space, following the general decomposition presented in Eq. (C.9) for a quartic operator $\hat{c}_{\mathbf{k}_1}^\dagger \hat{c}_{\mathbf{k}_2}^\dagger \hat{c}_{\mathbf{k}_3} \hat{c}_{\mathbf{k}_4}$, we distinguish contractions as follows:

- Hartree contractions:

$$\overbrace{c_{\mathbf{k}_1}^\dagger c_{\mathbf{k}_2}^\dagger c_{\mathbf{k}_3} c_{\mathbf{k}_4}}^{\text{Hartree}} \quad \text{and} \quad \overbrace{\hat{c}_{\mathbf{k}_1}^\dagger \hat{c}_{\mathbf{k}_2}^\dagger c_{\mathbf{k}_3} \hat{c}_{\mathbf{k}_4}}^{\text{Hartree}}.$$

- Fock contractions:

$$\overbrace{c_{\mathbf{k}_1}^\dagger c_{\mathbf{k}_2}^\dagger c_{\mathbf{k}_3} \hat{c}_{\mathbf{k}_4}}^{\text{Fock}} \quad \text{and} \quad \overbrace{\hat{c}_{\mathbf{k}_1}^\dagger \hat{c}_{\mathbf{k}_2}^\dagger \hat{c}_{\mathbf{k}_3} c_{\mathbf{k}_4}}^{\text{Fock}}.$$

- Cooper contractions:

$$\overbrace{c_{\mathbf{k}_1}^\dagger c_{\mathbf{k}_2}^\dagger \hat{c}_{\mathbf{k}_3} \hat{c}_{\mathbf{k}_4}}^{\text{Cooper}} \quad \text{and} \quad \overbrace{\hat{c}_{\mathbf{k}_1}^\dagger \hat{c}_{\mathbf{k}_2}^\dagger c_{\mathbf{k}_3} c_{\mathbf{k}_4}}^{\text{Cooper}}.$$

with appropriate sign factors.

3.4 Square lattice spatial symmetry channels

In this section, we discuss the decomposition in different spatial symmetry channels for expectation values defined on the square lattice. We define the set of functions $\varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma)}$, with $\gamma \in \{s, s^*, p_x, p_y, d\}$, defined in Tab. 3.4 and plotted in Fig. 3.4. We will often refer to them as “harmonics”. By construction, these functions obey an orthonormality condition:

$$\frac{1}{L_x L_y} \sum_{\mathbf{r}\mathbf{r}'} \varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma)} \varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma')*} = \delta_{\gamma\gamma'}, \quad (3.43)$$

which is proven by direct computation.

Channel	Form factor	Graph
s -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(s)} = \delta_{\mathbf{r}\mathbf{r}'}$	Fig. 3.4a
s^* -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(s^*)} = \frac{1}{2} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 3.4b
p_x -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(p_x)} = \frac{1}{\sqrt{2}} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}}]$	Fig. 3.4c
p_y -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(p_y)} = \frac{1}{\sqrt{2}} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 3.4d
$d_{x^2-y^2}$ -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(d)} = \frac{1}{2} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}} - \delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 3.4e

Table 3.4 | The harmonics $\varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma)}$. The notation of Fig. 3.3 is used. In the last column, we indicate a graphic representation given in the subfigures Fig. 3.4.




Figure 3.4 | The harmonics listed in Tab. 3.4. In figures five sites are represented: the hub site i and its four NNs. Solid lines connect the hub sites to a rim site where the harmonic is non-zero. Dashed lines connect to a rim site where the harmonic is zero.

Decomposition in symmetry channels. Consider a generic function of two variables $f(\mathbf{r}, \mathbf{r}')$. Suppose that, for a given centre $\mathbf{r}$, the function is non-zero only on the site itself ($\mathbf{r}' = \mathbf{r}$) and its NNs $\mathbf{r}' = \mathbf{r} \pm \boldsymbol{\delta}$, with $\boldsymbol{\delta} \in \{\underline{x}, \underline{y}\}$. In other words,

$$f(\mathbf{r}, \mathbf{r}') \neq 0 \iff |\mathbf{r} - \mathbf{r}'| = 0, 1, \quad (3.44)$$

in units of the lattice step. Using the harmonics $\varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma)}$, we get the following decomposition:

$$f(\mathbf{r}, \mathbf{r}') = \sum_{\gamma} f^{(\gamma)} \varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma)}, \quad (3.45)$$

where we defined:

$$f^{(\gamma)} = \frac{1}{L_x L_y} \sum_{\mathbf{r}, \mathbf{r}'} f(\mathbf{r}, \mathbf{r}') \varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma)}. \quad (3.46)$$

Eq. (3.45) will be referred to as “decomposition in symmetry channels”. The sum over γ is extended to the five symmetries of Tab. 3.4.

Proof 3.10 — Decomposition in symmetry channels of Eq. (3.45).

Consider a function $f(\mathbf{r}, \mathbf{r}')$ defined as in Eq. (3.44) and the harmonics of Tab. 3.4. By direct computation, the following identity holds

$$\begin{aligned} f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{x}}) &= \sum_{\mathbf{r}' \in \mathcal{L}} f(\mathbf{r}, \mathbf{r}') \delta_{\mathbf{r}' = \mathbf{r} + \underline{\mathbf{x}}} \\ &= \sum_{\mathbf{r}' \in \mathcal{L}} f(\mathbf{r}, \mathbf{r}') \left(\varphi_{\mathbf{r}\mathbf{r}'}^{(s^*)} + \varphi_{\mathbf{r}\mathbf{r}'}^{(d)} + \sqrt{2} \varphi_{\mathbf{r}\mathbf{r}'}^{(p_x)} \right). \end{aligned}$$

Similarly, also the following relations hold

$$\begin{aligned} f(\mathbf{r}, \mathbf{r}) &= \sum_{\mathbf{r}' \in \mathcal{L}} f(\mathbf{r}, \mathbf{r}') \varphi_{\mathbf{r}\mathbf{r}'}^{(s)}, \\ f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{x}}) &= \sum_{\mathbf{r}' \in \mathcal{L}} f(\mathbf{r}, \mathbf{r}') \left(\varphi_{\mathbf{r}\mathbf{r}'}^{(s^*)} + \varphi_{\mathbf{r}\mathbf{r}'}^{(d)} - \sqrt{2} \varphi_{\mathbf{r}\mathbf{r}'}^{(p_x)} \right), \\ f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{y}}) &= \sum_{\mathbf{r}' \in \mathcal{L}} f(\mathbf{r}, \mathbf{r}') \left(\varphi_{\mathbf{r}\mathbf{r}'}^{(s^*)} - \varphi_{\mathbf{r}\mathbf{r}'}^{(d)} + \sqrt{2} \varphi_{\mathbf{r}\mathbf{r}'}^{(p_y)} \right), \\ f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{y}}) &= \sum_{\mathbf{r}' \in \mathcal{L}} f(\mathbf{r}, \mathbf{r}') \left(\varphi_{\mathbf{r}\mathbf{r}'}^{(s^*)} - \varphi_{\mathbf{r}\mathbf{r}'}^{(d)} - \sqrt{2} \varphi_{\mathbf{r}\mathbf{r}'}^{(p_y)} \right). \end{aligned}$$

Hence, by defining

$$\begin{aligned} f^{(s)} &\equiv f(\mathbf{r}, \mathbf{r}), \\ f^{(s^*)} &\equiv f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{x}}) + f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{x}}) + f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{y}}) + f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{y}}), \\ f^{(p_x)} &\equiv f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{x}}) - f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{x}}), \\ f^{(p_y)} &\equiv f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{y}}) - f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{y}}), \\ f^{(d)} &\equiv f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{x}}) + f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{x}}) - f(\mathbf{r}, \mathbf{r} + \underline{\mathbf{y}}) - f(\mathbf{r}, \mathbf{r} - \underline{\mathbf{y}}). \end{aligned}$$

we get the decomposition:

$$f(\mathbf{r}, \mathbf{r}') = \sum_{\gamma} f^{(\gamma)} \varphi_{\mathbf{r}\mathbf{r}'}^{(\gamma)}.$$

The function f depends only on the difference of its two spatial variables. In other words, all the $f^{(\gamma)}$ components we defined above are independent of $\mathbf{r}$. Hence we can redefine them as in Eq. (3.46), averaging over $\mathbf{r}$. The decomposition obtained above still holds. With this redefinition of the $f^{(\gamma)}$ components, we recognise Eq. (3.45).

We decomposed the original function in its harmonics. Moving to reciprocal space, we get the $\mathcal{F}$ -transformed functions of Tab. 3.5, also satisfying the orthonormality condition:

$$\frac{1}{L_x L_y} \sum_{\mathbf{k}} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma')*} = \delta_{\gamma\gamma'}.$$

The functions of Tab. 3.5 are plotted in the BZ in Fig. 3.5.

A useful identity. Using the harmonics functions, it is useful to prove the following identity:

$$\cos(\delta k_x) + \cos(\delta k_y) = \frac{1}{2} \sum_{\gamma \neq s} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} \quad (3.47)$$

$$= \frac{1}{2} \sum_{\gamma \neq s} \varphi_{\mathbf{k}}^{(\gamma)*} \varphi_{\mathbf{k}'}^{(\gamma)}. \quad (3.48)$$

Structure	Structure factor	Graph
s -wave	$\varphi_{\mathbf{k}}^{(s)} = 1$	Fig. 3.4a
Extended s -wave	$\varphi_{\mathbf{k}}^{(s^*)} = \cos k_x + \cos k_y$	Fig. 3.4b
p_x -wave	$\varphi_{\mathbf{k}}^{(p_x)} = i\sqrt{2} \sin k_x$	Fig. 3.4c
p_y -wave	$\varphi_{\mathbf{k}}^{(p_y)} = i\sqrt{2} \sin k_y$	Fig. 3.4d
$d_{x^2-y^2}$ -wave	$\varphi_{\mathbf{k}}^{(d)} = \cos k_x - \cos k_y$	Fig. 3.4e

Table 3.5 | $\mathcal{F}$ -transforms of the harmonics of Tab. 3.4. These functions are plotted in Fig. 3.5.




Figure 3.5 | Representation of the various structure factors listed in Tab. 3.5 on the BZ. For the s^* and $d_{x^2-y^2}$ -wave factors, the real part is plotted; for the p_ℓ factors, the imaginary part is plotted.

Note that in both rows above the sum is intended for $\gamma \in \{s^*, p_x, p_y, d_{x^2-y^2}\}$, excluding s .

Proof 3.11 — Proof of Eq. (3.47).

Let $k_\ell \in \mathbb{R}$, with $\ell \in \{x, y\}$. We define:

$$c_\ell \equiv \cos k_\ell, \quad s_\ell \equiv \sin k_\ell, \quad c'_\ell \equiv \cos k'_\ell, \quad s'_\ell \equiv \sin k'_\ell.$$

Let $\delta k_\ell \equiv k_\ell - k'_\ell$. Using basic trigonometric identities we get:

$$\cos(\delta k_x) + \cos(\delta k_y) = \underbrace{(c_x c'_x + c_y c'_y)}_{\text{Symmetric}} + \underbrace{(s_x s'_x + s_y s'_y)}_{\text{Anti-symmetric}}. \quad (3.49)$$

All four terms are invariant under simultaneous inversion of both momenta, $(\mathbf{k}, \mathbf{k}') \rightarrow (-\mathbf{k}, -\mathbf{k}')$. Instead, if we invert just one momentum of the pair, the first two are symmetric for both arguments $\mathbf{k}, \mathbf{k}'$; the second two are anti-symmetric. Decoupling the symmetric part, we obtain

$$c_x c'_x + c_y c'_y = \frac{1}{2}(c_x + c_y)(c'_x + c'_y) + \frac{1}{2}(c_x - c_y)(c'_x - c'_y),$$

which finally gives:

$$\begin{aligned} \cos(\delta k_x) + \cos(\delta k_y) &= \frac{1}{2}(c_x + c_y)(c'_x + c'_y) && (s^*\text{-wave}) \\ &+ s_x s'_x && (p_x\text{-wave}) \\ &+ s_y s'_y && (p_y\text{-wave}) \\ &+ \frac{1}{2}(c_x - c_y)(c'_x - c'_y), && (d_{x^2-y^2}\text{-wave}) \end{aligned}$$

which is written in compact form as in Eq. (3.47).

Chapter 4

Mott localisation in the normal phase

In this chapter, we study the normal phase, namely the phase characterised by the absence of any spontaneous symmetry breaking mechanism.

We show that the non-local pairing induces a renormalisation of the hopping amplitude, leading to a Mott localisation effect. The chapter is organised as follows. In Sec. 4.1 we discuss the symmetries of the normal phase. In Secs. 4.2, 4.3 we derive the HF approximation to the EHM in the normal phase. In Sec. 4.4 we analyse the self-consistency equations and derive constraints on the self-consistent parameters identified by the Hartree-Fock method. In Sec. 4.5 we present the result of numeric simulations.

4.1 The normal phase and its symmetries

In the normal phase no symmetry is broken, as summarised in Tab. 3.1. Therefore, the HF approximation to the hamiltonian must exhibit the full $U(2) \otimes C_4 \otimes T_1^{(x)} \otimes T_1^{(y)}$ symmetry discussed in Sec. 3.1.

We anticipate here that we will actually allow for $C_4 \xrightarrow{\text{SSB}} C_2$, namely, breaking of planar rotational symmetry. As will result from our discussion, the normal phase does not break C_4 even if we allow it to. Nevertheless, we carry out this general derivation because breaking of planar rotational symmetry will be observed in Chap. 6 in the context of superconductivity, and the computations carried out for the normal phase will be re-used.

Translational invariance implies the following result:

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'} \rangle = \delta_{\mathbf{k}=\mathbf{k}'} \delta_{\sigma=\sigma'} \langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \rangle. \quad (4.1)$$

Proof 4.1 — Proof of Eq. (4.1).

Consider the generic EV on the left side of Eq. (4.1)

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'} \rangle,$$

for two momenta $\mathbf{k}, \mathbf{k}'$. Moving to real space through an $\mathcal{F}$ -transform, we get

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'} \rangle \stackrel{\mathcal{F}}{=} \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} \sum_{\mathbf{r}' \in \mathcal{L}} e^{-i(\mathbf{k} \cdot \mathbf{r} - \mathbf{k}' \cdot \mathbf{r}')} \langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}'\sigma'} \rangle,$$

where we used the convention of Eq. (3.34). Let now $\Delta \mathbf{r} \equiv \mathbf{r}_2 - \mathbf{r}_1$. Translational invariance implies that the function

$$g(\Delta \mathbf{r}, \sigma, \sigma') = \langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\Delta \mathbf{r}\sigma'} \rangle$$

Operator	Sector	Channel	Net effect
$\hat{H}_U$		Hartree	μ shift by $-nU$
$\hat{H}_V$	o.s.	Hartree	μ shift by nzV
		Hartree	μ shift by nzV
	s.s.	Fock	Bands renormalisation and resymmetrization

Table 4.1 | Wick's contractions for each interaction and channel in the HF framework, and their net effect in the normal phase.

is independent of $\mathbf{r}$. Thus we have:

$$\begin{aligned} \langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'} \rangle &\stackrel{\mathcal{F}}{=} \sum_{\Delta\mathbf{r} \in \mathcal{L}} e^{-i\mathbf{k}' \cdot \Delta\mathbf{r}} g(\Delta\mathbf{r}, \sigma, \sigma') \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} e^{-i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}} \\ &= \sum_{\Delta\mathbf{r} \in \mathcal{L}} e^{-i\mathbf{k}' \cdot \Delta\mathbf{r}} g(\Delta\mathbf{r}, \sigma, \sigma') \delta_{\mathbf{k}=\mathbf{k}'}, \end{aligned}$$

where we used the identity of Eq. (3.35). Then the EV of Eq. (4.1) vanishes for $\mathbf{k} \neq \mathbf{k}'$.

Recall now Eq. (3.32). Upon substituting $\delta \rightarrow \Delta\mathbf{r}$, we obtain that $g(\Delta\mathbf{r}, \sigma, \bar{\sigma}) = 0$ for a phase that is SU(2)-symmetric. Then, the EV of Eq. (4.1) is non-zero only if $\sigma = \sigma'$. This proves Eq. (4.1).

Translational invariance and spin rotational symmetry imply that the site occupation is independent of site and spin indices,

$$\langle \hat{n}_{i\sigma} \rangle = n, \quad (4.2)$$

with $n \in [0, 1]$.

We define the single-state occupation operator in reciprocal space:

$$\hat{n}_{\mathbf{k}\sigma} \equiv \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}$$

and its EV

$$u_{\mathbf{k}\sigma} \equiv \langle \hat{n}_{\mathbf{k}\sigma} \rangle, \quad u_{\mathbf{k}\sigma} = u_{\mathbf{k}\sigma}^*. \quad (4.3)$$

The relation on the right follows from the fact that $\hat{n}_{\mathbf{k}\sigma}$ is hermitian by construction. Note that $\hat{n}_{\mathbf{k}\sigma}$ is not the Fourier-transform of $\hat{n}_{i\sigma}$. Finally, using the functions $\varphi_{\mathbf{k}}^{(\gamma)}$ of Tab. 3.5, we define the harmonics

$$u_{\sigma}^{(\gamma)} \equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \rangle. \quad (4.4)$$

These harmonics are the HFPs of the normal phase, and Eq. (4.4) is their self-consistency equation.

4.2 Application of Hartree-Fock methods to the normal phase

Applying the HF methods to the interactions $\hat{H}_U$ and $\hat{H}_V$, from Tabs. 3.2 and 3.3, we know that in the normal phase the allowed contractions are:

$$\begin{aligned} \hat{H}_U & \quad \text{Hartree contractions,} \\ \hat{H}_V \text{ (o.s. sector)} & \quad \text{Hartree contractions,} \\ \hat{H}_V \text{ (s.s. sector)} & \quad \text{Hartree and Fock contractions.} \end{aligned}$$

As we will see in detail and is summarised in Tab. 4.1, Hartree contractions give a chemical potential shift, while the Fock contractions renormalise the bare bands.

4.2.1 Effects of the Hartree terms: chemical potential renormalisation

In this section we show that the Hartree channels of $\hat{H}_U$ and $\hat{H}_V$ reduce to multiples of the number operator. Applying HF methods, we get the decomposition of Eqs. (2.21), (2.22). Specifically, the Hartree channels are given by

$$\begin{aligned}\hat{H}_U &\stackrel{\text{HF}}{\simeq} U \sum_i \left(\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \langle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \rangle + \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} - \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle \langle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \rangle \right), \\ \hat{H}_V &\stackrel{\text{HF}}{\simeq} -V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle \right) + (\text{Fock channel}).\end{aligned}$$

Summing the two terms, we obtain

$$\hat{H}_U + \hat{H}_V \stackrel{\text{HF}}{\simeq} -n(2zV - U)\hat{N} - \left(E_{\text{H},U}^{(\text{N})} + E_{\text{H},V}^{(\text{N})} \right) + (\text{Fock channel}), \quad (4.5)$$

where we introduce the ‘‘Hartree contraction energy’’, namely a shift to energy coming from the scalar terms of Eqs. (2.21), (2.22):

$$E_{\text{H},U}^{(\text{N})} = -U \sum_{i \in \mathcal{L}} \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle \langle \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \rangle, \quad (4.6)$$

$$E_{\text{H},V}^{(\text{N})} = V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle. \quad (4.7)$$

From Eq. (4.5), we see that the chemical potential is shifted as

$$\tilde{\mu} \equiv \mu + n(2zV - U), \quad (4.8)$$

as is reported in Tab. 4.1. This result is valid for all the other phases discussed in this thesis.

Proof 4.2 — Proof of Eq. (4.5): HF approximation of $\hat{H}_U + \hat{H}_V$ in Hartree channel.

We analyse $\hat{H}_U$ and $\hat{H}_V$ separately.

Local repulsion. In the normal phase, the local repulsion $\hat{H}_U$ decomposes in the Hartree channel as is specified in Eq. (2.21),

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} U \sum_i (\langle \hat{n}_{i\uparrow} \rangle \hat{n}_{i\downarrow} + \hat{n}_{i\uparrow} \langle \hat{n}_{i\downarrow} \rangle - \langle \hat{n}_{i\downarrow} \rangle \langle \hat{n}_{i\uparrow} \rangle).$$

Using Eq. (4.2), we obtain:

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} nU \sum_i (\hat{n}_{i\uparrow} + \hat{n}_{i\downarrow}) - E_{\text{H},U}^{(\text{N})} \quad (4.9)$$

$$= nU \times \hat{N} - E_{\text{H},U}^{(\text{N})}. \quad (4.10)$$

Non-local attraction. The non-local attraction is divided in two sectors, as in Eq. (3.25). From Eq. (2.22), its Hartree-like terms are:

$$\begin{aligned}\hat{H}_V &\stackrel{\text{HF}}{\simeq} \overbrace{-V \sum_{\langle ij \rangle} \sum_{\sigma} (\langle \hat{n}_{i\sigma} \rangle \hat{n}_{j\sigma} + \hat{n}_{i\sigma} \langle \hat{n}_{j\sigma} \rangle - \langle \hat{n}_{i\sigma} \rangle \langle \hat{n}_{j\sigma} \rangle)}^{\text{s.s.}} \\ &\quad \underbrace{-V \sum_{\langle ij \rangle} \sum_{\sigma} (\langle \hat{n}_{i\sigma} \rangle \hat{n}_{j\bar{\sigma}} + \hat{n}_{i\sigma} \langle \hat{n}_{j\bar{\sigma}} \rangle - \langle \hat{n}_{i\sigma} \rangle \langle \hat{n}_{j\bar{\sigma}} \rangle)}_{\text{o.s.}} + (\text{Fock channel}).\end{aligned} \quad (4.11)$$

Using Eq. (4.2), we get

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} \underbrace{-nV \sum_{\langle ij \rangle} \sum_{\sigma} (\hat{n}_{j\sigma} + \hat{n}_{i\sigma})}_{\text{s.s.}} \underbrace{-nV \sum_{\langle ij \rangle} \sum_{\sigma} (\hat{n}_{j\bar{\sigma}} + \hat{n}_{i\sigma}) - E_{\text{H},V}^{(N)}}_{\text{o.s.}} + (\text{Fock channel}), \quad (4.12)$$

Both sums give a number operator. Note that we sum over NNs: each site is connected to $z = 4$ sites. Including this combinatorial factor, we have

$$\hat{H}_V \simeq -2nzV \times \hat{N} - E_{\text{H},V}^{(N)} + (\text{Fock channel}). \quad (4.13)$$

Summing Eqs. (4.10) and (4.13), we obtain Eq. (4.5).

The contraction energies of Eqs. (4.6), (4.7) read

$$E_{\text{H},U}^{(N)} = -n^2 U \quad (4.14)$$

$$E_{\text{H},V}^{(N)} = L_x L_y \times 2zn^2 V. \quad (4.15)$$

where $z = 4$ is the lattice coordination number.

Proof 4.3 — Proof of Eqs. (4.14), (4.15): Hartree contraction energies.

Using Eq. (4.2) in Eq. (4.6), we obtain

$$\begin{aligned} E_{\text{H},U}^{(N)} &= -U \sum_{\mathbf{r} \in \mathcal{L}} n \\ &= -n^2 U, \end{aligned}$$

which proves Eq. (4.14). Doing the same in Eq. (4.7), we get

$$\begin{aligned} E_{\text{H},V}^{(N)} &= V \sum_{\langle ij \rangle} \sum_{\sigma} (\langle \hat{n}_{i\sigma} \rangle \langle \hat{n}_{j\sigma} \rangle + \langle \hat{n}_{i\sigma} \rangle \langle \hat{n}_{j\bar{\sigma}} \rangle) \\ &= 2V \sum_{\langle ij \rangle} \sum_{\sigma} n^2 \\ &= L_x L_y \times 2zn^2 V, \end{aligned}$$

where in last line we used Eq. (3.21). We obtained Eq. (4.15).

4.2.2 Effects of the Fock terms: bare bands renormalisation

From the Wick's decomposition of $\hat{H}_V$ (see Eq. (2.22)), the only Fock term allowed in the normal phase comes from the same-spin sector. See Tab. 3.3 for a summary. Explicitly,

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + V \sum_{\langle ij \rangle} \sum_{\sigma} \left(\langle \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} \rangle \hat{c}_{j\sigma}^{\dagger} \hat{c}_{i\sigma} + \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} \langle \hat{c}_{j\sigma}^{\dagger} \hat{c}_{i\sigma} \rangle - \langle \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} \rangle \langle \hat{c}_{j\sigma}^{\dagger} \hat{c}_{i\sigma} \rangle \right). \quad (4.16)$$

We notice that the *plus* sign in front of the displayed term comes from the *minus* sign in Fock term of Eq. (2.1), applied to the interaction $-V$.

Using the Fourier transform of $\hat{H}_V$ given in Eq. (3.41), we obtain

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} & (\text{Hartree channel}) \\ & + \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \sum_{\sigma} (\cos(\delta k_x) + \cos(\delta k_y)) \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} - E_{\text{F},V}^{(N)}. \end{aligned} \quad (4.17)$$




Figure 4.1 Representation of the variable change employed to pass from Eq. (4.20) to (4.21). In Fig. 4.1a generic vectors are considered, cycling over all values of $\mathbf{K}, \mathbf{k} \in \text{BZ}$. In Fig. 4.1b is depicted the variables change to the new vectors $\mathbf{q}, \mathbf{q}'$.

Here, the 2 prefactor comes from the fact that the first two operators in square brackets in Eq. (4.16) are equal in reciprocal space. The contraction energy shift is given by

$$E_{\text{F},V}^{(\text{N})} \equiv \frac{V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \sum_{\sigma} (\cos(\delta k_x) + \cos(\delta k_y)) \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \rangle. \quad (4.18)$$

By recalling the definition of $u_{\mathbf{k}\sigma} = \langle \hat{n}_{\mathbf{k}\sigma} \rangle$, the non local attraction is approximated, in Fock channel, by:

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + V \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \left(u_{\sigma}^{(s^*)} \varphi_{\mathbf{k}}^{(s^*)*} + u_{\sigma}^{(d)} \varphi_{\mathbf{k}}^{(d)*} \right) \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} - E_{\text{F},V}^{(\text{N})}, \quad (4.19)$$

where $u_{\sigma}^{(\gamma)}$ are the harmonics of the occupation $u_{\mathbf{k}\sigma}$, defined in Eq. (4.4).

Proof 4.4 — Proof of Eq. (4.19): HF approximation of $\hat{H}_V$ in Fock channel.

Consider Eq. (4.17). Applying Eq. (4.1), it becomes:

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} & (\text{Hartree channel}) \\ & + \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} (\cos 2k_x + \cos 2k_y) \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma} \rangle - E_{\text{F},V}^{(\text{N})}. \end{aligned} \quad (4.20)$$

Let now $\mathbf{q} \equiv \mathbf{K} + \mathbf{k}$, $\mathbf{q}' \equiv \mathbf{K} - \mathbf{k}$. A representation of this variables change is given in Fig. 4.1. Being (for $\ell = x, y$)

$$\begin{aligned} \delta q_\ell & \equiv q_\ell - q'_\ell \\ & = (K_\ell + k_\ell) - (K_\ell - k_\ell) \\ & = 2k_\ell, \end{aligned}$$

Symmetry	Self-consistency equation	Graph
s^* -wave	$u^{(s^*)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})$	Fig. 3.4b
$d_{x^2-y^2}$ -wave	$u^{(d)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})$	Fig. 3.4e

Table 4.2 | Symmetry resolved self-consistency equations for the harmonics of $u_{\mathbf{k}}$.

we reduce Eq. (4.17) to

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + \frac{2V}{L_x L_y} \sum_{\mathbf{q}, \mathbf{q}'} \sum_{\sigma} [\cos(\delta q_x) + \cos(\delta q_y)] \underbrace{\langle \hat{c}_{\mathbf{q}\sigma}^\dagger \hat{c}_{\mathbf{q}\sigma} \rangle}_{u_{\mathbf{q}\sigma}} \hat{c}_{\mathbf{q}'\sigma}^\dagger \hat{c}_{\mathbf{q}'\sigma} - E_{\text{F},V}^{(\text{N})}, \quad (4.21)$$

where we recognised $u_{\mathbf{q}\sigma}$, as was defined in Eq. (4.3). Using Eq. (3.47), we obtain

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + \sum_{\gamma, \sigma} \left(\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} u_{\mathbf{q}\sigma} \right) \times V \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} \hat{c}_{\mathbf{q}'\sigma}^\dagger \hat{c}_{\mathbf{q}'\sigma} - E_{\text{F},V}^{(\text{N})}, \quad (4.22)$$

where we used the functions defined in Tab. 3.5. We recognise the harmonics $u_{\sigma}^{(\gamma)}$ from Eq. (4.4),

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + V \sum_{\mathbf{q}' \in \text{BZ}} \sum_{\gamma, \sigma} u_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}'}^{(\gamma)*} \hat{c}_{\mathbf{q}'\sigma}^\dagger \hat{c}_{\mathbf{q}'\sigma} - E_{\text{F},V}^{(\text{N})}, \quad (4.23)$$

which is almost Eq. (4.19).

To complete the proof, we recognise that in the normal phase inversion symmetry is not broken, thus

$$u_{\mathbf{q}\sigma} = u_{-\mathbf{q}\sigma}.$$

This follows from the fact that in the normal phase the EV of the operator $\hat{n}_{\mathbf{k}\sigma}$ must be inversion-symmetric. It follows that antisymmetric harmonics are zero,

$$u_{\sigma}^{(p_\ell)} = 0 \quad \text{for } \ell \in \{x, y\}. \quad (4.24)$$

Using this observation in Eq. (4.23), we obtain Eq. (4.19).

Bringing together Eq. (4.19) with Eqs. (4.10), (4.13) we obtain that the quadratic HF approximation to the EHM, for the normal phase, depends on the renormalised chemical potential $\tilde{\mu}$, and the occupation harmonics $u^{(s^*)}$ and $u^{(d)}$. The chemical potential $\tilde{\mu}$ is fixed by density and determined during computations. To find the ground-state, we must determine self-consistently the two HFPs $u_{\sigma}^{(s^*)}$, $u_{\sigma}^{(d)}$ through the self-consistency Eq. (4.4). The symmetry-resolved equations are reported in Tab. 4.2.

The operator of Eq. (4.19) has the same algebraic structure of the kinetic operator $\hat{H}_t$ in reciprocal space, given in Eq. (3.37). We can collapse the two into one effective kinetic operator with renormalised bands. Therefore,

$$\tilde{\epsilon}_{\mathbf{k}\sigma} \equiv \epsilon_{\mathbf{k}\sigma} + u_{\sigma}^{(s^*)} V (\cos k_x + \cos k_y) + u_{\sigma}^{(d)} V (\cos k_x - \cos k_y). \quad (4.25)$$

In Fock channel, we have:

$$\hat{H}_t + \hat{H}_V \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + \underbrace{\sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \tilde{\epsilon}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma}}_{\hat{H}_{\tilde{t}}} - E_{\text{F},V}^{(\text{N})}, \quad (4.26)$$

where we defined the effective kinetic operator, $\hat{H}_{\tilde{t}}$.

The renormalised bands of Eq. (4.25) are a sum of s^* and d -wave components. Instead, bare bands as defined Eq. (3.38) are purely s^* -symmetric functions,

$$\epsilon_{\mathbf{k}\sigma} = -2t(\cos k_x + \cos k_y) = -4t\varphi_{\mathbf{k}}^{(s^*)}. \quad (4.27)$$

We define the symmetry-resolved hoppings,

$$\tilde{t}_{\sigma}^{(s^*)} \equiv t - \frac{u_{\sigma}^{(s^*)}V}{2} \quad \text{and} \quad \tilde{t}_{\sigma}^{(d)} \equiv -\frac{u_{\sigma}^{(d)}V}{2}, \quad (4.28)$$

so that the bands renormalisation of Eq. (4.25) can be written as

$$\tilde{\epsilon}_{\mathbf{k}\sigma} \equiv -4\tilde{t}_{\sigma}^{(s^*)}\varphi_{\mathbf{k}}^{(s^*)} - 4\tilde{t}_{\sigma}^{(d)}\varphi_{\mathbf{k}}^{(d)}. \quad (4.29)$$

We notice the formal similarity with Eq. (4.27), showing how bands renormalisation reduces to a renormalisation of the hopping parameter t ,

$$t \text{ (} s^* \text{-wave)} \rightarrow \tilde{t}_{\sigma}^{(s^*)} \text{ (} s^* \text{-wave)} \text{ and } \tilde{t}_{\sigma}^{(d)} \text{ (} d \text{-wave)}.$$

As we will discuss in Sec. 4.3.1, the presence of $\tilde{t}^{(d)} \neq 0$ implies a nematic state that breaks C_4 .

We conclude this section by proving that the Fock contraction energy is given by:

$$E_{\text{F},V}^{(\text{N})} = L_x L_y \times \frac{V}{2} \sum_{\sigma} \left(|u_{\sigma}^{(s^*)}|^2 + |u_{\sigma}^{(d)}|^2 \right). \quad (4.30)$$

Later, using spin degeneracy, we will omit the sum on σ and the $1/2$ prefactor.

Proof 4.5 — Proof of Eq. (4.30): non-local Fock contraction energy.

Recall Eq. (4.18). Using Eq. (4.1), it reduces to:

$$E_{\text{F},V}^{(\text{N})} = \frac{V}{L_x L_y} \sum_{\mathbf{q}, \mathbf{q}'} \sum_{\sigma} [\cos(\delta q_x) + \cos(\delta q_y)] \langle \hat{c}_{\mathbf{q}\sigma}^{\dagger} \hat{c}_{\mathbf{q}\sigma} \rangle \langle \hat{c}_{\mathbf{q}'\sigma}^{\dagger} \hat{c}_{\mathbf{q}'\sigma} \rangle.$$

Inserting Eq. (4.3), we get:

$$\begin{aligned} E_{\text{F},V}^{(\text{N})} &= L_x L_y \times \frac{V}{2} \sum_{\gamma, \sigma} \left(\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} u_{\mathbf{q}\sigma} \right) \left(\frac{1}{L_x L_y} \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} u_{\mathbf{q}'\sigma}^* \right) \\ &= L_x L_y \times \frac{V}{2} \sum_{\gamma, \sigma} |u_{\sigma}^{(\gamma)}|^2, \end{aligned}$$

which, using the fact that $u_{\sigma}^{(p_{\ell})} = 0$, proves Eq. (4.30).

4.3 The HF hamiltonian and its properties

Collecting all the various terms, the HF approximation to the EHM hamiltonian in the normal phase is given by

$$\hat{H} - \mu \hat{N} \stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \tilde{\epsilon}_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} - \tilde{\mu} \hat{N} + E_c^{(\text{N})}, \quad (4.31)$$

where in first line we included the all contraction energies from Eqs. (4.14), (4.15) and (4.30) in $E_c^{(N)}$,

$$E_c^{(N)} \equiv - \left(E_{H,U}^{(N)} + E_{H,V}^{(N)} + E_{F,V}^{(N)} \right). \quad (4.32)$$

All dependence on the HFPs is inside the renormalised bands of Eq. (4.25) and the contributions to contraction energy.

Apart from energy shifts, this hamiltonian describes a perfectly degenerate Fermi gas whose bare bands are $\tilde{\epsilon}_{\mathbf{k}\sigma}$ as defined in Eq. (4.25), and whose chemical potential is $\tilde{\mu}$ as defined in Eq. (4.8). The latter is fixed self-consistently by setting the density; the former must be determined using self-consistency equations. The thermal state of the approximated hamiltonian obeys Fermi-Dirac statistics,

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'} \rangle = \delta_{\mathbf{k}=\mathbf{k}'} \delta_{\sigma=\sigma'} f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}), \quad (4.33)$$

being f the Fermi-Dirac distribution,

$$f(\epsilon; \beta, \mu) \equiv \frac{1}{e^{\beta(\epsilon-\mu)} + 1}.$$

In next subsections, we discuss in detail the renormalisation of the hopping parameter in real space, with a focus on the meaning of the d -wave component, and derive the free energy and entropy densities for the ground-state.

4.3.1 Nematicity: d -wave bands and anisotropic hopping

In Eq. (4.26) we defined $\hat{H}_{\tilde{t}}$ in reciprocal space. In real space, $\hat{H}_{\tilde{t}}$ takes the form

$$\begin{aligned} \hat{H}_{\tilde{t}} &= \hat{H}_t + V \sum_{\langle ij \rangle} \sum_{\sigma} \left[\langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} \rangle \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} + \text{h.c.} \right] \\ &= - \sum_{\langle ij \rangle} \sum_{\sigma} \left[\tilde{t}_{ij\sigma} \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} + \text{h.c.} \right]. \end{aligned}$$

where we introduce the renormalised hopping $\tilde{t}_{ij\sigma}$,

$$\tilde{t}_{ij\sigma} \equiv \begin{cases} \frac{1}{2} \left(t - V \langle \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} \rangle \right) & \text{if } i, j \text{ are NNs,} \\ 0 & \text{otherwise.} \end{cases}$$

Due to translational invariance, the renormalised hopping $\tilde{t}_{ij\sigma}$ depends only on the difference $i - j$. Therefore we can use the decomposition of Eq. (3.45),

$$\tilde{t}_{ij\sigma} = \sum_{\gamma} \tilde{t}_{\sigma}^{(\gamma)} \varphi_{ij}^{(\gamma)}.$$

The harmonics, given by Eq. (3.46), are identified with $\tilde{t}_{\sigma}^{(s^*)}$, $\tilde{t}_{\sigma}^{(d)}$ already introduced in Eq. (4.28). From the same argument we know that $\tilde{t}_{\sigma}^{(p_x)} = \tilde{t}_{\sigma}^{(p_y)} = 0$. Then, the renormalised hopping is explicitly given by

$$\begin{aligned} \tilde{t}_{\mathbf{r}\mathbf{r}'\sigma} &= \frac{\tilde{t}_{\sigma}^{(s^*)}}{2} \left[\delta_{\mathbf{r}'=\mathbf{r}+\underline{\mathbf{x}}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{\mathbf{x}}} + \delta_{\mathbf{r}'=\mathbf{r}+\underline{\mathbf{y}}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{\mathbf{y}}} \right] \\ &\quad + \frac{\tilde{t}_{\sigma}^{(d)}}{2} \left[\delta_{\mathbf{r}'=\mathbf{r}+\underline{\mathbf{x}}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{\mathbf{x}}} - \delta_{\mathbf{r}'=\mathbf{r}+\underline{\mathbf{y}}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{\mathbf{y}}} \right]. \end{aligned}$$

Let us define the direction-resolved renormalised hopping,

$$\tilde{t}_{\sigma}^{(x)} \equiv \frac{\tilde{t}_{\sigma}^{(s^*)} + \tilde{t}_{\sigma}^{(d)}}{2}, \quad (4.34)$$

$$\tilde{t}_{\sigma}^{(y)} \equiv \frac{\tilde{t}_{\sigma}^{(s^*)} - \tilde{t}_{\sigma}^{(d)}}{2}. \quad (4.35)$$




Figure 4.2 | Graphic rendition of anisotropic hopping, with a s^* -wave component (symmetric by $\pi/2$ rotations) and a d -wave component (antisymmetric).

This shows that, as anticipated, a finite $\tilde{t}^{(d)} \neq 0$ implies a nematic order characterised by the breaking of planar rotational symmetry, highlighted by the fact that kinetic operators in directions x and y differ. We can rewrite $\hat{H}_{\tilde{t}}$ as:

$$\hat{H}_{\tilde{t}} = - \sum_{\mathbf{r} \in \mathcal{L}} \sum_{\sigma} \left[\tilde{t}_{\sigma}^{(x)} \left(\hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}+\underline{x}\sigma} + \hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}-\underline{x}\sigma} \right) + \tilde{t}_{\sigma}^{(y)} \left(\hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}+\underline{y}\sigma} + \hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}-\underline{y}\sigma} \right) \right].$$

A schematics of anisotropic hopping is given in Fig. 4.2: for each bond linking two sites, the hopping parameter can differ, as the link direction is vertical or horizontal. We notice, however, that even in the presence of a non-zero d -wave hopping the density remains uniform.

Proof 4.6 — A d -wave hopping is compatible with uniform density.

Consider the on-site occupation operator $\hat{n}_{\mathbf{r}\sigma} = \hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}\sigma}$. Applying a Fourier transform as defined in Eq. (3.34), we obtain

$$\hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}\sigma} \stackrel{\mathcal{F}}{=} \frac{1}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} e^{i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}'\sigma}.$$

We compute the expectation value of the above operator and apply Eq. (4.33),

$$\begin{aligned} \langle \hat{n}_{\mathbf{r}\sigma} \rangle &= \frac{1}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} e^{i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}} \langle \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}'\sigma} \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}). \end{aligned}$$

The right-hand side of last line is independent of $\mathbf{r}$, therefore also $\langle \hat{n}_{\mathbf{r}\sigma} \rangle$ is.

Note that we never used the actual symmetry of the renormalised bands $\tilde{\epsilon}_{\mathbf{k}}$, but only the fact that the thermal state obeys the Fermi-Dirac distribution. A perfectly degenerate Fermi gas is always translational invariant and thus cannot exhibit non-uniform charge density.

From now on, we will omit the spin index σ from both the renormalised bands and the renormalised hopping,

$$\tilde{\epsilon}_{\mathbf{k}\sigma} \rightarrow \tilde{\epsilon}_{\mathbf{k}} \quad \forall \alpha : \quad \tilde{t}_{\sigma}^{(\alpha)} \rightarrow \tilde{t}^{(\alpha)}.$$

The normal phase does not break spin rotational invariance. Therefore the renormalised bands must be independent of the spin index. Inverting Eq. (4.28), we can express $\tilde{t}^{(x)}$ and $\tilde{t}^{(y)}$ in terms of the HFPs $u^{(s^*)}$ and $u^{(d)}$

$$\begin{aligned} \tilde{t}^{(x)} &\equiv t - \frac{u^{(s^*)} + u^{(d)}}{2} V, \\ \tilde{t}^{(y)} &\equiv t - \frac{u^{(s^*)} - u^{(d)}}{2} V. \end{aligned}$$

Inverting these relations we obtain:

$$u^{(s^*)} = \frac{(t - \tilde{t}^{(x)}) + (t - \tilde{t}^{(y)})}{V}, \quad (4.36)$$

$$u^{(d)} = \frac{\tilde{t}^{(x)} - \tilde{t}^{(y)}}{V}. \quad (4.37)$$

It follows that if $u^{(d)} > 0$, the effective hopping parameter is larger in direction x with respect to direction y . If $u^{(d)} < 0$, roles are inverted.

4.3.2 The entropy density of the normal ground-state

In this section, we compute the density of entropy of the ground state in the normal phase. Entropy density is given by

$$s^{(N)} = \frac{S^{(N)}}{L_x L_y},$$

being $S^{(N)}$ the total entropy.

The HF hamiltonian of Eq. (4.31) describes a non-interacting Fermi gas. The entropy density of a free Fermi gas in its ground state [8] is

$$s^{(N)} = -\frac{2k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} [(1 - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})) \ln(1 - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})) + f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \ln f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})]. \quad (4.38)$$

The 2 prefactor was included to account for spin degeneracy. The above equation can be written in another way:

$$s^{(N)} = -\frac{2k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} [\ln(1 - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})) - \beta(\tilde{\epsilon}_{\mathbf{k}} - \tilde{\mu})f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})]. \quad (4.39)$$

Proof 4.7 — Proof of Eq. (4.39): entropy density for the normal phase.

The following identity holds,

$$\begin{aligned} \ln\left(1 - \frac{1}{e^x + 1}\right) &= \ln\left(\frac{e^x}{e^x + 1}\right) \\ &= x + \ln\left(\frac{1}{e^x + 1}\right). \end{aligned} \quad (4.40)$$

Substituting $x = \beta(\tilde{\epsilon}_{\mathbf{k}} - \tilde{\mu})$ and using Eq. (4.40) in Eq. (4.38) we obtain

$$\begin{aligned} s^{(N)} &= -\frac{2k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[\ln(1 - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})) + f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \ln f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \right. \\ &\quad \left. - \beta(\tilde{\epsilon}_{\mathbf{k}} - \tilde{\mu})f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \ln f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \right]. \end{aligned}$$

Second and fourth terms cancel out. The remaining parts give Eq. (4.39).

4.3.3 The free energy density of the normal ground-state

In this section we compute the free energy density of the ground state, for the normal phase. The density of free energy is given by

$$f^{(N)} \equiv \frac{F^{(N)}}{L_x L_y}, \quad (4.41)$$

being $F^{(N)}$ the total free energy,

$$F^{(N)} = E_{\text{HF}}^{(N)} - \frac{S^{(N)}}{k_B \beta}. \quad (4.42)$$

Here $E_{\text{HF}}^{(\text{N})}$ is the ground-state energy for the HF hamiltonian. In the normal phase, we compute $f^{(\text{N})}$ using the following formula:

$$f^{(\text{N})} = \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln(1 - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})) - V \left(|u^{(s^*)}|^2 + |u^{(d)}|^2 \right) + \tilde{\mu} n. \quad (4.43)$$

Proof 4.8 — Proof of Eq. (4.43): free energy density for the normal phase.

Using Eqs. (4.41), (4.42), we have

$$f^{(\text{N})} = \frac{E_{\text{HF}}^{(\text{N})}}{L_x L_y} - \frac{s^{(\text{N})}}{k_B \beta},$$

with $s^{(\text{N})}$ the entropy density computed in Eq. (4.39). Then, $f^{(\text{N})}$ is obtained by summing the following terms:

1. The free particles ground state energy density,

$$e_g^{(\text{N})} = \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\tilde{\epsilon}_{\mathbf{k}} - \tilde{\mu}) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}).$$

The prefactor 2 accounts for spin degeneracy.

2. The contraction energy density,

$$\begin{aligned} e_c^{(\text{N})} &\equiv \frac{E_c^{(\text{N})}}{L_x L_y} \\ &= -n^2 U + V \left(2zn^2 - |u^{(s^*)}|^2 - |u^{(d)}|^2 \right), \end{aligned}$$

where we used Eq. (4.32).

3. The entropic contribution to free energy

$$\begin{aligned} e_s^{(\text{N})} &\equiv -\frac{s^{(\text{N})}}{k_B \beta} \\ &= \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln(1 - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})) - \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\tilde{\epsilon}_{\mathbf{k}} - \tilde{\mu}) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}). \end{aligned}$$

We used Eq. (4.39).

4. The chemical potential correction

$$e_n^{(\text{N})} = \mu n.$$

We explained at the end of Sec. 2.3.1 why this correction is needed. Note here that we are using the “physical” value of μ , not its RMP counterpart $\tilde{\mu}$.

Thus, composing everything

$$\begin{aligned} f^{(\text{N})} &\equiv e_g^{(\text{N})} + e_c^{(\text{N})} + e_s^{(\text{N})} + e_n^{(\text{N})} \\ &= \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln(1 - f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})) - V \left(|u^{(s^*)}|^2 + |u^{(d)}|^2 \right) + [\mu + n(2zV - U)] n. \end{aligned}$$

Using Eq. (4.8), we recognise

$$\mu + n(2zV - U) = \tilde{\mu}.$$

Hence Eq. (4.43) is proven.




Figure 4.3 | Graphic representation of the shrinking of renormalised band $\tilde{\epsilon}_{\mathbf{k}} = -4t\varphi_{\mathbf{k}}^{(s^*)}$, as the hopping parameter is decreased by renormalisation.

4.4 Constraints on the self-consistent solution

In this section, we collect some constraints on the HFPs $u^{(s^*)}$, $u^{(d)}$, that follow from a formal analysis of the self-consistency problem for the normal phase. It is possible to prove that:

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} = 0, \quad (4.44)$$

for any choice of $U, V > 0$ and doping δ . The proof is reported in App. E. Eq. (4.44) implies that in the normal phase there is no nematic effect, and the renormalised hopping parameter $\tilde{t}_{ij}$ is isotropic with an amplitude smaller than t ,

$$0 < \tilde{t}^{(s^*)} < t.$$

Here we used Eq. (4.28). Then, bands renormalisation

$$\epsilon_{\mathbf{k}} \rightarrow \tilde{\epsilon}_{\mathbf{k}},$$

is actually a flattening. We denote this effect as “bands shrinking”, and show a schematics in Fig. 4.3. The flattening of electronic bands is a signature of a Mott transition, as we discussed in Chap. 1.

From Eq. (4.28) we notice that, if the converged value of $u^{(s^*)}$ approaches its boundaries, we have

$$\begin{aligned} u^{(s^*)} \rightarrow 0 & \implies \tilde{t}^{(s^*)} \rightarrow t, \\ u^{(s^*)} \rightarrow \frac{2t}{V} & \implies \tilde{t}^{(s^*)} \rightarrow 0. \end{aligned}$$

In the second line, we see that the upper boundary can be interpreted as a critical value for Mott transition. Hence we define

$$u_c^{(s^*)}(V) \equiv \frac{2t}{V}. \quad (4.45)$$

If $u^{(s^*)} \rightarrow u_c^{(s^*)}(V)$ (from below), bands are completely flattened.

We conclude this section by proving some analytic properties of $u^{(s^*)}$. Its self-consistency equation, from Tab. 4.2, reads

$$u^{(s^*)}(\beta, \tilde{\mu}) = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}), \quad (4.46)$$

where we explicitly highlighted the dependence of $u^{(s^*)}$ on the chemical potential and the temperature. We also notice that the renormalised bands, inside the Fermi-Dirac factor on the right-hand side, depend on $u^{(s^*)}$. We now prove that in the low-temperature limit the above equation implies that $u^{(s^*)}$ is a monotonically descendent function of doping,

$$\frac{\partial}{\partial \delta} u^{(s^*)}(\beta, \tilde{\mu}) \leq 0, \quad (4.47)$$

with the equality being verified when bands are completely flattened. The fact that $u^{(s^*)}$ cannot increase as we increase δ implies that the bands shrinking effect must be suppressed as we move away from half-filling.

Proof 4.9 — Proof of Eq. (4.47): $u^{(s^*)}$ decreases with doping at low temperature.

Whatever the shape of the renormalised bands, the chemical potential cannot decrease with doping,

$$\frac{\partial \tilde{\mu}}{\partial \delta} \geq 0,$$

the equality holding only when adding electrons the Fermi level does not rise – namely, when the band is flat. Moreover,

$$\frac{\partial}{\partial \delta} u^{(s^*)}(\beta, \tilde{\mu}) = \frac{\partial \tilde{\mu}}{\partial \delta} \frac{\partial}{\partial \tilde{\mu}} u^{(s^*)}(\beta, \tilde{\mu}).$$

Then, if we shows that $\partial u^{(s^*)} / \partial \tilde{\mu} < 0$, Eq. (4.47) is proven.

Upon taking the derivative of Eq. (4.46), we obtain

$$\begin{aligned} \frac{\partial}{\partial \tilde{\mu}} u^{(s^*)}(\beta, \tilde{\mu}) &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) \frac{\partial}{\partial \tilde{\mu}} f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \\ &= -\frac{\beta}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) \frac{e^{\beta(\tilde{\epsilon}_{\mathbf{k}} - \tilde{\mu})}}{[e^{\beta(\tilde{\epsilon}_{\mathbf{k}} - \tilde{\mu})} + 1]^2} < 0 \end{aligned}$$

which ends the proof.

This is the first expected behaviour of the self-consistent functional $u^{(s^*)}$: if there exists a region of parametric space where the shrinking effect is not strong enough to flatten the band, a rise in the doping level must lower $u^{(s^*)}$. Such a region must exist: at least all the segment $(V = 0, \delta)$ is included, since on said region of parametric space the HF hamiltonian reduces to the that of the free Fermi gas.

We conclude this section with the analytic computation of $u^{(s^*)}$ at half filling and unitary filling, both at low temperature:

$$\lim_{\beta \rightarrow \infty} u^{(s^*)}[\beta, 0] = \left(\frac{2}{\pi}\right)^2 \approx 0.405, \quad (4.48)$$

$$\lim_{\beta \rightarrow \infty} u^{(s^*)}[\beta, 2\tilde{t}^{(s^*)}] = 0, \quad (4.49)$$

where, at unitary filling, we set the chemical potential equal to the maximum energy reached by a band of amplitude $4\tilde{t}^{(s^*)}$. All the properties hereby discussed are independent of the sign of δ . Namely, all we derived is symmetric as we add or remove electrons.

In order to prove Eq. (4.48), it is necessary to introduce the Magnetic Brillouin Zone (MBZ):

$$\text{MBZ} \equiv \left\{ \mathbf{k} \in \text{BZ} \mid \varphi_{\mathbf{k}}^{(s^*)} \geq 0 \right\}. \quad (4.50)$$

In Chap. 5 the reason for such a name will be clarified. The MBZ coincides with the set of points (k_x, k_y) that satisfy the relation

$$|k_x| + |k_y| \leq \pi.$$




Figure 4.4 | Representation of the MBZ. The dashed lines represent the boundaries of the BZ.

We draw the MBZ in Fig. 4.4. The renormalised bands satisfy

$$\tilde{\epsilon}_{\mathbf{k}} = -4\tilde{t}^{(s^*)}\varphi_{\mathbf{k}}^{(s^*)}.$$

Hence, $\tilde{\epsilon}_{\mathbf{k}} < 0$ inside the MBZ and $\tilde{\epsilon}_{\mathbf{k}} > 0$ outside. Along the boundary ∂MBZ we have $\tilde{\epsilon}_{\mathbf{k}} = 0$.

Proof 4.10 — Proof of Eqs. (4.48), (4.49).

The two proofs are the following:

1. At half-filling, the Fermi surface coincides with the boundary of the MBZ. Hence, being the Fermi-Dirac distribution a step-function at low temperatures, Eq. (4.46) reduces to

$$\lim_{\beta \rightarrow \infty} u^{(s^*)}(\beta, 0) \simeq \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x + \cos k_y).$$

In thermodynamic limit, performing integration on the MBZ shown in Fig. 4.4, we obtain

$$\begin{aligned} \lim_{\beta \rightarrow \infty} u^{(s^*)}(\beta, 0) &\simeq \frac{1}{(2\pi)^2} \int_{-\pi}^{\pi} dk_y \int_{-\pi+|k_y|}^{\pi-|k_y|} dk_x \cos k_x + \frac{1}{(2\pi)^2} \int_{-\pi}^{\pi} dk_x \int_{-\pi+|k_x|}^{\pi-|k_x|} dk_y \cos k_y \\ &= \frac{1}{2\pi^2} \int_{-\pi}^0 dk_y \int_{-\pi-k_y}^{\pi+k_y} dk_x \cos k_x + \frac{1}{2\pi^2} \int_0^{\pi} dk_y \int_{-\pi+k_y}^{\pi-k_y} dk_x \cos k_x \\ &= \frac{1}{\pi^2} \int_{-\pi}^0 dk_y \sin(\pi + k_y) + \frac{1}{\pi^2} \int_0^{\pi} dk_y \sin(\pi - k_y) \\ &= \left(\frac{2}{\pi}\right)^2, \end{aligned}$$

which proves Eq. (4.48).

2. At unitary filling we have

$$\lim_{\beta \rightarrow \infty} u^{(s^*)}(\beta, 2\tilde{t}^{(s^*)}) = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) = 0,$$

which vanishes due to the structure factor periodicity by π . This proves Eq. (4.49).




Figure 4.5 | Numerical results for $u^{(s^*)}$. On the left: colour plot in the (δ, V) plane. On the right: line plot at doubled lattice size, varying δ . The theoretical value for $u^{(s^*)}$ computed in Eqs. (4.48) at half-filling is plotted as a green dashed line.

4.5 Discussion of numeric results

In this section we present and discuss the numeric results we obtained for the normal phase. In the end we present a concrete scenario where careful tuning of the algorithm is necessary to ensure convergence. As is discussed above, in the normal phase the HFPs vector (defined in Sec. 2.3.2) has just one component:

$$\mathbf{v} = [u^{(s^*)}] .$$

We have two RMPs,

$$\begin{aligned} \tilde{t}_{ij} &\equiv \left(t - \frac{u^{(s^*)}V}{2} \right) \varphi_{ij}^{(s^*)} , \\ \tilde{\mu} &\equiv \mu + n(2zV - U) , \end{aligned}$$

the second one being only dependent on the model parameters and not on HFPs. Finally, the phase free energy is given by Eq. (4.43). The normal phase is algebraically independent of U . Therefore, we present the results varying the other parameters, keeping everywhere $U = 0$. We ran simulations in the (V, δ) plane, with the setup:

$$U = 0, \quad \Delta n = 10^{-2}, \quad \Delta \mathbf{v} = [u^{(s^*)} : 5 \times 10^{-4}] , \quad \beta = 100.0 .$$


Here $\Delta \mathbf{v}$ is the tolerance vector, defined as in Sec. 2.3.2. We have set $\beta = 100.0$ as an (arbitrary) approximation of zero temperature. We measure temperature in adimensional units,

$$\frac{\beta}{\beta_0} \rightarrow \beta$$

being $\beta_0 = 1/(1\text{K} \times k_B)$.

The trend of Eq. (4.47) is confirmed in Fig. 4.5, where we show $u^{(s^*)}$ as a function of V and varying δ , with $\beta = 100$. We see that $u^{(s^*)}$ decreases as we increase δ while keeping V fixed. In the colour plot of Fig. 4.5a we observe that in the top-left corner (low- δ and high- V limit) there is a brighter region in which, increasing δ while keeping V fixed, the converged value of $u^{(s^*)}$ is approximately constant. This effect is more visible in Fig. 4.5b, where we doubled L and ran simulations with a smaller δ resolution. At small values of V , the convergence value of $u^{(s^*)}$ is approximately independent of V . Note that, as was expected from the discussion of Sec. 4.4, the half-filling convergence value of $u^{(s^*)}$ in the V -independent region is approximately given by Eq. (4.48), which we report in Fig. 4.5b as an horizontal dashed green line. We did not find significant discrepancies in the converged data for $L = 128$ and $L = 256$.

For larger values of V , $u^{(s^*)}$ starts to decrease monotonically, consistently with the constraints of Eq. (4.44). We sketch the upper boundary $u_c^{(s^*)}(V) = 2t/V$, defined in Eq. (4.45), in Fig. 4.5b



a | Numerical results for $\tilde{t}^{(s^*)}$ b | Critical Mott boundary.
 Figure 4.6 | Plots of $\tilde{t}^{(s^*)}$ and the Mott-boundary V_c , computed as in Eq. (4.51).

as a black dashed line. We observe that when the red lines (numerical results) reach the black line (theoretical upper boundary), we have

$$u^{(s^*)} \lesssim \frac{2t}{V} \quad \Rightarrow \quad \tilde{t}^{(s^*)} \approx 0$$

Interestingly, this would correspond to a complete flattening of the renormalised bands.

4.5.1 Mott localisation

In Sec. 1.1.1 we introduced Mott insulation, and argued that the flattening of bands is a signature of a Mott transition. Interestingly, our results suggest that Mott localisation is induced by the non-local attraction $-V$, instead of the usual Mott localisation induced by the local repulsion $+U$.

In Fig. 4.5b, we see that the converged value of $u^{(s^*)}$ is approximately horizontal up to the boundary $u_c^{(s^*)} = 2t/V$, represented as a dashed black line. Therefore, in the region where it is constant, we have

$$u^{(s^*)}[\beta, \tilde{\mu}(\delta)]|_V = u^{(s^*)}[\beta, \tilde{\mu}(\delta)]|_{V=0}.$$

We define the critical Mott boundary as:

$$V_c[\beta, \delta] \equiv \frac{2t}{u^{(s^*)}[\beta, \tilde{\mu}(\delta)]|_{V=0}}. \quad (4.51)$$

When V becomes larger than this critical value, due to the constraint of Eq. (4.44), $u^{(s^*)}$ is forced to decrease while $\tilde{t}^{(s^*)}$ drops to 0. In Fig. 4.6a the converged values of $\tilde{t}^{(s^*)}$, extracted from the same data of Fig. 4.5a, are plotted in parametric space.

The localisation effect is particularly effective at half-filling, where the kinetic energy vanishes for $V/t \approx 5$. This value needs to be compared with the critical value of the local repulsion U_c for Mott localisation. In DMFT, $U_c \simeq 3W$, with W half the bare bandwidth [13]. In Fig. 4.6b we report, superimposed, the convergence values of the RMP $\tilde{t}^{(s^*)}$ and $V_c[\beta = 100, \delta]$ given by Eq. (4.51). Our prediction of the critical value for V approximates reasonably the localisation region.

To extract the value of $V_c[\beta, \delta]$ is, in general, a non-trivial analytic problem due to the insurgence of elliptic integrals and the smearing effects of temperature. At low temperature, however, we can use Eq. (4.48) and obtain the asymptotic critical boundary at half-filling

$$V_c[\beta \rightarrow \infty, 0] = \frac{2t}{(2/\pi)^2} \approx 4.93t, \quad (4.52)$$

which we report in Fig. 4.6b as a tick of the vertical axis.




Figure 4.7 | Comparative plots of the HFP $u^{(s^*)}$ (top panel) and RMP $\tilde{t}^{(s^*)}$ (bottom panel) in null-doped (left plots), “medium-doped” ($\delta = 0.2$, central plots) and “heavily-doped” ($\delta = 0.4$, right plots) contexts. Increasing temperature progressively destroys Mott localisation, reducing the bands shrinking effect.

Increasing temperature, Mott localisation is suppressed. In Fig. 4.7a we report results for the HFP $u^{(s^*)}$ computed at decreasingly lower temperature and three example doping levels,

$$1.0 \leq \beta \leq 100.0, \quad \delta = 0.0, 0.2, 0.4.$$

Increasing temperature, the constant $u^{(s^*)}$ region progressively disappears in favour of a monotonically decreasing behaviour. In Fig. 4.7a the same dataset is plotted with focus on the RMP $\tilde{t}^{(s^*)}$. Temperature increase disrupts localisation effects, at least at small values of V . However, the bands shrinking effect is still relevant up to $\beta = 1.0$, with the hopping parameter approximately halving at $V \approx 5 \div 6$.

4.5.2 Solution free energy and entropy densities

In order to compare phases, we need an estimation of free energy of each. The plots of Fig. 4.8 report our findings on the normal phase free energy, computed based on Eq. (4.43). The free energy is intended in unities of t . For a non-flat band, doping increases energy because the band fills up and electrons populate higher energy levels.

Looking at the data of Fig. 4.8a at fixed δ , free energy density remains approximately constant up to the Mott boundary. Indeed, as long as a band with non-zero amplitude is present, the HFP correction of Eq. (4.43) compensates the free energy increase due to shrinking of bands,

$$\begin{aligned} -V|u^{(s^*)}|^2 &= -\frac{Vu^{(s^*)*}}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\epsilon_{\mathbf{k}} - \tilde{\epsilon}_{\mathbf{k}}) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}), \end{aligned}$$

having we used in the second passage

$$\tilde{\epsilon}_{\mathbf{k}} = \epsilon_{\mathbf{k}} + u^{(s^*)*} V (\cos k_x + \cos k_y).$$




Figure 4.8 | Free energy density of the normal phase. On the left-side panel, the dashed black line has the same meaning as the dotted line of Fig. 4.6b, indicating the Mott transition.




Figure 4.9 | Entropy density of the normal phase (left panel) and theoretical contribution to entropy for a state with occupation f . On the left-side panel, the dashed white line has the same meaning as the dotted line of Fig. 4.6b, indicating the Mott transition.

Recalling the first point of the proof given in Sec. 4.3.3, to add the HFP correction to free energy is the same as removing the *tilde* sign from single-state energies, thus using the free model ($V = 0$) free energy density expression.

Mott localisation is energetically expensive. This is visible in both panels of Fig. 4.8. The V -dependent energy increase is an entropic effect: indeed, for a flat band,

$$f(0; \beta, \tilde{\mu}(\delta)) = \frac{1}{2}$$

at any temperature. The expression for the contribution to entropy of a single particle state with occupation f [8] is given by

$$s(f) = -k_B [f \log f + (1 - f) \log(1 - f)] ,$$

and is represented in Fig. 4.9b. The maximum is located at $f = 1/2$, then a flat band realises the maximally entropic condition. The Mott insulating region can be seen as a degeneration of the FS interface to the entire BZ, a process that maximises entropy.

All of this holds, as all the rest, at low temperature: by lowering β , finite-temperature effects become more and more relevant and the statements above need to be corrected.

4.5.3 Convergence problems and mixing fine-tuning

In this subsection we describe some of the typical convergence problems arising in HF iterative schemes anticipated in Sec. 2.3.2. Bands flattening is a source of algorithmic instability: numerics




Figure 4.10 | Comparison of three example tracks for the single-valued HFPs vector of the normal phase, simulated at a “Mott-deep” point.

can lead to values of $u^{(s^*)}$, during iterations, slightly out of bounds with respect to Eq. (4.44). Even if the numeric effect is small, the consequence can be vast. To heal this effect and extract the actual convergence values, it is necessary to optimise the mixing parameter $g \in [0, 1]$.

We defined g in Sec. 2.3.2, and operatively in Eq. (2.25). This parameter controls deterministically how, from an initialiser (the HFPs vector we plug in the algorithm) and an output (the HFPs vector we get back from the algorithm) the new initialiser for the following step is generated. We show in Fig. 4.10 the evolutions at different mixing parameters of a simulation at

$$V = 6.0, \quad \delta = 0.1, \quad \beta = 100.0.$$

This point is inside the Mott region, as can be checked in Fig. 4.5a. The HFP $u^{(s^*)}$ needs to converge at $2t/V = t/3$ in order to give complete localisation. The initialiser

$$\text{Step 1} \quad : \quad u^{(s^*)} = 0.1,$$

was arbitrarily chosen. In each panel of Fig. 4.10, the red track represents the value of $u^{(s^*)}$ we plug in the self-consistency equations, the blue the result of the self-consistent equation, the green one the mixed value of the two based on different g s. The gray dashed line is the theoretical convergence target.

In the first two panels, there is an evident infinite loop occurring whenever the green line crosses the threshold $2t/V$. At those steps, the green point is out of bounds with respect to Eq. (4.44). When this happens, the successive red point gives a band with inverted sign, a feature that “breaks” the self-consistency equation and generates a blue point with no physical significance.

To overcome this problem it is necessary to move “slowly”. As is shown in the right panel, the choice of a small enough mixing parameter ensures convergence. In the track, each blue dot is above the target and would produce a band with inverted sign. Mixing blue and red dots with a 95% unbalance on the red ones, we ensure that the green track approaches the target from below.

This kind of problems can turn out to be pretty common for iHF schemes, and to understand how to knob the algorithmic parameters without sacrificing accurateness and precision, as well as avoiding computationally costly and frustrating infinite loops, was an essential part when designing this algorithm.

Chapter 5

Antiferromagnetic phase

In this chapter, we study antiferromagnetism in the extended Hubbard model, limiting ourselves to commensurate antiferromagnetism at half-filling. This phase is well known and studied on the square Hubbard lattice. Here we show that the insertion of the non-local attraction $\hat{H}_V$ leads to an enhancement of the antiferromagnetic (AF) instability, as a result of two cooperative effects. First, the direct amplification of the AF gap, and, second, the shrinking of electronic bandwidth that lead to the aforementioned Mott localisation.

This chapter is organised as follows. In Sec. 5.1, we present the commensurate antiferromagnetic (AF) phase and discuss its symmetries. In Secs. 5.2, 5.3, 5.4, we discuss the HF approximation in the AF phase, starting from the conventional Hubbard model and highlighting the modifications in the presence of the non-local attraction $\hat{H}_V$. In Sec. 5.5, we analyse the self-consistency equations and derive a set of constraints on the HFPs that describe the self-consistent solution. In Sec. 5.6, we present the result of numeric simulations.

Disclaimer. Sec. 5.3, where we carry out all technical computations to derive the HF approximation to the EHM, is *very* detailed and cumbersome. It can be skipped, jumping directly to Sec. 5.4, where we summarise all results.

5.1 The AF phase and its symmetries

Long-range antiferromagnetic ordering is defined by the insurgence of a staggered pattern of spin density with a given periodicity, with zero net magnetisation. In Fig. 5.1, we show three (of many) possible AF orderings, with the blue/red arrows indicating an abundance of spin-up/down




Figure 5.1 | Three possible antiferromagnetic structures. Dots and arrows indicate spin abundance on a given site, while empty sites are intended as spin-balanced. Within this text we deal with the lowest-periodicity commensurate phase, the antiferromagnet on left panel.

electrons on-site. The left panel of Fig. 5.1 shows an alternating structure with two-sites periodicity in both directions. We denote this configuration as “commensurate”. This structure is periodic by shifts of an amount $(1, 1)$ in lattice units. Therefore, in reciprocal space this phase has a non-zero component at wavevector

$$\boldsymbol{\pi} = (\pi, \pi). \quad (5.1)$$

which we denote, from now on, the “AF vector”. The central panel of Fig. 5.1 presents another staggered configuration with doubled periodicity, where sites with a given polarisation are inter-mediated by sites where spin populations are balanced (gray dots). The right panel shows a possible “striped” antiferromagnet, where AF ordering is established along one particular direction. In our example, in along direction y we have AF ordering with periodicity of 2 sites, while along x we have ferromagnetic ordering. Striped antiferromagnetism is studied in HM physics [1] as a possible long-range ordering energetically competing with anisotropic superconductivity. Nematic antiferromagnets, establishing long-range magnetic ordering along one particular direction, are interesting due to competition and interplay against superconducting ordering at finite doping [2]. Geometrically non-trivial magnetic ordering are at the core of SDW formation in copper oxides at finite doping [3, 4].

In this thesis, we focus on the commensurate configuration on the left panel of Fig. 5.1, which is known to be favourite at half-filling [5]. As we anticipated in Sec. 3.2 and summarised in Tab. 3.1, the AF phase is described by the breaking of global spin rotation and translational symmetries:

$$\begin{aligned} \text{SU}(2) &\xrightarrow{\text{SSB}} \text{U}^z(1), \\ \text{T}_1^{(x)} \otimes \text{T}_1^{(y)} &\xrightarrow{\text{SSB}} \text{T}_2^{(x)} \otimes \text{T}_2^{(y)}. \end{aligned}$$

As we did for the normal phase, we also allow for

$$\text{C}_4 \xrightarrow{\text{SSB}} \text{C}_2.$$

We will show that the commensurate AF phase actually leaves C_4 unbroken.

Commensurate antiferromagnetism is specified by the Ansatz:

$$\langle \hat{n}_{\mathbf{r}\sigma} \rangle = n - (-1)^{x+y+\delta_{\sigma=\uparrow}} m \quad \text{and} \quad n, m \in [0, 1]. \quad (5.2)$$

where $\mathbf{r} = (x, y) \in \mathcal{L}$. Eq. (5.2) represents the staggered distribution of spin populations shown in the left panel of Fig. 5.1. At site $\mathbf{r}$ the two spin populations average to n and differ by $2m$. Note that $\langle \hat{n}_{\mathbf{r}\sigma} \rangle \leq 1$ due to Pauli exclusion principle.

5.1.1 Symmetries of the commensurate AF Ansatz

The Ansatz of Eq. (5.2) breaks translational invariance in each spin sector, but preserves $\text{U}^z(1)$ and $\text{U}^c(1)$ symmetries. Eq. (5.2) is formulated equivalently as

$$\langle \hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow} \rangle = (-1)^{x+y} 2m. \quad (5.3)$$

Antiferromagnetism establishes when $m \neq 0$.

Proof 5.1 — The Ansatz of Eq. (5.2) is $\text{U}^z(1)$ -symmetric.

We use the spinors $\hat{\Psi}_{\mathbf{r}}$ defined in Eq. (3.8)

$$\hat{\Psi}_{\mathbf{r}} = \begin{bmatrix} \hat{c}_{\mathbf{r}\uparrow} \\ \hat{c}_{\mathbf{r}\downarrow} \end{bmatrix},$$

and the local spin operators of Eq. (3.7),

$$\hat{S}_{\mathbf{r}}^+ = \frac{\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}}{2}, \quad \hat{S}_{\mathbf{r}}^- = \frac{\hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\uparrow}}{2}, \quad \hat{S}_{\mathbf{r}}^z = \frac{\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} - \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}}{2}.$$

Consider Eq. (3.23) and its conjugate,

$$[\hat{c}_{\mathbf{r}\sigma}^\dagger, \hat{n}_{\mathbf{r}\sigma}] = -\hat{c}_{\mathbf{r}\sigma}^\dagger \quad \text{and} \quad [\hat{c}_{\mathbf{r}\sigma}, \hat{n}_{\mathbf{r}\sigma}] = \hat{c}_{\mathbf{r}\sigma}.$$

The operators $\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}$ and $\hat{S}_{\mathbf{r}}^\pm$ do not commute. Indeed,

$$\begin{aligned} [\hat{S}_{\mathbf{r}}^+, \hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}] &= \frac{1}{2} [\hat{c}_{\mathbf{r}\uparrow}^\dagger, \hat{n}_{\mathbf{r}\uparrow}] \hat{c}_{\mathbf{r}\downarrow} - \frac{1}{2} \hat{c}_{\mathbf{r}\uparrow}^\dagger [\hat{c}_{\mathbf{r}\downarrow}, \hat{n}_{\mathbf{r}\downarrow}] \\ &= -\hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}, \\ [\hat{S}_{\mathbf{r}}^-, \hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}] &= \frac{1}{2} \hat{c}_{\mathbf{r}\downarrow}^\dagger [\hat{c}_{\mathbf{r}\uparrow}, \hat{n}_{\mathbf{r}\uparrow}] - \frac{1}{2} [\hat{c}_{\mathbf{r}\downarrow}^\dagger, \hat{n}_{\mathbf{r}\downarrow}] \hat{c}_{\mathbf{r}\uparrow} \\ &= \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\uparrow}. \end{aligned}$$

Instead, $\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}$ and $\hat{S}_{\mathbf{r}}^z$ commute, because all on-site number operators commute among themselves. Then, in order to have $m \neq 0$ in Eq. (5.3), SU(2) must be broken while $U^z(1)$ does not. Therefore the commensurate AF phase breaks global spin rotational invariance, reducing it to the smaller symmetry of rotations around the magnetisation axis.

Proof 5.2 — The Ansatz of Eq. (5.2) is $U^c(1)$ -symmetric.

From Eq. (3.5) we know that on-site density operators are gauge-invariant separately in both spin sectors,

$$\hat{n}_{\mathbf{r}\sigma} \rightarrow e^{i(\eta-\eta)} \hat{n}_{\mathbf{r}\sigma} = \hat{n}_{\mathbf{r}\sigma}.$$

Then no selection rule applies to $\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}$: its EV can be non-zero without breaking $U^z(1)$.

Eq. (5.2) contains two physical observables: the average density n and the staggered magnetisation m . These two quantities obey the self-consistency equations

$$n = \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle + \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2}, \quad (5.4)$$

$$m = \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle - \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2}, \quad (5.5)$$

where the expectation values on the right-hand side depend on n and m . Since we work at fixed density, Eq. (5.4) is used to determine self-consistently the chemical potential.

Proof 5.3 — Proof of Eqs. (5.4) and (5.5).

The on-site averaged density is given by:

$$\forall \mathbf{r} \in \mathcal{L} \quad : \quad n \equiv \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle + \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2}, \quad (5.6)$$

while on-site magnetisation is, inverting Eq. (5.3),

$$\forall \mathbf{r} = (x, y) \in \mathcal{L} \quad : \quad m \equiv (-1)^{x+y} \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle - \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2}. \quad (5.7)$$

By construction, both n and m are translationally invariant quantities. Hence

$$\begin{aligned} n &= \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle + \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2}, \\ m &= \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle - \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2}, \end{aligned}$$

which ends the proof.

5.1.2 The AF Ansatz in reciprocal space

Translational symmetry is not completely broken in the AF phase; rather, is reduced. Therefore, it is useful to move to reciprocal space and transform Eq. (5.2). We define

$$\hat{n}_{\mathbf{k}\sigma} \equiv \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}, \quad (5.8)$$

$$\hat{\psi}_{\mathbf{k}\sigma} \equiv \hat{c}_{\mathbf{k}+\boldsymbol{\pi}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}, \quad (5.9)$$

and their expectation values (EVs),

$$u_{\mathbf{k}\sigma} \equiv \langle \hat{n}_{\mathbf{k}\sigma} \rangle, \quad (5.10)$$

$$v_{\mathbf{k}\sigma} \equiv i \langle \hat{\psi}_{\mathbf{k}\sigma} \rangle. \quad (5.11)$$

In the definition of $v_{\mathbf{k}\sigma}$ we included a i prefactor that is useful in later calculations. Using the functions of Tab. 3.5, we define the harmonics

$$u_\sigma^{(\gamma)} \equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} u_{\mathbf{k}\sigma}, \quad (5.12)$$

$$v_\sigma^{(\gamma)} \equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} v_{\mathbf{k}\sigma}. \quad (5.13)$$

As will be discussed, these harmonics are HFPs in the AF phase and must be determined self-consistently. Eqs. (5.12) and (5.13) are their self-consistency equations.

A finite staggered magnetisation ($m \neq 0$) implies, at least for a subset of momenta $\{\mathbf{k}\}$,

$$m \neq 0 \implies \exists \mathbf{k} \in \text{BZ} : |\langle \hat{\psi}_{\mathbf{k}\sigma} \rangle| > 0. \quad (5.14)$$

Recalling Eq. (5.9), we see that removing one electron at $(\mathbf{k}, \sigma)$ from the ground-state and creating another at $(\mathbf{k} + \boldsymbol{\pi}, \sigma)$ produces a state with a finite overlap.

Proof 5.4 — If $m \neq 0$, for some $\mathbf{k}$ it must hold $\langle \hat{\psi}_{\mathbf{k}\sigma} \rangle \neq 0$.

We assume $m \neq 0$. From Eq. (5.2) we get

$$\begin{aligned} (-1)^{\delta_{\sigma=\downarrow} m} &= \overbrace{\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} n}^{=0} + \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{\delta_{\sigma=\downarrow} m} \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} (n + (-1)^{x+y+\delta_{\sigma=\downarrow} m}) \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} \langle \hat{n}_{\mathbf{r}\sigma} \rangle, \end{aligned} \quad (5.15)$$

where we used $1 = ((-1)^{x+y})^2$. $\mathcal{F}$ -transforming the operator and its sign factor inside the sum in last line, we get

$$\begin{aligned} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} \hat{n}_{\mathbf{r}\sigma} &= \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma} \\ &\stackrel{\mathcal{F}}{=} \sum_{\mathbf{r} \in \mathcal{L}} e^{i\boldsymbol{\pi} \cdot \mathbf{r}} \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma}^\dagger \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k}' \in \text{BZ}} e^{-i\mathbf{k}' \cdot \mathbf{r}} \hat{c}_{\mathbf{k}'\sigma} \\ &= \sum_{\mathbf{k} \in \text{BZ}} \sum_{\mathbf{k}' \in \text{BZ}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma} \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} e^{-i[\mathbf{k}' - (\mathbf{k} + \boldsymbol{\pi})] \cdot \mathbf{r}}. \end{aligned}$$

Using Eq. (3.35), last line is reduced to

$$\sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} \hat{n}_{\mathbf{r}\sigma} = \sum_{\mathbf{k} \in \text{BZ}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}+\boldsymbol{\pi}\sigma} \quad (5.16)$$

$$= \sum_{\mathbf{k} \in \text{BZ}} \hat{\psi}_{\mathbf{k}\sigma}. \quad (5.17)$$

We insert the last line in Eq. (5.15),

$$(-1)^{\delta_{\sigma=\downarrow}} m = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\psi}_{\mathbf{k}\sigma} \rangle. \quad (5.18)$$

Since $m \neq 0$, at least for some momentum $\mathbf{k}$ it must hold $\langle \hat{\psi}_{\mathbf{k}\sigma} \rangle \neq 0$.

The self-consistency equation for m reads, in reciprocal space,

$$m = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\psi}_{\mathbf{k}\uparrow} - \hat{\psi}_{\mathbf{k}\downarrow} \rangle. \quad (5.19)$$

This is the transformed version of Eq. (5.5).

Proof 5.5 — Proof of Eq. (5.19): $\mathcal{F}$ -transform of Eq. (5.5).

From Eq. (5.18), we have

$$m = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\psi}_{\mathbf{k}\uparrow} \rangle \quad (5.20)$$

$$= -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\psi}_{\mathbf{k}\downarrow} \rangle. \quad (5.21)$$

Summing the two lines, we obtain Eq. (5.19). From Eq. (5.13), we notice that $v_\sigma^{(s)} = i(-1)^{\delta_{\sigma=\downarrow}} m$.

The operator $\hat{\psi}_{\mathbf{k}\sigma}$ satisfies the relation

$$\varphi_{\mathbf{k}+\boldsymbol{\pi}}^{(\gamma)} \hat{\psi}_{\mathbf{k}+\boldsymbol{\pi}\sigma}^\dagger = -\varphi_{\mathbf{k}}^{(\gamma)} \hat{\psi}_{\mathbf{k}\sigma}, \quad (5.22)$$

where we used the harmonics $\varphi_{\mathbf{k}}^{(\gamma)}$ defined in Tab. 3.5.

Proof 5.6 — Proof of Eq. (5.22).

The following identity holds

$$\begin{aligned} \hat{\psi}_{\mathbf{k}+\boldsymbol{\pi}\sigma}^\dagger &= \left(\hat{c}_{\mathbf{k}+2\boldsymbol{\pi}\sigma}^\dagger \hat{c}_{\mathbf{k}+\boldsymbol{\pi}\sigma} \right)^\dagger \\ &= \hat{c}_{\mathbf{k}+\boldsymbol{\pi}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \\ &= \hat{\psi}_{\mathbf{k}\sigma}. \end{aligned} \quad (5.23)$$

Moreover, being combinations of sines and cosines, the functions $\varphi_{\mathbf{k}+\boldsymbol{\pi}}^{(\gamma)}$ are antisymmetric under $\boldsymbol{\pi}$ shifts,

$$\varphi_{\mathbf{k}+\boldsymbol{\pi}}^{(\gamma)} = -\varphi_{\mathbf{k}}^{(\gamma)}. \quad (5.24)$$

Combining Eq. (5.23) and (5.24), we obtain Eq. (5.22).

5.2 The AF phase in the conventional HM

In this section, we set $V = 0$ and describe antiferromagnetism in the conventional HM, $\hat{H}_t + \hat{H}_U$. We start by deriving the analytic expression of the HF hamiltonian for the AF phase, and then diagonalise it.

5.2.1 Derivation of the HF hamiltonian

The two-body repulsive interaction is

$$\hat{H}_U = U \sum_{\mathbf{r} \in \mathcal{L}} \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow}.$$

We know from Tab. 3.2 that the only contribution to the HF hamiltonian comes from Hartree channel, while Fock and Cooper channel are identically zero in the AF phase.

Local Hartree terms. In Eq. (2.21) we decomposed $\hat{H}_U$ through Wick's theorem. The decoupling in the Hartree channel reads

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} U \sum_{\mathbf{r} \in \mathcal{L}} [\hat{n}_{\mathbf{r}\uparrow} \langle \hat{n}_{\mathbf{r}\downarrow} \rangle + \langle \hat{n}_{\mathbf{r}\uparrow} \rangle \hat{n}_{\mathbf{r}\downarrow} - \langle \hat{n}_{\mathbf{r}\uparrow} \rangle \langle \hat{n}_{\mathbf{r}\downarrow} \rangle]. \quad (5.25)$$

We define the MBZ as in Eq. (4.50), as the set of points in reciprocal space that satisfy the inequality $|k_x| + |k_y| \leq \pi$. We sketched it in Fig. 4.4. Applying the Ansatz of Eq. (5.2), we obtain the HF approximation

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} - \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \left(\Delta_{\mathbf{k}\sigma} \hat{\psi}_{\mathbf{k}\sigma} + \Delta_{\mathbf{k}\sigma}^* \hat{\psi}_{\mathbf{k}\sigma}^\dagger \right) + nU \hat{N} - E_{\text{H},U}^{(\text{AF})}, \quad (5.26)$$

where we defined the gap function

$$\Delta_{\mathbf{k}\sigma} = (-1)^{\delta_{\sigma=\downarrow}} mU, \quad (5.27)$$

and the energy shift

$$E_{\text{H},U}^{(\text{AF})} \equiv U \sum_{\mathbf{r} \in \mathcal{L}} \langle \hat{n}_{\mathbf{r}\uparrow} \rangle \langle \hat{n}_{\mathbf{r}\downarrow} \rangle. \quad (5.28)$$

We observe, in Eq. (5.26), the presence of a number operator, meaning that chemical potential is shifted in the Hartree channel by

$$\mu \rightarrow \mu - nU. \quad (5.29)$$

We report these results in the summary of Tab. 5.1, provided in next section.

Proof 5.7 — Proof of Eq. (5.26): HF approximation of $\hat{H}_U$ in Hartree channel.

Consider Eq. (5.25). Using Eqs. (5.4), (5.5) we can re-write

$$\begin{aligned} \hat{n}_{\mathbf{r}\uparrow} \langle \hat{n}_{\mathbf{r}\downarrow} \rangle + \langle \hat{n}_{\mathbf{r}\uparrow} \rangle \hat{n}_{\mathbf{r}\downarrow} &= \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle + \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2} (\hat{n}_{\mathbf{r}\uparrow} + \hat{n}_{\mathbf{r}\downarrow}) - \frac{\langle \hat{n}_{\mathbf{r}\uparrow} \rangle - \langle \hat{n}_{\mathbf{r}\downarrow} \rangle}{2} (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}) \\ &= n (\hat{n}_{\mathbf{r}\uparrow} + \hat{n}_{\mathbf{r}\downarrow}) - m (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}). \end{aligned}$$

Inserting this decomposition and the expression for $E_{\text{H},U}^{(\text{AF})}$ given in Eq. (5.28) into Eq. (5.25), we obtain

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} nU \sum_{\mathbf{r} \in \mathcal{L}} (\hat{n}_{\mathbf{r}\uparrow} + \hat{n}_{\mathbf{r}\downarrow}) - mU \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}) - E_{\text{H},U}^{(\text{AF})}. \quad (5.30)$$

The first term gives a number operator,

$$nU \sum_{\mathbf{r} \in \mathcal{L}} (\hat{n}_{\mathbf{r}\uparrow} + \hat{n}_{\mathbf{r}\downarrow}) = nU \times \hat{N}.$$

Using Eq. (5.16), the second term of Eq. (5.28) becomes

$$mU \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}) = mU \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} (-1)^{\delta_{\sigma=\downarrow}} \hat{\psi}_{\mathbf{k}\sigma}.$$

Recall the definition of the MBZ given in Eq. (4.50). The full BZ can be seen as the sum of two MBZs shifted by π . Then, from Eq. (5.23), we get

$$mU \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}) = mU \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} (-1)^{\delta_{\sigma=\downarrow}} \left(\hat{\psi}_{\mathbf{k}\sigma} + \hat{\psi}_{\mathbf{k}\sigma}^{\dagger} \right).$$

Composing everything, Eq. (5.30) becomes

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} -mU \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} (-1)^{\delta_{\sigma=\downarrow}} \left(\hat{\psi}_{\mathbf{k}\sigma} + \hat{\psi}_{\mathbf{k}\sigma}^{\dagger} \right) + nU\hat{N} - E_{\text{H},U}^{(\text{AF})}.$$

By defining the gap function $\Delta_{\mathbf{k}\sigma}$ as in Eq. (5.27), we obtain the HF approximation of Eq. (5.26).

The contraction energy, defined in Eq. (5.28), is given by

$$E_{\text{H},U}^{(\text{AF})} = L_x L_y \times U (n^2 - m^2). \quad (5.31)$$

Compared with that for the normal phase given in Eq. (4.14), we observe the presence of an additional term dependent on m^2 .

Proof 5.8 — Proof of Eq. (5.31): local Hartree contraction energy.

Inserting Eq. (5.2) in Eq. (5.28), we obtain

$$\begin{aligned} E_{\text{H},U}^{(\text{AF})} &\equiv U \sum_{\mathbf{r} \in \mathcal{L}} \langle \hat{n}_{\mathbf{r}\uparrow} \rangle \langle \hat{n}_{\mathbf{r}\downarrow} \rangle \\ &= U \sum_{\mathbf{r} \in \mathcal{L}} (n + (-1)^{x+y} m) (n - (-1)^{x+y} m). \end{aligned}$$

Expanding the product above we get:

$$\begin{aligned} (n + (-1)^{x+y} m) (n - (-1)^{x+y} m) &= n^2 + ((-1)^{x+y} - (-1)^{x+y}) nm - ((-1)^{x+y})^2 m^2 \\ &= n^2 - m^2. \end{aligned}$$

The sum over $\mathcal{L}$ gives $L_x L_y$ identical terms. Hence we obtain Eq. (5.31).

5.2.2 Diagonalisation of the HF hamiltonian

In this section, we diagonalise the HF hamiltonian identified by Eqs. (5.26) and (5.31).

Matrix representation. For the AF phase, the HF hamiltonian can be written as

$$\hat{H}_t + \hat{H}_U \stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} h_{\mathbf{k}\sigma} \hat{\Psi}_{\mathbf{k}\sigma} + nU\hat{N} - E_{\text{H},U}^{(\text{AF})}, \quad (5.32)$$

where we defined the 2×2 matrix:

$$h_{\mathbf{k}\sigma} \equiv \begin{bmatrix} \epsilon_{\mathbf{k}\sigma} & -\Delta_{\mathbf{k}\sigma}^* \\ -\Delta_{\mathbf{k}\sigma} & -\epsilon_{\mathbf{k}\sigma} \end{bmatrix}, \quad (5.33)$$

and the spinors

$$\hat{\Psi}_{\mathbf{k}\sigma} \equiv \begin{bmatrix} \hat{c}_{\mathbf{k}\sigma} \\ \hat{c}_{\mathbf{k}+\pi\sigma} \end{bmatrix}. \quad (5.34)$$

Proof 5.9 — Proof of Eq. (5.32).

Recall Eq. (3.37): the kinetic sector is represented in reciprocal space by

$$\begin{aligned}\hat{H}_t &= \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \epsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} \\ &= \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \epsilon_{\mathbf{k}} \left(\hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} - \hat{c}_{\mathbf{k}+\boldsymbol{\pi}\sigma}^{\dagger} \hat{c}_{\mathbf{k}+\boldsymbol{\pi}\sigma} \right),\end{aligned}$$

where we used antiperiodicity by $\boldsymbol{\pi}$ of the bare bands $\epsilon_{\mathbf{k}}$ defined in Eq. (3.38),

$$\epsilon_{\mathbf{k}} = -\epsilon_{\mathbf{k}+\boldsymbol{\pi}}.$$

The kinetic operator can be written in terms of the spinors $\hat{\Psi}_{\mathbf{k}\sigma}$ defined in Eq. (5.34),

$$\hat{H}_t = \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \begin{bmatrix} \epsilon_{\mathbf{k}\sigma} & \\ & -\epsilon_{\mathbf{k}\sigma} \end{bmatrix} \hat{\Psi}_{\mathbf{k}\sigma}.$$

The local repulsion $\hat{H}_U$ of Eq. (5.26) can also be written in terms of these spinors:

$$\hat{H}_U = \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \begin{bmatrix} & -\Delta_{\mathbf{k}\sigma}^* \\ -\Delta_{\mathbf{k}\sigma} & \end{bmatrix} \hat{\Psi}_{\mathbf{k}\sigma} + nU\hat{N} - E_{\text{H},U}^{(\text{AF})}.$$

Summing $\hat{H}_t + \hat{H}_U$ and defining $h_{\mathbf{k}\sigma}$ as in Eq. (5.33), we obtain the matricial representation of Eq. (5.32).

Pseudospin representation. The matrix $h_{\mathbf{k}\sigma}$ can be written in term of the Pauli matrices τ^{α} ,

$$h_{\mathbf{k}\sigma} = \epsilon_{\mathbf{k}\sigma} \tau^z - \Delta_{\mathbf{k}\sigma} \tau^x.$$

Let us define the pseudospin operators $\hat{\mathbf{s}}_{\mathbf{k}\sigma}$ as

$$\hat{s}_{\mathbf{k}\sigma}^{\alpha} \equiv \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \tau^{\alpha} \hat{\Psi}_{\mathbf{k}\sigma} \quad \text{with} \quad \hat{\mathbf{s}}_{\mathbf{k}\sigma} = \begin{bmatrix} \hat{s}_{\mathbf{k}\sigma}^x \\ \hat{s}_{\mathbf{k}\sigma}^y \\ \hat{s}_{\mathbf{k}\sigma}^z \end{bmatrix}. \quad (5.35)$$

These operators satisfy a $\mathfrak{su}(2)$ algebra up to a constant C ,

$$[\hat{s}_{\mathbf{k}\sigma}^{\alpha}, \hat{s}_{\mathbf{k}\sigma}^{\beta}] = C \times i \sum_{\gamma} \epsilon_{\alpha\beta\gamma} \hat{s}_{\mathbf{k}\sigma}^{\gamma}. \quad (5.36)$$

The proof of Eq. (5.36) is identical to the one presented in Sec. 3.1.2. Let us express the pseudospin and identity operators in terms of the operators of Eqs. (5.8), (5.9):

$$\hat{s}_{\mathbf{k}\sigma}^x = \hat{\psi}_{\mathbf{k}\sigma} + \hat{\psi}_{\mathbf{k}\sigma}^{\dagger}, \quad (5.37)$$

$$\hat{s}_{\mathbf{k}\sigma}^y = i \left(\hat{\psi}_{\mathbf{k}\sigma} - \hat{\psi}_{\mathbf{k}\sigma}^{\dagger} \right), \quad (5.38)$$

$$\hat{s}_{\mathbf{k}\sigma}^z = \hat{n}_{\mathbf{k}\sigma} - \hat{n}_{\mathbf{k}+\boldsymbol{\pi}\sigma}, \quad (5.39)$$

$$\begin{aligned}\hat{\mathbb{I}}_{\mathbf{k}\sigma} &\equiv \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \mathbb{I}_{2 \times 2} \hat{\Psi}_{\mathbf{k}\sigma} \\ &= \hat{n}_{\mathbf{k}\sigma} + \hat{n}_{\mathbf{k}+\boldsymbol{\pi}\sigma}.\end{aligned} \quad (5.40)$$

We define the pseudofield $\mathbf{b}_{\mathbf{k}\sigma}$:

$$\mathbf{b}_{\mathbf{k}\sigma} \equiv \begin{bmatrix} -\Delta_{\mathbf{k}\sigma} \\ 0 \\ \epsilon_{\mathbf{k}\sigma} \end{bmatrix}.$$

In pseudospin representation, the hamiltonian of Eq. (5.32) can be expressed as

$$\hat{H}_t + \hat{H}_U \stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \mathbf{b}_{\mathbf{k}\sigma} \cdot \hat{\mathbf{s}}_{\mathbf{k}\sigma} + nU\hat{N} - E_{\text{H},U}^{(\text{AF})}, \quad (5.41)$$




Figure 5.2 | Sketch of the diagonalisation procedure for the HM antiferromagnet. To diagonalise the HF hamiltonian, we align the z axis with the pseudo-field.

describing a set of (pseudo)spins subject to local (pseudo)magnetic fields. The field $\mathbf{b}_{\mathbf{k}\sigma}$ is orthogonal to the y plane, and is tilted in the z plane by an angle $2\theta_{\mathbf{k}\sigma}$, where

$$\cos 2\theta_{\mathbf{k}\sigma} = \frac{\epsilon_{\mathbf{k}\sigma}}{E_{\mathbf{k}\sigma}}, \quad \sin 2\theta_{\mathbf{k}\sigma} = \frac{\Delta_{\mathbf{k}\sigma}}{E_{\mathbf{k}\sigma}}, \quad E_{\mathbf{k}\sigma} \equiv \sqrt{\epsilon_{\mathbf{k}\sigma}^2 + |\Delta_{\mathbf{k}\sigma}|^2}.$$

A sketch of these angles and the pseudofield is given in Fig. 5.2.

We observe that we could have redefined the pseudospin operators of Eq. (5.35) dividing them by an appropriate constant, in order to satisfy exactly the $\mathfrak{su}(2)$ algebra, and at the same time multiply the pseudo-magnetic field by the same constant. The product $\mathbf{b}_{\mathbf{k}\sigma} \cdot \hat{\mathbf{s}}_{\mathbf{k}\sigma}$ would be unchanged, and the diagonalisation matrix $W_{\mathbf{k}\sigma}$ would be the same. Because of this, the constant C in Eq. (5.36) is irrelevant.

Diagonalisation. In order to diagonalise the HF hamiltonian, Eq. (5.41), we perform a unitary operation on the $\hat{\Psi}_{\mathbf{k}\sigma}$ spinors which reduces the 2×2 matrix $h_{\mathbf{k}\sigma}$ of Eq. (5.33) to a diagonal form. This means to diagonalise each $h_{\mathbf{k}\sigma}$. This is obtained through a rotation around the y axis:

$$d_{\mathbf{k}\sigma} = W_{\mathbf{k}\sigma} h_{\mathbf{k}\sigma} W_{\mathbf{k}\sigma}^\dagger \quad \text{with} \quad d_{\mathbf{k}\sigma} \equiv \begin{bmatrix} E_{\mathbf{k}\sigma} & \\ & -E_{\mathbf{k}\sigma} \end{bmatrix}. \quad (5.42)$$

The diagonal values of $d_{\mathbf{k}\sigma}$ are $\pm E_{\mathbf{k}\sigma}$, with

$$E_{\mathbf{k}\sigma} = \sqrt{\epsilon_{\mathbf{k}\sigma}^2 + |\Delta_{\mathbf{k}\sigma}|^2}. \quad (5.43)$$

In Eq. (5.42), the matrix $W_{\mathbf{k}\sigma}$ is unitary, and gives a counter-clockwise rotation by an angle $2\theta_{\mathbf{k}\sigma}$ around the y axis:

$$\begin{aligned} W_{\mathbf{k}\sigma} &\equiv e^{-i\theta_{\mathbf{k}\sigma}\tau^y} \\ &= \cos \theta_{\mathbf{k}\sigma} - i \sin \theta_{\mathbf{k}\sigma} \tau^y \\ &= \begin{bmatrix} \cos \theta_{\mathbf{k}\sigma} & -\sin \theta_{\mathbf{k}\sigma} \\ \sin \theta_{\mathbf{k}\sigma} & \cos \theta_{\mathbf{k}\sigma} \end{bmatrix}. \end{aligned} \quad (5.44)$$

In Fig. 5.2 we sketch the pseudo-magnetic field in the xz plane, and the diagonalisation procedure. $W_{\mathbf{k}\sigma}$ is a rotation by an angle $2\theta_{\mathbf{k}\sigma}$ around the y axis, and in the end aligns the original z axis with the pseudofield axis. We notice that $E_{\mathbf{k}\sigma}$ is actually spin independent: only the gap depends on σ through a sign factor, which is irrelevant in Eq. (5.43).

Applying the rotation matrix to the spinors we obtain the “tilted” spinors,

$$\hat{\Gamma}_{\mathbf{k}\sigma} \equiv W_{\mathbf{k}\sigma} \hat{\Psi}_{\mathbf{k}\sigma} \quad \text{with} \quad \hat{\Gamma}_{\mathbf{k}\sigma} \equiv \begin{bmatrix} \hat{\gamma}_{\mathbf{k}\sigma}^{(+)} \\ \hat{\gamma}_{\mathbf{k}\sigma}^{(-)} \end{bmatrix}, \quad (5.45)$$

where we defined a new set of fermionic operators $\hat{\gamma}_{\mathbf{k}\sigma}^{(\pm)}$ (see App. B for details on the unitary mapping). Bringing all together, we obtain the non-interacting hamiltonian:

$$\begin{aligned}\hat{H}_t + \hat{H}_U &\stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} h_{\mathbf{k}\sigma} \hat{\Psi}_{\mathbf{k}\sigma} + (\dots) \\ &= \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} W_{\mathbf{k}\sigma}^{\dagger} W_{\mathbf{k}\sigma} h_{\mathbf{k}\sigma} W_{\mathbf{k}\sigma}^{\dagger} W_{\mathbf{k}\sigma} \hat{\Psi}_{\mathbf{k}\sigma} + (\dots) \\ &= \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \hat{\Gamma}_{\mathbf{k}\sigma}^{\dagger} d_{\mathbf{k}\sigma} \hat{\Gamma}_{\mathbf{k}\sigma} + (\dots) \\ &= \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} E_{\mathbf{k}\sigma} \left(\hat{\gamma}_{\mathbf{k}\sigma}^{(+)\dagger} \hat{\gamma}_{\mathbf{k}\sigma}^{(+)} - \hat{\gamma}_{\mathbf{k}\sigma}^{(-)\dagger} \hat{\gamma}_{\mathbf{k}\sigma}^{(-)} \right) + (\dots),\end{aligned}$$

where we intend $(\dots) = nU\hat{N} - E_{\text{H},U}^{(\text{AF})}$. The final non-interacting hamiltonian describes a free Fermi gas of fermions described by the operators $\hat{\gamma}_{\mathbf{k}\sigma}^{(\pm)}$, with energies $\pm E_{\mathbf{k}\sigma}$.

Pseudospin EVs. The expectation value of the original pseudospin operators in the various directions can be expressed in terms of the tilted pseudospins $\hat{\Gamma}_{\mathbf{k}\sigma}^{\dagger} \tau^{\alpha} \hat{\Gamma}_{\mathbf{k}\sigma}$,

$$\begin{aligned}\langle \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \tau^x \hat{\Psi}_{\mathbf{k}\sigma} \rangle &= -\sin 2\theta_{\mathbf{k}\sigma} \langle \hat{\Gamma}_{\mathbf{k}\sigma}^{\dagger} \tau^z \hat{\Gamma}_{\mathbf{k}\sigma} \rangle, \\ \langle \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \tau^y \hat{\Psi}_{\mathbf{k}\sigma} \rangle &= 0, \\ \langle \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \tau^z \hat{\Psi}_{\mathbf{k}\sigma} \rangle &= \cos 2\theta_{\mathbf{k}\sigma} \langle \hat{\Gamma}_{\mathbf{k}\sigma}^{\dagger} \tau^z \hat{\Gamma}_{\mathbf{k}\sigma} \rangle,\end{aligned}$$

where we used the angular relations of Fig. 5.2. Moving to the tilted frame, the z component of the pseudo-spin reduces to the thermal occupation of “up” and “down” states

$$\langle \hat{\Gamma}_{\mathbf{k}\sigma}^{\dagger} \tau^z \hat{\Gamma}_{\mathbf{k}\sigma} \rangle = f(E_{\mathbf{k}\sigma}; \beta, \tilde{\mu}) - f(-E_{\mathbf{k}\sigma}; \beta, \tilde{\mu}). \quad (5.46)$$

Let us define the thermal factors:

$$\text{Th}_{\mathbf{k}\sigma}^{(\pm)}[\beta, \mu] \equiv f(-E_{\mathbf{k}\sigma}; \beta, \mu) \pm f(E_{\mathbf{k}\sigma}; \beta, \mu), \quad (5.47)$$

which will often appear in computations. Being $E_{\mathbf{k}\sigma}$ spin independent, so are the thermal factors. We have:

$$\langle \hat{s}_{\mathbf{k}\sigma}^x \rangle = \frac{\Delta_{\mathbf{k}\sigma}}{E_{\mathbf{k}\sigma}} \text{Th}_{\mathbf{k}\sigma}^{(-)}[\beta, \mu], \quad (5.48)$$

$$\langle \hat{s}_{\mathbf{k}\sigma}^y \rangle = 0, \quad (5.49)$$

$$\langle \hat{s}_{\mathbf{k}\sigma}^z \rangle = -\frac{\epsilon_{\mathbf{k}\sigma}}{E_{\mathbf{k}\sigma}} \text{Th}_{\mathbf{k}\sigma}^{(-)}[\beta, \mu]. \quad (5.50)$$

Magnetisation self-consistency equation. We gave the self-consistency equation for the magnetisation m in Eq. (5.19). By further elaboration, it can be written equivalently as

$$m = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\Delta_{\mathbf{k}\uparrow}}{E_{\mathbf{k}\uparrow}} \text{Th}_{\mathbf{k}\uparrow}^{(-)}[\beta, \mu]. \quad (5.51)$$

This equation is the “operational” version of the self-consistency equation for the HFP m .

Proof 5.10 — Proof of Eq. (5.51): self-consistency equation for m .

Consider Eq. (5.19). Restricting the sum to the MBZ and using Eq. (5.23), we get

$$m = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \left\langle \left(\hat{\psi}_{\mathbf{k}\uparrow} + \hat{\psi}_{\mathbf{k}\uparrow}^{\dagger} \right) - \left(\hat{\psi}_{\mathbf{k}\downarrow} + \hat{\psi}_{\mathbf{k}\downarrow}^{\dagger} \right) \right\rangle.$$

From Eq. (5.37), we recognise a pseudospin EV in direction x . From Eq. (5.48)

$$\begin{aligned} m &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\langle \hat{s}_{\mathbf{k}\uparrow}^x \rangle - \langle \hat{s}_{\mathbf{k}\downarrow}^x \rangle) \\ &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \left(\frac{\text{Re}\{\Delta_{\mathbf{k}\uparrow}\}}{E_{\mathbf{k}\uparrow}} \text{Th}_{\mathbf{k}\uparrow}^{(-)}[\beta, \mu] - \frac{\text{Re}\{\Delta_{\mathbf{k}\downarrow}\}}{E_{\mathbf{k}\downarrow}} \text{Th}_{\mathbf{k}\downarrow}^{(-)}[\beta, \mu] \right) \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\Delta_{\mathbf{k}\uparrow}}{E_{\mathbf{k}\uparrow}} \text{Th}_{\mathbf{k}\uparrow}^{(-)}[\beta, \mu], \end{aligned}$$

which proves Eq. (5.51). In last line we used $\Delta_{\mathbf{k}\uparrow} = -\Delta_{\mathbf{k}\downarrow}$, which follows from Eq. (5.27), and the fact that $E_{\mathbf{k}\sigma}$ and $\text{Th}_{\mathbf{k}\sigma}^{(-)}[\beta, \mu]$ are spin actually spin-independent.

Density. The averaged density obeys the equations

$$n = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \text{Th}_{\mathbf{k}\uparrow}^{(+)}[\beta, \mu] \quad (5.52)$$

$$= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \text{Th}_{\mathbf{k}\uparrow}^{(+)}[\beta, \mu]. \quad (5.53)$$

The two are equivalent, but the second one is handier to implement in numeric computations.

Proof 5.11 — Proof of Eq. (5.52): self-consistency equation for n .

The $\mathcal{F}$ -transform of the on-site density operator satisfies

$$\begin{aligned} \sum_{\mathbf{r} \in \mathcal{L}} \hat{n}_{\mathbf{r}\sigma} &= \sum_{\mathbf{r} \in \mathcal{L}} \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma} \\ &\stackrel{\mathcal{F}}{=} \sum_{\mathbf{r} \in \mathcal{L}} \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma}^\dagger \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k}' \in \text{BZ}} e^{-i\mathbf{k}' \cdot \mathbf{r}} \hat{c}_{\mathbf{k}'\sigma}, \end{aligned} \quad (5.54)$$

where we used Eq. (3.34). By further elaboration, we obtain

$$\begin{aligned} \sum_{\mathbf{r} \in \mathcal{L}} \hat{n}_{\mathbf{r}\sigma} &= \sum_{\mathbf{k} \in \text{BZ}} \sum_{\mathbf{k}' \in \text{BZ}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma} \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} e^{-i(\mathbf{k}' - \mathbf{k}) \cdot \mathbf{r}} \\ &= \sum_{\mathbf{k} \in \text{BZ}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \\ &= \sum_{\mathbf{k} \in \text{BZ}} \hat{n}_{\mathbf{k}\sigma}, \end{aligned} \quad (5.55)$$

where we used Eq. (3.35). Hence, applying Eq. (5.2),

$$\begin{aligned} n &= \frac{1}{2} \sum_{\sigma} \left(\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} n + \overbrace{\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} (-1)^{x+y} m}^{=0} \right) \\ &= \frac{1}{2L_x L_y} \sum_{\sigma} \sum_{\mathbf{r} \in \mathcal{L}} \langle \hat{n}_{\mathbf{r}\sigma} \rangle \\ &= \frac{1}{2L_x L_y} \sum_{\sigma} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{n}_{\mathbf{k}\sigma} \rangle. \end{aligned} \quad (5.56)$$




Figure 5.3 Left panel: electronic density in the AF phase as a function of the chemical potential, computed through Eq. (5.53), at various temperatures. Right panel: example doped bands, with $\mu/\Delta = 1.5$ set arbitrarily and temperature set to $\beta = 10$ to enhance smearing effects around the FS. Single-state filling is proportional to the colour intensity.

Restricting the sum of Eq. (5.56) to the MBZ, we get

$$\begin{aligned}
 n &= \frac{1}{2L_x L_y} \sum_{\sigma} \sum_{\mathbf{k} \in \text{MBZ}} \langle \hat{n}_{\mathbf{k}\sigma} + \hat{n}_{\mathbf{k}+\pi\sigma} \rangle \\
 &= \frac{1}{2L_x L_y} \sum_{\sigma} \sum_{\mathbf{k} \in \text{MBZ}} \langle \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \hat{\Psi}_{\mathbf{k}\sigma} \rangle.
 \end{aligned} \tag{5.57}$$

The mapping $\hat{\Psi} \rightarrow \hat{\Gamma}$ is unitary, so

$$\hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \hat{\Psi}_{\mathbf{k}\sigma} = \hat{\Gamma}_{\mathbf{k}\sigma}^{\dagger} \hat{\Gamma}_{\mathbf{k}\sigma}.$$

Therefore, Eq. (5.57) reduces to

$$\begin{aligned}
 n &= \frac{1}{2L_x L_y} \sum_{\sigma} \sum_{\mathbf{k} \in \text{MBZ}} \langle \hat{\Gamma}_{\mathbf{k}\sigma}^{\dagger} \hat{\Gamma}_{\mathbf{k}\sigma} \rangle \\
 &= \frac{1}{2L_x L_y} \sum_{\sigma} \sum_{\mathbf{k} \in \text{MBZ}} \left(\langle \hat{\gamma}_{\mathbf{k}\sigma}^{(+)\dagger} \hat{\gamma}_{\mathbf{k}\sigma}^{(+)} \rangle + \langle \hat{\gamma}_{\mathbf{k}\sigma}^{(-)\dagger} \hat{\gamma}_{\mathbf{k}\sigma}^{(-)} \rangle \right) \\
 &= \frac{1}{2L_x L_y} \sum_{\sigma} \sum_{\mathbf{k} \in \text{MBZ}} \text{Th}_{\mathbf{k}\sigma}^{(+)}[\beta, \mu].
 \end{aligned}$$

In last line we used Eq. (5.47). The thermal factor is spin-independent, hence we can set $\sigma = \uparrow$ and remove the sum over spin. This proves Eq. (5.52).

In Fig. 5.3a we sketch density as a function of the chemical potential, computed using Eq. (5.52) at $U/t = 10$ and $L = 256$. As we increase β , we observe that a sort of plateau emerges at $\mu \in [-\Delta, \Delta]$, becoming sharper as we decrease temperature. For an insulating phase, any infinitesimal shift to μ does not increase n . Indeed, in that case the chemical potential lies within a gap and no states can be filled by incrementing a little the chemical potential. On the contrary, for a metallic phase n changes as we vary infinitesimally μ . In Fig. 5.3a, we observe that the plateau is located at $n = 1/2$, i.e. half-filling, while for $|\mu| > \Delta$ the density varies with μ . Therefore, the commensurate AF phase is insulating only at half-filling.

In Fig. 5.3b we sketch the bands $\pm E_{\mathbf{k}\sigma}$ at $\mu/\Delta = 1.5$. The intense colour indicates an almost filled single-particle state, while the lighter gray colour an almost empty single-particle state. The band are gapped at zero energy, while μ crosses the upper band. This can be explained algebraically as follows. Consider the gran canonical hamiltonian $\hat{H} - \mu \hat{N}$, and use Eq. (5.40). The resulting

operator is diagonalised by the same matrix given in Eq. (5.44), being

$$-\mu\hat{N} = \sum_{\mathbf{k} \in \text{MBZ}} \begin{bmatrix} -\mu & \\ & -\mu \end{bmatrix} \implies W_{\mathbf{k}\sigma} [h_{\mathbf{k}\sigma} - \mu\mathbb{I}_{2 \times 2}] W_{\mathbf{k}\sigma}^\dagger = d_{\mathbf{k}\sigma} - \mu\mathbb{I}_{2 \times 2}, \quad (5.58)$$

thus for non-null chemical potential (i.e. finite doping) the eigenvalues are just shifted, $\pm E_{\mathbf{k}} - \mu$. This means that the chemical potential for the new fermions, described by the operators $\hat{\gamma}_{\mathbf{k}\sigma}$, is the same we had for the electrons. From Fig. 5.3a we know that at finite doping the chemical potential becomes larger than the gap, therefore μ crosses the upper/lower band for electron/hole doping. The resulting phase is metallic.

Instability of the doped commensurate antiferromagnet. As we said at the beginning of this chapter, there exist several ways to realise an Hubbard antiferromagnet. For the conventional HM, it is known [6, 7] that the commensurate AF phase is unstable at any finite doping. This is discussed briefly in App. D. The key takeaway is that insulating antiferromagnets are stable with respect to spin fluctuations, while metallic antiferromagnets are not.

In fact, the mere existence of a self-consistent solution does not guarantee stability: indeed, when in Sec. 2.2.2 we derived the variational solution, we only required it to be a stationary point of the HF energy. There is no need for the solution to be a minimum. In particular, the commensurate AF phase is a saddle point of HF energy and it can be shown that any finite doping disrupts commensurate ordering towards uncommensurate phases [6].

5.3 Generalisation to the EHM

We now include the non-local interaction $\hat{H}_V$. We anticipate here that we are going to obtain the same HF hamiltonian of Eq. (5.26) with renormalised chemical potential, bands and gap,

$$\mu \rightarrow \tilde{\mu}, \quad \epsilon_{\mathbf{k}\sigma} \rightarrow \tilde{\epsilon}_{\mathbf{k}\sigma}, \quad \Delta_{\mathbf{k}\sigma} \rightarrow \tilde{\Delta}_{\mathbf{k}\sigma}.$$

The explicit expressions for these renormalised quantities will be given, respectively, in Eqs. (5.72), (5.86) and (5.103). Due to this renormalisation of parameters, the diagonalisation procedure of Sec. 5.2.2 can be generalised.

In diagonal form, the bands $\pm \tilde{E}_{\mathbf{k}\sigma}$ read

$$\tilde{E}_{\mathbf{k}\sigma} \equiv \sqrt{\tilde{\epsilon}_{\mathbf{k}\sigma}^2 + |\tilde{\Delta}_{\mathbf{k}\sigma}|^2}.$$

We allow $\tilde{\Delta}_{\mathbf{k}\sigma}$ to be a complex function of $\mathbf{k}$ and σ ,

$$\tilde{\Delta}_{\mathbf{k}\sigma} = \text{Re}\{\tilde{\Delta}_{\mathbf{k}\sigma}\} + i \text{Im}\{\tilde{\Delta}_{\mathbf{k}\sigma}\}.$$

We define the renormalised angle $\tilde{\theta}_{\mathbf{k}\sigma}$, $\tilde{\zeta}_{\mathbf{k}\sigma}$ through the relations:

$$\sin 2\tilde{\theta}_{\mathbf{k}\sigma} = \frac{|\tilde{\Delta}_{\mathbf{k}\sigma}|}{\tilde{E}_{\mathbf{k}\sigma}}, \quad \cos 2\tilde{\theta}_{\mathbf{k}\sigma} = \frac{\tilde{\epsilon}_{\mathbf{k}\sigma}}{\tilde{E}_{\mathbf{k}\sigma}}, \quad (5.59)$$

and

$$\sin 2\tilde{\zeta}_{\mathbf{k}\sigma} = \frac{\text{Re}\{\tilde{\Delta}_{\mathbf{k}\sigma}\}}{|\tilde{\Delta}_{\mathbf{k}\sigma}|}, \quad \cos 2\tilde{\zeta}_{\mathbf{k}\sigma} = \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}\sigma}\}}{|\tilde{\Delta}_{\mathbf{k}\sigma}|}. \quad (5.60)$$

A sketch of these angles and the generalised diagonalisation procedure is given in Fig. 5.4. For a complex gap function, the new diagonalisation matrix is given by:

$$\begin{aligned} \tilde{W}_{\mathbf{k}\sigma} &\equiv e^{-i\tilde{\theta}_{\mathbf{k}\sigma}\tau^y} e^{-i\tilde{\zeta}_{\mathbf{k}\sigma}\tau^z} \\ &= \left(\cos \tilde{\theta}_{\mathbf{k}\sigma} - i \sin \tilde{\theta}_{\mathbf{k}\sigma} \tau^y \right) \left(\cos \tilde{\zeta}_{\mathbf{k}\sigma} - i \sin \tilde{\zeta}_{\mathbf{k}\sigma} \tau^z \right) \\ &= \begin{bmatrix} \cos \tilde{\theta}_{\mathbf{k}\sigma} & -\sin \tilde{\theta}_{\mathbf{k}\sigma} \\ \sin \tilde{\theta}_{\mathbf{k}\sigma} & \cos \tilde{\theta}_{\mathbf{k}\sigma} \end{bmatrix} \begin{bmatrix} e^{-i\tilde{\zeta}_{\mathbf{k}\sigma}} & \\ & e^{i\tilde{\zeta}_{\mathbf{k}\sigma}} \end{bmatrix} \\ &= \begin{bmatrix} \cos \tilde{\theta}_{\mathbf{k}\sigma} e^{-i\tilde{\zeta}_{\mathbf{k}\sigma}} & -\sin \tilde{\theta}_{\mathbf{k}\sigma} e^{i\tilde{\zeta}_{\mathbf{k}\sigma}} \\ \sin \tilde{\theta}_{\mathbf{k}\sigma} e^{-i\tilde{\zeta}_{\mathbf{k}\sigma}} & \cos \tilde{\theta}_{\mathbf{k}\sigma} e^{i\tilde{\zeta}_{\mathbf{k}\sigma}} \end{bmatrix}, \end{aligned}$$

and “updates” the diagonalisation matrix of Eq. (5.44).

Pseudospin EVs. The following relations hold (*tilda* sign indicates the renormalised quantity):

$$\begin{aligned}\langle \hat{\Psi}_{\mathbf{k}\sigma}^\dagger \tau^x \hat{\Psi}_{\mathbf{k}\sigma} \rangle &= -\sin 2\tilde{\theta}_{\mathbf{k}\sigma} \cos 2\tilde{\zeta}_{\mathbf{k}\sigma} \langle \hat{\Gamma}_{\mathbf{k}\sigma}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}\sigma} \rangle, \\ \langle \hat{\Psi}_{\mathbf{k}\sigma}^\dagger \tau^y \hat{\Psi}_{\mathbf{k}\sigma} \rangle &= -\sin 2\tilde{\theta}_{\mathbf{k}\sigma} \sin 2\tilde{\zeta}_{\mathbf{k}\sigma} \langle \hat{\Gamma}_{\mathbf{k}\sigma}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}\sigma} \rangle, \\ \langle \hat{\Psi}_{\mathbf{k}\sigma}^\dagger \tau^z \hat{\Psi}_{\mathbf{k}\sigma} \rangle &= \cos 2\tilde{\theta}_{\mathbf{k}\sigma} \langle \hat{\Gamma}_{\mathbf{k}\sigma}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}\sigma} \rangle.\end{aligned}$$

Let us extend Eq. (5.46) and define the renormalised thermal factors,

$$\tilde{\text{Th}}_{\mathbf{k}\sigma}^{(\pm)}[\beta, \tilde{\mu}] \equiv f\left(-\tilde{E}_{\mathbf{k}\sigma}; \beta, \tilde{\mu}\right) \pm f\left(\tilde{E}_{\mathbf{k}\sigma}; \beta, \tilde{\mu}\right), \quad (5.61)$$

which give, in the tilted frame, the z component of the pseudospin

$$\begin{aligned}\langle \hat{\Gamma}_{\mathbf{k}\sigma}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}\sigma} \rangle &= f\left(\tilde{E}_{\mathbf{k}\sigma}; \beta, \tilde{\mu}\right) - f\left(-\tilde{E}_{\mathbf{k}\sigma}; \beta, \tilde{\mu}\right) \\ &= -\tilde{\text{Th}}_{\mathbf{k}\sigma}^{(-)}[\beta, \tilde{\mu}].\end{aligned} \quad (5.62)$$

At half-filling, we have:

$$\tilde{\text{Th}}_{\mathbf{k}\sigma}^{(+)}[\beta, 0] = 1, \quad (5.63)$$

$$\tilde{\text{Th}}_{\mathbf{k}\sigma}^{(-)}[\beta, 0] = \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}\sigma}}{2}\right). \quad (5.64)$$

We will use these expression only in the last part, carrying out computation at generic $\tilde{\mu}$.

Proof 5.12 — Proof of Eqs. (5.63), (5.64).

At half-filling, $\mu = 0$. For a null chemical potential, the Fermi-Dirac distribution is given by

$$f(\epsilon; \beta, 0) = \frac{1}{e^{\beta\epsilon} + 1}.$$

Therefore, the following identity holds:

$$\begin{aligned}f(-\epsilon; \beta, 0) &= \frac{1}{e^{-\beta\epsilon} + 1} \\ &= \frac{e^{\beta\epsilon}}{1 + e^{\beta\epsilon}} \\ &= 1 - \frac{1}{e^{\beta\epsilon} + 1} \\ &= 1 - f(\epsilon; \beta, 0).\end{aligned}$$

Hence,

$$f(-\tilde{E}_{\mathbf{k}\sigma}; \beta, 0) + f(\tilde{E}_{\mathbf{k}\sigma}; \beta, 0) = 1$$

which proves Eq. (5.63).

Moreover, we have:

$$\begin{aligned}f(-\epsilon; \beta, 0) - f(\epsilon; \beta, 0) &= \frac{1}{e^{-\beta\epsilon} + 1} - \frac{1}{e^{\beta\epsilon} + 1} \\ &= \frac{e^{\beta\epsilon}}{e^{\beta\epsilon} + 1} - \frac{1}{e^{\beta\epsilon} + 1} \\ &= \frac{e^{\beta\epsilon/2} - e^{-\beta\epsilon/2}}{e^{\beta\epsilon/2} + e^{-\beta\epsilon/2}}.\end{aligned}$$

Therefore,

$$f(-\tilde{E}_{\mathbf{k}\sigma}; \beta, 0) - f(\tilde{E}_{\mathbf{k}\sigma}; \beta, 0) = \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}\sigma}}{2}\right)$$




Figure 5.4 Sketch of the diagonalisation procedure for the EHM antiferromagnet. To diagonalise the HF hamiltonian, we align the z axis with the pseudo-field. This is an extension of Fig. 5.2.

which proves Eq. (5.64).

Pseudospin EVs in the three directions are:

$$\langle \hat{s}_{\mathbf{k}\sigma}^x \rangle = \frac{\text{Re}\{\tilde{\Delta}_{\mathbf{k}\sigma}\}}{\tilde{E}_{\mathbf{k}\sigma}} \tilde{\text{Th}}_{\mathbf{k}\sigma}^{(-)}[\beta, \tilde{\mu}], \quad (5.65)$$

$$\langle \hat{s}_{\mathbf{k}\sigma}^y \rangle = \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}\sigma}\}}{\tilde{E}_{\mathbf{k}\sigma}} \tilde{\text{Th}}_{\mathbf{k}\sigma}^{(-)}[\beta, \tilde{\mu}], \quad (5.66)$$

$$\langle \hat{s}_{\mathbf{k}\sigma}^z \rangle = -\frac{\tilde{\epsilon}_{\mathbf{k}\sigma}}{\tilde{E}_{\mathbf{k}\sigma}} \tilde{\text{Th}}_{\mathbf{k}\sigma}^{(-)}[\beta, \tilde{\mu}]. \quad (5.67)$$

These equations are an extension of Eqs. (5.48), (5.49), (5.50) for the EHM.

Magnetisation and density self-consistency equations. By renormalising all quantities, the self-consistency equation for magnetisation must stay formally the same of Eq. (5.51),

$$m = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\tilde{\Delta}_{\mathbf{k}\uparrow}}{\tilde{E}_{\mathbf{k}\uparrow}} \tilde{\text{Th}}_{\mathbf{k}\uparrow}^{(-)}[\beta, \tilde{\mu}]. \quad (5.68)$$

We will be able to prove this statement rigorously in Sec. 5.4. For the density n an identical argument holds, and Eq. (5.52) is extended to

$$n = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \tilde{\text{Th}}_{\mathbf{k}\uparrow}^{(+)}[\beta, \tilde{\mu}]. \quad (5.69)$$

Contributions of each channel. We recall now Tab. 3.3. The contractions that, through Wick's theorem, possibly give non-zero contributions are, for $\hat{H}_V$,

$$\begin{aligned} \hat{H}_V \text{ (o.s. sector)} & \quad \text{Hartree contractions,} \\ \hat{H}_V \text{ (s.s. sector)} & \quad \text{Hartree and Fock contractions.} \end{aligned}$$

Operator	Sector	Channel	Net effect
$\hat{H}_U$		Hartree	μ shift by $-nU$, real part of the AF gap
	o.s.	Hartree	μ shift by nzV
$\hat{H}_V$		Hartree	μ shift by nzV
	s.s.	Fock	Bands renormalisation and resymmetrization, imaginary part of the AF gap

Table 5.1 | Wick's contractions for each interaction and channel in the HF framework, and their net effect in the AF phase.

Tab. 5.1 summarises the roles of the various contraction channels (including those from the local repulsion), which are described in next sections.

5.3.1 Effects of the non-local Hartree terms: chemical potential renormalisation

The approximation to $\hat{H}_V$ in the non-local Hartree channel is given by

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} -2znV \times \hat{N} - E_{\text{H},V}^{(\text{AF})} + (\text{Fock channel}), \quad (5.70)$$

with the contraction energy

$$E_{\text{H},V}^{(\text{AF})} \equiv V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma, \sigma'} \langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}\sigma} \rangle \langle \hat{c}_{\mathbf{r}'\sigma'}^\dagger \hat{c}_{\mathbf{r}'\sigma'} \rangle. \quad (5.71)$$

From Eq. (5.70), we see that the Hartree channel of the non-local attraction contributes to the chemical potential shift. Combining Eq. (5.26) with Eq. (5.70), the full shift to chemical potential reads

$$\tilde{\mu} \equiv \mu + n(2zV - U), \quad (5.72)$$

which is identical to what we found for the normal phase, in Eq. (4.8).

Proof 5.13 — Proof of Eq. (5.70).

In Eq. (4.11) we separated the Hartree channel approximation to $\hat{H}_V$ in s.s. and o.s. channels. The same equation is valid in this context. Using the definition of $E_{\text{H},V}^{(\text{AF})}$ given in Eq. (5.71), we obtain

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} & -V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (\langle \hat{n}_{\mathbf{r}\sigma} \rangle \hat{n}_{\mathbf{r}'\sigma} + \hat{n}_{\mathbf{r}\sigma} \langle \hat{n}_{\mathbf{r}'\sigma} \rangle) \\ & - V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (\langle \hat{n}_{\mathbf{r}\sigma} \rangle \hat{n}_{\mathbf{r}'\bar{\sigma}} + \hat{n}_{\mathbf{r}\sigma} \langle \hat{n}_{\mathbf{r}'\bar{\sigma}} \rangle) - E_{\text{H},V}^{(\text{AF})} + (\text{Fock channel}). \end{aligned} \quad (5.73)$$

Inserting the Ansatz of Eq. (5.2), we obtain

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} & \overbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (\hat{n}_{\mathbf{r}'\sigma} + \hat{n}_{\mathbf{r}\sigma}) + mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{\delta_{\sigma=\uparrow}} \left((-1)^{x+y} \hat{n}_{\mathbf{r}'\sigma} + (-1)^{x'+y'} \hat{n}_{\mathbf{r}\sigma} \right)}^{\text{s.s.}} \\ & \underbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{n}_{\mathbf{r}'\bar{\sigma}} + \hat{n}_{\mathbf{r}\sigma}] + mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \left((-1)^{x+y+\delta_{\sigma=\uparrow}} \hat{n}_{\mathbf{r}'\bar{\sigma}} + (-1)^{x'+y'+\delta_{\bar{\sigma}=\uparrow}} \hat{n}_{\mathbf{r}\sigma} \right)}_{\text{o.s.}} \\ & - E_{\text{H},V}^{(\text{AF})} + (\text{Fock channel}). \end{aligned} \quad (5.74)$$

For a square lattice, if $\mathbf{r} = (x, y)$ and $\mathbf{r}' = (x', y')$ are NNs, both the following identities are true

$$\begin{aligned} (-1)^{x'+y'} &= (-1)^{x+y+1}, \\ (-1)^{\delta_{\bar{\sigma}=\uparrow}} &= (-1)^{\delta_{\sigma=\uparrow}+1}. \end{aligned}$$

Therefore, Eq. (5.74) becomes

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} & \overbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\sigma})}^{\text{s.s. (density)}} - \overbrace{mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{x+y+\delta_{\sigma=\uparrow}} (\hat{n}_{\mathbf{r}\sigma} - \hat{n}_{\mathbf{r}'\sigma})}^{\text{s.s. (magnetisation)}} \\ & \underbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\bar{\sigma}})}_{\text{o.s. (density)}} + \underbrace{mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{x+y+\delta_{\sigma=\uparrow}} (\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\bar{\sigma}})}_{\text{o.s. (magnetisation)}} \\ & - E_{\text{H},V}^{(\text{AF})} + (\text{Fock channel}). \end{aligned} \quad (5.75)$$

In the above expressions we separated in “density” and “magnetisation” terms. Let us deal with them separately.

Density terms. In the sum of s.s. and o.s. density terms of Eq. (5.75) we recognise the operator of Eq. (4.12), obtained for the normal phase. Hence we obtain the same result of Eq. (4.13),

$$-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\sigma}) - nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\bar{\sigma}}) = -2znV \times \hat{N}. \quad (5.76)$$

Magnetisation terms. The magnetisation s.s. and o.s. terms of Expr. (5.75) perfectly cancel out:

$$mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{x+y+\delta_{\sigma=\uparrow}} [(\hat{n}_{\mathbf{r}\sigma} - \hat{n}_{\mathbf{r}'\sigma}) - (\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\bar{\sigma}})] = 0.$$

Therefore the only contribution to the HF approximation of $\hat{H}_V$, in the Hartree channel, comes from Eq. (5.76). Inserting the latter in Eq. (5.75), Eq. (5.70) is proven.

The contraction energy of Eq. (5.71) reduces to

$$E_{\text{H},V}^{(\text{AF})} = -L_x L_y \times 2n^2, \quad (5.77)$$

which coincides with that for the normal phase, given in Eq. (4.15). This is coherent with the fact that the shift to chemical potential is identical.

Proof 5.14 — Proof of Eq. (5.77): non-local Hartree contraction energy.

Separate Eq. (5.71) in s.s. and o.s. sectors,

$$E_{\text{H},V}^{(\text{AF})} = V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \left(\overbrace{\langle \hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}\sigma} \rangle \langle \hat{c}_{\mathbf{r}'\sigma}^{\dagger} \hat{c}_{\mathbf{r}'\sigma} \rangle}^{\text{s.s.}} + \overbrace{\langle \hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}\sigma} \rangle \langle \hat{c}_{\mathbf{r}'\bar{\sigma}}^{\dagger} \hat{c}_{\mathbf{r}'\bar{\sigma}} \rangle}^{\text{o.s.}} \right). \quad (5.78)$$

s.s. sector contraction energy. The s.s. sector of Eq. (5.78) reduces to

$$V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \langle \hat{n}_{\mathbf{r}\sigma} \rangle \langle \hat{n}_{\mathbf{r}'\sigma} \rangle = V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \overbrace{\left[(n + (-1)^{x+y}m)(n + (-1)^{x'+y'}m) \right]}^{\sigma=\uparrow} + V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \underbrace{\left[(n - (-1)^{x+y}m)(n - (-1)^{x'+y'}m) \right]}_{\sigma=\downarrow}.$$

The mixed terms (those of order $\mathcal{O}(n) \times \mathcal{O}(m)$) cancel out, and since sign factors necessarily alternate on neighbouring sites, the remainder is just

$$V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \langle \hat{n}_{\mathbf{r}\sigma} \rangle \langle \hat{n}_{\mathbf{r}'\sigma} \rangle = V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} 2(n^2 - m^2). \quad (5.79)$$

o.s. sector contraction energy. Proceeding identically on the o.s. sector of Eq. (5.78), we get

$$V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \langle \hat{n}_{\mathbf{r}\sigma} \rangle \langle \hat{n}_{\mathbf{r}'\bar{\sigma}} \rangle = V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \overbrace{\left[(n + (-1)^{x+y}m)(n - (-1)^{x'+y'}m) \right]}^{\sigma=\uparrow} + V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \underbrace{\left[(n - (-1)^{x+y}m)(n + (-1)^{x'+y'}m) \right]}_{\sigma=\downarrow},$$

which gives

$$V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \langle \hat{n}_{\mathbf{r}\sigma} \rangle \langle \hat{n}_{\mathbf{r}'\bar{\sigma}} \rangle = V \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} 2(n^2 + m^2). \quad (5.80)$$

Then, summing Eqs. (5.79) and (5.80), we get

$$E_{\text{H},V}^{(\text{AF})} = -2V \sum_{\langle ij \rangle} \sum_{\sigma} n^2 = -4n^2V \times \frac{z}{2} L_x L_y.$$

which coincides with Eq. (5.77).

5.3.2 Effects of the Fock terms: gap and bare bands renormalisation

In this subsection we discuss the Fock contributions to the HF approximated hamiltonian, in the AF phase. We start by Eq. (4.17), derived in the context of the normal phase,

$$\hat{H}_V^{\text{HF}} \simeq (\text{Hartree channel}) + \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \sum_{\sigma} (\cos(\delta k_x) + \cos(\delta k_y)) \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \rangle - E_{\text{F},V}^{(\text{AF})}, \quad (5.81)$$

where $E_{\text{F},V}^{(\text{AF})}$, see Eq. (4.18), reads

$$E_{\text{F},V}^{(\text{AF})} \equiv \frac{V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \sum_{\sigma} (\cos(\delta k_x) + \cos(\delta k_y)) \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \rangle, \quad (5.82)$$

As we specified in Sec. 5.1.2, the commensurate AF phase is described by two EVs being in non-zero in reciprocal space,

$$|\langle \hat{n}_{\mathbf{k}\sigma} \rangle| \geq 0 \quad \text{and} \quad |\langle \hat{\psi}_{\mathbf{k}\sigma} \rangle| \geq 0,$$

This implies only two contributions in Eq. (5.81) are non-zero:

$$\mathbf{k} = -\mathbf{k}' \quad \text{and} \quad \mathbf{k} + \boldsymbol{\pi} = -\mathbf{k}'.$$

Then Eq. (5.81) becomes

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \\ \times \left(\underbrace{\langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma} \rangle}_{\text{Diagonal terms}} - \underbrace{\langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}+\boldsymbol{\pi}\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}-\boldsymbol{\pi}\sigma} \rangle}_{\text{Off-diagonal terms}} \right) - E_{\text{F},V}^{(\text{AF})}, \end{aligned} \quad (5.83)$$

where we separated the hamiltonian in *diagonal* and *off-diagonal* terms.

Diagonal terms. The diagonal terms of Eq. (5.83) reduce to

$$\begin{aligned} \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma} \rangle \\ = V \sum_{\mathbf{k}, \sigma} \left(u_{\sigma}^{(s^*)} (\cos q_x + \cos q_y) + u_{\sigma}^{(d)} (\cos q_x - \cos q_y) \right) \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}, \end{aligned} \quad (5.84)$$

where we used Eq. (5.12), where the harmonics of $u_{\mathbf{k}\sigma}$ were defined.

Proof 5.15 — Proof of Eq. (5.84).

We define new variables as in Fig. 4.1, $\mathbf{q} \equiv \mathbf{K} + \mathbf{k}$, $\mathbf{q}' \equiv \mathbf{K} - \mathbf{k}$. Applying this change of variables to the diagonal terms of Eq. (5.83), we obtain

$$\begin{aligned} \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}\sigma} \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma} \rangle \\ = \frac{2V}{L_x L_y} \sum_{\mathbf{q}, \mathbf{q}'} \sum_{\sigma} [\cos(\delta q_x) + \cos(\delta q_y)] \langle \hat{c}_{\mathbf{q}\sigma}^\dagger \hat{c}_{\mathbf{q}\sigma} \rangle \langle \hat{c}_{\mathbf{q}'\sigma}^\dagger \hat{c}_{\mathbf{q}'\sigma} \rangle \\ = \sum_{\gamma, \sigma} \left(\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{n}_{\mathbf{q}\sigma} \rangle \right) \times V \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} \hat{n}_{\mathbf{q}'\sigma} \\ = \sum_{\gamma, \sigma} u_{\sigma}^{(\gamma)} \times V \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} \hat{n}_{\mathbf{q}'\sigma}. \end{aligned} \quad (5.85)$$

We used the decomposition of Eq. (3.47). In the last line, we recognised the harmonics given in Eq. (5.12).

We can re-use the argument that, for the normal phase, lead us to Eq. (4.24). We are not breaking inversion symmetry (and as a consequence, neither C_2): this implies that, whatever the shape of the renormalised bands, it must be

$$\tilde{\epsilon}_{\mathbf{q}\sigma} = \tilde{\epsilon}_{-\mathbf{q}\sigma} \implies u_{\sigma}^{(p\ell)} = 0 \quad \text{with} \quad \ell \in \{x, y\}.$$

Renormalised bare bands need to be symmetric functions of the moment. Hence for $\gamma \in \{p_x, p_y\}$ the above sum vanishes. Using this selection rule, Eq. (5.85) reduces to Eq. (5.84).

Apart from the explicit form of the self-consistency equations, this discussion is formally the same of the normal phase, carried out in Sec. 4.2.2. Therefore the bands renormalisation, discussed in Sec. 4.3.1, applies also for the AF phase. Specifically, the renormalised bands are

$$\tilde{\epsilon}_{\mathbf{k}\sigma} \equiv \epsilon_{\mathbf{k}\sigma} + u_{\sigma}^{(s^*)} V (\cos k_x + \cos k_y) + u_{\sigma}^{(d)} V (\cos k_x - \cos k_y). \quad (5.86)$$

The d -wave harmonic $u^{(d)}$ is zero if C_4 is unbroken, otherwise is finite.

Symmetry	Self-consistency equation	Graph
s^* -wave	$u^{(s^*)} = -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x + \cos k_y) \frac{\tilde{\epsilon}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]$	Fig. 3.4b
$d_{x^2-y^2}$ -wave	$u^{(d)} = -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x - \cos k_y) \frac{\tilde{\epsilon}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]$	Fig. 3.4e

Table 5.2 | Symmetry and spin resolved self-consistency equations for the harmonics of $u_{\mathbf{k}}$.

The self-consistency equation for the harmonics of $u_{\mathbf{k}\sigma}$, given in Eq. (5.12), can be written as

$$u_{\sigma}^{(\gamma)} = -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\tilde{\epsilon}_{\mathbf{k}\sigma}}{\tilde{E}_{\mathbf{k}\sigma}} \tilde{\text{Th}}_{\mathbf{k}\sigma}^{(-)}[\beta, \tilde{\mu}]. \quad (5.87)$$

The symmetry-resolved self-consistency equations are reported explicitly in Tab. 5.2. Since in Eq. (5.87) all quantities are spin-independent, we omit spin index on these HFPs.

Proof 5.16 — Proof of Eq. (5.87): self-consistency equations for $u_{\sigma}^{(\gamma)}$.

Recall the definition of $u_{\mathbf{k}\sigma}$,

$$u_{\mathbf{k}\sigma} = \langle \hat{n}_{\mathbf{k}\sigma} \rangle,$$

given in Eq. (5.10). We expand the harmonics decomposition of Eq. (5.12),

$$\begin{aligned} u_{\sigma}^{(\gamma)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{n}_{\mathbf{k}\sigma} \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \left(\varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{n}_{\mathbf{k}\sigma} \rangle + \varphi_{\mathbf{k}+\pi}^{(\gamma)} \langle \hat{n}_{\mathbf{k}+\pi\sigma} \rangle \right) \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{n}_{\mathbf{k}\sigma} - \hat{n}_{\mathbf{k}+\pi\sigma} \rangle. \end{aligned}$$

In last line we used Eq. (5.24). From Eq. (5.39), we recognise the z component of the pseudospin operator

$$\begin{aligned} u_{\sigma}^{(\gamma)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{s}_{\mathbf{k}\sigma}^z \rangle \\ &= -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\tilde{\epsilon}_{\mathbf{k}\sigma}}{\tilde{E}_{\mathbf{k}\sigma}} \tilde{\text{Th}}_{\mathbf{k}\sigma}^{(-)}[\beta, \tilde{\mu}], \end{aligned} \quad (5.88)$$

where, in the last line, Eq. (5.67) has been used. This proves Eq. (5.87).

Similarly to the normal phase, also for the AF phase we can define the symmetry-resolved components of the renormalised hopping amplitude,

$$\tilde{t}_{\sigma}^{(s^*)} \equiv t - \frac{u_{\sigma}^{(s^*)} V}{2} \quad \text{and} \quad \tilde{t}_{\sigma}^{(d)} \equiv -\frac{u_{\sigma}^{(d)} V}{2}. \quad (5.89)$$

We re-used the definitions of Eq. (4.28).

Off-diagonal terms. The off-diagonal sector of Eq. (5.83) is reduced to

$$\begin{aligned} & -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}+\pi\sigma}^{\dagger} \hat{c}_{\mathbf{K}+\mathbf{k}\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{K}-\mathbf{k}-\pi\sigma} \\ & = V \sum_{\sigma} \sum_{\gamma \neq s} \sum_{\mathbf{k} \in \text{MBZ}} \left(i v_{\sigma}^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} \hat{\psi}_{\mathbf{k}\sigma} + \text{h.c.} \right), \end{aligned} \quad (5.90)$$

where we used the harmonics of $v_{\mathbf{k}\sigma}$, defined in Eq. (5.13), and the functions of Tab. 3.5.

Proof 5.17 — Proof of Eq. (5.90).

We define new variables as in Fig. 4.1, $\mathbf{q} \equiv \mathbf{K} + \mathbf{k}$, $\mathbf{q}' \equiv \mathbf{K} - \mathbf{k}$. Applying this change of variables to the off-diagonal terms of Eq. (5.83) and using the decomposition of Eq. (3.47), we obtain

$$\begin{aligned} & -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}+\boldsymbol{\pi}\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}-\boldsymbol{\pi}\sigma} \\ & = -\sum_{\sigma} \sum_{\gamma \neq s} \left(\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{q}\sigma}^\dagger \rangle \right) \times V \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} \hat{\psi}_{\mathbf{q}'\sigma}^\dagger. \end{aligned} \quad (5.91)$$

We shift the first sum by an AF vector $\boldsymbol{\pi}$ defining $\mathbf{k} + \boldsymbol{\pi} \equiv \mathbf{q}$. Using Eq. (5.22) as well as reciprocal space periodicity by a shift $\boldsymbol{\pi}$,

$$\begin{aligned} \frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{q}\sigma}^\dagger \rangle &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}+\boldsymbol{\pi}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{k}+\boldsymbol{\pi}\sigma}^\dagger \rangle \\ &= -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{k}\sigma} \rangle. \end{aligned}$$

Recall the definition $v_{\mathbf{k}\sigma} = i \langle \hat{\psi}_{\mathbf{k}\sigma} \rangle$ of Eq. (5.11). Inserting the above equation in Eq. (5.91), we obtain

$$\begin{aligned} & -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}+\boldsymbol{\pi}\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}-\boldsymbol{\pi}\sigma} \\ & = \sum_{\sigma} \sum_{\gamma \neq s} \left(\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{q}\sigma} \rangle \right) \times V \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} \hat{\psi}_{\mathbf{q}'\sigma}^\dagger \\ & = -i \sum_{\sigma} \sum_{\gamma \neq s} v_{\sigma}^{(\gamma)} \times V \sum_{\mathbf{q}'} \varphi_{\mathbf{q}'}^{(\gamma)*} \hat{n}_{\mathbf{q}'\sigma}, \end{aligned} \quad (5.92)$$

where we used Eq. (5.13). We restrict the sum to the MBZ,

$$\begin{aligned} & -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}+\boldsymbol{\pi}\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}-\boldsymbol{\pi}\sigma} \\ & = -iV \sum_{\sigma} \sum_{\gamma \neq s} \sum_{\mathbf{q} \in \text{MBZ}} \left(v_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}}^{(\gamma)*} \hat{\psi}_{\mathbf{q}\sigma}^\dagger + v_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}+\boldsymbol{\pi}}^{(\gamma)*} \hat{\psi}_{\mathbf{q}+\boldsymbol{\pi}\sigma}^\dagger \right), \end{aligned}$$

and use Eq. (5.22), exchanging terms positions

$$\begin{aligned} & -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}+\boldsymbol{\pi}\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}-\boldsymbol{\pi}\sigma} \\ & = iV \sum_{\sigma} \sum_{\gamma \neq s} \sum_{\mathbf{q} \in \text{MBZ}} \left(v_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}}^{(\gamma)*} \hat{\psi}_{\mathbf{q}\sigma} - v_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}}^{(\gamma)*} \hat{\psi}_{\mathbf{q}\sigma}^\dagger \right). \end{aligned} \quad (5.93)$$

Recalling Tab. 3.5, for the four symmetries we are considering, the functions $\varphi_{\mathbf{k}}^{(\gamma)}$ are real-valued for $\gamma = s^*, d$, and purely imaginary for $\gamma = p_x, p_y$. We anticipate the result of Eq. (5.96),

$$v_{\sigma}^{(s^*)}, v_{\sigma}^{(d)} \in \mathbb{R}, \quad v_{\sigma}^{(p_x)}, v_{\sigma}^{(p_y)} \in i\mathbb{R},$$

Symmetry	Self-consistency equation	Graph
s^* -wave	$v^{(s^*)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x + \cos k_y) \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]$	Fig. 3.4b
p_x -wave	$v^{(p_x)} = \frac{i\sqrt{2}}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \sin k_x \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]$	Fig. 3.4c
p_y -wave	$v^{(p_y)} = \frac{i\sqrt{2}}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \sin k_y \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]$	Fig. 3.4d
$d_{x^2-y^2}$ -wave	$v^{(d)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x - \cos k_y) \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]$	Fig. 3.4e

Table 5.3 | Symmetry and spin resolved self-consistency equations for the harmonics of $v_{\mathbf{k}}$.

that will be obtained later. Then

$$iv_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}}^{(\gamma)*} \in i\mathbb{R}, \quad (5.94)$$

and to conjugate a purely imaginary number it suffices to put a minus sign in front of it,

$$-iv_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}}^{(\gamma)*} = \left(iv_{\sigma}^{(\gamma)} \varphi_{\mathbf{q}}^{(\gamma)*} \right)^*.$$

Therefore, substituting this identity in Eq. (5.93), we obtain Eq. (5.90).

The HFPs $v_{\sigma}^{(\gamma)}$ obey the self-consistency equations

$$\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{MBZ}} \varphi_{\mathbf{q}}^{(\gamma)} \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{q}\sigma}\}}{\tilde{E}_{\mathbf{q}\sigma}} \tilde{\text{Th}}_{\mathbf{q}\sigma}^{(-)}[\beta, \tilde{\mu}]. \quad (5.95)$$

These equations are reported, symmetry-resolved, in Tab. 5.3. By direct inspection of the equations thereby reported, we see that

$$v_{\sigma}^{(s^*)}, v_{\sigma}^{(d)} \in \mathbb{R}, \quad v_{\sigma}^{(p_x)}, v_{\sigma}^{(p_y)} \in i\mathbb{R}. \quad (5.96)$$

We have used this result before, in Eq. (5.94).

In Tab. 5.3 the four derived self-consistency equations are reported, ignoring spin index. Indeed, all quantities in Eq. (5.95) are independent of σ .

Proof 5.18 — Proof of Eq. (5.95): self-consistency equations for $v_{\sigma}^{(\gamma)}$.

We expand Eq. (5.13),

$$\begin{aligned} v_{\sigma}^{(\gamma)} &= \frac{i}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{q}\sigma} \rangle \\ &= \frac{i}{L_x L_y} \sum_{\mathbf{q} \in \text{MBZ}} \left(\varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{q}\sigma} \rangle + \varphi_{\mathbf{q}+\boldsymbol{\pi}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{q}+\boldsymbol{\pi}\sigma} \rangle \right) \\ &= \frac{i}{L_x L_y} \sum_{\mathbf{q} \in \text{MBZ}} \varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{\psi}_{\mathbf{q}\sigma} - \hat{\psi}_{\mathbf{q}\sigma}^{\dagger} \rangle. \end{aligned}$$

In last line we used Eq. (5.24). From Eq. (5.38), we recognise the y component of the pseudospin

operator

$$\begin{aligned} v_{\sigma}^{(\gamma)} &= \frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{MBZ}} \varphi_{\mathbf{q}}^{(\gamma)} \langle \hat{\Psi}_{\mathbf{q}\sigma}^{\dagger} \tau^y \hat{\Psi}_{\mathbf{q}\sigma} \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{MBZ}} \varphi_{\mathbf{q}}^{(\gamma)} \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{q}\sigma}\}}{\tilde{E}_{\mathbf{q}\sigma}} \tilde{\text{Th}}_{\mathbf{q}\sigma}^{(-)}[\beta, \tilde{\mu}], \end{aligned} \quad (5.97)$$

where we used Eq. (5.66). Last line coincides with Eq. (5.95).

The contraction energy of Eq. (5.82) reduces to

$$E_{\text{F},V}^{(\text{AF})} = L_x L_y \times \frac{V}{2} \sum_{\sigma} \sum_{\gamma \neq s} \left(|u_{\sigma}^{(\gamma)}|^2 + |v_{\sigma}^{(\gamma)}|^2 \right), \quad (5.98)$$

Note that, due to spin independency, we can omit the spin index, remove the sum over σ and the $1/2$ prefactor.

Proof 5.19 — Proof of Eq. (5.98): Fock contraction energy.

Consider Eq. (5.82). Using the changing variables of Fig. 4.1, we get

$$E_{\text{F},V}^{(\text{AF})} = \frac{V}{L_x L_y} \sum_{\mathbf{q}, \mathbf{q}'} \sum_{\sigma} [\cos(\delta q_x) + \cos(\delta q_y)] \left(\underbrace{\langle \hat{n}_{\mathbf{q}\sigma} \rangle \langle \hat{n}_{\mathbf{q}'\sigma} \rangle}_{\text{Diagonal terms}} - \underbrace{\langle \hat{\psi}_{\mathbf{q}\sigma} \rangle \langle \hat{\psi}_{\mathbf{q}'\sigma} \rangle}_{\text{Off-diagonal terms}} \right).$$

Applying the decomposition of Eq. (3.47) and using Eqs. (5.10), (5.11), we obtain:

$$E_{\text{F},V}^{(\text{AF})} = \underbrace{\frac{V}{2L_x L_y} \sum_{\sigma} \sum_{\gamma \neq s} \sum_{\mathbf{q}, \mathbf{q}'} \varphi_{\mathbf{q}}^{(\gamma)} \varphi_{\mathbf{q}'}^{(\gamma)*} u_{\mathbf{q}\sigma} u_{\mathbf{q}'\sigma}}_{\text{Diagonal terms}} + \underbrace{\frac{V}{2L_x L_y} \sum_{\sigma} \sum_{\gamma \neq s} \sum_{\mathbf{q}, \mathbf{q}'} \varphi_{\mathbf{q}}^{(\gamma)} \varphi_{\mathbf{q}'}^{(\gamma)*} v_{\mathbf{q}\sigma} v_{\mathbf{q}'\sigma}}_{\text{Off-diagonal terms}}. \quad (5.99)$$

Diagonal terms. Let us start from the diagonal part of Eq. (5.99). It holds

$$u_{\mathbf{q}\sigma} = \langle \hat{n}_{\mathbf{q}\sigma} \rangle \implies u_{\mathbf{q}\sigma} = u_{\mathbf{q}\sigma}^*,$$

and then

$$\begin{aligned} \frac{1}{(L_x L_y)^2} \sum_{\mathbf{q}, \mathbf{q}'} \varphi_{\mathbf{q}}^{(\gamma)} \varphi_{\mathbf{q}'}^{(\gamma)*} u_{\mathbf{q}\sigma} u_{\mathbf{q}'\sigma} &= \left(\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} u_{\mathbf{q}\sigma} \right) \left(\frac{1}{L_x L_y} \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} u_{\mathbf{q}'\sigma}^* \right) \\ &= |u_{\sigma}^{(\gamma)}|^2. \end{aligned} \quad (5.100)$$

Off-diagonal terms. Consider the off-diagonal part of Eq. (5.99). Using Eq. (5.23), we get

$$v_{\mathbf{q}\sigma} = i \langle \hat{\psi}_{\mathbf{q}\sigma} \rangle \implies v_{\mathbf{q}+\pi\sigma} = -v_{\mathbf{q}\sigma}^*,$$

Since the form factors satisfy $\varphi_{\mathbf{q}+\pi}^{(\gamma)} = -\varphi_{\mathbf{q}}^{(\gamma)}$, we obtain

$$\begin{aligned} \frac{1}{(L_x L_y)^2} \sum_{\mathbf{q}, \mathbf{q}'} \varphi_{\mathbf{q}}^{(\gamma)} \varphi_{\mathbf{q}'}^{(\gamma)*} v_{\mathbf{q}\sigma} v_{\mathbf{q}'\sigma} &= \left(\frac{1}{L_x L_y} \sum_{\mathbf{q} \in \text{BZ}} \varphi_{\mathbf{q}}^{(\gamma)} v_{\mathbf{q}\sigma} \right) \left(\frac{1}{L_x L_y} \sum_{\mathbf{q}' \in \text{BZ}} \varphi_{\mathbf{q}'}^{(\gamma)*} v_{\mathbf{q}'\sigma} \right) \\ &= v_{\sigma}^{(\gamma)} \left(\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}+\pi}^{(\gamma)*} v_{\mathbf{k}+\pi\sigma} \right), \end{aligned}$$

where in the second passage we employed the change of variables $\mathbf{q}' = \mathbf{k} + \boldsymbol{\pi}$ and used BZ periodicity to map the original sum onto a new sum for $\mathbf{k} \in \text{BZ}$.

$$\begin{aligned} \frac{1}{(L_x L_y)^2} \sum_{\mathbf{q}, \mathbf{q}'} \varphi_{\mathbf{q}}^{(\gamma)} \varphi_{\mathbf{q}'}^{(\gamma)*} v_{\mathbf{q}\sigma} v_{\mathbf{q}'\sigma} &= v_{\sigma}^{(\gamma)} \left(\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)*} v_{\mathbf{k}\sigma}^* \right) \\ &= |v_{\sigma}^{(\gamma)}|^2. \end{aligned} \quad (5.101)$$

Inserting Eqs. (5.100), (5.101) in Eq. (5.99), we obtain the expression of Eq. (5.98).

We examined all contribution to the HF hamiltonian. In next section we collect all results and describe the features of the HF ground-state.

5.4 The HF hamiltonian and its properties

We recollect all results into a single hamiltonian approximation. Summing Eqs. (5.26), (5.84) and (5.90), we obtain the HF hamiltonian

$$\begin{aligned} \hat{H} - \mu \hat{N} &\stackrel{\text{HF}}{\simeq} \sum_{\sigma} \sum_{\mathbf{k} \in \text{MBZ}} \tilde{\epsilon}_{\mathbf{k}\sigma} (\hat{n}_{\mathbf{k}\sigma} - \hat{n}_{\mathbf{k}+\boldsymbol{\pi}\sigma}) \\ &\quad - \sum_{\sigma} \sum_{\mathbf{k} \in \text{MBZ}} \left(\tilde{\Delta}_{\mathbf{k}\sigma} \hat{\psi}_{\mathbf{k}\sigma} + \tilde{\Delta}_{\mathbf{k}\sigma}^* \hat{\psi}_{\mathbf{k}\sigma}^{\dagger} \right) - \tilde{\mu} \hat{N} + E_c^{(\text{AF})}. \end{aligned} \quad (5.102)$$

We defined renormalised bands, see Eq. (5.86), as

$$\tilde{\epsilon}_{\mathbf{k}\sigma} = \epsilon_{\mathbf{k}\sigma} + V \sum_{\gamma \in \{s^*, d\}} u_{\sigma}^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*},$$

the renormalised gap function as

$$\tilde{\Delta}_{\mathbf{k}\sigma} \equiv \Delta_{\mathbf{k}\sigma} - iV \sum_{\gamma} v_{\sigma}^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*}, \quad (5.103)$$

and total contraction energy as the (negative) sum of all contraction energies, see Eqs. (5.31), (5.77) and (5.98),

$$E_c^{(\text{AF})} \equiv - \left(E_{\text{H},U}^{(\text{AF})} + E_{\text{H},V}^{(\text{AF})} + E_{\text{F},V}^{(\text{AF})} \right). \quad (5.104)$$

5.4.1 Gap complexification and AF flavours

Recalling Eq. (5.94), we observe that the correction to the gap function in Eq. (5.103) is purely imaginary. Instead, the HM gap is purely real, $\Delta_{\mathbf{k}\sigma} = (-1)^{\delta_{\sigma=\downarrow}} mU \in \mathbb{R}$. We denote this effect as “gap complexification”. A direct consequence of the fact that the correction is purely imaginary is that Eq. (5.68), which expresses the self-consistency condition for m , which we postulated, is proven.

Proof 5.20 — Proof of Eq. (5.68): self-consistency equation for m .

We know from Eq. (5.65) that the real part of the gap is connected to the x projection of the pseudospin expectation value. Instead, from Eq. (5.66) we know that the imaginary part is related to the y projection. It follows

$$\text{Re}\{\tilde{\Delta}_{\mathbf{k}\sigma}\} = (-1)^{\delta_{\sigma=\downarrow}} mU \in \mathbb{R}$$

Then, the x projection of the pseudospin gives a spin-antisymmetric EV,

$$\langle \hat{s}_{\mathbf{k}\uparrow}^x \rangle = - \langle \hat{s}_{\mathbf{k}\downarrow}^x \rangle$$

Phase	HFPs
sAF	$\{m, u^{(s*)}, u^{(d)}, v^{(s*)}, v^{(d)}\}$
aAF	$\{m, u^{(s*)}, u^{(d)}, -iv^{(p_x)}, -iv^{(p_y)}\}$

Table 5.4 | Set of HFP for the two possible realisations of the AF phase, respectively with a symmetric and an antisymmetric imaginary part of the gap.

Then Eq. (5.68) holds identically, because the proof that lead us to Eq. (5.51) for the HM was based on this property.

In this analysis, antiferromagnetism is realised without breaking inversion symmetry. This translates in the requirement that:

$$\tilde{E}_{\mathbf{k}\sigma} \stackrel{!}{=} \tilde{E}_{-\mathbf{k}\sigma}.$$

We know that:

$$\tilde{E}_{\mathbf{k}\sigma} = \sqrt{\tilde{\epsilon}_{\mathbf{k}\sigma}^2 + \text{Re}^2\{\tilde{\Delta}_{\mathbf{k}\sigma}\} + \text{Im}^2\{\tilde{\Delta}_{\mathbf{k}\sigma}\}}.$$

Renormalised bare bands as well as the real part of the gap satisfy said inversion symmetry requirement,

$$\tilde{\epsilon}_{\mathbf{k}\sigma} = \tilde{\epsilon}_{-\mathbf{k}\sigma} \quad \text{Re}\{\tilde{\Delta}_{\mathbf{k}\sigma}\} = \text{Re}\{\tilde{\Delta}_{-\mathbf{k}\sigma}\},$$

hence inversion symmetry is respected only if

$$\text{Im}\{\tilde{\Delta}_{\mathbf{k}\sigma}\} = \pm \text{Im}\{\tilde{\Delta}_{-\mathbf{k}\sigma}\}.$$

Either the imaginary part of the gap is inversion symmetric, or is antisymmetric. Based on this consideration, we are able to discard two of the four self-consistency equations for the harmonics $v_{\sigma}^{(\gamma)}$, listed in Tab. 5.3. If $\text{Im}\{\tilde{\Delta}_{\mathbf{k}\sigma}\}$ is an even function of momentum, $v_{\sigma}^{(p_x)} = v_{\sigma}^{(p_y)} = 0$. If $\text{Im}\{\tilde{\Delta}_{\mathbf{k}\sigma}\}$ is an odd function of momentum, $v_{\sigma}^{(s*)} = v_{\sigma}^{(d)} = 0$. The two options cannot coexist, and separate simulations can be implemented in the two symmetry sectors.

We name the two possible symmetries “sAF” (symmetric antiferromagnet) and “aAF” (antisymmetric antiferromagnet). The set of HFPs then restricts to these AF structures, and is reported for both cases in Tab. 5.4. As we will discuss in Sec. 5.5, the two phases actually coincide.

5.4.2 The entropy density of the AF ground-state

The approximated ground-state in the AF phase is given by a gas of free $\hat{\gamma}^{(\pm)}$ fermions populating the two bands $\pm\tilde{E}_{\mathbf{k}}$ following a Fermi-Dirac distribution. Hence, the total entropy density of the ground state is given by that of a Fermi gas [8]:

$$s^{(\text{AF})} = s_{-}^{(\text{AF})} + s_{+}^{(\text{AF})},$$

where the superscript indicates the band,

$$s_{\pm}^{(\text{AF})} \equiv -\frac{k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \left[\left(1 - f(\pm\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu})\right) \ln \left(1 - f(\pm\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu})\right) + f(\pm\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) \ln f(\pm\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) \right]. \quad (5.105)$$

Using Eq. (4.40), we can give an alternate formula for the entropy density

$$s^{(\text{AF})} = -\frac{k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu})\right) - \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\tilde{E}_{\mathbf{k}} - \tilde{\mu}) f(\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) - \frac{k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \ln \left(1 - f(-\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu})\right) + \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\tilde{E}_{\mathbf{k}} + \tilde{\mu}) f(-\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}). \quad (5.106)$$

The proof is identical to the one provided for the normal phase in Sec. 4.3.2. Note that, as the gap closes the above expression reduces to the entropy density for the normal phase reported in Eq. (4.39). Indeed, we are summing over the MBZ, and the bare bands $\tilde{\epsilon}_{\mathbf{k}}$ change sign when crossing its boundary. Thus, substituting $\tilde{E}_{\mathbf{k}} \rightarrow \tilde{\epsilon}_{\mathbf{k}}$, the first term coincides with half the sum of Eq. (4.39), the second term with the other half.

5.4.3 The free energy density of the AF ground-state

We compute the density of free energy, following the same procedure we used for the normal phase in Sec. 4.3.3. The free energy density is given by

$$f^{(\text{AF})} \equiv \frac{F^{(\text{AF})}}{L_x L_y}, \quad (5.107)$$

being $F^{(\text{AF})}$ the total free energy,

$$F^{(\text{AF})} = E_{\text{HF}}^{(\text{AF})} - \frac{S^{(\text{AF})}}{k_B \beta}, \quad (5.108)$$

and $S^{(\text{AF})} \equiv L_x L_y \times s^{(\text{AF})}$ the total entropy. Here $E_{\text{HF}}^{(\text{AF})}$ is the ground-state energy for the HF hamiltonian. In the AF phase, we compute $f^{(\text{AF})}$ using the following formula:

$$f^{(\text{AF})} = \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \left[\ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) \right) + \ln \left(1 - f(-\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) \right) \right] \\ + U m^2 - V \sum_{\gamma \neq s} \left(|u^{(\gamma)}|^2 + |v^{(\gamma)}|^2 \right) + \tilde{\mu} n. \quad (5.109)$$

Proof 5.21 — Proof of Eq. (5.109): free energy density for the AF phase.

Using Eqs. (5.107), (5.108), we have

$$f^{(\text{AF})} = \frac{E_{\text{HF}}^{(\text{AF})}}{L_x L_y} - \frac{s^{(\text{AF})}}{k_B \beta},$$

with $s^{(\text{AF})}$ the entropy density computed in Eq. (5.106). Then, $f^{(\text{AF})}$ is obtained by summing the following terms:

1. Quasiparticles energy, taking care of spin degeneracy,

$$e_g^{(\text{AF})} = \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\tilde{E}_{\mathbf{k}} - \tilde{\mu}) f(\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) - \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\tilde{E}_{\mathbf{k}} + \tilde{\mu}) f(-\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}).$$

2. Contraction energy,

$$e_c^{(\text{AF})} = \frac{E_c^{(\text{AF})}}{L_x L_y} \\ = U(m^2 - n^2) + V \left[2zn^2 - \sum_{\gamma \neq s} \left(|u^{(\gamma)}|^2 + |v^{(\gamma)}|^2 \right) \right].$$

The total contraction energy $E_c^{(\text{AF})}$ was given in Eq. (5.104). The above result was obtained by using Eqs. (5.31), (5.77) and (5.98).

3. Entropic energy,

$$e_s^{(\text{AF})} = \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) \right) - \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\tilde{E}_{\mathbf{k}} - \tilde{\mu}) f(\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) \\ + \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \ln \left(1 - f(-\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}) \right) + \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\tilde{E}_{\mathbf{k}} + \tilde{\mu}) f(-\tilde{E}_{\mathbf{k}}; \beta, \tilde{\mu}),$$

obtained by multiplying Eq. (5.106) by $-2/\beta k_B$.

4. Chemical potential correction,

$$e_n^{(\text{AF})} = \mu n.$$

Composing everything,

$$f^{(\text{AF})} \equiv e_g^{(\text{AF})} + e_c^{(\text{AF})} + e_s^{(\text{AF})} + e_n^{(\text{AF})},$$

we obtain Eq. (5.109).

5.5 Analysis of the solution

In this section, we present some constraints on the solution for the AF phase, showing that gap complexification is absent. Then we discuss how the renormalisation of bands can induce magnetisation. In the last paragraph, we use these constraints to approximate and interpret the formula for free energy of Eq. (5.109).

Self-consistency constraints. In Tab. 5.4 we collected the two sets HFPs for the two flavours of the AF phase. Each set contains five parameters. The scenario appears quite complicated, but it is possible to show that

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} = 0, \quad (5.110)$$

which implies that the hopping amplitude is reduced by renormalisation and that C_4 is unbroken, identically to the normal phase (see Sec. 4.4). Moreover,

$$\forall \gamma \quad : \quad v^{(\gamma)} = 0. \quad (5.111)$$

The proofs of Eqs. (5.110) and (5.111) are given in App. E. As a consequence, there is no need to distinguish two AF phases,

$$\text{sAF} = \text{aAF}.$$

It follows that the AF gap coincides with the one given by Eq. (5.27),

$$\tilde{\Delta}_{\mathbf{k}} = mU.$$

Therefore, the Bogoliubov bands are given by

$$\tilde{E}_{\mathbf{k}} = \sqrt{\tilde{\epsilon}_{\mathbf{k}}^2 + (Um)^2} \geq Um. \quad (5.112)$$

At any finite magnetisation m , the bands are gapped.

Mott localisation-enhanced magnetisation. The AF phase is described by two HFPs: $u^{(s^*)}$ and m . Moreover, we know that the HFP $u^{(s^*)}$ takes part in the renormalisation of the hopping amplitude,

$$\tilde{t}^{(s^*)} = t - \frac{u^{(s^*)}V}{2},$$

where we used Eq. (5.89). From the constraint of Eq. (5.110), we see that the hopping parameter is reduced in amplitude,

$$t \rightarrow 0 < \tilde{t}^{(s^*)} < t.$$

Bands are effectively shrunk by the non-local attraction, as happened for the normal phase.

From the analysis of antiferromagnetism in the Hubbard model, we know that the local repulsion $\hat{H}_U$ induces magnetisation. We now show that the bands shrinking enhances it. Consider the Bogoliubov bands $\tilde{E}_{\mathbf{k}}$ given in Eq. (5.43),

$$\tilde{E}_{\mathbf{k}} = \sqrt{\tilde{\epsilon}_{\mathbf{k}}^2 + |mU|^2}.$$

If $mU \gg \tilde{t}^{(s^*)}$, these are approximated by

$$\begin{aligned} \tilde{E}_{\mathbf{k}} &= |mU| \sqrt{1 + \left(\frac{\tilde{\epsilon}_{\mathbf{k}}}{mU}\right)^2} \\ &\simeq mU \left[1 + \frac{1}{2} \left| \frac{\tilde{\epsilon}_{\mathbf{k}}}{mU} \right|^2 + \mathcal{O}(r^4) \right], \end{aligned}$$

where we defined the ratio

$$r \equiv \frac{\tilde{t}^{(s^*)}}{mU}, \quad (5.113)$$

being $\tilde{\epsilon}_{\mathbf{k}} = \mathcal{O}(\tilde{t}^{(s^*)})$.

By inserting this result in the self consistency equation for m , Eq. (5.68), we obtain

$$\begin{aligned} m &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{mU}{\tilde{E}_{\mathbf{k}}} \tilde{\text{T}}\text{h}_{\mathbf{k}}^{(-)}[\beta, 0] \\ &\simeq \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \left[1 - \frac{1}{2} \left| \frac{\tilde{\epsilon}_{\mathbf{k}}}{mU} \right|^2 + \mathcal{O}(r^4) \right] \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right). \end{aligned} \quad (5.114)$$

In last line we used Eq. (5.64), valid at half-filling. Let $m \neq 0$. From Eq. (5.112) we know that $\tilde{E}_{\mathbf{k}} \neq 0$. The hyperbolic tangent is, at low temperature, approximated by,

$$\lim_{\beta \rightarrow \infty} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) = 1.$$

Therefore, Eq. (5.114) becomes

$$m \simeq \frac{1}{2} - \underbrace{\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \left| \frac{\tilde{\epsilon}_{\mathbf{k}}}{mU} \right|^2}_{\mathcal{O}(r^2)} + \mathcal{O}(r^4). \quad (5.115)$$

The second order correction can be made small in two ways:

1. By increasing U . From Eq. (5.113), we see that r decreases as U grows. This is the “conventional” way of magnetising the Hubbard lattice, by taking advantage of local repulsion which induces direct exchange (see App. A).
2. By shrinking bands. From Eq. (5.113), we see that as the renormalised bandwidth goes to zero, so does r . This is the “unconventional” way of magnetising the Hubbard lattice, where localisation is not necessarily due to a strong on-site repulsion, rather a low electronic mobility on the lattice.

Hence, as we wanted to prove, a strong Mott flattening can induce commensurate antiferromagnetism.

Approximation of free energy at half-filling. At the end of Sec. 5.2.2, we discussed the fact that the ground-state we find for the AF phase is stable only at half-filling, where $\tilde{\mu} = 0$. We can derive an approximation to free energy in this condition.

$$f^{(\text{AF})} \simeq -\frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \tilde{E}_{\mathbf{k}} + Um^2 - V \sum_{\gamma \neq s} |u^{(\gamma)}|^2. \quad (5.116)$$

This approximation is valid when $\tilde{E}_{\mathbf{k}} \neq 0$ everywhere. This is always realised at finite magnetisation, as is stated in Eq. (5.112). At half filling and low temperature, the system is well approximated by a completely filled electronic band $-\tilde{E}_{\mathbf{k}}$.

From a computational point of view, Eq. (5.116) can be useful when calculating the free energy of the ground-state, since the logarithms of Eq. (5.109) were removed.

Proof 5.22 — Proof of Eq. (5.116).

We start with the identity:

$$\ln\left(1 - \frac{1}{e^x + 1}\right) + \ln\left(1 - \frac{1}{e^{-x} + 1}\right) = \ln\left[\frac{e^x}{(e^x + 1)^2}\right] \\ = x - 2\ln(1 + e^x).$$

At $x \rightarrow \infty$, we have

$$x - 2\ln(1 + e^x) \simeq -x \quad (x \rightarrow \infty).$$

Let now $x = \beta\tilde{E}_{\mathbf{k}}$. At low temperature for a band with no nodes, we can approximate $\forall \mathbf{k} : \beta\tilde{E}_{\mathbf{k}} \rightarrow +\infty$ on the entire BZ. Hence

$$\ln\left(1 - \frac{1}{e^{\beta\tilde{E}_{\mathbf{k}}} + 1}\right) + \ln\left(1 - \frac{1}{e^{-\beta\tilde{E}_{\mathbf{k}}} + 1}\right) \simeq -\beta\tilde{E}_{\mathbf{k}}.$$

Substituting in Eq. (5.109), we obtain Eq. (5.116).

5.6 Discussion of numeric results

In this section, we present numeric results. For the AF phase, the HFPs vector has two components:

$$\mathbf{v} = \begin{bmatrix} m \\ u^{(s^*)} \end{bmatrix} \quad \text{with} \quad \mathbf{v} = \mathbb{R}^2.$$

The RMPs are, as for the normal phase:

$$\tilde{t}_{ij} \equiv \left(t - \frac{u^{(s^*)}V}{2}\right) \varphi_{ij}^{(s^*)}, \\ \tilde{\mu} \equiv \mu + n(2zV - U).$$

We simulate in the ranges:

$$0 \leq U \leq 8, \quad 0 \leq V \leq 8,$$

and limit to $\delta = 0$, since we restrict to the commensurate AF phase which is stable only at half-filling (see Sec. 5.2.2, last paragraph). Tolerances on density and HFPs have been set to

$$\Delta n = 10^{-2}, \quad \Delta \mathbf{v} = \begin{bmatrix} m : 5 \times 10^{-4} \\ u^{(s^*)} : 5 \times 10^{-4} \end{bmatrix}.$$

We will show heatmaps in the (U, V) plane. For each heatmap, we will superimpose six arbitrarily chosen contour lines, in order to highlight how iso valued domains are shaped. These contour lines will be coloured in gray scale.

5.6.1 Mott localisation in the AF phase

In this subsection, we present results for the HFP $u^{(s^*)}$ and the RMP $\tilde{t}^{(s^*)}$.

Numerical results for $u^{(s^*)}$. In Fig. 5.5, we show the converged values of the HFP $u^{(s^*)}$ in the (U, V) plane at $t = 1.0$. We see that $u^{(s^*)}$ is maximised in the bottom-left corner (low- U ,




Figure 5.5 | Numerical results for the HFP $u^{(s*)}$. The gray curves indicate isovalued lines.




Figure 5.6 | Numerical results for the HFP $u^{(s*)}$ as we vary U (left) and V (right), keeping the other variable fixed.

low- V limit). As we see from the colour plot, $u^{(s*)}$ exhibits a sort of triangle-shaped plateau in this region, where it reaches the value

$$u^{(s*)} \lesssim 0.405.$$

The above converged value coincides with that computed at half-filling for the normal phase, in Eq. (4.48). This is expected at $U = 0$, where magnetisation is necessarily null and we reduce to the normal phase. The approximate plateau in the bottom-left corner of Fig. 5.5 extends to $U/t \lesssim 2$, and indicates the region of parameters space where we expect magnetisation to be low enough to retrieve, approximately, the normal phase. We indicated the tick $V = 4.93t$ on the vertical axis, where $U = 0$. We know from Eq. (4.52) that, at half-filling, it is the critical value for the Mott transition. Indeed, we see that the plateau of $u^{(s*)}$ has the point

$$(U/t, V/t) = (0, 4.93)$$

as one of its “corners”.

In Fig. 5.5, we see that isovalued domains are, specifically, straight lines. Moreover, we see that as we move to the top-right corner (high- U , high- V) the isovalued lines become approximately parallel and $u^{(s*)}$ decreases steadily.

The fact that $u^{(s*)}$ is a decreasing function of both U and V is confirmed by looking at Fig. 5.6. In the left panel (Fig. 5.6a), we see that $u^{(s*)}$ decreases as a function of U . For small V (brighter colour), the curves show a plateau for $U \rightarrow 0$, which disappears upon increasing V . An analogous behaviour is observed in the right panel (Fig. 5.6b), by varying V at fixed values of U . We notice that, since $m = 0$ for $U = 0$, the $U = 0$ curve of Fig. 5.6b coincides with the one at $\delta = 0$ for the normal phase, reported in Fig. 4.5b.





Figure 5.7 | Numerical results for the RMP $\tilde{t}^{(s^*)}$. The gray curves indicate isovalues.



a | U -dependency of $t^{(s^*)}$ varying V .

b | V -dependency of $t^{(s^*)}$ varying U .

Figure 5.8 | Numerical results for the RMP $\tilde{t}^{(s^*)}$ as we vary U (left) and V (right), keeping the other variable fixed.

Hopping renormalisation. In Fig. 5.7, we show the the corresponding values for the renormalised hopping amplitude, defined as

$$\tilde{t}^{(s^*)} = t - \frac{u^{(s^*)}V}{2}.$$

As is clear from the definition, isovalues curves of $\tilde{t}^{(s^*)}$ and $u^{(s^*)}$ do not coincide. Specifically, we see that the top-left (low- U , high- V) and bottom-right (high- U , low- V) corners differ significantly (while, for $u^{(s^*)}$, they were similar enough). In the first case, $\tilde{t}^{(s^*)}$ is minimised; in the second, is maximised. The two corners correspond, respectively, to an “strong shrinking” and a “weak shrinking” of bare bands. The region of strong shrinking, namely the brighter region of Fig. 5.7, extends from the critical value $(U/t, V/t) = (0, 4.93)$ to small values of the local repulsion, approximately $U \lesssim 1$. While isovalues lines in the weak shrinking region are approximately parallel to the horizontal axis, moving towards the strong shrinking region they bend progressively.

Apart from the vertical axis, $U = 0$, the converged value of $\tilde{t}^{(s^*)}$ never vanishes (within numerical tolerance). This is better appreciated in Fig. 5.8, where we show data for $\tilde{t}^{(s^*)}$ as we vary U and V . In the right panel (Fig. 5.8b) we see that increasing V while U is fixed makes $\tilde{t}^{(s^*)}$ decrease monotonically, but at any $U \neq 0$ we do not reach in our simulation range a condition of total flattening. In the left panel (Fig. 5.8a), we see that as we increase U , the renormalised hopping amplitude gets larger.

In Fig. 5.9 we show a comparative plot of the converged values for $\tilde{t}^{(s^*)}$ for the normal phase (red surface) and the AF phase (blue surface). Being the normal phase independent of U , the data for the normal phase are a concatenation at various values of the non-local repulsion of the $\delta = 0.0$ data shown in Fig. 4.6. We see that the blue surface reduces to zero only at $U = 0$, whereas the red surface is consistently null at $V \gtrsim 4.93t$. As we mentioned before, in the AF phase we




Figure 5.9 | Comparative plot of the converged values for $\tilde{t}^{(s*)}$ for the normal phase and the AF phase.

do not observe a complete flattening of bare bands. On the contrary, the presence of U reduces significantly the bands shrinking effect.

For the AF phase, we conclude that the non-local attraction $\hat{H}_V$ induces the flattening of bare bands, while the local repulsion $\hat{H}_U$ does the opposite. When we discussed Mott localisation in the context of the normal phase (see Sec. 4.5.1), we argued that bands flattening was induced by V , while U was irrelevant. Here we see that in the AF phase, the local repulsion opposes Mott localisation. This is somewhat the opposite of what we would expect, based on the phenomenology of Mott localisation, which is in general enhanced by the repulsion of two electrons on the same site (see Sec. 1.1.1).

The above observation need to be addressed in the context of antiferromagnetic ordering, where magnetisation m is a macroscopic observable. Recall the discussion of Sec. 5.5: we claimed that the flattening of bare bands could induce magnetisation. Indeed, if by increasing U we are “by force” enhancing the AF gap (even at low magnetisation), by increasing V we are reducing the bare bandwidth. This suggests that we can obtain the same macroscopic magnetisation even if the underlying physics is much different. Indeed, as we observe from Fig. 5.8, it is only by increasing the non-local attraction that the actual hopping amplitude of electrons is reduced. From the above considerations, we observe that either increasing U or V we should expect approximately the same magnetic ordering.

5.6.2 Mott-induced enhancement of magnetisation

In Fig. 5.10 we show the converged results for m as a function of U and V . In the bottom-left corner (low- U , low- V), a bright triangle-shaped region is present in this area, where magnetisation is low. While we cannot assess precisely if this low-magnetisation region has the same shape of the plateau observed in Fig. 5.5, we see that they share the same top-left corner at $(U/t, V/t) = (0, 4.93)$. Indeed, consider a path starting from the low- m region and moving vertically, namely increasing V while keeping U fixed. If U is small enough, the transition of m to approximately zero to its saturation value happens as soon as we reach the strong shrinking region highlighted in Fig. 5.7.

In Fig. 5.10, we see that isomagnetic curves are actually straight lines, and become approximately parallel as we move towards the top-right corner of the plot. This is similar to what we




Figure 5.10 | Numerical results for the HFP m . The gray curves indicate isomagnetic lines.




Figure 5.11 | Numerical results for the HFP m as we vary U (left) and V (right), keeping the other variable fixed.

found for the HFP $u^{(s*)}$. The fact that iso-valued curves are linearly shaped suggests that possibly some kind of critical ratio among the interactions exists, such that the net sum of local repulsion and Mott physics lead to the same macroscopic magnetic ordering. This relation should be, roughly, something like

$$\frac{V}{V_0} = \frac{U_0 - U}{U_0},$$

where V_0 is the intersection of the isomagnetic line with the vertical axis ($U = 0$) and U_0 with the horizontal axis ($V = 0$).

In Fig. 5.11 we show the converged data for m as we vary, alternatively, U (left panel) and V (right panel). As we already commented, magnetisation is induced by both interactions, but the general shape of the curves is slightly different. In Fig. 5.11a we see that m saturates rapidly, but as we lower U all curves go to zero. In Fig. 5.11b, instead, lowering V the curves reach a finite intersection with the vertical axis. The local repulsion is necessary in order to establish antiferromagnetic ordering, while the non-local attraction acts – at fixed U – as a sort of magnetisation boost.

5.6.3 Solution free energy density

In Fig. 5.12 the free energy density is plotted. In the right panel, Fig. 5.12b, we see that free energy is minimised in the bottom-right corner (low- V and high- U). In the top-left corner of Fig. 5.12a free energy grows, eventually reaching zero in the strong shrinking region.

It is then evident how, even if the macroscopic magnetic properties are the same along one isomagnetic line of those plotted in Fig. 5.10, and thus the realised state exhibits an identical value of m , much different is the energetic content of the state itself. Increasing U while keeping V fixed, we amplify the gap. Hence, Bogoliubov bands $\pm \tilde{E}_{\mathbf{k}}$ are shifted away, namely the lower band is




Figure 5.12 | Free energy density of the AF phase.

pushed towards more negative energies and the upper band towards more positive energies. At half-filling and low temperature, the lower band is roughly completely occupied while the upper band is roughly completely empty. Hence, quasiparticles lower their energy by the amplification of the gap. On the other hand, increasing V while keeping U fixed does the opposite. As bands get flattened, their distance lowers and quasiparticles increase their energies (namely, the lower band gets closer to zero). Hence, even if we observe isomagnetic domains, this does not mean that the content in energy of the HF ground state is identical.

The AF phase is always energetically favoured at half-filling, with respect to the normal phase. In Fig. 5.13 we plot the energy difference

$$f^{(N)} - f^{(AF)}.$$

The data for the normal phase, similarly to Fig. 5.9, are a concatenation of those of Fig. 4.8 at $\delta = 0.0$. We see that, on the entire $(U/t, V/t)$ region of our simulations, the free energy difference is positive. Hence the AF phase is consistently more stable with respect to the normal phase, except for the segment at $U = 0$, where the two coincide. Looking at Fig. 5.13, we see that the energy difference is enhanced as we increase V at fixed U .

At low temperature and half filling, it is not worth plotting entropy density. Half-filled commensurate antiferromagnets are insulators, with two gapped Bogoliubov bands. At low temperature, electronic distribution fills completely the lower band and empties the upper one. Because of this, insulating gapped systems inherently are low entropic and our results are compatible with zero in the entire region of parametric space we explored.

5.6.4 Fixed-point interpretation of HF solution

We end this chapter, by presenting a final remark regarding the numerical structure of the algorithm. As we anticipated in Sec. 2.3.2, our convergence scheme can be seen as an effective search for the singular fixed point of a nonlinear map. A physical solution is represented as a HFP vector $\mathbf{v}$ such that the algorithm is a local identity,

$$A(\mathbf{v}) = \mathbf{v}.$$

To predict, for a given initialiser $\mathbf{v}_0$, the path of the algorithm in HFPs space is in general non-trivial due to the intrinsic nonlinearity of the problem. An example of such nonlinearity is given in the derivation of Secs. E.2.2, E.2.3.

In Fig. 5.14 we sketch this concept in a practical situation. In figure are plotted various algorithmic paths through AF HFPs space, with operational setup at

$$U = 3, \quad V = 6, \quad L = 128, \quad \beta = 100, \quad \delta = 0.$$

At this point, the self-consistent HF “fixed-point” solution is given by

$$m \approx 0.449, \quad u^{(s^*)} \approx 0.212,$$




Figure 5.13 | Difference of free energies for the normal phase minus the AF phase.




Figure 5.14 | Quiver plot of algorithmic evolution in HFPs space at fixed model parametrization. The black dot $\mathbf{v}$ represents the nonlinear map fixed point, while the colored paths the sequences of HFPs coordinates simulated via iterative HF procedure, starting from different HFPs initialisers (colored dots).

indicated in panels as the black dot. The three panels show different map evolutions varying the mixing parameter within three choices, $g \in \{0.25, 0.1, 0.05\}$. We chose various initialisers in the set:

$$\mathbf{v}_0 = \begin{bmatrix} m_0 \\ u_0^{(s^*)} \end{bmatrix} : m_0 \in \{0.2, 0.3, \dots, 0.6\} \quad \text{and} \quad u_0^{(s^*)} \in \{0.0, 0.1, \dots, 0.4\}.$$

In figures, different colours indicate different evolutions, while the tip size indicates the magnitude of the algorithmic step in passing from one iteration to the following.

All paths are able to converge to $\mathbf{v}$, and the quantity of steps necessary to reach the fixed point, a sort of attractor, increases as we lower g . To select the correct g turns out to be the real algorithm fine-tuning knob: as we saw for the normal phase, some model parametrisation could lead to unavoidable algorithmic instabilities. The problem is that we do not know beforehand where the instabilities will occur, and how. The correct g is the compromise between keeping low the number of steps (larger mixing) and moving slow enough to not diverge when encountering unstable points (smaller mixing). By recursively fine-tuning g repeating non-converged simulation with more care, we are essentially making the nonlinear map “softer” and curvier, as is in the right panel of Fig. 5.14. It’s not always resource convenient to do so, so for most simulations we stick to a rougher map (like the left panel one) by setting $g = 0.5$.

Chapter 6

Superconducting instability

In this chapter, we study the translationally invariant superconducting (SC) phase. We derive the self-consistent HF hamiltonian in both the singlet and triplet channels, and simulate only in the singlet channel. We will observe various effects: bare bands shrinking, insurgence of mixed-symmetry superconductivity and nematicity.

This chapter is organised as follows. In Sec. 6.1 we present the translationally invariant superconducting phase and its symmetries. In Secs. 6.2, 6.3, 6.4 we derive the approximated HF hamiltonian for the SC phase, starting from the conventional Hubbard model and then adding the non-local attraction $\hat{H}_V$. In Sec. 6.5 we derive a constraint on the HFPs by analysing their self-consistency equations. In Sec. 6.6 we present the result of numeric simulations.

Disclaimer. Sec. 6.3, where we carry out all technical computations to derive the HF approximation to the EHM, is *very* detailed and cumbersome. It can be skipped, jumping directly to Sec. 6.4, where we summarise all results.

6.1 The SC phase and its symmetries

In Sec. 3.2 we anticipated that the translationally invariant SC phase is described by the two SSBs

$$\begin{aligned} \text{U}^z(1) &\xrightarrow{\text{SSB}} 0, \\ \text{C}_4 &\xrightarrow{\text{SSB}} \text{C}_2. \end{aligned}$$

Translational invariance is unbroken. Therefore, Eqs. (4.1) and (4.2), derived in the context of the normal phase, hold equivalently,

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'} \rangle = \delta_{\mathbf{k}=\mathbf{k}'} \delta_{\sigma=\sigma'} \langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \rangle, \quad (6.1)$$

$$\langle \hat{n}_{i\sigma} \rangle = n. \quad (6.2)$$

We define the operators

$$\hat{n}_{\mathbf{k}\sigma} \equiv \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}, \quad (6.3)$$

$$\hat{\phi}_{\mathbf{k}} \equiv \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow}, \quad (6.4)$$

and their expectation values (EVs),

$$u_{\mathbf{k}\sigma} \equiv \langle \hat{n}_{\mathbf{k}\sigma} \rangle, \quad (6.5)$$

$$w_{\mathbf{k}} \equiv \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle. \quad (6.6)$$

Using the functions of Tab. 3.5, we define the harmonics

$$u_{\sigma}^{(\gamma)} \equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} u_{\mathbf{k}\sigma}, \quad (6.7)$$

$$w^{(\gamma)} \equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} w_{\mathbf{k}}. \quad (6.8)$$

As was for the AF phase, these harmonics are HFPs in the SC phase and must be determined self-consistently. Eqs. (6.7) and (6.8) are their self-consistency equations.

6.2 The SC phase in the conventional HM

The conventional HM is given by $\hat{H}_t + \hat{H}_U$, thus its unique two-body interaction is the local repulsion. In this section, we study superconductivity in the conventional HM, and prove that it is impossible at HF level.

6.2.1 Derivation of the HF hamiltonian

Following Tab. 3.2 and Eqs. (2.21), (2.22), we know the local repulsion decomposes in Hartree and Cooper channels,

$$\begin{aligned} \hat{H}_U &= U \sum_{\mathbf{r} \in \mathcal{L}} \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}\downarrow} \\ &\stackrel{\text{HF}}{\simeq} U \sum_{\mathbf{r} \in \mathcal{L}} \left(\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \rangle + \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} - \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}\downarrow}^\dagger \hat{c}_{\mathbf{r}\downarrow} \rangle \right. \\ &\quad \left. + \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \langle \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle + \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \rangle \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} - \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle \right), \end{aligned} \quad \begin{array}{l} \text{(Hartree)} \\ \text{(Cooper)} \end{array}$$

while the Fock channel is identically zero. We deal with each channel separately.

Local Hartree terms. For the Hartree channel, being the density uniform across sites, we can re-use the entirety of Sec. 4.2.1 for the normal phase. Substituting Eq. (6.2), we get the approximation of Eq. (4.10),

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} nU \times \hat{N} - E_{\text{H},U}^{(\text{SC})} + (\text{Cooper channel}), \quad (6.9)$$

thus chemical potential renormalisation,

$$\tilde{\mu} = \mu - nU \quad (6.10)$$

is present also in the SC phase. Note that $\tilde{\mu}$ will be renormalised again, later, and we will redefine it accordingly. The contraction energy is the same of Eq. (4.14)

$$E_{\text{H},U}^{(\text{SC})} = L_x L_y \times n^2 U. \quad (6.11)$$

Local Cooper terms. We move to reciprocal space, through the transformation of Eq. (3.39). Consider the Cooper channel of Eq. (2.21). In reciprocal space we get:

$$\begin{aligned} \hat{H}_U \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + \frac{U}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \left(\langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow} \right. \\ \left. + \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \langle \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow} \rangle - \langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow} \rangle \right). \end{aligned} \quad (6.12)$$

We are not breaking translational invariance, therefore,

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'}^\dagger \rangle = \langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'}^\dagger \rangle \delta_{\mathbf{k}=-\mathbf{k}'}. \quad (6.13)$$

Proof 6.1 — Proof of Eq. (6.13).

Consider the EV

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'}^\dagger \rangle,$$

for two momenta $\mathbf{k}, \mathbf{k}'$. Moving to real space via an $\mathcal{F}$ -transform, we get

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'}^\dagger \rangle \stackrel{\mathcal{F}}{=} \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} \sum_{\mathbf{r}' \in \mathcal{L}} e^{-i(\mathbf{k} \cdot \mathbf{r} + \mathbf{k}' \cdot \mathbf{r}')} \langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}'\sigma'}^\dagger \rangle,$$

where we used the convention of Eq. (3.34). Let now $\mathbf{r} \equiv \mathbf{r}$ and $\Delta \mathbf{r} \equiv \mathbf{r} - \mathbf{r}'$. Translational invariance implies that the function

$$g(\Delta \mathbf{r}, \sigma, \sigma') = \langle \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}+\Delta \mathbf{r}\sigma'}^\dagger \rangle$$

is independent of $\mathbf{r}$. Thus, we have:

$$\begin{aligned} \langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma'}^\dagger \rangle &\stackrel{\mathcal{F}}{=} \sum_{\Delta \mathbf{r} \in \mathcal{L}} e^{-i\mathbf{k}' \cdot \Delta \mathbf{r}} g(\Delta \mathbf{r}, \sigma, \sigma') \frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{L}} e^{-i(\mathbf{k} + \mathbf{k}') \cdot \mathbf{r}} \\ &= \sum_{\Delta \mathbf{r} \in \mathcal{L}} e^{-i\mathbf{k}' \cdot \Delta \mathbf{r}} g(\Delta \mathbf{r}, \sigma, \sigma') \delta_{\mathbf{k}=\mathbf{k}'}, \end{aligned}$$

where we used the identity of Eq. (3.35). Then, the above EV vanishes for $\mathbf{k} \neq \mathbf{k}'$. This proves Eq. (6.13).

Using Eq. (6.13) in Eq. (6.12), only $\mathbf{K} = \mathbf{0}$ gives a non-zero contribution, and we obtain

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) - \sum_{\mathbf{k} \in \text{BZ}} \left(\Delta^{(s)} \hat{\phi}_{\mathbf{k}} + \Delta^{(s)*} \hat{\phi}_{\mathbf{k}}^\dagger \right) - E_{C,U}^{(\text{SC})}, \quad (6.14)$$

where we defined

$$\Delta^{(s)} \equiv -U w^{(s)}, \quad (6.15)$$

and the contraction energy is given by

$$E_{C,U}^{(\text{SC})} = -\frac{U}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \langle \hat{\phi}_{\mathbf{k}'} \rangle. \quad (6.16)$$

Proof 6.2 — Proof of Eq. (6.14): HF approximation of $\hat{H}_U$ in Cooper channel.

Using Eqs. (6.4), (6.12) and (6.13), we obtain

$$\hat{H}_U \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + \frac{U}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} \left(\langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \hat{\phi}_{\mathbf{k}'} + \hat{\phi}_{\mathbf{k}}^\dagger \langle \hat{\phi}_{\mathbf{k}'} \rangle \right) - E_{C,U}^{(\text{SC})}, \quad (6.17)$$

From Eq. (6.6), we have

$$\begin{aligned} \hat{H}_U \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + \left(\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} w_{\mathbf{k}} \right) \times U \sum_{\mathbf{k}' \in \text{BZ}} \hat{\phi}_{\mathbf{k}'} \\ + \left(\frac{1}{L_x L_y} \sum_{\mathbf{k}' \in \text{BZ}} w_{\mathbf{k}'}^* \right) \times U \sum_{\mathbf{k} \in \text{BZ}} \hat{\phi}_{\mathbf{k}}^\dagger. \end{aligned}$$

Therefore, from Eq. (6.8), we see that the parenthesised terms coincide, respectively, with $w^{(s)}$ and $w^{(s)*}$. Applying Eq. (6.15), we obtain Eq. (6.14).

The contraction energy of Eq. (6.16) reduces to

$$E_{C,U}^{(\text{SC})} = L_x L_y \times U |w^{(s)}|^2. \quad (6.18)$$

Proof 6.3 — Proof of Eq. (6.18): local Cooper contraction energy.

Consider Eq. (6.16). Substituting Eqs. (6.6), (6.8) we get:

$$E_{C,U}^{(\text{SC})} = L_x L_y \times U \left(\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} w_{\mathbf{k}}^* \right) \left(\frac{1}{L_x L_y} \sum_{\mathbf{k}' \in \text{BZ}} w_{\mathbf{k}'} \right) \quad (6.19)$$

$$= L_x L_y \times U |w^{(s)}|^2, \quad (6.20)$$

which proves Eq. (6.18).

In general, $\Delta^{(s)}$ as defined in Eq. (6.15) is a complex number. However, we can always choose $\Delta^{(s)}$ as purely real. This is due to the fact that the hamiltonian is gauge-invariant. Indeed, if we set in Eq. (3.4)

$$\eta \equiv -\frac{\arg w^{(s)}}{2}, \quad (6.21)$$

we have $\Delta^{(s)} \in \mathbb{R}$.

Proof 6.4 — Choosing the gauge of Eq. (6.21), we obtain a real gap.

Inserting Eq. (6.21) in Eq. (3.4), we obtain

$$\hat{c}_{\mathbf{k}\sigma}^\dagger \rightarrow e^{-i \arg w^{(s)}/2} \hat{c}_{\mathbf{k}\sigma}^\dagger.$$

Therefore,

$$\hat{\phi}_{\mathbf{k}}^\dagger \rightarrow \hat{\phi}_{\mathbf{k}}^\dagger e^{-i \arg w^{(s)}} \implies w_{\mathbf{k}} \rightarrow w_{\mathbf{k}} e^{-i \arg w^{(s)}},$$

where we used Eq. (6.6). It follows, from Eq. (6.8),

$$w^{(s)} \rightarrow w^{(s)} e^{-i \arg w^{(s)}}.$$

For any complex number z , we have $z e^{-i \arg z} \in \mathbb{R}$. Then, $w^{(s)}$ is purely real under this gauge transform, and so is $\Delta^{(s)}$.

6.2.2 Diagonalisation of the HF hamiltonian

We proceed here as we did for the AF phase in Sec. 5.2.2. We start by giving a matrix representation of the approximated hamiltonian, then map it to a pseudospin problem and finally specify how to diagonalise it to retrieve its ground state.

Matrix representation. Let $\Delta_{\mathbf{k}} = \Delta^{(s)}$. The HM gran-canonical hamiltonian is given by $\hat{H}_t + \hat{H}_U - \mu \hat{N}$. Its HF approximation for the SC phase can be written as

$$\hat{H}_t + \hat{H}_U - \mu \hat{N} \stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \hat{\Phi}_{\mathbf{k}}^\dagger h_{\mathbf{k}} \hat{\Phi}_{\mathbf{k}} + E_a^{(\text{SC})} - \left(E_{\text{H},U}^{(\text{SC})} + E_{\text{C},U}^{(\text{SC})} \right), \quad (6.22)$$

where we defined the 2×2 matrix:

$$h_{\mathbf{k}} \equiv \begin{bmatrix} \xi_{\mathbf{k}} & -\Delta_{\mathbf{k}}^* \\ -\Delta_{\mathbf{k}} & -\xi_{\mathbf{k}} \end{bmatrix} \quad \text{with} \quad \xi_{\mathbf{k}} \equiv \epsilon_{\mathbf{k}} - \tilde{\mu}. \quad (6.23)$$

In Eq. (6.22), we defined the Nambu spinors

$$\hat{\Phi}_{\mathbf{k}} \equiv \begin{bmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{bmatrix}, \quad (6.24)$$

and the “anticommutation energy” $E_a^{(\text{SC})}$,

$$E_a^{(\text{SC})} \equiv \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}}. \quad (6.25)$$

Proof 6.5 — Proof of Eq. (6.22).

Summing Eq. (3.37) and Eq. (6.14), we obtain

$$\begin{aligned} \hat{H}_t + \hat{H}_U &\stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \epsilon_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} \\ &\quad - \sum_{\mathbf{k} \in \text{BZ}} \left(\Delta^{(s)} \hat{\phi}_{\mathbf{k}} + \Delta^{(s)*} \hat{\phi}_{\mathbf{k}}^{\dagger} \right) + nU \times \hat{N} - \left(E_{\text{H},U}^{(\text{SC})} + E_{\text{C},U}^{(\text{SC})} \right). \end{aligned} \quad (6.26)$$

Inversion symmetry is not broken, then $\xi_{\mathbf{k}} = \xi_{-\mathbf{k}}$. Using Fermi anticommutation rules, the kinetic operator can be written as:

$$\begin{aligned} \hat{H}_t - \mu \hat{N} &= \sum_{\sigma} \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}\sigma} \hat{n}_{\mathbf{k}\sigma} \\ &= \sum_{\mathbf{k} \in \text{BZ}} (\xi_{\mathbf{k}} \hat{n}_{\mathbf{k}\uparrow} + \xi_{-\mathbf{k}} \hat{n}_{-\mathbf{k}\downarrow}) \\ &= \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}} \left(\hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} + 1 \right) \\ &= \sum_{\mathbf{k} \in \text{BZ}} \xi_{\mathbf{k}} \left(\hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \right) + E_{\text{a}}^{(\text{SC})}. \end{aligned}$$

In last line we used Eq. (6.25). Inserting in Eq. (6.26), we obtain

$$\begin{aligned} \hat{H}_t + \hat{H}_U - \mu \hat{N} &\stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{BZ}} \epsilon_{\mathbf{k}} \left(\hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\downarrow}^{\dagger} \right) + E_{\text{a}}^{(\text{SC})} \\ &\quad - \sum_{\mathbf{k} \in \text{BZ}} \left(\Delta_{\mathbf{k}} \hat{\phi}_{\mathbf{k}} + \Delta_{\mathbf{k}} \hat{\phi}_{\mathbf{k}}^{\dagger} \right) - \tilde{\mu} \hat{N} - \left(E_{\text{H},U}^{(\text{SC})} + E_{\text{C},U}^{(\text{SC})} \right), \end{aligned} \quad (6.27)$$

where $\tilde{\mu} = \mu - nU$ and $\Delta^{(s)} = \Delta_{\mathbf{k}}$. The energies $E_{\text{H},U}^{(\text{SC})}$, $E_{\text{C},U}^{(\text{SC})}$ are given respectively in Eqs. (6.11) and (6.18). Using the Nambu spinors defined in Eq. (6.24) and the 2×2 matrix $h_{\mathbf{k}}$ of Eq. (6.22), the HM hamiltonian is reduced to the matrix form of Eq. (6.22).

Pseudospin representation. Let now the pseudospin operator $\hat{\mathbf{s}}_{\mathbf{k}}$ and the pseudofield $\mathbf{b}_{\mathbf{k}}$

$$\hat{s}_{\mathbf{k}}^{\alpha} \equiv \hat{\Phi}_{\mathbf{k}}^{\dagger} \tau^{\alpha} \hat{\Phi}_{\mathbf{k}} \quad \text{with} \quad \hat{\mathbf{s}}_{\mathbf{k}} = \begin{bmatrix} \hat{s}_{\mathbf{k}}^x \\ \hat{s}_{\mathbf{k}}^y \\ \hat{s}_{\mathbf{k}}^z \end{bmatrix}, \quad \mathbf{b}_{\mathbf{k}} \equiv \begin{bmatrix} -\text{Re}\{\Delta_{\mathbf{k}}\} \\ -\text{Im}\{\Delta_{\mathbf{k}}\} \\ \xi_{\mathbf{k}} \end{bmatrix}.$$

As was for the AF phase, these operators $\hat{s}_{\mathbf{k}}^{\alpha}$ satisfy a $\mathfrak{su}(2)$ algebra up to a constant, irrelevant in the context of this discussion. They satisfy the identities:

$$\hat{s}_{\mathbf{k}}^x = \hat{\phi}_{\mathbf{k}} + \hat{\phi}_{\mathbf{k}}^{\dagger}, \quad (6.28)$$

$$\hat{s}_{\mathbf{k}}^y = i \left(\hat{\phi}_{\mathbf{k}} - \hat{\phi}_{\mathbf{k}}^{\dagger} \right), \quad (6.29)$$

$$\hat{s}_{\mathbf{k}}^z = \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{\mathbf{k}\downarrow} - 1. \quad (6.30)$$

Proceeding precisely as for the AF phase, in pseudospin representation the HF hamiltonian of Eq. (6.22) can be reduced to:

$$\hat{H}_t + \hat{H}_U - \mu \hat{N} \stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{BZ}} \mathbf{b}_{\mathbf{k}} \cdot \hat{\mathbf{s}}_{\mathbf{k}} + E_{\text{a}}^{(\text{SC})} - \left(E_{\text{H},U}^{(\text{SC})} + E_{\text{C},U}^{(\text{SC})} \right). \quad (6.31)$$

Diagonalisation. Let us define the angles $\theta_{\mathbf{k}}$, $\zeta_{\mathbf{k}}$ through the relations

$$\cos 2\theta_{\mathbf{k}} = \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}}, \quad \sin 2\theta_{\mathbf{k}} \cos 2\zeta_{\mathbf{k}} = \frac{\text{Re}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}}, \quad \sin 2\theta_{\mathbf{k}} \sin 2\zeta_{\mathbf{k}} = \frac{\text{Im}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}}.$$

For the SC phase, diagonalisation of the HF hamiltonian follows the same procedure as was for the AF phase. Indeed, the diagonalising matrix is given by

$$W_{\mathbf{k}} \equiv e^{-i\theta_{\mathbf{k}}\tau^y} e^{-i\zeta_{\mathbf{k}}\tau^z}, \quad (6.32)$$

and the graphic procedure for the diagonalisation is the same of Fig. 5.2, with $\epsilon \rightarrow \xi$. We get:

$$d_{\mathbf{k}} = W_{\mathbf{k}} h_{\mathbf{k}} W_{\mathbf{k}}^\dagger \quad \text{with} \quad d_{\mathbf{k}} \equiv \begin{bmatrix} E_{\mathbf{k}} & \\ & -E_{\mathbf{k}} \end{bmatrix}, \quad (6.33)$$

where we defined the Bogoliubov bands $E_{\mathbf{k}}$,

$$E_{\mathbf{k}} \equiv \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}. \quad (6.34)$$

In the tilted frame we have the expression for the Nambu spinors:

$$\hat{\Gamma}_{\mathbf{k}} = W_{\mathbf{k}} \hat{\Psi}_{\mathbf{k}} \quad \text{with} \quad \hat{\Gamma}_{\mathbf{k}} \equiv \begin{bmatrix} \hat{\gamma}_{\mathbf{k}}^{(+)} \\ \hat{\gamma}_{\mathbf{k}}^{(-)} \end{bmatrix}. \quad (6.35)$$

Applying the rotation of Eq. (6.33) to Eq. (6.31), we obtain the free Fermi gas representation of the HF hamiltonian,

$$\hat{H}_t + \hat{H}_U - \tilde{\mu}\hat{N} \stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} E_{\mathbf{k}} \left(\hat{\gamma}_{\mathbf{k}}^{(+)\dagger} \hat{\gamma}_{\mathbf{k}}^{(+)} - \hat{\gamma}_{\mathbf{k}}^{(-)\dagger} \hat{\gamma}_{\mathbf{k}}^{(-)} \right) + (\dots),$$

with $(\dots) = E_{\text{a}}^{(\text{SC})} - (E_{\text{H},U}^{(\text{SC})} + E_{\text{C},U}^{(\text{SC})})$ the various energy shifts.

Pseudospin EVs. The EVs of pseudospin operators in the various directions are given by:

$$\begin{aligned} \langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^x \hat{\Psi}_{\mathbf{k}} \rangle &= -\sin 2\theta_{\mathbf{k}} \cos 2\zeta_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle, \\ \langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^y \hat{\Psi}_{\mathbf{k}} \rangle &= -\sin 2\theta_{\mathbf{k}} \sin 2\zeta_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle, \\ \langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^z \hat{\Psi}_{\mathbf{k}} \rangle &= \cos 2\theta_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle. \end{aligned}$$

Let us define the thermal factors:

$$\text{Th}_{\mathbf{k}}^{(\pm)}[\beta] \equiv f(-E_{\mathbf{k}}; \beta, 0) \pm f(E_{\mathbf{k}}; \beta, 0). \quad (6.36)$$

The only difference with Eq. (6.36) is in the fact that the chemical potential is zero in the Fermi-Dirac distributions, because it is already inside the definition of $E_{\mathbf{k}}$ given in Eq. (6.34). The thermal factors satisfy, at any density,

$$\text{Th}_{\mathbf{k}}^{(+)}[\beta] = 1, \quad (6.37)$$

$$\text{Th}_{\mathbf{k}}^{(-)}[\beta] = \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right). \quad (6.38)$$

The proofs are identical to the ones provided for Eqs. (5.63), (5.63).

In the tilted frame, the averaged z projection of pseudospin is:

$$\begin{aligned} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle &= f(E_{\mathbf{k}}; \beta, 0) - f(-E_{\mathbf{k}}; \beta, 0) \\ &= -\tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right). \end{aligned} \quad (6.39)$$

Bringing everything together, we have the EVs for the pseudospin operator:

$$\langle \hat{s}_{\mathbf{k}}^x \rangle = \frac{\text{Re}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right), \quad (6.40)$$

$$\langle \hat{s}_{\mathbf{k}}^y \rangle = \frac{\text{Im}\{\Delta_{\mathbf{k}}\}}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right), \quad (6.41)$$

$$\langle \hat{s}_{\mathbf{k}}^z \rangle = -\frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right). \quad (6.42)$$

***s*-wave gap self-consistency equation.** The only HFP that we obtained for the SC phase is $w^{(s)}$. It obeys the self-consistency equation

$$w^{(s)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\Delta_{\mathbf{k}}}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right). \quad (6.43)$$

which is the “operational” version of Eq. (6.8), for $\gamma = s$.

Proof 6.6 — Proof of Eq. (6.43): self-consistency equation for $w^{(s)}$.

Let $\tau^{\pm} \equiv (\tau^x \pm i\tau^y)/2$. From Eq. (6.8), we obtain

$$\begin{aligned} w^{(s)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\phi}_{\mathbf{k}}^{\dagger} \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{\Phi}_{\mathbf{k}}^{\dagger} \tau^+ \hat{\Phi}_{\mathbf{k}} \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\langle \hat{s}_{\mathbf{k}}^x \rangle + i \langle \hat{s}_{\mathbf{k}}^y \rangle}{2} \\ &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\Delta_{\mathbf{k}}}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right). \end{aligned}$$

The last line coincides with Eq. (6.43).

Impossibility of *s*-wave superconductivity. At HF level, the HM cannot host superconductivity, i.e.

$$w^{(s)} = 0, \quad (6.44)$$

therefore no superconducting gap can develop at HF level. This does not hold for $U < 0$: an attractive model can host *s*-wave superconductivity.

Proof 6.7 — Proof of Eq. (6.44): impossibility of *s*-wave superconductivity in the conventional HM.

Let

$$U_c[\beta] \equiv \left[\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{1}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right) \right]^{-1}. \quad (6.45)$$

By construction $U_c[\beta] > 0$. Using Eq. (6.45) Eq. (6.43) can be rewritten as:

$$w^{(s)} = -\frac{U}{U_c[\beta]} w^{(s)}.$$

The unique possible self-consistent solution for the equation above is $w^{(s)} = 0$. This implication terminates the argument and proves Eq. (6.44): superconductivity cannot establish self-consistently at HF level in the repulsive HM.

Density. Density n can be computed as

$$n = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right) \right]. \quad (6.46)$$

Proof 6.8 — Proof of Eq. (6.46).

In order to calculate density, we can use the z projection of the pseudospin operators. From Eq. (6.30) we know

$$\hat{s}_{\mathbf{k}}^z + 1 = \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{\mathbf{k}\downarrow}.$$

Then, density can be compute as:

$$\begin{aligned} n &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \langle \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{-\mathbf{k}\downarrow} \rangle \\ &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\langle \hat{s}_{\mathbf{k}}^z \rangle + 1) \\ &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right) \right], \end{aligned}$$

which proves Eq. (6.46).

6.3 Generalisation to the EHM

In this section we add to the hamiltonian the non-local attraction $\hat{H}_V$. As we did for the AF phase, we look for a solution with an identical mathematical structure as the one here described for the conventional HM. We define the renormalised quantities

$$\xi_{\mathbf{k}} \rightarrow \tilde{\xi}_{\mathbf{k}}, \quad \Delta_{\mathbf{k}} \rightarrow \tilde{\Delta}_{\mathbf{k}}, \quad \mu \rightarrow \tilde{\mu}.$$

The chemical potential was already renormalised once in the HM derivation in Sec. 6.3.1. Renormalisation of Bogoliubov bands follows that of bare bands and gap function,

$$\tilde{E}_{\mathbf{k}} \equiv \sqrt{\tilde{\xi}_{\mathbf{k}}^2 + |\tilde{\Delta}_{\mathbf{k}}|^2}.$$

Angles are changed too. We define operatively the angles $\tilde{\theta}_{\mathbf{k}}$ and $\tilde{\zeta}_{\mathbf{k}}$ as

$$\sin 2\tilde{\theta}_{\mathbf{k}} = \frac{|\tilde{\Delta}_{\mathbf{k}}|}{\tilde{E}_{\mathbf{k}}}, \quad \cos 2\tilde{\theta}_{\mathbf{k}} = \frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}}, \quad (6.47)$$

and

$$\sin 2\tilde{\zeta}_{\mathbf{k}} = \frac{\text{Re}\{\tilde{\Delta}_{\mathbf{k}}\}}{|\tilde{\Delta}_{\mathbf{k}}|}, \quad \cos 2\tilde{\zeta}_{\mathbf{k}} = \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{|\tilde{\Delta}_{\mathbf{k}}|}. \quad (6.48)$$

Pseudospin EVs. Using Eqs. (6.47) and (6.48), we compute the pseudospin EVs

$$\langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^x \hat{\Phi}_{\mathbf{k}} \rangle = -\sin 2\tilde{\theta}_{\mathbf{k}} \sin 2\tilde{\zeta}_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle, \quad (6.49)$$

$$\langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^y \hat{\Phi}_{\mathbf{k}} \rangle = -\sin 2\tilde{\theta}_{\mathbf{k}} \cos 2\tilde{\zeta}_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle, \quad (6.50)$$

$$\langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^z \hat{\Phi}_{\mathbf{k}} \rangle = \cos 2\tilde{\theta}_{\mathbf{k}} \langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle. \quad (6.51)$$

Renormalisation of the thermal factors and the z pseudospin EV in the tilted frame gives, similarly to Eqs. (6.36), (6.39),

$$\tilde{\text{Th}}_{\mathbf{k}}^{(\pm)}[\beta, \tilde{\mu}] \equiv f(-\tilde{E}_{\mathbf{k}}; \beta, 0) \pm f(\tilde{E}_{\mathbf{k}}; \beta, 0). \quad (6.52)$$

$$\langle \hat{\Gamma}_{\mathbf{k}}^\dagger \tau^z \hat{\Gamma}_{\mathbf{k}} \rangle = f(\tilde{E}_{\mathbf{k}}; \beta, 0) - f(-\tilde{E}_{\mathbf{k}}; \beta, 0), \quad (6.53)$$

Operator	Sector	Channel	Net effect
$\hat{H}_U$		Hartree	μ shift by $-nU$
		Cooper	Isotropic pairing in singlet channel
$\hat{H}_V$	o.s.	Hartree	μ shift by nzV
		Cooper	Anisotropic pairing in singlet and triplet channels
	s.s.	Hartree	μ shift by nzV
		Fock	Band renormalisation and resymmetrisation

Table 6.1 | Wick's contractions for each interaction and channel in the HF framework, and their net effect in the SC phase.

Eqs. (6.37), (6.38) hold identically. Hence, pseudospin expectation values are:

$$\langle \hat{s}_{\mathbf{k}}^x \rangle = \frac{\text{Re}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right), \quad (6.54)$$

$$\langle \hat{s}_{\mathbf{k}}^y \rangle = \frac{\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right), \quad (6.55)$$

$$\langle \hat{s}_{\mathbf{k}}^z \rangle = -\frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right). \quad (6.56)$$

These equations are an extension of Eqs. (6.54), (6.55), (6.56) for the EHM.

$w^{(s)}$ and density self-consistency equations. We can rewrite the equations for $w^{(s)}$ and n accordingly,

$$w^{(s)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right), \quad (6.57)$$

$$n = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[1 - \cos 2\tilde{\theta}_{\mathbf{k}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right], \quad (6.58)$$

where we used Eqs. (6.43), (6.46).

Contributions of each channel. Recall Tab. 3.3, collecting the selection rules for the non-local attraction $\hat{H}_V$. From Eq. (2.22), for the SC phase we need to account for the following contractions:

$$\begin{aligned} \hat{H}_V \text{ (o.s. sector)} & \quad \text{Hartree and Cooper contractions,} \\ \hat{H}_V \text{ (s.s. sector)} & \quad \text{Hartree and Fock contractions.} \end{aligned}$$

Note that Cooper contractions in the non-local s.s. sector are neglected, because of the selection rule discussed at the end of Sec. 3.2.2. We specify the role of each contraction channel in Tab. 6.1, where also those from $\hat{H}_U$ are included.

6.3.1 Effects of non-local Hartree terms: chemical potential renormalisation

We start by analysing the non-local Hartree channel,

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} -V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} & \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle \right) \\ & + (\text{Fock channel}) + (\text{Cooper channel}), \end{aligned}$$

where we used Eq. (2.22). We have already carried out this computation in Sec. 4.2.1, which lead to the renormalisation of the chemical potential. The non-local attraction $\hat{H}_V$ is reduced in the Hartree channel to

$$\hat{H}_V \stackrel{\text{HF}}{\simeq} -2nzV \times \hat{N} - E_{\text{H},V}^{(\text{SC})} + (\text{Fock channel}) + (\text{Cooper channel}), \quad (6.59)$$

This, together Eq. (6.9), gives the chemical potential full renormalisation, identical to Eq. (4.8),

$$\tilde{\mu} \equiv \mu + n(2zV - U). \quad (6.60)$$

The contraction energy for the non-local sector was derived in Eq. (4.15),

$$E_{\text{H},V}^{(\text{SC})} = -L_x L_y \times 2zn^2V. \quad (6.61)$$

6.3.2 Effects of Fock terms: bare bands renormalisation

From Eq. (2.22), we have

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} & (\text{Hartree channel}) \\ & V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rangle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} \rangle \right) + (\text{Cooper channel}). \end{aligned}$$

Repeating the same derivation of Sec. 4.2.2, with the averages now being computed on the SC thermal state, $\hat{H}_V$ is approximated in the Fock channel by

$$\begin{aligned} \hat{H}_V \stackrel{\text{HF}}{\simeq} & (\text{Hartree channel}) \\ & + V \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \left(u_{\sigma}^{(s^*)} \varphi_{\mathbf{k}}^{(s^*)*} + u_{\sigma}^{(d)} \varphi_{\mathbf{k}}^{(d)*} \right) \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} - E_{\text{F},V}^{(\text{SC})} \\ & + (\text{Cooper channel}). \end{aligned} \quad (6.62)$$

where, as was for the normal phase,

$$E_{\text{F},V}^{(\text{SC})} = L_x L_y \times \frac{V}{2} \sum_{\sigma} \left(|u_{\sigma}^{(s^*)}|^2 + |u_{\sigma}^{(d)}|^2 \right). \quad (6.63)$$

The proofs are identical to those presented in Sec. 4.2.2, because the SC phase is translationally invariant as was the normal phase. Renormalisation of bands reads

$$\tilde{\epsilon}_{\mathbf{k}\sigma} \equiv \epsilon_{\mathbf{k}} + u_{\sigma}^{(s^*)} V (\cos k_x + \cos k_y) + u_{\sigma}^{(d)} V (\cos k_x - \cos k_y). \quad (6.64)$$

From now on we omit spin index, using spin rotational invariance. Each harmonic $u^{(\gamma)}$ satisfies the self-consistency equation:

$$u^{(\gamma)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \left[1 - \cos 2\tilde{\theta}_{\mathbf{k}} \tanh \left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2} \right) \right]. \quad (6.65)$$

In Tab. 6.2 we report these equations, writing explicitly the harmonics from Tab. 3.5 and using the angular relations of Eq. (6.47).

Proof 6.9 — Proof of Eq. (6.65): self-consistency equations for $u^{(\gamma)}$.

Symmetry	Self-consistency equation	Graph
s^* -wave	$u^{(s^*)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) \left[1 - \frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right]$	Fig. 3.4b
$d_{x^2-y^2}$ -wave	$u^{(d)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) \left[1 - \frac{\tilde{\xi}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right]$	Fig. 3.4e

Table 6.2 | Symmetry resolved self-consistency equations for the harmonics of $u_{\mathbf{k}}$. The shown equations are reformulations of Eq. (6.65).

We start by computing explicitly $u_{\mathbf{k}\sigma}$. For $\sigma = \uparrow$, we get the chain of equalities:

$$\begin{aligned}
u_{\mathbf{k}\uparrow} &= \langle \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} \rangle \\
&= \left\langle \hat{\Phi}_{\mathbf{k}}^\dagger \frac{\mathbb{I} + \tau^z}{2} \hat{\Phi}_{\mathbf{k}} \right\rangle \\
&= \frac{1}{2} \left[\langle \hat{\Gamma}_{\mathbf{k}}^\dagger \hat{\Gamma}_{\mathbf{k}} \rangle + \langle \hat{s}_{\mathbf{k}}^z \rangle \right] \\
&= \frac{1}{2} \left[\tilde{\text{Th}}_{\mathbf{k}}^{(+)}[\beta] - \cos 2\tilde{\theta}_{\mathbf{k}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta] \right] \\
&= \frac{1}{2} \left[1 - \cos 2\tilde{\theta}_{\mathbf{k}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right], \tag{6.66}
\end{aligned}$$

having we used Eq. (6.52) and sequent. Inserting this result in Eq. (6.7), we get Eq. (6.65) in this spin sector. For $\sigma = \downarrow$,

$$\begin{aligned}
u_{-\mathbf{k}\downarrow} &= \langle \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow} \rangle \\
&= 1 - \langle \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^\dagger \rangle \\
&= 1 - \left\langle \hat{\Phi}_{\mathbf{k}}^\dagger \frac{\mathbb{I} - \tau^z}{2} \hat{\Phi}_{\mathbf{k}} \right\rangle \\
&= 1 - \frac{1}{2} \left[\langle \hat{\Gamma}_{\mathbf{k}}^\dagger \hat{\Gamma}_{\mathbf{k}} \rangle - \langle \hat{s}_{\mathbf{k}}^z \rangle \right] \\
&= 1 - \frac{1}{2} \left[\tilde{\text{Th}}_{\mathbf{k}}^{(+)}[\beta] + \cos 2\tilde{\theta}_{\mathbf{k}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta] \right] \\
&= 1 - \frac{1}{2} \left[1 + \cos 2\tilde{\theta}_{\mathbf{k}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \right], \tag{6.67}
\end{aligned}$$

We observe that the last line equals Eq. (6.66). Hence

$$u_{\mathbf{k}\uparrow} = u_{-\mathbf{k}\downarrow}.$$

Inserting this result in Eq. (6.7), we get Eq. (6.65) for both spin sectors.

6.3.3 Effects of non-local Cooper terms: generalised pairing

In this section we discuss the effects of Cooper contractions in the HF decomposition of $\hat{H}_V$, shown in Eq. (2.22),

$$\begin{aligned}
\hat{H}_V &\stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) + (\text{Fock channel}) \\
&= V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \langle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle + \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \rangle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} - \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \rangle \langle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle \right).
\end{aligned}$$

The HF approximation to $\hat{H}_V$ in Cooper channel is given by

$$\hat{H}_V^{\text{HF}} \simeq (\text{Hartree channel}) + (\text{Fock channel}) - V \sum_{\gamma \neq s} \sum_{\mathbf{k} \in \text{BZ}} \left[w^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} \hat{\phi}_{\mathbf{k}} + \text{h.c.} \right] - E_{C,V}^{(\text{SC})}, \quad (6.68)$$

where the harmonics $w^{(\gamma)}$ have been defined in Eq. (6.8). The contraction energy is given by:

$$E_{C,V}^{(\text{SC})} \equiv -\frac{V}{L_x L_y} \sum_{\gamma \neq s} \sum_{\mathbf{k}, \mathbf{k}'} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} w_{\mathbf{k}} w_{\mathbf{k}'}^*. \quad (6.69)$$

In both the above equations, the sum is intended for $\gamma \in \{s^*, p_x, p_y, d_{x^2-y^2}\}$.

Proof 6.10 — Proof of Eq. (6.68): HF approximation of $\hat{H}_V$ in Cooper channel.

We know from Eq. (3.25) that the non-local attraction can be separated in o.s. and s.s. channels. As we reported in Tab. 3.3, the only non-zero contribution comes from the o.s. sector.

Let us move to reciprocal space. Applying Eq. (3.40), for the o.s. sector we get

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) \\ - \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \times \left[\langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \right. \\ \left. + \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \langle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \rangle - \langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \rangle \right]. \end{aligned} \quad (6.70)$$

We know that Cooper pairs have zero net momentum for translationally invariant SC phases. Hence, only $\mathbf{K} = \mathbf{0}$ contributes. Therefore, the above operator is reduced to:

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) \\ - \frac{2V}{L_x L_y} \sum_{\mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \times \left[\langle \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \rangle \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} \right. \\ \left. + \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \langle \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} \rangle - \langle \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger \rangle \langle \hat{c}_{-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{k}'\uparrow} \rangle \right]. \end{aligned}$$

Recall now Eq. (6.6), where we defined $w_{\mathbf{k}} \equiv \langle \phi_{\mathbf{k}}^\dagger \rangle$. Using the decomposition of Eqs. (3.47), (3.48), we end up with:

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) - \frac{V}{L_x L_y} \sum_{\gamma \neq s} \sum_{\mathbf{k}, \mathbf{k}'} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} w_{\mathbf{k}} \hat{\phi}_{\mathbf{k}'} \\ - \frac{V}{L_x L_y} \sum_{\gamma \neq s} \sum_{\mathbf{k}, \mathbf{k}'} \varphi_{\mathbf{k}}^{(\gamma)*} \varphi_{\mathbf{k}'}^{(\gamma)} w_{\mathbf{k}}^* \hat{\phi}_{\mathbf{k}'}^\dagger - E_{C,V}^{(\text{SC})}. \end{aligned} \quad (6.71)$$

Sum is intended for $\gamma \in \{s^*, p_x, p_y, d_{x^2-y^2}\}$. We recognised $E_{C,V}^{(\text{SC})}$ as defined in Eq. (6.69). Eq. (6.71) is reduced to

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} \stackrel{\text{HF}}{\simeq} (\text{Hartree channel}) - \sum_{\gamma \neq s} \left(\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} w_{\mathbf{k}} \right) \times V \sum_{\mathbf{k}' \in \text{BZ}} \varphi_{\mathbf{k}'}^{(\gamma)*} \hat{\phi}_{\mathbf{k}'} \\ - \sum_{\gamma \neq s} \left(\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)*} w_{\mathbf{k}}^* \right) \times V \sum_{\mathbf{k}' \in \text{BZ}} \varphi_{\mathbf{k}'}^{(\gamma)} \hat{\phi}_{\mathbf{k}'} - E_{C,V}^{(\text{SC})}. \end{aligned} \quad (6.72)$$

Inserting the expression for $w^{(\gamma)}$ of Eq. (6.8), since we have no s.s. contribution, we obtain Eq. (6.68).

Symmetry	Self-consistency equation	Graph
s -wave	$w^{(s)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$	Fig. 3.4a
s^* -wave	$w^{(s^*)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$	Fig. 3.4b
p_x -wave	$w^{(p_x)} = \frac{i\sqrt{2}}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \sin k_x \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$	Fig. 3.4c
p_y -wave	$w^{(p_y)} = \frac{i\sqrt{2}}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \sin k_y \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$	Fig. 3.4d
$d_{x^2-y^2}$ -wave	$w^{(d)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)$	Fig. 3.4e

Table 6.3 | Symmetry resolved self-consistency equations for the harmonics of $w_{\mathbf{k}}$. The shown equations are reformulations of Eq. (6.74)

The contraction energy of Eq. (6.69) can be written as follows:

$$E_{C,V}^{(\text{SC})} = -L_x L_y \times V \sum_{\gamma \neq s} |w^{(\gamma)}|^2. \quad (6.73)$$

with $\gamma \in \{s^*, p_x, p_y, d_{x^2-y^2}\}$.

Proof 6.11 — Proof of Eq. (6.73): non-local Cooper contraction energy..

The contraction energy is formally given by Eq. (6.69). Substituting the harmonics decomposition of Eq. (6.8), we get:

$$\begin{aligned} E_{C,V}^{(\text{SC})} &= -L_x L_y \times V \sum_{\gamma \neq s} \left(\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} w_{\mathbf{k}} \right) \left(\frac{1}{L_x L_y} \sum_{\mathbf{k}' \in \text{BZ}} \varphi_{\mathbf{k}'}^{(\gamma)*} w_{\mathbf{k}'} \right) \\ &= -L_x L_y \times V \sum_{\gamma \neq s} w^{(\gamma)} w^{(\gamma)*} \\ &= -L_x L_y \times V \sum_{\gamma \neq s} |w^{(\gamma)}|^2, \end{aligned}$$

which proves Eq. (6.73).

Each harmonic, including s -wave, satisfies the self-consistency equation:

$$w^{(\gamma)} = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \sin 2\tilde{\theta}_{\mathbf{k}} e^{i2\tilde{\zeta}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right). \quad (6.74)$$

In Tab. 6.3 we report these equations, writing explicitly the harmonics from Tab. 3.5 and using the angular relations of Eq. (6.47).

Proof 6.12 — Proof of Eq. (6.74): self-consistency equations for $w^{(\gamma)}$.

Consider Eq. (6.8). For the γ -wave component of the superconducting order parameter, the

following chain holds:

$$\begin{aligned}
w^{(\gamma)} &\equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \\
&= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \langle \hat{\Phi}_{\mathbf{k}}^\dagger \tau^+ \hat{\Phi}_{\mathbf{k}} \rangle \\
&= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\langle \hat{s}_{\mathbf{k}}^x \rangle + i \langle \hat{s}_{\mathbf{k}}^y \rangle}{2} \\
&= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right),
\end{aligned}$$

where $\tau^+ = (\tau^x + i\tau^y)/2$. We used Eqs. (6.49), (6.50) and (6.53). Use now Eqs. (6.47), (6.48),

$$\begin{aligned}
\frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} &= \frac{|\tilde{\Delta}_{\mathbf{k}}|}{\tilde{E}_{\mathbf{k}}} \left(\frac{\text{Re}\{\tilde{\Delta}_{\mathbf{k}}\}}{|\tilde{\Delta}_{\mathbf{k}}|} + \frac{i \text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}}{|\tilde{\Delta}_{\mathbf{k}}|} \right) \\
&= \sin 2\tilde{\theta}_{\mathbf{k}} \left(\cos 2\tilde{\zeta}_{\mathbf{k}} + i \sin 2\tilde{\zeta}_{\mathbf{k}} \right) \\
&= \sin 2\tilde{\theta}_{\mathbf{k}} e^{i2\tilde{\zeta}_{\mathbf{k}}},
\end{aligned}$$

and insert in previous equation. This gives Eq. (6.74). The derivation above holds for all symmetries, including s -wave.

With this proof we conclude this cumbersome algebraic derivation. In next section we summarise the obtained results, give the final form of the HF hamiltonian and discuss the symmetries of the gap function.

6.4 The HF hamiltonian and its properties

In this section we take together everything we said so far. Adding $\hat{H}_V$ to the conventional HM, we have seen, produces renormalisation of the bare bands, the gap function and the chemical potential.

We compose Eqs. (6.27), (6.59), (6.62) and (6.68). The final form of the HF gran-canonical hamiltonian reads

$$\hat{H} - \mu \hat{N} \stackrel{\text{HF}}{\simeq} \sum_{\mathbf{k} \in \text{BZ}} \tilde{\xi}_{\mathbf{k}} \left(\hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^\dagger \right) - \sum_{\mathbf{k} \in \text{BZ}} \left(\tilde{\Delta}_{\mathbf{k}} \hat{\phi}_{\mathbf{k}} + \tilde{\Delta}_{\mathbf{k}}^* \hat{\phi}_{\mathbf{k}}^\dagger \right) + E_a^{(\text{SC})} + E_c^{(\text{SC})},$$

having we included the anticommutation energy defined in Eq. (6.25), and the total contraction energy,

$$E_c^{(\text{SC})} \equiv - \left(E_{\text{H},U}^{(\text{SC})} + E_{\text{C},U}^{(\text{SC})} + E_{\text{H},V}^{(\text{SC})} + E_{\text{F},V}^{(\text{SC})} + E_{\text{C},V}^{(\text{SC})} \right) \quad (6.75)$$

which include all the shifts to energy we encountered. These are given, respectively, in Eqs. (6.11), (6.18), (6.61), (6.63) and (6.73).

6.4.1 Gap symmetries and anisotropic pairing

We focus on the renormalised quantities. The bare bands are given by:

$$\tilde{\xi}_{\mathbf{k}} = \epsilon_{\mathbf{k}} + V \sum_{\gamma \in \{s^*, d\}} u^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} - \tilde{\mu}, \quad (6.76)$$

and the full gap function by

$$\tilde{\Delta}_{\mathbf{k}} = -U w^{(s)} + V \sum_{\gamma} w^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*}. \quad (6.77)$$

We can identify the gap harmonics as

$$\tilde{\Delta}^{(s)} = -U w^{(s)}, \quad \tilde{\Delta}^{(\gamma)} = V w^{(\gamma)}, \quad \text{being } \gamma \in \{s^*, p_x, p_y, d\}.$$

We will investigate competition among these components.

We anticipated in Tab. ?? that Cooper pairing in the singlet channel requires for the gap wavefunction to be inversion-symmetric. Triplet pairing, instead, requires inversion anti-symmetry. This gives a symmetry selection rule for the symmetries of the gap function: if we work in singlet channel, the gap must be symmetric, thus only three harmonics are allowed:

$$\text{Singlet channel} \quad : \quad \tilde{\Delta}_{\mathbf{k}} = \tilde{\Delta}^{(s)} + \tilde{\Delta}^{(s^*)}(\cos k_x + \cos k_y) + \tilde{\Delta}^{(d)}(\cos k_x - \cos k_y).$$

An inverse statement holds for the triplet pairing, for which

$$\text{Triplet channel} \quad : \quad \tilde{\Delta}_{\mathbf{k}} = i\sqrt{2} \left(\tilde{\Delta}^{(p_x)} \sin k_x + \tilde{\Delta}^{(p_y)} \sin k_y \right).$$

6.4.2 The entropy density of the SC ground-state

The approximated ground-state in the SC phase is given by a gas of free fermions populating the two bands $\pm \tilde{E}_{\mathbf{k}}$, with occupations specified by the Fermi-Dirac distribution. Hence, the total entropy density of the ground state is given by that of a Fermi gas [8]:

$$s^{(\text{SC})} = s_-^{(\text{SC})} + s_+^{(\text{SC})},$$

where the superscript indicates the band,

$$s_{\pm}^{(\text{SC})} \equiv -\frac{k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[\left(1 - f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \right) \ln \left(1 - f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \right) + f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \ln f(\pm \tilde{E}_{\mathbf{k}}; \beta, 0) \right]. \quad (6.78)$$

Note the fact that the Fermi-Dirac distributions are computed at null chemical potential. Because of this, we get:

$$s_-^{(\text{SC})} = s_+^{(\text{SC})} \implies s^{(\text{SC})} \equiv s_-^{(\text{SC})} + s_+^{(\text{SC})} = 2s_+^{(\text{SC})}. \quad (6.79)$$

Proof 6.13 — Proof of Eq. (6.79): Bogoliubov bands $\pm \tilde{E}_{\mathbf{k}}$ give equal entropy densities..

Consider the entropy of the lower band,

$$s_-^{(\text{SC})} \equiv -\frac{k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[\left(1 - f(-\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \ln \left(1 - f(-\tilde{E}_{\mathbf{k}}; \beta, 0) \right) + f(-\tilde{E}_{\mathbf{k}}; \beta, 0) \ln f(-\tilde{E}_{\mathbf{k}}; \beta, 0) \right],$$

and Eq. (6.37), which implies

$$f(x; \beta, 0) = 1 - f(-x; \beta, 0).$$

Inserting the latter for $x = \tilde{E}_{\mathbf{k}}$ in the expression for $s_-^{(\text{SC})}$, we get

$$s_-^{(\text{SC})} \equiv -\frac{k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[f(\tilde{E}_{\mathbf{k}}; \beta, 0) \ln f(\tilde{E}_{\mathbf{k}}; \beta, 0) + \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \right],$$

therefore, comparing with Eq. (6.78), we get Eq. (6.79): the two bands give an identical contribution to entropy.

The expression for the entropy density can be simplified by using Eq. (4.40), which implies

$$\begin{aligned} & \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \\ &= \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \left(\tilde{E}_{\mathbf{k}} \beta + \ln f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right). \end{aligned}$$

It follows that $s^{(\text{SC})} = 2s_+^{(\text{SC})}$ can be written as:

$$s^{(\text{SC})} \equiv -\frac{2k_B}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) + \frac{2k_B \beta}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} f(\tilde{E}_{\mathbf{k}}; \beta, 0). \quad (6.80)$$

We note that, as the gap closes and $\tilde{E}_{\mathbf{k}} \rightarrow \tilde{\xi}_{\mathbf{k}}$, the above expression reduces to the entropy density for the normal phase reported in Eq. (4.39).

6.4.3 The free energy density of the SC ground-state

We compute the density of free energy, following the same procedure we used for the normal phase in Sec. 4.3.3. The free energy density is given by

$$f^{(\text{SC})} \equiv \frac{F^{(\text{SC})}}{L_x L_y}, \quad (6.81)$$

being $F^{(\text{SC})}$ the total free energy,

$$F^{(\text{SC})} = E_{\text{HF}}^{(\text{SC})} - \frac{S^{(\text{SC})}}{k_B \beta}, \quad (6.82)$$

and $S^{(\text{SC})} \equiv L_x L_y \times s^{(\text{SC})}$ the total entropy. Here $E_{\text{HF}}^{(\text{SC})}$ is the ground-state energy for the HF hamiltonian. In the SC phase, we compute $f^{(\text{SC})}$ using the following formula:

$$f^{(\text{SC})} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left(\tilde{\xi}_{\mathbf{k}} - \tilde{E}_{\mathbf{k}} \right) + \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) - U |w^{(s)}|^2 - V \left(\sum_{\gamma} |u^{(\gamma)}|^2 - \sum_{\gamma} |w^{(\gamma)}|^2 \right) + \tilde{\mu} n, \quad (6.83)$$

which gives the analytic expression for the free energy. Note that, as the gap closes and $\tilde{E}_{\mathbf{k}} \rightarrow \tilde{\xi}_{\mathbf{k}}$, the above expression reduces to the entropy density for the normal phase reported in Eq. (4.39).

Proof 6.14 — Proof of Eq. (6.83): free energy density for the SC phase.

Free energy density of the SC phase is derived following the procedure used for the other two phases. The five contributions to free energy are:

1. Quasiparticles energy:

$$\begin{aligned} e_g^{(\text{SC})} &= -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta] \\ &= -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} \tanh \left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2} \right). \end{aligned}$$

2. Anticommutation energy, as defined in Eq. (6.25):

$$\begin{aligned} e_a^{(\text{SC})} &\equiv \frac{E_a^{(\text{SC})}}{L_x L_y} \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{\xi}_{\mathbf{k}}. \end{aligned}$$

3. Contraction energy:

$$\begin{aligned} e_c^{(\text{SC})} &= \frac{E_c^{(\text{SC})}}{L_x L_y} \\ &= -U \left(n^2 + |w^{(s)}|^2 \right) + V \left(2zn^2 - \sum_{\gamma} |u^{(\gamma)}|^2 + \sum_{\gamma} |w^{(\gamma)}|^2 \right), \end{aligned}$$

where we used Eq. (6.75).

4. Entropic energy

$$\begin{aligned} e_s^{(\text{SC})} &= -\frac{s^{(\text{SC})}}{k_B \beta} \\ &= \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) - \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} f(\tilde{E}_{\mathbf{k}}; \beta, 0), \end{aligned}$$

where we used Eq. (6.80).

5. Chemical potential correction

$$e_n^{(\text{SC})} = \mu n.$$

Now, bringing everything together,

$$f^{(\text{SC})} \equiv e_g^{(\text{SC})} + e_a^{(\text{SC})} + e_c^{(\text{SC})} + e_s^{(\text{SC})} + e_n^{(\text{SC})}.$$

In particular, note that $e_g^{(\text{SC})}$ and the second term of $e_s^{(\text{SC})}$ can be composed nicely,

$$-\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} \tanh \left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2} \right) - \frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}} f(\tilde{E}_{\mathbf{k}}; \beta, 0) = -\frac{2}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{E}_{\mathbf{k}}.$$

Indeed, this is proven by noting that

$$\tanh \left(\frac{x}{2} \right) + \frac{2}{e^x + 1} = 1.$$

which follows from basic algebra.

Taking everything together, in the end we get the following expression for free energy density:

$$\begin{aligned} f^{(\text{SC})} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[\tilde{\xi}_{\mathbf{k}} - \tilde{E}_{\mathbf{k}} \right] + \frac{2}{\beta L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \ln \left(1 - f(\tilde{E}_{\mathbf{k}}; \beta, 0) \right) \\ &\quad - U \left(n^2 + |w^{(s)}|^2 \right) + V \left(2zn^2 - \sum_{\gamma} |u^{(\gamma)}|^2 + \sum_{\gamma} |w^{(\gamma)}|^2 \right) + \mu n \end{aligned}$$

As was for the normal and AF phases, we recognise the chemical potential shift of Eq. (6.60),

$$\tilde{\mu} = \mu + n(2zV - U).$$

Upon substituting this result, we obtain Eq. (6.83).

6.5 Analysis of the solution

In this section, we derive one constraint on the solution for the SC phase, which “regulates” the competition among various symmetries of the superconducting gap. We specify that, differently from AF and normal phases, for the singlet SC phase we do not have any theoretical constraint on the HFPs $u^{(\gamma)}$, $w^{(\gamma)}$.

Energetic constraints. We derived the expression of Eq. (6.83), which contains in its second row the HFPs $w^{(\gamma)}$ and $w^{(s)}$. We have:

$$-U|w^{(s)}|^2 + V \sum_{\gamma} |w^{(\gamma)}|^2 = \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{|\tilde{\Delta}_{\mathbf{k}}|^2}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right). \quad (6.84)$$

The right-hand side of Eq.(6.84) is a sum of positive terms, hence is positive (or zero, if the gap is null). So must be the left-hand side. Then,

$$-U|w^{(s)}|^2 + V \sum_{\gamma} |w^{(\gamma)}|^2 \geq 0.$$

If the superconducting gap is null, the above inequality is satisfied, because $w^{(\gamma)} = 0$ for all γ s. If the gap is finite, the above requirement translates to:

$$\sum_{\gamma} |w^{(\gamma)}|^2 \geq M^{(\text{SC})} \quad \text{and} \quad \gamma \neq s \quad (6.85)$$

where we defined the radius:

$$M^{(\text{SC})} \equiv |w^{(s)}|^2 \frac{U}{V}.$$

For the conventional HM, we know from Sec. 6.2.2 that $w^{(s)} = 0$. For the EHM, it can assume non-zero values, as long as Eq. (6.85) is satisfied. In other words, $w^{(s)}$ cannot be non-zero without at least one of the other two singlet harmonics – namely, s^* -wave and d -wave – being non-zero.

Proof 6.15 — Proof of Eq. (6.84).

Recall the explicit form of the conjugate gap function from Eq. (6.77),

$$\tilde{\Delta}_{\mathbf{k}}^* = -U w^{(s)*} + V \sum_{\gamma} w^{(\gamma)*} \varphi_{\mathbf{k}}^{(\gamma)}.$$

Expand the right-hand side of Eq. (6.84) and insert the above equation,

$$\begin{aligned} \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{|\tilde{\Delta}_{\mathbf{k}}|^2}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \tilde{\Delta}_{\mathbf{k}}^* \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) \\ &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \left[-U w^{(s)*} + V \sum_{\gamma \neq s} w^{(\gamma)*} \varphi_{\mathbf{k}}^{(\gamma)} \right] \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right). \end{aligned}$$

Using the self-consistency equations of Tab. 6.3 to recognise the various harmonics. We reduce the right-hand side of Eq. (6.84) to:

$$\begin{aligned} \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{|\tilde{\Delta}_{\mathbf{k}}|^2}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right) &= -U w^{(s)*} \times \overbrace{\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)}^{w^{(s)}} \\ &\quad + V \sum_{\gamma \neq s} w^{(\gamma)*} \times \underbrace{\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \varphi_{\mathbf{k}}^{(\gamma)} \frac{\tilde{\Delta}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tanh\left(\frac{\beta \tilde{E}_{\mathbf{k}}}{2}\right)}_{w^{(\gamma)}}, \end{aligned}$$

which coincides with the left-hand side of Eq. (6.84).

This proof ends the theoretical discussion of the SC phase. Differently from the other phases, we are not able to provide any more constraints on the HFPs.

6.6 Discussion of numeric results for the singlet channel

The last part of this chapter is devoted to presenting and discussing the numeric results we extracted from the iHF procedure. **We limited simulations to the singlet channel, because [I wanted to see the end of this project!]**. The SC is described by five HFPs

$$\mathbf{v} = \begin{bmatrix} u^{(s^*)} \\ u^{(d)} \\ w^{(s)} \\ w^{(s^*)} \\ w^{(d)} \end{bmatrix} \quad \text{with} \quad \mathbf{v} = \mathbb{R}^5,$$

and three RMPs:

$$\begin{aligned} \tilde{t}_{ij}^{(s^*)} &\equiv \left(t - \frac{u^{(s^*)}V}{2} \right) \varphi_{ij}^{(s^*)}, \\ \tilde{t}_{ij}^{(d)} &\equiv -\frac{u^{(s^*)}V}{2} \varphi_{ij}^{(d)}, \\ \tilde{\mu} &\equiv \mu + n(2zV - U). \end{aligned}$$

For AF and normal phase, we had two HFPs and no d -wave component of the renormalised hopping. We ran simulations at $\beta = 100.0$ within the boundaries:

$$0 \leq U \leq 12 \quad 0 \leq V \leq 8 \quad 0 \leq \delta \leq 0.45$$

For the normal phase we did not vary U ; for the AF phase we did not vary δ . Here, we need to vary both in order to explore the entire parametric space.

6.6.1 Multi-symmetry singlet superconductivity

In singlet channel, the most general gap function is a combination of s , s^* and d components. The resulting gap function $\tilde{\Delta}_{\mathbf{k}}$ emerges from a competition among these channels. Running unconstrained simulations, we are able to see directly the result of the competition in the self-consistent solution. We discuss the three channels separately.

s -wave gap. Consider the isotropic harmonic of the gap function, namely the s -wave component. In Fig. 6.1, we show the data for $w^{(s)}$. It is given by:

$$\tilde{\Delta}^{(s)} = -Uw^{(s)}.$$


In the conventional HM we know that s -wave superconductivity is impossible at HF level. This was discussed in Sec. (6.2.2). Instead in the EHM there is no symmetry selection rule for $w^{(s)}$. The only constraint comes from Eq. (6.85): $w^{(s)}$ can be non-zero as long as at least one $w^{(\gamma)}$ with $\gamma = s^*, d$ is non-zero. This is a necessary but insufficient condition.

In Fig. 6.1a we see that $w^{(s)}$ is null at $\delta = 0$ (left panel). Hence


$$\forall U/t, V/t \quad : \quad \tilde{\Delta}^{(s)}|_{\delta=0} = 0, \quad (6.86)$$

at least within the range of parameters we explored. Increasing doping (central and right panels), $w^{(s)}$ is null up to a critical value $V_c^{(s)}$, which depends also on U . For $V \geq V_c^{(s)}$, $w^{(s)}$ is negative and the shape of the curve depends strongly on U . In particular, the amplitude $|w^{(s)}|$ exhibits a local maximum. At almost unitary filling, ratios V/t comparable with 1 are sufficient to develop a finite convergence value for $w^{(s)}$. Therefore, doping lowers the critical value of the non-local attraction that is needed in order to establish an isotropic gap in the local channel.

The dependence of $w^{(s)}$ on doping is shown in Fig. 6.1b. As anticipated, $w^{(s)} = 0$ at half-filling. Varying U/t and V/t we see that $w^{(s)} = 0$ reaches some negative minimum whose absolute value is lower at higher U s.




a | Dependency of $w^{(s)}$ on V at various U s, compared at three values of δ .




b | Dependency of $w^{(s)}$ on δ at various V s, compared at three values of U .

Figure 6.1 | Dependency of the s -wave HFP $w^{(s)}$ on V at fixed U and δ (top panel, compared figures) and on δ at fixed V and U (bottom panel, compared figures).



a | Dependency of $w^{(s^*)}$ on V at various U s, compared at three values of δ .



b | Dependency of $w^{(s^*)}$ on δ at various V s, compared at three values of U .

Figure 6.2 | Dependency of the s^* -wave HFP $w^{(s^*)}$ on V at fixed U and δ (top panel, compared figures) and on δ at fixed V and U (bottom panel, compared figures).




Figure 6.3 | Dependency of the d -wave HFP $w^{(d)}$ on V at fixed U and δ (top panel, compared figures) and on δ at fixed V and U (bottom panel, compared figures).

s^* -wave gap. The results for the non-local isotropic HFP $w^{(s^*)}$ are shown in Fig. 6.2. This component of the full gap is given by

$$\tilde{\Delta}^{(s^*)} = V w^{(s^*)}.$$

Consider first Fig. 6.2a. At null and finite fillings, $w^{(s^*)}$ is null up to a critical value $V_c^{(s^*)}$. Its position depends on δ and weakly on U/t . The half-filling curve (left panel) is independent of U/t . Differently from $w^{(s)}$, we have

$$\forall U/t \quad \wedge \quad \forall V < V_c^{(s^*)} \quad : \quad \tilde{\Delta}^{(s^*)}|_{\delta=0} = 0,$$

The local isotropic gap harmonic $\tilde{\Delta}^{(s)}$ was always null at half filling. Its non-local counterpart $\tilde{\Delta}^{(s^*)}$ is null up to the critical value delimiting the phase transition. Therefore, superconductivity can establish at half-filling with a gap whose isotropic part is purely non-local. Note that we are not saying that pure s^* -wave superconductivity is possible. As we discuss later, at half-filling the dominant harmonic of the gap function has d -wave symmetry.

In Fig. 6.2b we see that the dependence on doping of $w^{(s^*)}$ is peculiar. At low doping ($\delta \lesssim 0.3$), $w^{(s^*)}$ is small for small values of V (left panel). At higher doping (central and right panels) we see that a local maximum emerges, indicating that $w^{(s^*)}$ converges to higher values at smaller V s.

d -wave gap. In Fig. 6.3 we report our findings for the HFP $w^{(d)}$, recalling that the d -wave gap is given by:

$$\tilde{\Delta}^{(d)} = V w^{(d)}.$$

We see from the top panel (Fig. 6.3a) that $w^{(d)}$ depends weakly on U . At half-filling (left panel), the d -wave gap is zero only at $V = 0$:

$$\forall U/t \quad \wedge \quad V = 0 \quad : \quad \tilde{\Delta}^{(d)}|_{\delta=0} = 0.$$

Differently from the other components, $w^{(d)}$ grows to finite values already at infinitesimal V/t . Recalling what we said in the above paragraph, we note that when the strength of the non-local


Singlet gap $\tilde{\Delta}_k$ harmonics ($t = 1.0$, $U = 4.0$, $\beta = 100.0$)

Figure 6.4 | Plot of the singlet gap harmonics $\tilde{\Delta}^{(\gamma)}$, with $\gamma = s, s^*, d$ at $U/t = 4$. The panels on the left show the three harmonics separately as heatmaps, while the surface plot on the right shows them comparatively.

attraction grows enough, the s^* -wave part sets in and symmetries mix. At finite filling (central and right panels) a critical value $V_c^{(d)}$ emerges such that the d -wave gap appears at $V > V_c^{(d)}$.

The dependence of $w^{(d)}$ on doping at different values of U is shown in the bottom row (Fig. 6.3b). For any V/t , the HFP $w^{(d)}$ is a monotonically decreasing function of δ . As we said, the critical value $V_c^{(d)}$ increases with doping. Because of this, we see that the lighter curves of Fig. 6.3b, corresponding to lower values of V , reach zero quite rapidly. The curves for larger values of V , instead, reach zero at almost unitary filling. In all three panels of Fig. 6.3b we see that at doping $\delta \approx 0.3$ various lines cross at $w^{(d)} \approx 0.3$. Finally, at null doping, there exists a finite value of V that maximises $w^{(d)}$: it is visible in Fig. 6.3a.

Competition among the channels. To get a valid understanding of the competition among channels, we need to compare the values of the harmonics $\tilde{\Delta}^{(\gamma)}$ for $\gamma = s, s^*, d$. Competition is reflected in the relative ratio between different components. We report a comparison of such in Fig. 6.4. **The presented data are limited to the case $U/t = 4$ for graphic reasons, but data and plots for other values of U are openly accessible at our repository.**

On the left column we plot three heatmaps, one for each harmonic in singlet SC phase. The same data are shown, superimposed, in the surface plot on the right. The harmonics $\tilde{\Delta}^{(\gamma)}$ cross if V/t gets large enough, with the d -wave component reaching the largest values in the parametric region we investigated, followed closely by the s^* -wave harmonic. Isotropic harmonics (s and s^*) are significantly non-zero in the same region, as we see in the first two rows of the left column. The two channel differ in amplitude significantly: the maximum value reached by the local gap harmonic $\tilde{\Delta}^{(s)}$ is approximately an order of magnitude smaller than the maximum of the non-local harmonic $\tilde{\Delta}^{(s^*)}$. Moreover, we see that at $\delta = 0.0$ the local isotropic harmonic $\tilde{\Delta}^{(s)}$ is completely suppressed, as we expressed in Eq. (6.86).

All harmonics share a quite strange boundary around $\delta \approx 0.34$, $V/t \approx 2.5$, visible as a sharp vertical segment in the three heatmaps on the left column of Fig. 6.4. The isotropic harmonics suddenly change from zero to large values moving from left to right, while the d -wave part does the opposite. Whether this behaviour underlies some deeper structure we were not able to tackle




Figure 6.5 | Depiction of the normal state region and the three singlet superconducting regions, at $U/t = 0, 4, 12$ from left to right. Lighter blue indicates the pure d -wave SC phase, blue the isotropic $s \oplus s^*$ phase, deeper blue the mixed region. In text is described the convention we used to identify the boundaries.

with our coarse-grained simulations, or is a numeric artifact, we do not know.

Four phases can be distinguished: the normal state and three superconducting regions (pure d -wave, pure $s \oplus s^*$ -wave, mixed). In Fig. 6.5 we give a schematic depiction of how the plane (δ, V) divides in the four regions at $U/t = 0$ (left), $U/t = 4$ (centre) and $U/t = 12$ (right). The boundaries were computed by setting an arbitrary threshold $\varepsilon = 10^{-2}$, with the following rule:

$$\begin{aligned}
 \text{Normal} & \quad |\tilde{\Delta}^{(s)}| < \varepsilon \quad \wedge \quad |\tilde{\Delta}^{(s^*)}| < \varepsilon \quad \wedge \quad |\tilde{\Delta}^{(d)}| < \varepsilon, \\
 d\text{-wave} & \quad |\tilde{\Delta}^{(s)}| < \varepsilon \quad \wedge \quad |\tilde{\Delta}^{(s^*)}| < \varepsilon \quad \wedge \quad |\tilde{\Delta}^{(d)}| \geq \varepsilon, \\
 s \oplus s^*\text{-wave} & \quad [|\tilde{\Delta}^{(s)}| \geq \varepsilon \quad \vee \quad |\tilde{\Delta}^{(s^*)}| \geq \varepsilon] \quad \wedge \quad |\tilde{\Delta}^{(d)}| < \varepsilon, \\
 \text{Mixed} & \quad [|\tilde{\Delta}^{(s)}| \geq \varepsilon \quad \vee \quad |\tilde{\Delta}^{(s^*)}| \geq \varepsilon] \quad \wedge \quad |\tilde{\Delta}^{(d)}| \geq \varepsilon.
 \end{aligned}$$

We consider s and s^* as part of the same isotropic channel: just one harmonic being larger than the threshold is sufficient for us to assert that the gap function has an isotropic harmonic.

Comparing the three panels of Fig. 6.5, we see that increasing the value of the local repulsion U the shapes of the four regions change. In particular, the d -wave region expands to the right (larger δ s); the normal- $s \oplus s^*$ SC boundary moves up (larger V s). Note that on the left panel the isotropic region is labeled as purely s^* -wave: this is because if $U = 0$, the s -wave gap harmonic is null, even if $w^{(s)}$ converges to a non-zero value. The growth of the local repulsion U has two effects:

1. turning on the s -wave gap harmonic;
2. expanding the d -wave region and restricting the isotropic and mixed region.

From the last point we understand that increasing U we allow for the gap function to have a finite s -wave component, but this acts as a sort of internal competition within the isotropic region. Indeed, from the constraint of Eq. (6.85), we know that $w^{(s)} \neq 0$ only if $w^{(\gamma)} \neq 0$, with $\gamma \neq s$. In the $s \oplus s^*$ region the constraint reads:

$$|w^{(s^*)}|^2 \geq |w^{(s)}|^2 \frac{U}{V}, \quad (\text{Isotropic region})$$

Suppose we work at fixed δ and $V > 0$ in the isotropic region at $U = 0$. There is no constraint whatsoever on $w^{(s^*)}$. Also, $w^{(s)}$ is free to converge at some finite value (we saw this in the left panel of Fig. 6.1b): the gap still has no s -wave component. Suppose now we switch on $U > 0$. Now, in order for $w^{(s)}$ to reach the same value as before, $w^{(s^*)}$ has to respect the constraint posed by the above equation. As U increases, the right-hand side gets bigger and $w^{(s^*)}$ needs to converge to larger and larger values to sustain $w^{(s)}$, otherwise to drop to zero. This is why increasing U the isotropic boundary moves up: $w^{(s^*)}$ isn't "free" anymore and must respect the constraint of Eq. (6.85).

In the three panels of Fig. 6.5, the mixing region is pretty much unchanged. At low doping, it is bounded from below by an horizontal line at $V/t \approx 3$. Increasing doping at fixed U , we see that the lower boundary reaches a cusp-like minimum around $\delta \approx 0.3 \div 0.4$ and then grows. This cusp coincides with the aforementioned sharp vertical segment, visible in Fig. 6.4. If we work at fixed doping and U , and choose δ such that we are aligned with the horizontal position of the cusp on the (δ, V) plane, the critical value of V/t for transitioning to a mixed SC state is minimised. At high values of V/t the mixing region extends to almost unitary doping, being limited on the right by a vertical line at $\delta \gtrsim 0.4$. We remind that we are discerning the SC phases by applying an arbitrary threshold ε to each gap component: we imagine that by using a more refined technique and by increasing the resolution of the simulation the boundary would appear less straight. Anyway, apart from a vertical stripe at almost unitary filling, at high V/t the dominant singlet SC phase is described by a mix of d -wave and $s \oplus s^*$ -wave component. Recalling the surface plot of Fig. 6.4, we see that at $V/t \gg 1$ the s -wave component gets weaker with respect to the other two. In this limit, if $U \ll V$, the s -wave component is negligible, giving an approximate $d \oplus s^*$ mixed phase.

The mixing region is of particular interest also due its connection with the planar rotational SSB,

$$\text{Symmetry mixing} \implies C_4 \xrightarrow{\text{SSB}} C_2.$$

As we show in next section, the mixed-symmetry region coincides with the portion of the (δ, V) plane where planar rotational symmetry is broken.

6.6.2 Rotational SSB and nematic superconductivity

In the context of this thesis, nematicity is established whenever $\tilde{t}^{(d)}$ becomes non-zero. This was discussed in detail in the context of the normal phase. If the renormalised hopping parameter has a non-zero d -wave component, this means that bare hopping is effectively boosted in one direction and damped in the other. Indeed, from Eq. (4.37), we know that

$$\tilde{t}^{(d)} = \frac{\tilde{t}^{(y)} - \tilde{t}^{(x)}}{2}.$$

As we anticipated at the end of Sec. 4.3.1, if $\tilde{t}^{(d)} > 0$ the renormalised hopping is larger in direction y , if $\tilde{t}^{(d)} < 0$ instead it's larger in direction x . This interpretation serves to grasp the general idea behind nematicity, but is to be taken carefully. Let us get more rigorous. In the SC phase, the many body state is a gas of Bogoliubov quasiparticles which do not “hop on neighbouring sites”. If $\tilde{t}^{(d)}$ is non-zero, the superconductor is nematic in the sense that Bogoliubov quasibands are not rotationally invariant,

$$\tilde{E}_{\mathbf{k}} \neq \tilde{E}_{\mathcal{R}\mathbf{k}},$$

being $\mathcal{R}$ a $\pi/2$ rotation. We connect this effect to hopping renormalisation *a posteriori* following the formal discussion of Sec. 4.3.1.

For the normal and AF phases we were able to prove that $u^{(d)} = 0$ for any choice of model parameters. For the SC phase, instead, $u^{(d)}$ converges to non-zero values. For the sake of conciseness, we skip plotting the curves for $u^{(d)}$ **(which are, anyway, openly accessible at our repository)**, and show directly the curves for the quantity

$$\tilde{t}^{(d)} = -\frac{Vu^{(d)}}{2}.$$

This is done in Fig. 6.6: nematicity is undoubtedly present. In the top panel (Fig. 6.6a), we see that a critical value of V exists over which nematicity is established. Moreover, the $\delta = 0.0$ case (left panel) is independent of U . In all cases, the convergence values for $\tilde{t}^{(d)}$ are two order of magnitudes smaller than $t = 1.0$: in the region of parametric space we explore, nematicity is a sharp but small effect. Note that at $\delta = 0.2$ (central panel) $\tilde{t}^{(d)} \geq 0$, while at $\delta = 0.4$ (right panel) $\tilde{t}^{(d)} \leq 0$. Increasing the doping we change the sign of $\tilde{t}^{(d)}$.

This sign change is most evident in the first two figures of the bottom panel (Fig. 6.6b), where we vary the doping level while keeping fixed the local repulsion U . The behaviour of $\tilde{t}^{(d)}$ is similar at low doping in the three cases, but changes significantly at larger doping. Indeed, at $U/t = 0$




Figure 6.6 | Dependency of the d -wave renormalised hopping $\tilde{t}^{(d)}$ varying the doping δ and the local repulsion U . In the top panel we compare three doping levels $\delta = 0.0, 0.2, 0.4$ varying $0 \leq U \leq 8$. In the bottom panel we compare three repulsion values $U = 0.0, 5.0, 10.0$ varying $0 \leq \delta \leq 0.45$.




Figure 6.7 | Heatmap of the anisotropic component of renormalised hopping, $\tilde{t}^{(d)}$, at $U/t = 0, 4, 12$ from left to right. The dataset is the same of Fig. 6.5, and the mixing boundary is the set of points separating the mixed region from the rest of the phases.

(left figure) and $U/t = 5$ (central figure) at some point $\tilde{t}^{(d)}$ changes sign. At $U/t = 10$ (right figure), all our data points are positive (possibly the sign flip occurs at higher V s). Moreover, as U increases, the amplitude of the oscillation of $\tilde{t}^{(d)}$ as a function of δ reduces. We conclude that both the local repulsion and electronic doping disrupt nematicity, which is instead most effective for a purely NNs attractive model.

Nematic effects are directly related to symmetry mixing in the superconducting order parameter. Consider Fig. 6.7, where we plot the convergence values for $\tilde{t}^{(d)}$ in the (δ, V) plane for $U/t = 0, 4, 12$. The used dataset is the same of Fig. 6.5. The superimposed black lines are the mixing boundaries, the set of extremal points of the deeper blue regions of Fig. 6.5. Despite a bit of roughness related to the gridded nature of our data, it is evident that these boundaries encircle the regions where $\tilde{t}^{(d)} \neq 0$. Note that the set of points composing the mixing regions are those satisfying the threshold rule exposed in last section; choosing another threshold, the black curves

change. Nevertheless, the key point is that there is an evident connection between symmetry mixing and breaking of planar rotational symmetry.

The mixing region coincides with the nematic region. As we said at the beginning of this section, nematicity can be established in two ways, based on the direction along which hopping is boosted. Hence it must hold

$$M \simeq N_x \cup N_y$$

with the following convention on notation:

M = Region of (δ, V) plane where mixing occurs,

N_x = Region of (δ, V) plane where $\tilde{t}^{(d)} < 0$,

N_y = Region of (δ, V) plane where $\tilde{t}^{(d)} > 0$.

The notation N_ℓ has the following meaning: the index ℓ indicates the direction along which renormalised hopping is larger. We indicate in Fig. 6.7 these regions for $U/t = 0, 4, 12$.

Note that the map were colour-corrected in order to enhance the visibility of the blue region. The intensity of the colouring there suggests that the amplitude is comparable with the red region, but is not. Indeed, we set up the colormap such that gray is aligned with $\tilde{t}^{(d)} = 0.0$, and intensity of colour grows asymmetrically in the negative and positive direction.

The three panels of Fig. 6.7 differ crucially in the fact that the data for $U/t = 12$ (right) are all positive, while those for $U/t = 0, 4$ (left, centre) are negative in a portion of space. In other words, as we increase U/t , N_y expands and N_x shrinks. Moving from $U/t = 0$ (left) to $U/t = 4$ (centre) we see that the amplitude of $\tilde{t}^{(d)}$ decreases in N_x and not in N_y . We already observed this in Fig. 6.6b. Moreover, for $U/t = 12$ (right) N_x is apparently absent and N_y takes up the entirety of the mixing region. Whether there exists some critical value of U/t above which N_x disappears in the entirety of the (δ, V) plane, we do not know.

Breaking of planar rotational symmetry and the insurgence of a d -wave harmonic for the renormalised hopping is certainly the most peculiar effect for the singlet SC phase. It adds to the isotropic bands shrinking, which is, the s^* -wave renormalisation of the hopping parameter. We discuss it in next section.

6.6.3 Isotropic bands shrinking

The last HFP to be checked is $u^{(s^*)}$, responsible of bands shrinking. Its physical role is the isotropic renormalisation of the hopping parameter


$$\tilde{t}^{(s^*)} = t - V u^{(s^*)} 2,$$

again, we skip plotting $u^{(s^*)}$ and plot directly $\tilde{t}^{(s^*)}$ in Fig. 6.8. In the top panel (Fig. 6.8a), we show the dependence of $t^{(s^*)}$ on V . The three curves, obtained at three doping levels, are approximately independent of U . The common feature is that $\tilde{t}^{(s^*)}$ decreases somewhat linearly with V up to some critical value, above which it flattens to an horizontal asymptote. Electronic doping changes primarily the height of this asymptote


As was for the AF and normal phases, electronic doping disrupts the isotropic bands shrinking effect. At half-filling (left figure) we see that bands shrinking is most effective, reducing the hopping parameter to less than half its $V = 0$ value. In the bottom panel (Fig. 6.8b) we show $t^{(s^*)}$ as a function of δ , varying V and U . As we already noted, the dependence on U is very weak and the three figures are a lot alike. As we already noted, $t^{(s^*)}$ increases with doping.

In Fig. 6.9 we compare $t^{(s^*)}$ for the normal phase (blue surface) and the SC phase (green surface). The two surfaces share a general feature: $t^{(s^*)}$ is approximately unitary at low V/t and almost unitary doping, and decreases significantly in the low- δ and high- V/t limit. In this region, for the normal phase we observe the Mott-localisation plateau discussed in Sec. 4.5.1, where $t^{(s^*)} \approx 0$. In the SC phase, looking at the same region, $t^{(s^*)}$ is minimised but no plateau is formed. More importantly, isotropic hopping is reduced to roughly half the value of t and not nullified. We do not observe a Mott-like area as for the normal phase.

Recalling the plots of Fig. 6.5, the low- δ and high- V/t limit (the corner where $t^{(s^*)}$ is minimised) falls deep within the mixed region. Recalling also Fig. 6.4, in this corner the d -wave and s^* -wave



a | Dependency of $\tilde{t}^{(s^*)}$ on V at various U s, compared at three values of δ .



b | Dependency of $\tilde{t}^{(s^*)}$ on δ at various V s, compared at three values of U .

Figure 6.8 | Dependency of the s^* -wave renormalised hopping $t^{(s^*)}$ varying the doping δ and the local repulsion U . In the top panel we compare three doping levels $\delta = 0.0, 0.2, 0.4$ varying $0 \leq U \leq 8$. In the bottom panel we compare three repulsion values $U = 0.0, 5.0, 10.0$ varying $0 \leq \delta \leq 0.45$.




Figure 6.9 | Comparison of the isotropic part of the renormalised hopping $\tilde{t}^{(s^*)}$ for the SC (green) and normal (blue) phases on the (δ, V) plane. The data from the normal phase are the same of Fig. 4.6.




Figure 6.10 | Heatmaps of the free energy and entropy densities at $U/t = 4$. The black dotted lines represent the internal boundaries of the singlet SC phase, extracted from Fig. 6.5.

harmonics of the gap function are large and comparable. Finally, we know from Fig. 6.7 (central panel) that here nematic effects are small, but significantly non-zero. In particular, considering that

$$\lim_{\delta \rightarrow 0} \lim_{V/t \gg 1} \frac{\tilde{t}(s^*)}{t} \approx 0.4 \quad \text{and} \quad \lim_{\delta \rightarrow 0} \lim_{V/t \gg 1} \frac{\tilde{t}(d)}{t} \approx 0.06,$$

we see that the two components differ just by one order of magnitude. Nematic effects account for a rather large portion of hopping renormalisation.

With this consideration, we end the presentation of HFPs and RMPs and move to the last section, where we discuss the results regarding free energy and entropy densities.

6.6.4 Solution free energy and entropy densities

For the SC phase, free energy density is given by Eq. (6.83), while entropy density is given by Eq. (6.78). In Fig. 6.10 we plot, sided, the heatmaps for f and s at $U/t = 4$ in the (δ, V) plane. Superimposed, we show the boundaries between the four regions identified in Fig. 6.5. Consider first the free energy, on the left: our data suggest that free energy is maximised in the d -wave phase, followed by the mixed phase. At high doping, the $s \oplus s^*$ -wave phase is low-energetic. Overall, f varies smoothly across phase boundaries, and we do not identify the black lines with some isoenergetic line separating phases.

This is not the case for entropy density. Indeed, the entropic region on the right panel of Fig. 6.10 coincides with the normal phase. The three superconducting regions are low-entropic. This is explained by the following argument:

- In the normal phase, the Bogoliubov band reduces to $\tilde{E}_{\mathbf{k}} \rightarrow \tilde{\xi}_{\mathbf{k}}$. The chemical potential crosses the bare band and a Fermi surface exists. Hence, at finite temperature, the Fermi-Dirac occupation of single-particle states varies continuously from almost unitary (deep in the Fermi sea) to almost zero (far outside). All those states in the middle, around the Fermi level, are highly entropic: we have shown this in Fig. 4.9b. It is the intrinsic metallic nature of the normal phase that makes it entropic.
- In the SC phase, we open a gap at the chemical potential, not at zero energy (as instead was in the AF phase). Computing entropy as in Sec. 6.4.2, we apply the Fermi-Dirac distribution on the two bands $\pm \tilde{E}_{\mathbf{k}}$. As the gap opens, the lower one fills ($f \approx 1$), while the upper empties ($f \approx 0$). As we lower temperature, the above statement gets exact. Recalling Fig. 4.9b, we see that entropy is minimised at $f = 0, 1$. Hence, the SC phase is low-entropic.

This is the reason behind the sharp separation of the normal region from the SC regions in the right panel of Fig. 6.10.

In Fig. 6.11 we show f and s as functions of V , varying δ and comparing for three values of U . On both rows, moving from left to right we do not observe significant differences: f and s depend weakly on U . We already observed the presence of a maximum of the free energy at ratios




Figure 6.11 | Dependency of free energy (top) and entropy (bottom) densities on V varying the doping δ and the local repulsion U .




Figure 6.12 | Surface plots of the energies among the three phases. For the normal phase, we use that fact that $f^{(\text{SC})}$ depends weakly on U , while $f^{(\text{N})}$ does not depend on it at all, to limit to the case $U/t = 4$. For the AF phase, we limit to the half-filling condition and show data in the (U, V) plane.

V/t comparable with 1. In Fig. 6.11a we see this maximum around $V/t \lesssim 4$. Comparing with the data of Fig. 6.10, we can imagine that this maximum could be located at the boundary between d -wave and mixed regions. Our data and methods are too rough to assert with certainty.

In the bottom panel, entropy density (Fig. 6.11b) shows the aforementioned sharp separation between normal and SC regions, and is maximised at half-filling and $V = 0$. That is the case of the conventional HM. We already argued that superconductivity is impossible at HF level, while commensurate antiferromagnetism is favoured at null doping with respect to the normal phase.

To conclude, we compare the free energy of the SC phase with that of the normal phase and the AF phase. We said that f depends rather weakly on U for the singlet SC phase, so we carry

out the comparison with normal phase at $U/t = 4$ in the (δ, V) plane. In Fig. 6.12a we plot the quantity

$$f^{(N)} - f^{(SC)},$$

and observe that it is greater or equal to zero over the entire explored region. The energy difference is null only at $V/t \lesssim 2$, where the superconducting gap is null. We plot, superimposed, the boundaries of the normal region extracted from Fig. 6.5. In the rest of the (δ, V) plane the SC phase is energetically favoured with respect to the normal phase.

In Fig. 6.12b we show the energy difference between AF and SC phases,

$$f^{(AF)} - f^{(SC)}.$$

We colour-corrected the surface plot to align the gray with the $f = 0$ level. Colour separates positive and negative regions: the SC phase is dominant where $f^{(AF)} > f^{(SC)}$ (green region), in the rest of the plot the AF phase dominates (blue region). We indicate the contour level at $f = 0$ as a black dotted line, obtained as the set of data points whose energy difference is closest to zero. At half-filling, commensurate antiferromagnetism is almost everywhere energetically favoured. Singlet superconductivity prevails in the $U/t \ll 1$, $V/t \gg 1$ limit. The fact that the phase boundary has somewhat a linear shape suggests the fact that possibly a critical ratio exists,

$$\left(\frac{V}{U}\right)_c = k, \quad (\text{Critical ratio})$$

delimiting a phase transition from antiferromagnetic to superconducting ordering at half-filling.

6.7 A final note

We conclude this chapter with a note of method. We performed simulations on the singlet channel, but also gathered data for the triplet channel. We omitted to show them in order to avoid unnecessary complications. Indeed, we observed that triplet superconductivity (as we formulated it at HF level) is suppressed everywhere in the region of parametric space we are exploring. This means that

$$w^{(p_x)} = w^{(p_y)} = 0. \quad (\text{Observed numerically})$$

Although non present in this discussions, **the plots for these HFPs are openly accessible at our repository.**

We did not provide any theoretical constraint on the five HFPs for the singlet SC phase. We tried to obtain a proof similar to those presented for the normal and AF phases, but we did not find any. It would be interesting, if possible, to work out a method to express analytically the position of the various boundaries we identified.

Appendix A

Superexchange and virtual hopping in Hubbard lattices

Superexchange is a key mechanism in the formation of AF ordering in Hubbard lattices. It is an extension of the direct exchange effect, that we describe in Sec. A.2, that favours antiferromagnetic ordering and takes advantage of the chemical structure of cuprates. The mechanism becomes clear enough if we start with a 2-sites Hubbard toy model.

A.1 Singlet pairing in Hubbard lattices

Consider the 2 sites Hubbard model of Eq. (1.2),

$$\hat{H} = -t \left(\hat{c}_{1\uparrow}^\dagger \hat{c}_{2\uparrow} + \hat{c}_{1\downarrow}^\dagger \hat{c}_{2\downarrow} + \text{h.c.} \right) + U \left(\hat{n}_{1\uparrow} \hat{n}_{1\downarrow} + \hat{n}_{2\uparrow} \hat{n}_{2\downarrow} \right),$$

with $i = 1, 2$ the site index. For an half-filled system (i.e. with two electrons) the Hilbert space is six-dimensional. We use the notation $|n_{1\uparrow} n_{1\downarrow} n_{2\uparrow} n_{2\downarrow}\rangle$ to indicate the six states of the computational basis:

$$\begin{aligned} |\psi_1\rangle &\equiv |1010\rangle, & |\psi_3\rangle &\equiv |1001\rangle, & |\psi_5\rangle &\equiv |0011\rangle, \\ |\psi_2\rangle &\equiv |1100\rangle, & |\psi_4\rangle &\equiv |0110\rangle, & |\psi_6\rangle &\equiv |0101\rangle. \end{aligned} \quad (\text{A.1})$$

We give a pictorial representation of these states in Fig. A.1. We ordered the states in such a way that the two states with total spin different from zero, $|\psi_1\rangle$ and $|\psi_6\rangle$, are at the extrema of the set.

The matrix elements of the hamiltonian can be computed using this basis,

$$H_{ij} = \langle \psi_i | \hat{H} | \psi_j \rangle \quad \Rightarrow \quad H = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & U & -t & -t & 0 & 0 \\ 0 & -t & 0 & 0 & -t & 0 \\ 0 & -t & 0 & 0 & -t & 0 \\ 0 & 0 & -t & -t & U & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}. \quad (\text{A.2})$$

In the above matrix, rows and columns are ordered following the states indices of Eq. (A.1). The states $|\psi_1\rangle$ (both spins up) and $|\psi_6\rangle$ (both spins down) are zero-energy eigenstates. Indeed, they do not interact via Coulomb repulsion (are on different sites) and neither through NNs hopping, due to Pauli exclusion principle.




Figure A.1 | Computational basis of Eq. (A.1) for the 2-sites Hubbard model.

Structure	Eigenstate	Energy
Spin-1/2 singlet	$\frac{ \psi_3\rangle - \psi_4\rangle}{\sqrt{2}}$	$E^- = \frac{U}{2} - \sqrt{\frac{U^2}{4} + 4t^2}$
Spin-1/2 triplet	$ \psi_1\rangle, \frac{ \psi_3\rangle + \psi_4\rangle}{\sqrt{2}}, \psi_6\rangle$	0
	$\frac{ \psi_2\rangle - \psi_5\rangle}{\sqrt{2}}$	U
	$\frac{ \psi_2\rangle + \psi_5\rangle}{\sqrt{2}}$	$E^+ = \frac{U}{2} + \sqrt{\frac{U^2}{4} + 4t^2}$

Table A.1 | List of exact eigenstates and relative energies for the 2-sites half-filled Hubbard model.

All states with null total spin projection, namely $\{|\psi_i\rangle\}$ for $2 \leq i \leq 5$, give a null matrix elements when computed with the aforementioned states, but non-zero matrix elements among themselves. Note that this set includes both the states with one doubly occupied site and one empty site ($|\psi_2\rangle, |\psi_5\rangle$) and those with two singly occupied sites ($|\psi_3\rangle, |\psi_4\rangle$). These states are connected by the central 4×4 matrix in Eq. (A.2). The latter is diagonalised by the means of a change of basis M ,

$$M \begin{bmatrix} U & -t & -t & \\ -t & & & -t \\ -t & & & -t \\ & -t & -t & U \end{bmatrix} M^\dagger = \begin{bmatrix} E^- & & & \\ & 0 & & \\ & & U & \\ & & & E^+ \end{bmatrix}.$$

where blank slots indicate zeros. In Tab. A.1 we report the eigenvectors and relative eigenvalues obtained from diagonalisation.

The ground-state is the singlet state,

$$\frac{|\psi_3\rangle - |\psi_4\rangle}{\sqrt{2}} = \frac{|1010\rangle - |0101\rangle}{\sqrt{2}} = \frac{|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle}{\sqrt{2}},$$

whose energy is

$$E^- = \frac{U}{2} - \sqrt{\frac{U^2}{4} + 4t^2} \simeq -\frac{4t^2}{U}.$$

The latter approximation is true if $U \gg t$ (strong coupling limit). The singlet state pairs with a spatially-symmetric (node-less) wavefunction, while triplet states with an antisymmetric wavefunction (which, instead, has a node). The entire triplet (second row of Tab. A.1) has zero energy. Excited states are anti-symmetrised and symmetrised version of the polarised states $|\psi_1\rangle$ and $|\psi_6\rangle$.

A.2 Direct exchange and virtual hopping

The fact that we limited to two sites allowed for an exact diagonalisation of the toy hamiltonian of Eq. (1.2). When moving to lattices, the above procedure cannot be extended easily. By using perturbation theory, instead, we can obtain the same result and easily extend to larger lattices.

Nevertheless, let us obtain the same result of Sec. A.1 by the means of perturbation theory. Consider Eq. (1.2) separating two parts,

$$\hat{H} = \hat{H}_0 + \hat{H}_1,$$

where

$$\begin{aligned} \hat{H}_0 &\equiv U (\hat{n}_{1\uparrow}\hat{n}_{1\downarrow} + \hat{n}_{2\uparrow}\hat{n}_{2\downarrow}), \\ \hat{H}_1 &\equiv -t (\hat{c}_{1\uparrow}^\dagger \hat{c}_{2\uparrow} + \hat{c}_{1\downarrow}^\dagger \hat{c}_{2\downarrow} + \text{h.c.}). \end{aligned}$$



Figure A.2 | Schematisation of a second order virtual hopping. Note that this is a virtual process induced by the kinetic perturbation, not an actual physical process.

At zeroth order in perturbation theory, thus considering just the Coulomb repulsion $\hat{H}_0$, the Hilbert space separates in two manifolds

$$\begin{aligned} \text{Span}(|\psi_1\rangle, |\psi_3\rangle, |\psi_4\rangle, |\psi_6\rangle) & \quad \text{with} \quad E = 0, \\ \text{Span}(|\psi_2\rangle, |\psi_5\rangle) & \quad \text{with} \quad E = U. \end{aligned}$$

We limit to the ground manifold, thus “excluding from the Hilbert space” the states with double occupation $|\psi_2\rangle, |\psi_5\rangle$.

First order corrections to energy are null:

$$\forall i, j \quad : \quad \langle \psi_i | \hat{H}_1 | \psi_j \rangle = 0,$$

because the hopping operator $\hat{H}_1$ produces a doubly occupied site when applied to $|\psi_3\rangle, |\psi_4\rangle$ and zero when applied to $|\psi_1\rangle, |\psi_6\rangle$ due to Pauli exclusion principle.

We need to move to second order corrections. Let us change basis, and define the singlet state

$$|s\rangle \equiv \frac{|\psi_3\rangle - |\psi_4\rangle}{\sqrt{2}},$$

and the triplet states

$$|t_+\rangle \equiv |\psi_1\rangle, \quad |t_0\rangle \equiv \frac{|\psi_3\rangle + |\psi_4\rangle}{\sqrt{2}}, \quad |t_-\rangle \equiv |\psi_6\rangle.$$

Let also $E_\psi^{(0)}$ be the zeroth-order energy associated to state $|\psi\rangle$. Second order correction to the singlet state energy is given by

$$\begin{aligned} E_s^{(2)} &= \sum_{\psi \neq s} \frac{\langle s | \hat{H}_1 | \psi \rangle \langle \psi | \hat{H}_1 | s \rangle}{E_s^{(0)} - E_\psi^{(0)}} \\ &= \frac{2t^2}{E_s^{(0)} - E_{\psi_2}^{(0)}} + \frac{2t^2}{E_s^{(0)} - E_{\psi_5}^{(0)}} = -\frac{4t^2}{U}. \end{aligned}$$

The above result can be obtained easily by direct computation. Note that the + sign in the middle is obtained as a combination of the internal – sign in the definition of the singlet state, and another – sign arising from Fermi anticommutation rules.

Instead, for the central triplet state

$$\begin{aligned} E_{t_0}^{(2)} &= \sum_{\psi \neq t_0} \frac{\langle t_0 | \hat{H}_1 | \psi \rangle \langle \psi | \hat{H}_1 | t_0 \rangle}{E_{t_0}^{(0)} - E_\psi^{(0)}} \\ &= \frac{2t^2}{E_{t_0}^{(0)} - E_{\psi_2}^{(0)}} - \frac{2t^2}{E_{t_0}^{(0)} - E_{\psi_5}^{(0)}} = 0. \end{aligned}$$

This time, the central – sign is present precisely because of the internal sign of the state $|t_0\rangle$. Finally,

$$E_{t_\pm}^{(2)} = \sum_{\psi \neq t_0} \frac{\langle t_\pm | \hat{H}_1 | \psi \rangle \langle \psi | \hat{H}_1 | t_\pm \rangle}{E_{t_\pm}^{(0)} - E_\psi^{(0)}} = 0,$$

again, because of Pauli exclusion principle.

What makes $|s\rangle$ and $|t_0\rangle$ fundamentally different? The singlet state takes advantage of the second-order virtual process where one spin hops to the neighbouring site and returns back. This

process is schematised in Fig. A.2, and is referred to as “virtual hopping”. But each one of the two electrons can do this. The antisymmetry properties of many-body electronic wavefunctions imply, in the end, that these two virtual processes interfere constructively. For the triplet state $|t_0\rangle$, the two processes are perfectly plausible but interfere destructively. The coupling of electronic spin on neighbouring atoms is known as “direct exchange”.

This perturbative scheme scales up to larger lattices. It can be shown analytically that the low-energy sector of the strong-coupled Hubbard hamiltonian of Eq. (1.1) is approximated by an effective Heisenberg hamiltonian with antiferromagnetic spin-spin coupling [28].

A.3 Superexchange in cuprates

The scheme of direct exchange explains the insurgence of antiferromagnetism in the Hubbard model at strong coupling, $U \gg t$. The physics of cuprates is slightly more complicated.

Indeed, in the copper oxides active layers, the Cu^{2+} ions are surrounded by oxygens in a closed shell configuration. Hence, copper $d_{x^2-y^2}$ -wave orbitals (where conduction electrons reside) do not overlap. Instead, they overlap with p -wave orbitals of the neighbouring oxygens. This enables a mechanism known as “superexchange”: even if the copper spins cannot interact directly, they use the intermediate p -orbitals as a sort of “bridge” and establish an direct exchange mechanism with the intermediate spins. The resulting interaction, as is “seen” by the copper electrons, is again an effective antiferromagnetic coupling [45].

Appendix B

Gaussian states

In this appendix we discuss the properties of the many-body states we use to approximate the real ground-state of the system within HF method. We introduce Slater determinants to grasp the general idea behind HF variational method, and then extend to a larger set of states.

B.1 Slater determinants

A many-body wavefunction describing a collective electronic state needs to be antisymmetric under interchange of coordinates and spins $(\mathbf{r}_i, \sigma_i)$, $(\mathbf{r}_j, \sigma_j)$ of any couple of electrons. The simplest way to build a many-body wavefunction satisfying this rule is “to anti-symmetrise” a set of orthonormal N one-body spin-orbitals, being N the total number of electrons and D the total number of spin-orbitals. Let $\{\psi_i\}$ be said set,

$$\frac{1}{2\mathcal{V}} \sum_{\sigma \in \{\uparrow, \downarrow\}} \int d\mathbf{r} \psi_i(\mathbf{r}, \sigma) \psi_j(\mathbf{r}, \sigma) = \delta_{ij} ,$$

being $\mathcal{V}$ the total volume.

Let S_N be the $N!$ elements symmetric group of permutations of N objects and $P \in S_N$ a specific permutation of a given N -uple,

$$P : (x_1, x_2, \dots, x_N) \rightarrow (x_{P(1)}, x_{P(2)}, \dots, x_{P(N)}) .$$

Let F_P be the operator that implements the permutation P on the N variables of a function f ,

$$F_P [f(x_1, x_2, \dots, x_N)] = f(x_{P(1)}, x_{P(2)}, \dots, x_{P(N)}) .$$

We define the anti-symmetrisation operator:

$$\mathcal{A}[f] \equiv \frac{1}{\sqrt{N!}} \sum_{P \in S_N} (-1)^{\sigma(P)} F_P[f] ,$$

being $\sigma(P)$ the “parity factor” of the permutation $P \in S_N$. The factor $\sigma(P)$ counts the number of exchanges needed to obtain P : this way, an even number of exchanges gives a total sign $+1$, an odd number gives -1 . By construction, the function $\mathcal{A}[f]$ is antisymmetric under exchange of a pair of coordinates.

Consider now the many-body wavefunction

$$\Psi(\mathbf{r}_1, \sigma_1; \mathbf{r}_2, \sigma_2; \dots; \mathbf{r}_N, \sigma_N) = \mathcal{A} \left[\prod_{i=1}^N \psi_i(\mathbf{r}_i, \sigma_i) \right] .$$

This class of wavefunctions, obtained as antisymmetric combinations of product states, are representable as so-called Slater determinants,

$$\Psi(\mathbf{r}_1, \sigma_1; \mathbf{r}_2, \sigma_2; \dots; \mathbf{r}_N, \sigma_N) = \frac{1}{\sqrt{N!}} \det \begin{bmatrix} \psi_1(\mathbf{r}_1, \sigma_1) & \psi_1(\mathbf{r}_2, \sigma_2) & \cdots & \psi_1(\mathbf{r}_N, \sigma_N) \\ \psi_2(\mathbf{r}_1, \sigma_1) & \psi_2(\mathbf{r}_2, \sigma_2) & \cdots & \psi_2(\mathbf{r}_N, \sigma_N) \\ \vdots & \vdots & & \vdots \\ \psi_N(\mathbf{r}_1, \sigma_1) & \psi_N(\mathbf{r}_2, \sigma_2) & \cdots & \psi_N(\mathbf{r}_N, \sigma_N) \end{bmatrix} .$$

This representation includes automatically Pauli exclusion principle: determinants of matrices with two equal rows or two equal columns are null. Thus, all spin-orbitals must differ, as well as no single spin-orbital ψ_i can be doubly occupied (which is, all spin-coordinate couples must be different).

Any normalised combinations of Slater determinants gives an antisymmetric wavefunction, although not necessarily representable as a single Slater determinant. In HF theory, we search for the best possible choice of orthonormal one-body spin orbitals $\{\psi_1, \psi_2, \dots, \psi_N\}$ to approximate the true ground-state of the model through a single Slater determinant. The space of possible “Slater-like” many-body wavefunctions is the parametric space where we perform the variational constrained search.

The EHM is formulated in second-quantisation. It is then useful to represent Slater-like states in this language. Let us map all the spin-orbitals from the discussion above to pairs of annihilation-creation fermionic operators,

$$\psi_i \rightarrow \hat{c}_i, \hat{c}_i^\dagger.$$

Any Slater state can be expressed in second-quantisation as:

$$|\Psi\rangle = \prod_{i=1}^N \hat{c}_i^\dagger |\Omega\rangle, \quad (\text{B.1})$$

being $|\Omega\rangle$ the vacuum. Antisymmetry is guaranteed by Fermi anticommutation rules.

B.2 Wick’s theorem applied to Slater determinants

The concept of vacuum state needs to be interpreted in relation to an appropriate choice of second-quantisation annihilation and creation operators. For a given set of fermion operators, the vacuum is that state mapped to zero by any annihilation operator. As we will see, to correctly define the vacuum state is crucial in order to apply Wick’s theorem and develop an HF approximation of EHM. Before moving to generalisation, let us discuss a playground setup.

Consider some unspecified model whose ground state at null temperature is given in Eq. (B.1) for a given set of electronic operators $\hat{c}_i, \hat{c}_i^\dagger$. Let $\hat{c}_i$ with $N < i < D$ identify all spin-orbitals not already included in the definition of $|\Psi\rangle$ of Eq. (B.1) (thus, non populated). Then it holds:

$$\begin{aligned} 0 \leq i \leq N & \quad \hat{c}_i^\dagger |\Psi\rangle = 0, \\ i > N & \quad \hat{c}_i |\Psi\rangle = 0. \end{aligned}$$

Let:

$$\hat{a}_i \equiv \begin{cases} \hat{c}_i^\dagger & \text{for } 0 \leq i \leq N, \\ \hat{c}_i & \text{for } i > N. \end{cases}$$

These new operators satisfy a Fermi algebra $\{\hat{a}_i, \hat{a}_j^\dagger\} = \delta_{ij}$ and annihilate the “vacuum” state $|\Psi\rangle$. In other words, any Slater state like Eq. (B.1) can be seen as a vacuum state for a suitably chosen set of fermionic operators.

This property is handy for evaluation of expectation values (EV). Let $\hat{O}^{(\ell)}$ be a linear combination of $\hat{c}_i$ or $\hat{c}_i^\dagger$,

$$\hat{O}^{(\ell)} \equiv \sum_{i=1}^D \left(u_i^{(\ell)} \hat{c}_i + v_i^{(\ell)} \hat{c}_i^\dagger \right) \quad \text{where} \quad u_i^{(\ell)}, v_i^{(\ell)} \in \mathbb{C}. \quad (\text{B.2})$$

Consider the product operator:

$$\hat{O} \equiv \hat{O}^{(1)} \hat{O}^{(2)} \dots \hat{O}^{(2k)} \quad \text{where} \quad k \in \mathbb{N}.$$

Inserting the above decomposition, we get a large sum of many terms, each containing some sequence of annihilations and creations. Using linearity we can concentrate on one specific term of these, derive results and then recompose them for the original operator. Consider one of these terms,

$$\hat{O} = \dots + (\text{coefficient}) \times \hat{C}_1 \hat{C}_2 \dots \hat{C}_{2k} + \dots,$$

with $\hat{C}_i$ some fermionic operator, either $\hat{c}_j$ or $\hat{c}_j^\dagger$ for some j . Now, Wick's theorem states that

$$\begin{aligned} \hat{C}_1 \hat{C}_2 \cdots \hat{C}_{2k} &= N \left\{ \hat{C}_1 \hat{C}_2 \cdots \hat{C}_{2k} \right\} \\ &+ \sum_{(\alpha\beta)} (-1)^{\zeta_{\alpha\beta}} D_{\alpha\beta} N \left\{ \prod_{i \neq \alpha, \beta} \hat{C}_i \right\} \\ &+ \sum_{(\alpha\beta)(\gamma\delta)} (-1)^{\zeta_{\alpha\beta}} D_{\alpha\beta} (-1)^{\zeta_{\gamma\delta}} D_{\gamma\delta} N \left\{ \prod_{i \neq \alpha, \beta, \gamma, \delta} \hat{C}_i \right\} \\ &+ \cdots, \end{aligned}$$

where $N\{\cdots\}$ indicates normal ordering, $D_{\alpha\beta}$ is the contraction of the pair $\hat{C}_\alpha \hat{C}_\beta$ and $\zeta_{\alpha\beta}$ is a symmetry factor included to take care of the number of fermionic algebra. In this context contractions are just expectation values over $|\Psi\rangle$. Sums are performed over pairs of indices, implicitly taken to be different from each other. To take averages of this last operator over $|\Psi\rangle$ is a not-so-hard task, given the simple expression of the state in terms of electronic annihilations and creations. However it can be made simpler: if we change basis and express all operators in terms of $\hat{a}$, $\hat{a}^\dagger$,

$$\hat{C}_i \rightarrow \hat{A}_i,$$

the above decomposition holds identically, with slight change of notation. Now, taking a zero temperature average on $|\Psi\rangle$ (the vacuum with respect to the new operators) all normal-ordered terms disappear,

$$\langle \Psi | N\{\hat{A}_i \hat{A}_j \cdots\} | \Psi \rangle = 0,$$

thus in the decomposition above only fully contracted terms survive. Going back to $\hat{c}$ operators, we can re-express fully contracted terms in the original basis. Thus:

$$\langle \hat{C}_1 \hat{C}_2 \cdots \hat{C}_{2k} \rangle = \sum (\text{total sign factor}) \times (\text{product of complete contractions}),$$

where sum is intended, implicitly, over all possible ways of choosing k couples of indices from a starting set of $2k$ elements.

Taking another step back, due to linearity we can apply this result to all terms of $\mathcal{O}$ and conclude that the EV of $\mathcal{O}$ decomposes as a sum of 2-points EVs. This can be seen as the defining property of gaussian states. In this procedure, the only key property of Slater states we used was the fact that they are vacuum to some fermionic basis. Finite-temperature extension of Wick's theorem as well as more complicated change of basis are possible, but the concept remains the same: on a Slater state, which is essentially a non-interacting state for some set of fermions, Wick's theorem is particularly well-behaved.

From the point above we can conclude that:

$$\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta \rangle = \langle \hat{c}_\alpha^\dagger \hat{c}_\delta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\gamma \rangle - \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \langle \hat{c}_\gamma^\dagger \hat{c}_\delta \rangle,$$

being the average intended on a Slater state. In this context, only contractions with an equal number of creation and annihilation operators contribute.

B.3 Generalisation to Bogoliubov-Valatin states

In HF theory we look for a good approximation of the true many-body wavefunction constraining search to a class of states. Introducing Slater determinants was useful to grasp the concrete idea behind the method. However, in order to treat some physical phenomena we need to extend the variational procedure to a other sets of states. For example, Bardeen-Cooper-Shrieffer (BCS) theory of superconductivity gives a many-body ground-state which is not representable as a Slater determinant of "standard" electronic operators. In this section we extend the notions introduced before and set grounds for the generalised Hartree-Fock-Bogoliubov (HFB) method we use in this thesis.

Consider the Nambu spinor

$$\hat{\Phi} \equiv \begin{bmatrix} \hat{\mathbf{c}} \\ \hat{\mathbf{c}}^\dagger \end{bmatrix},$$

being $\hat{\mathbf{c}}$ the vector of all annihilation operators, ordered, and $\hat{\mathbf{c}}^\dagger$ of all creation operators. Let us assume the vectors have D entries. Consider an unitary map M , such that

$$\hat{\Gamma} \equiv M\hat{\Phi} \quad \text{being} \quad \hat{\Gamma} \equiv \begin{bmatrix} \hat{\gamma} \\ \hat{\gamma}^\dagger \end{bmatrix}.$$

The new Nambu vectors $\hat{\gamma}$ and $\hat{\gamma}^\dagger$ contain our new, generalised fermionic operators in second quantisation. Let:

$$M \equiv \begin{bmatrix} U & V \\ U^\dagger & V^\dagger \end{bmatrix}.$$

If M is unitary, it follows:

$$\begin{aligned} \mathbb{I}_{2D \times 2D} &= MM^\dagger \\ &= \begin{bmatrix} U & V \\ U^\dagger & V^\dagger \end{bmatrix} \begin{bmatrix} U & U^\dagger \\ V^\dagger & V \end{bmatrix} \\ &= \begin{bmatrix} UU^\dagger + VV^\dagger & U^2 + V^2 \\ (U^\dagger)^2 + (V^\dagger)^2 & U^\dagger U + V^\dagger V \end{bmatrix}, \end{aligned}$$

thus, among other conditions, $UU^\dagger + VV^\dagger = \mathbb{I}_{D \times D}$.

The new operators in $\hat{\Gamma}$ satisfy Fermi anticommutation rules. In fact, writing the mapping explicitly for any of the new operators and some index $i \in [1, D]$,

$$\hat{\gamma}_i \equiv \sum_{j=1}^D U_{ij} \hat{c}_j + \sum_{j=1}^D V_{ij} \hat{c}_j^\dagger \quad \text{and} \quad \hat{\gamma}_i^\dagger \equiv \sum_{j=1}^D U_{ji}^* \hat{c}_j^\dagger + \sum_{j=1}^D V_{ji}^* \hat{c}_j,$$

and substituting in the anti-commutator

$$\begin{aligned} \{\hat{\gamma}_i, \hat{\gamma}_j^\dagger\} &= \sum_{i'=1}^D U_{ii'} \sum_{j'=1}^D U_{j'j}^* \{\hat{c}_{i'}, \hat{c}_{j'}^\dagger\} + \sum_{i'=1}^D V_{ii'} \sum_{j'=1}^D V_{j'j}^* \{\hat{c}_{i'}^\dagger, \hat{c}_{j'}\} \\ &= \sum_{i'=1}^D U_{ii'} \sum_{j'=1}^D U_{j'j}^* \delta_{i'j'} + \sum_{i'=1}^D V_{ii'} \sum_{j'=1}^D V_{j'j}^* \delta_{i'j'} \\ &= [UU^\dagger + VV^\dagger]_{ij} \\ &= \delta_{ij}, \end{aligned}$$

being M unitary.

Assume now $|\Psi\rangle$ to be the vacuum state for the new operators,

$$\forall i \quad : \quad \hat{\gamma}_i |\Psi\rangle = 0.$$

Suppose we want to evaluate, with identical meaning of the notation as in previous section, something like

$$\langle \hat{C}_1 \hat{C}_2 \cdots \hat{C}_{2k} \rangle.$$

We can apply the result obtained above. Indeed, inverse maps linking old and new operators read:

$$\hat{c}_i \equiv \sum_{j=1}^D U_{ij} \hat{\gamma}_j + \sum_{j=1}^D U_{ji}^* \hat{\gamma}_j^\dagger \quad \text{and} \quad \hat{c}_i^\dagger \equiv \sum_{j=1}^D V_{ji}^* \hat{\gamma}_j^\dagger + \sum_{j=1}^D V_{ij} \hat{\gamma}_j.$$

In Sec. B.2 we argued that, when averaging a product of operators like the one of Eq. (B.2), only 2-points EVs contribute. This time the idea is the same, substituting $\mathcal{O}^{(\ell)} \rightarrow \hat{C}_\ell$ and $\hat{c}_i, \hat{c}_i^\dagger \rightarrow \hat{\gamma}_i, \hat{\gamma}_i^\dagger$.

Thus, also here we can decompose the product operator, apply Wick's theorem, discard all non-fully-contracted terms and recompose the results. All we used was just the unitary mapping between $\hat{\Phi}$ and $\hat{\Gamma}$, the fact that $\gamma_i, \hat{\gamma}_i^\dagger$ obey Fermi anticommutation rules and “see” $|\Psi\rangle$ as a vacuum state.

As in Sec. B.2, we conclude that:

$$\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\gamma \hat{c}_\delta \rangle = \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\delta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\gamma \rangle}_{\text{Hartree}} - \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \langle \hat{c}_\gamma^\dagger \hat{c}_\delta \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \langle \hat{c}_\gamma \hat{c}_\delta \rangle}_{\text{Cooper}}. \quad (\text{B.3})$$

The subscripts indicate the name we give to these terms throughout the text. Note that, differently from “pure Slater states” EVs with two annihilations or two creations do not vanish, in general.

All of this holds when averaging is performed on a state, $|\Psi\rangle$, which is vacuum for a set of operators linked to fermionic Nambu spinor $\hat{\Phi}$ by some unitary operation. This set of states is the domain of our variational constrained search. This kind of unitary mapping is often referred to as Bogoliubov-Valatin transformation, and the method hereby presented as Hartree-Fock-Bogoliubov. For simplicity, in this thesis we stick to the name “HF method” referring to this discussion. This argument has been carried out at zero temperature, but can be extended to finite temperature.

Appendix C

Connection with mean-field theories

In this appendix we bridge the gap between HF methods and mean-field theory (MFT). In general, in MFT approaches we substitute the two-body (or more) interactions of the hamiltonian with a single-body interaction with a mean field, determined locally via self-consistency equations. This is, for example, the Weiß mean-field solution to the quantum Ising model. The HF method we sketched here does not work like that. In the context of Hubbard models, the analogous of Weiß MFT would be dynamical mean-field theory (DMFT) [13].

However, there are some points of connection. Let us sketch first the general scheme behind MFT.

C.1 Mean-field theory

Suppose we have some operator $\hat{A} = \hat{B}\hat{C}$. Let

$$\delta\hat{O} \equiv \hat{O} - \langle\hat{O}\rangle \quad \text{with} \quad \hat{O} \in \{\hat{A}, \hat{B}, \hat{C}\}.$$

Then we have:

$$\begin{aligned} \hat{A} &= (\langle\hat{B}\rangle + \delta\hat{B}) (\langle\hat{C}\rangle + \delta\hat{C}) \\ &= \langle\hat{B}\rangle\langle\hat{C}\rangle + \langle\hat{B}\rangle\delta\hat{C} + \delta\hat{B}\langle\hat{C}\rangle + \delta\hat{B}\delta\hat{C} \\ &= \langle\hat{B}\rangle\langle\hat{C}\rangle + \langle\hat{B}\rangle(\hat{C} - \langle\hat{C}\rangle) + (\hat{B} - \langle\hat{B}\rangle)\langle\hat{C}\rangle + \delta\hat{B}\delta\hat{C} \\ &= -\langle\hat{B}\rangle\langle\hat{C}\rangle + \langle\hat{B}\rangle\hat{C} + \hat{B}\langle\hat{C}\rangle + \delta\hat{B}\delta\hat{C}. \end{aligned}$$

Up to now, this relation is exact. MFT relies on the idea that the last term, of order $\mathcal{O}(\delta^2)$, can be neglected. This amounts to say that the operator fluctuations about its average are “small”. This gives the approximation:

$$\hat{A} \simeq \langle\hat{B}\rangle\hat{C} + \hat{B}\langle\hat{C}\rangle - \langle\hat{B}\rangle\langle\hat{C}\rangle. \quad (\text{C.1})$$

Within the assumption that independently $\hat{B}$ and $\hat{C}$ fluctuations are “small”, we are able to simplify $\hat{A}$ with a sum of three terms: the product of each operator with the averaged value of the other, minus the product of the averages.

In the context of HF methods, we are in an slightly different situation. We are not making assumptions on the fluctuations of the quartic sectors of the hamiltonian. We are restricting the space of wavefunctions and minimising energy there.

C.2 Hartee-Fock methods

Let us restart from the interacting hamiltonian of Eq. (2.2). In this thesis we deal with the EHM, thus (limitedly to this section) let us restrict to the two-body generic quartic interaction:

$$\hat{V} = \sum_{\alpha,\beta} V_{\alpha\beta} \hat{c}_{\alpha}^{\dagger} \hat{c}_{\beta}^{\dagger} \hat{c}_{\beta} \hat{c}_{\alpha},$$

with $\forall \alpha, \beta: V_{\alpha\beta} = V_{\beta\alpha}^*$. In terms of the general hamiltonian of Eq. (2.2), we have:

$$B_{\alpha\beta\gamma\delta} = 2V_{\alpha\beta}\delta_{\alpha\delta}\delta_{\beta\gamma}. \quad (\text{C.2})$$

We work with HF states. Thus Eq. (2.1) holds exactly (adapted for two indices instead of four)

$$\alpha \neq \beta \quad : \quad \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\beta \hat{c}_\alpha \rangle = \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle}_{\text{Hartree}} - \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \langle \hat{c}_\beta \hat{c}_\alpha \rangle}_{\text{Cooper}}.$$

This gives:

$$\begin{aligned} \langle \hat{V} \rangle &= \sum_{\alpha, \beta} V_{\alpha\beta} \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\beta \hat{c}_\alpha \rangle \\ &= \sum_{\alpha, \beta} V_{\alpha\beta} \left(\langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle - \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \langle \hat{c}_\beta \hat{c}_\alpha \rangle \right), \end{aligned} \quad (\text{C.3})$$

from which we have the equations:

$$\frac{\partial \langle \hat{V} \rangle}{\partial \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle} = \delta_{\alpha\beta} \left[\sum_{\gamma} V_{\alpha\gamma} \langle \hat{c}_\gamma^\dagger \hat{c}_\gamma \rangle \right] - V_{\alpha\beta} \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle, \quad (\text{C.4})$$

$$\frac{\partial \langle \hat{V} \rangle}{\partial \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle} = V_{\alpha\beta} \langle \hat{c}_\beta \hat{c}_\alpha \rangle. \quad (\text{C.5})$$

Now, our aim is to approximate $\hat{V}$ as:

$$\hat{V} \simeq \sum_{\alpha\beta} \left[x_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta + y_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger + y_{\beta\alpha}^* \hat{c}_\alpha \hat{c}_\beta \right] + (\text{some constant}), \quad (\text{C.6})$$

where $x_{\alpha\beta} = x_{\beta\alpha}^*$ to ensure hermiticity. Note that we are actively ignoring terms like $\hat{c}\hat{c}^\dagger$. Anti-commuting these, we get sums of operators like $\hat{c}^\dagger \hat{c}$ and constants. The above expression is the most general. Take its EV:

$$\langle \hat{V} \rangle \simeq \sum_{\alpha\beta} \left[x_{\alpha\beta} \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle + y_{\alpha\beta} \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle + y_{\beta\alpha}^* \langle \hat{c}_\alpha \hat{c}_\beta \rangle \right] + (\text{some constant}).$$

Deriving the equation above, we obtain:

$$x_{\alpha\beta} = \frac{\partial \langle \hat{V} \rangle}{\partial \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle} \quad \text{and} \quad y_{\alpha\beta} = \frac{\partial \langle \hat{V} \rangle}{\partial \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle}.$$

Applying Eqs. (C.4) and (C.5), we get the identities:

$$x_{\alpha\beta} = \delta_{\alpha\beta} \left[\sum_{\gamma} V_{\alpha\gamma} \langle \hat{c}_\gamma^\dagger \hat{c}_\gamma \rangle \right] - V_{\alpha\beta} \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle, \quad (\text{C.7})$$

$$y_{\alpha\beta} = V_{\alpha\beta} \langle \hat{c}_\beta \hat{c}_\alpha \rangle. \quad (\text{C.8})$$

These equations are a rephrasing of Eqs. (2.15), (2.16) for the fields $h_{\alpha\beta}^{pq}$, setting $A_{\alpha\beta} = 0$. This can be checked by using Eq. (C.2).

Now: consider the special case where $\forall \alpha: V_{\alpha\alpha} = 0$. This is the case for the EHM. Then the above equations imply:

$$\begin{aligned} \hat{V} &= \sum_{\alpha \neq \beta} V_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\beta \hat{c}_\alpha \\ &\simeq \sum_{\alpha \neq \beta} V_{\alpha\beta} \left(\underbrace{\hat{c}_\alpha^\dagger \hat{c}_\alpha \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \hat{c}_\beta^\dagger \hat{c}_\beta}_{\text{Hartree}} - \underbrace{\hat{c}_\alpha^\dagger \hat{c}_\beta \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle - \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \hat{c}_\beta^\dagger \hat{c}_\alpha}_{\text{Fock}} \right. \\ &\quad \left. + \underbrace{\hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \langle \hat{c}_\beta \hat{c}_\alpha \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \hat{c}_\beta \hat{c}_\alpha}_{\text{Cooper}} \right) + (\text{some constant}). \end{aligned}$$

To determine the constant, we take the EV of the above equation. The result should equal Eq. (C.3). Take e.g. the Hartree sector:

$$\langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle + C = \langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle \implies C = -\langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle,$$

with C a constant. A similar implication holds for the Fock and the Cooper sectors. Then:

$$\begin{aligned} \hat{V} &= \sum_{\alpha \neq \beta} V_{\alpha\beta} \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \hat{c}_\beta \hat{c}_\alpha & (C.9) \\ &\stackrel{\text{HF}}{\simeq} \sum_{\alpha \neq \beta} V_{\alpha\beta} \left(\hat{c}_\alpha^\dagger \hat{c}_\alpha \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \hat{c}_\beta^\dagger \hat{c}_\beta - \langle \hat{c}_\alpha^\dagger \hat{c}_\alpha \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\beta \rangle \right. & (\text{Hartree}) \\ &\quad - \hat{c}_\alpha^\dagger \hat{c}_\beta \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle - \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \hat{c}_\beta^\dagger \hat{c}_\alpha + \langle \hat{c}_\alpha^\dagger \hat{c}_\beta \rangle \langle \hat{c}_\beta^\dagger \hat{c}_\alpha \rangle & (\text{Fock}) \\ &\quad \left. + \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \langle \hat{c}_\beta \hat{c}_\alpha \rangle + \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \hat{c}_\beta \hat{c}_\alpha - \langle \hat{c}_\alpha^\dagger \hat{c}_\beta^\dagger \rangle \langle \hat{c}_\beta \hat{c}_\alpha \rangle \right). & (\text{Cooper}) \end{aligned}$$

Passing from the first row to the following ones, we decomposed the original quartic operator to a sum of three decomposition much similar to Eq. (C.1). This is the main connection with MFT. In HF theory, we do not write the quartic operator as a product of two quadratic operators and decompose it as we would do in MFT. Instead, a very similar decomposition (summed over the three sectors) is given by the fact that EVs are computed over HF states.

Appendix D

Antiferromagnetism in the HM

In this appendix we give some details about the antiferromagnetic phase in the conventional Hubbard model,

$$\hat{H} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \left[\hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + \text{h.c.} \right] + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \quad \text{where } t, U > 0, \quad (\text{D.1})$$

approximated by HF methods. We use the results discussed in Sec. ??.

D.1 The AF ground-state for the conventional HM

Let the Nambu spinors:

$$\hat{\Psi}_{\mathbf{k}\sigma} \equiv \begin{bmatrix} \hat{c}_{\mathbf{k}\sigma} \\ \hat{c}_{\mathbf{k}+\pi\sigma} \end{bmatrix}.$$

Let also the gap function:

$$\Delta_{\mathbf{k}\sigma} = (-1)^{\delta_{\sigma=\downarrow}} mU.$$

We know from Eq. (5.41) that the approximated hamiltonian is:

$$\hat{H}_t + \hat{H}_U \simeq \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \mathbf{b}_{\mathbf{k}\sigma} \cdot \hat{\mathbf{s}}_{\mathbf{k}\sigma} + nU\hat{N} - E_{\text{H}/U}^{(\text{AF})},$$

where:

$$\mathbf{b}_{\mathbf{k}\sigma} \equiv \begin{bmatrix} -\Delta_{\mathbf{k}\sigma} \\ 0 \\ \epsilon_{\mathbf{k}\sigma} \end{bmatrix} \quad \hat{s}_{\mathbf{k}\sigma}^{\alpha} \equiv \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} \tau^{\alpha} \hat{\Psi}_{\mathbf{k}\sigma} \quad \hat{\mathbf{s}}_{\mathbf{k}\sigma} \equiv \begin{bmatrix} \hat{s}_{\mathbf{k}\sigma}^x \\ \hat{s}_{\mathbf{k}\sigma}^y \\ \hat{s}_{\mathbf{k}\sigma}^z \end{bmatrix}.$$

Through the unitary matrix $W_{\mathbf{k}\sigma}$ given in Eq. (5.44), the hamiltonian is mapped to a gas of non-interacting fermions collected in the $\hat{\Gamma}$ spinors of Eq. (5.45),

$$\hat{\Gamma}_{\mathbf{k}\sigma} = W_{\mathbf{k}\sigma} \hat{\Psi}_{\mathbf{k}\sigma} = \begin{bmatrix} \hat{\gamma}_{\mathbf{k}\sigma}^{(+)} \\ \hat{\gamma}_{\mathbf{k}\sigma}^{(-)} \end{bmatrix}.$$

The final non-interacting hamiltonian is:

$$\hat{H}_t + \hat{H}_U \simeq \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} E_{\mathbf{k}\sigma} \left(\hat{\gamma}_{\mathbf{k}\sigma}^{(+)\dagger} \hat{\gamma}_{\mathbf{k}\sigma}^{(+)} - \hat{\gamma}_{\mathbf{k}\sigma}^{(-)\dagger} \hat{\gamma}_{\mathbf{k}\sigma}^{(-)} \right) + nU\hat{N} - E_{\text{H}/U}^{(\text{AF})},$$

with $E_{\mathbf{k}\sigma}$ the Bogoliubov bands of Eq. (5.43). In next section we give the explicit form of the many-body ground-state at zero temperature and half-filling. From now on, limitedly to this appendix, we use the simplified notation:

$$\Delta_{\mathbf{k}\uparrow} \rightarrow \Delta,$$

with $\Delta_{\mathbf{k}\uparrow} = -\Delta_{\mathbf{k}\downarrow}$, and $\Delta = mU$.

D.2 Explicit AF ground state at half-filling and $\beta \rightarrow \infty$

Let $|\uparrow_{\mathbf{k}\sigma}\rangle$ and $|\downarrow_{\mathbf{k}\sigma}\rangle$ be respectively the “up” and “down” $\mathbf{k}$ -th pseudo-spin states. Let $|\overline{\downarrow}_{\mathbf{k}\sigma}\rangle$ be the “down” state in the tilted frame. The ground state is given by

$$|\Psi\rangle \equiv \bigotimes_{\sigma} \bigotimes_{\mathbf{k} \in \text{MBZ}} |\overline{\downarrow}_{\mathbf{k}\sigma}\rangle. \quad (\text{D.2})$$

We can link the “down” states via an inverse rotation,

$$\begin{aligned} |\overline{\downarrow}_{\mathbf{k}\sigma}\rangle &= e^{-i\theta_{\mathbf{k}\sigma} \hat{s}_{\mathbf{k}\sigma}^y} |\downarrow_{\mathbf{k}\sigma}\rangle \\ &= \cos \theta_{\mathbf{k}\sigma} |\downarrow_{\mathbf{k}\sigma}\rangle - i \sin \theta_{\mathbf{k}\sigma} \hat{s}_{\mathbf{k}\sigma}^y |\downarrow_{\mathbf{k}\sigma}\rangle \\ &= \cos \theta_{\mathbf{k}\sigma} |\downarrow_{\mathbf{k}\sigma}\rangle - \sin \theta_{\mathbf{k}\sigma} (\hat{s}_{\mathbf{k}\sigma}^+ - \hat{s}_{\mathbf{k}\sigma}^-) |\downarrow_{\mathbf{k}\sigma}\rangle \\ &= \cos \theta_{\mathbf{k}\sigma} |\downarrow_{\mathbf{k}\sigma}\rangle - \sin \theta_{\mathbf{k}\sigma} |\uparrow_{\mathbf{k}\sigma}\rangle, \end{aligned} \quad (\text{D.3})$$

having we used

$$\hat{s}_{\mathbf{k}\sigma}^{\pm} \equiv \frac{\hat{s}_{\mathbf{k}\sigma}^x \pm i \hat{s}_{\mathbf{k}\sigma}^y}{2}.$$

From Eqs. (5.40) and (5.39),

$$\frac{\hat{\mathbb{I}}_{\mathbf{k}\sigma} + \hat{s}_{\mathbf{k}\sigma}^z}{2} = \hat{n}_{\mathbf{k}\sigma}, \quad \frac{\hat{\mathbb{I}}_{\mathbf{k}\sigma} - \hat{s}_{\mathbf{k}\sigma}^z}{2} = \hat{n}_{\mathbf{k}+\pi\sigma}.$$

Evaluating the above operators on the “up” and “down” states, we get

$$\begin{aligned} \frac{1}{2} \langle \uparrow_{\mathbf{k}\sigma} | \hat{\mathbb{I}}_{\mathbf{k}\sigma} + \hat{s}_{\mathbf{k}\sigma}^z | \uparrow_{\mathbf{k}\sigma} \rangle &= 1, & \frac{1}{2} \langle \uparrow_{\mathbf{k}\sigma} | \hat{\mathbb{I}}_{\mathbf{k}\sigma} - \hat{s}_{\mathbf{k}\sigma}^z | \uparrow_{\mathbf{k}\sigma} \rangle &= 0, \\ \frac{1}{2} \langle \downarrow_{\mathbf{k}\sigma} | \hat{\mathbb{I}}_{\mathbf{k}\sigma} + \hat{s}_{\mathbf{k}\sigma}^z | \downarrow_{\mathbf{k}\sigma} \rangle &= 0, & \frac{1}{2} \langle \downarrow_{\mathbf{k}\sigma} | \hat{\mathbb{I}}_{\mathbf{k}\sigma} - \hat{s}_{\mathbf{k}\sigma}^z | \downarrow_{\mathbf{k}\sigma} \rangle &= 1. \end{aligned}$$

Thus, up to a phase

$$|\uparrow_{\mathbf{k}\sigma}\rangle \sim \hat{c}_{\mathbf{k}\sigma}^{\dagger} |\Omega_{\mathbf{k}\sigma}\rangle, \quad |\downarrow_{\mathbf{k}\sigma}\rangle \sim \hat{c}_{\mathbf{k}+\pi\sigma}^{\dagger} |\Omega_{\mathbf{k}\sigma}\rangle,$$

being $|\Omega_{\mathbf{k}\sigma}\rangle$ the $(\mathbf{k}, \sigma)$ Hilbert subspace vacuum.

It follows that composing Eqs. (D.2) and (D.3) the ground state $|\Psi\rangle$ can be written as:

$$|\Psi\rangle = \bigotimes_{\sigma} \bigotimes_{\mathbf{k} \in \text{MBZ}} \left[\sin \theta_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma}^{\dagger} + e^{2i\zeta_{\mathbf{k}\sigma}} \cos \theta_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}+\pi\sigma}^{\dagger} \right] |\Omega_{\mathbf{k}\sigma}\rangle, \quad (\text{D.4})$$

having we absorbed in the relative phase $e^{2i\zeta_{\mathbf{k}\sigma}}$ the two states phases, as well as the original sign difference of Eq. (D.3). Now, inserting the explicit form of the ground state from equation above:

$$\begin{aligned} \langle \overline{\downarrow}_{\mathbf{k}\sigma} | \hat{\psi}_{\mathbf{k}\sigma} + \hat{\psi}_{\mathbf{k}\sigma}^{\dagger} | \overline{\downarrow}_{\mathbf{k}\sigma} \rangle &= (e^{2i\zeta_{\mathbf{k}\sigma}} + e^{-2i\zeta_{\mathbf{k}\sigma}}) \sin \theta_{\mathbf{k}\sigma} \cos \theta_{\mathbf{k}\sigma} \\ &= \cos 2\zeta_{\mathbf{k}\sigma} \sin 2\theta_{\mathbf{k}\sigma}. \end{aligned}$$

From Eq. (5.37)

$$\begin{aligned} \langle \overline{\downarrow}_{\mathbf{k}\sigma} | \hat{\psi}_{\mathbf{k}\sigma} + \hat{\psi}_{\mathbf{k}\sigma}^{\dagger} | \overline{\downarrow}_{\mathbf{k}\sigma} \rangle &= \langle \overline{\downarrow}_{\mathbf{k}\sigma} | \hat{s}_{\mathbf{k}\sigma}^x | \overline{\downarrow}_{\mathbf{k}\sigma} \rangle \\ &= \sin 2\theta_{\mathbf{k}\sigma}, \end{aligned}$$

since the pseudofield has no y component. This implies necessarily

$$\cos 2\zeta_{\mathbf{k}\sigma} \implies \zeta_{\mathbf{k}\sigma} = k\pi \quad \text{with } k \in \mathbb{Z}.$$

Using Eq. (D.4), we conclude

$$|\Psi\rangle = \bigotimes_{\sigma} \bigotimes_{\mathbf{k} \in \text{MBZ}} \left[\sin \theta_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}\sigma}^{\dagger} + \cos \theta_{\mathbf{k}\sigma} \hat{c}_{\mathbf{k}+\pi\sigma}^{\dagger} \right] |\Omega_{\mathbf{k}\sigma}\rangle. \quad (\text{D.5})$$

For low-temperature physics, we can expect the ground-state density matrix to be highly dominated by this BCS-like zero-temperature ground-state, which highlights the essential features of the system.

Finally, using Eq. (5.45), we check that correctly the ground-state describes a perfect degenerate $\gamma^{(-)}$ quasiparticles Fermi gas,

$$|\Psi\rangle = \bigotimes_{\sigma} \bigotimes_{\mathbf{k} \in \text{MBZ}} \gamma_{\mathbf{k}\sigma}^{(-)\dagger} |\Omega_{\mathbf{k}\sigma}\rangle. \quad (\text{D.6})$$

The AF phase can be thought of as a long-range coordination between fermionic states related by a π shift in reciprocal space, producing a new free gas where all interactions have been “condensed” in these AF pairs.

D.3 Instability of the commensurate AF phase

The commensurate AF phase is unstable at any finite doping. This result is discussed in [6, 7], and here we recall just the main passages. Consider the local spin operator defined in Eq. (3.7),

$$\hat{S}_j^\alpha = \frac{1}{2} \hat{\Psi}_j^\dagger \tau^\alpha \hat{\Psi}_j, \quad \hat{S}_j^+ \equiv \frac{\hat{S}_j^x + i\hat{S}_j^y}{2}, \quad \hat{S}_j^- \equiv \frac{\hat{S}_j^x - i\hat{S}_j^y}{2}.$$

Let us take a continuum limit, $L_x \rightarrow \infty$ and $L_y \rightarrow \infty$, while the lattice spacing goes to zero.

Continuous SSBs are connected to the presence of a stable gapless Goldstone mode. Such a Goldstone mode is essentially a sharp peak at zero frequency for some dressed response function. For the commensurate AF phase, the low-energy collective excitations are transverse spin fluctuations about the antiferromagnetic ground-state, describing SDWs. In order to see if the Goldstone mode is stable for a given doping δ , we need to investigate whether the respective response function diverges at zero frequency. If it does, we have a gapless Goldstone mode and the phase is stable; if not, the phase is unstable. We work at $\omega = 0$ in order to capture gapless modes.

We work in interaction representation, and indicate the spin operators by

$$\hat{\mathbf{S}}(\mathbf{r}, t)$$

The retarded transverse spin-spin response is given by:

$$\chi^{-+}(\mathbf{r}, t; \mathbf{r}', t') \equiv iH(t - t') \langle \Psi | [\hat{S}^-(\mathbf{r}, t), \hat{S}^+(\mathbf{r}', t')] | \Psi \rangle,$$

being $H(x)$ the Heaviside function. The argument carried by Singh and Tešanović relies on random phase approximation (RPA). Let χ_0^{-+} be the non-interacting transverse response function. Then

$$\chi^{-+} \simeq \frac{\chi_0^{-+}}{1 - U\chi_0^{-+}}, \quad (\text{RPA})$$

is the random phase approximation (RPA) proper response function.

We move to momentum-frequency space. Within RPA approximation, low-lying spin waves collective excitations at wavevector $\mathbf{q}$ give a stable gapless Goldstone mode if:

$$1 - U\chi_0^{-+}(\mathbf{q}, 0) = 0. \quad (\text{D.7})$$

Commensurate antiferromagnetism is stable if the above equation is satisfied at $\mathbf{q} \rightarrow \pi$. Indeed, the wavevector π expresses two-sites periodicity in both directions.

Now, as we specified in Sec. 5.1, commensurate antiferromagnetism is described by the simultaneous SSB:

$$\begin{aligned} \text{SU}(2) &\xrightarrow{\text{SSB}} \text{U}^z(1), \\ \text{T}_1^{(x)} \otimes \text{T}_1^{(y)} &\xrightarrow{\text{SSB}} \text{T}_2^{(x)} \otimes \text{T}_2^{(y)}. \end{aligned}$$

Thus translational invariance is halved in both directions. The lattice $\mathcal{L}$ is separated in the two sublattices $\mathcal{S}_1, \mathcal{S}_2$ of Fig. 3.1. As a consequence, the non-interacting response function χ_0^{+-} does not depend strictly on $\mathbf{r}, \mathbf{r}'$ themselves. Instead, it depends only on which sublattice $\mathbf{r}$ and $\mathbf{r}'$ belong to. Because of this χ_0^{+-} is representable as a 2×2 matrix χ_0^{+-} . Then Eq. (D.7) is actually an eigenvalue equation,

$$\mathbb{I} - U\chi_0^{+-}(\mathbf{q}, 0) = \mathbf{0},$$

with $\mathbf{0}$ the 2×2 zero matrix. Diagonalising $\chi_0^{+-}(\mathbf{q}, 0)$ (which is symmetric due to lattice exchange symmetry) we can express the above equation in terms of the diagonal matrix containing its eigenvalues.

Consider the largest eigenvalue $\lambda_\delta(\mathbf{k}, \omega)$ of the matrix $[\chi_0^{+-}]$ at doping δ . We work in momentum-frequency space. The phase is unstable if

$$1 - U\lambda_\delta \neq 0, \quad (\text{D.8})$$

because, as we said, in that case we do not have a pole for the transverse spin-spin response function. Singh and Tešanović [6] found

$$\lambda_0(\mathbf{k} \rightarrow \boldsymbol{\pi}, 0) = \frac{1}{U}, \quad (\text{half filling}) \quad (\text{D.9})$$

$$\lambda_\delta(\mathbf{k} \rightarrow \boldsymbol{\pi}, 0) = \frac{1}{U} + \frac{2\epsilon_F}{\Delta} N(\epsilon_F). \quad (\text{finite doping}) \quad (\text{D.10})$$

The reason for the distinction above is the intrinsic metallic nature of the system for infinitesimal doping. The half-filled AF phase is insulating, because the chemical potential lies within the gap. For an insulator Fermi energy is not defined. Instead, at finite filling the phase is metallic because the chemical potential crosses the upper band.

From Eq. (D.8) and (D.9), we see that the half filled antiferromagnet is stable towards transverse spin fluctuations, because χ^{+-} develops a divergence at zero frequency. At any finite doping, since we are describing a metal, a FS exists and thus $N(\epsilon_F)$ is finite. This makes the phase unstable, and this is the reason we do not discuss the AF phase outside of half-filling.

As we discussed at the beginning of Chap. 5, antiferromagnetism can establish in many ways. For instance, striped antiferromagnets are of particular interest [1]. In order to investigate the most stable physical realisation of an antiferromagnet on the conventional HM, on a pure HF level, we should implement a variational search for the minimum of free energy while varying the AF wavevector $\mathbf{k}$. Notice that at incommensurate wavevectors, $\mathbf{k} \neq \boldsymbol{\pi}$, the unit cell is not as simple as a pair of sites, but can become much larger. Hence, the HF analytical solution would be more complicated than our pseudospin representation of the HM hamiltonian in the commensurate AF channel.

D.4 iHF results

For the conventional HM, antiferromagnetism is described by a single HFP, the magnetisation m . Its self-consistency Eq. (5.51) reads:

$$m_{i+1} = \frac{U}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{m_i}{2E_{\mathbf{k}}[\Delta_i]} [f(-E_{\mathbf{k}}[m_i]; \beta, \mu) - f(E_{\mathbf{k}}[m_i]; \beta, \mu)], \quad (\text{D.11})$$

where we indicated as m_i the value of magnetisation at step i . We ran iHF simulations in the ranges:

$$0 \leq U \leq 16, \quad 0 \leq \delta \leq 0.48.$$

We considered also $\delta > 0$, although – as we specified in previous section – a doped commensurate antiferromagnet is actually unstable.

In Fig. D.1 magnetisation m at $\beta = 100$ is plotted. As is visible in Fig. D.1a, ignoring instability issues, doping disrupts AF ordering; the horizontal asymptote lowers as δ gets bigger. Also, for $\delta > 0$, a finite magnetisation m exists only for $U \geq U_c[\delta]$ (a critical value parametrically dependent on doping).

Fig. D.1b shows the dependence of the magnetisation on doping varying the local repulsion U . Identical analysis have been performed for different temperatures, leading to analogous results as long as $\beta \gtrsim 10$ and to $m \simeq 0$ for higher temperatures.




Figure D.1 | Plots of the zero-temperature AF instability order parameter m as a function of both the local repulsion U/t (Fig. D.1a) and the doping $\delta = n - 1/2$ (Fig. D.1b).

D.5 Analytic expression of self-consistent magnetisation at half-filling

In this subsection we provide an analytic expression to compute m self-consistently at half-filling. Multiply Eq. (5.51) on both sides by U . Since due to symmetry $\mu = 0$, we have the finite-temperature self-consistent equation

$$\Delta = \frac{U}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\Delta}{\sqrt{\epsilon_{\mathbf{k}}^2 + \Delta^2}} \tanh\left(\frac{\beta}{2} \sqrt{\epsilon_{\mathbf{k}}^2 + \Delta^2}\right),$$

where we used the relation

$$\text{Th}_{\mathbf{k}}^{(-)}[\beta, 0] = \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right).$$

In thermodynamic limit $L \rightarrow \infty$

$$\frac{1}{U} = \frac{1}{2} \int_{\text{BZ}} \frac{d\mathbf{k}}{(2\pi)^2} \frac{1}{\sqrt{\epsilon_{\mathbf{k}}^2 + \Delta^2}} \tanh\left(\frac{\beta}{2} \sqrt{\epsilon_{\mathbf{k}}^2 + \Delta^2}\right).$$

Substitute the momentum integral with an integral over energies:

$$\frac{2}{U} = \int_{-4t}^{4t} \frac{d\epsilon \rho(\epsilon)}{\sqrt{\epsilon^2 + \Delta^2}} \tanh\left(\frac{\beta}{2} \sqrt{\epsilon^2 + \Delta^2}\right),$$

being $\rho(\epsilon)$ the 2D square lattice DoS. Its explicit form [27] is given by

$$\rho(\epsilon) = \frac{1}{2\pi^2} K\left(\sqrt{1 - \left(\frac{\epsilon}{4t}\right)^2}\right) \quad K(x) \equiv \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - x^2 \sin^2 \theta}},$$

where $K(x)$ is the complete elliptic integral of the first kind, shown in Fig. D.2a.

As is plotted in Fig. D.2b, the resulting DoS is logarithmically peaked at $\epsilon = 0$, on the MBZ. A common approximation is to substitute the actual DoS with a step function,

$$\rho(\epsilon) \simeq \rho_0 H(\epsilon_0 - |\epsilon|) \quad \text{with} \quad 2\rho_0 \epsilon_0 = 1.$$

Here, the peak parameters ϵ_0 and ρ_0 are chosen in order to maintain a correct DoS integral, considering also the 2 pre-factor of spin degeneracy. Within this extremely rough approximation, we have

$$\frac{2}{U} \simeq \rho_0 \int_{-\epsilon_0}^{\epsilon_0} \frac{d\epsilon}{\sqrt{\epsilon^2 + \Delta^2}} \tanh\left(\frac{\beta}{2} \sqrt{\epsilon^2 + \Delta^2}\right).$$




Figure D.2 | Plots of the complete elliptic integral of the first kind $K(x)$ and the density of states for the planar square lattice.




Figure D.3 | Sketch of the analytic expression for m . The left panel shows a family of coloured curves described by Eq. (D.12). The right panel shows a family of coloured curves described by Eq. (D.13), only reasonable at $U \rightarrow 0$.

Suppose the system is magnetised, $\Delta > 0$. At low temperature, $\beta \gg 1$, as a first approximation we can substitute the hyperbolic tangent with unity, obtaining

$$\begin{aligned}
 \frac{2}{\rho_0 U} &\simeq \int_{-\epsilon_0}^{\epsilon_0} \frac{d\epsilon}{\sqrt{\epsilon^2 + \Delta^2}} \\
 &= \int_{-\epsilon_0/\Delta}^{\epsilon_0/\Delta} \frac{dx}{\sqrt{1+x^2}} \\
 &= \sinh^{-1} \left(\frac{\epsilon_0}{\Delta} \right) - \sinh^{-1} \left(-\frac{\epsilon_0}{\Delta} \right) \\
 &= 2 \sinh^{-1} \left(\frac{\epsilon_0}{\Delta} \right),
 \end{aligned}$$

thus

$$m \simeq \frac{\epsilon_0}{U \sinh(1/\rho_0 U)}, \quad (\text{D.12})$$

having we used $\Delta = mU$. In the left panel of Fig. D.3 we sketch this function of U varying ρ_0 , ϵ_0 . The same data of Fig. D.1a, $\delta = 0$ are plotted as grey dots. The shown curves are not fits.

Now: at large U ,

$$\lim_{U \rightarrow +\infty} m \simeq \epsilon_0 \rho_0 = \frac{1}{2},$$

which gives the correct saturation limit at high local repulsion. In the opposite limit,

$$\begin{aligned}\lim_{U \rightarrow 0} m &\simeq \lim_{U \rightarrow 0} \frac{2\epsilon_0/U}{\exp(1/\rho_0 U) - \exp(-1/\rho_0 U)} \\ &\simeq \lim_{U \rightarrow 0} \frac{2\epsilon_0}{U} \exp\left(-\frac{1}{\rho_0 U}\right),\end{aligned}\tag{D.13}$$

giving a near-0 exponential suppression of magnetisation, typical of MFT self-consistent solutions. In the right panel of Fig. [D.3](#) we sketch this approximation.

Appendix E

Theoretical constraints on the HFPs

In this appendix we derive some theoretical constraints on the HFPs of the normal and AF phase. The derivations presented here are used in the discussions of Chap. 4 and Chap. 5.

E.1 Normal phase

The normal phase is described by two HFPs:

$$\text{Normal HFPs} \quad : \quad u^{(s^*)}, u^{(d)},$$

which obey the self-consistency equations of Tab. 4.2. In this section we show that Eq. (4.44) holds,

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} = 0. \quad (\text{E.1})$$

The discussion is divided in three parts: in Sec. E.1.1, we discuss the spatial geometry of bands with a mixed symmetry $s^* \oplus d$. In Sec. E.1.2 we derive Eq. (E.1) by considering the simpler case of unbroken C_4 symmetry (hence setting $u^{(d)} = 0$ “by hand”). In Sec. E.1.3 we break rotational symmetry, $C_4 \xrightarrow{\text{SSB}} C_2$, and terminate the proof.

E.1.1 Mixed $s^* \oplus d$ bands

Before starting, it is useful to spend a few words on the spatial structure of a mixed $s^* \otimes d$ -wave band. Let $\tilde{\epsilon}_{\mathbf{k}}$ be a generic mixed band,

$$\tilde{\epsilon}_{\mathbf{k}} = \epsilon^{(s^*)}(\cos k_x + \cos k_y) + \epsilon^{(d)}(\cos k_x - \cos k_y).$$

It is the balance between the two components $\epsilon^{(s^*)}$ and $\epsilon^{(d)}$ that gives the main features of the renormalised band. Consider the s^* -wave and d -wave structure factors already presented in Fig. 3.5, and define the Magnetic Brillouin Zone (MBZ) and Nematic Brillouin Zone (NBZ) as follows:

$$\text{MBZ} \equiv \left\{ \mathbf{k} \in \text{BZ} \mid \varphi_{\mathbf{k}}^{(s^*)} \geq 0 \right\}, \quad (\text{E.2})$$

$$\text{NBZ} \equiv \left\{ \mathbf{k} \in \text{BZ} \mid \varphi_{\mathbf{k}}^{(d)} \geq 0 \right\}, \quad (\text{E.3})$$

represented in the top-left panel of Fig. E.1. In the rest of the panels of Fig. E.1 are reported examples of mixed symmetry bands, for positive components. The respective opposite sign plots are obtained by the following rules:

$$\epsilon^{(s^*)} < 0 \quad \text{and} \quad \epsilon^{(d)} < 0 \quad \implies \quad \text{Invert colors in Fig. E.1}$$

$$\epsilon^{(s^*)} > 0 \quad \text{and} \quad \epsilon^{(d)} < 0 \quad \implies \quad \text{Rotate plots by } \pi/2 \text{ in Fig. E.1}$$

$$\epsilon^{(s^*)} < 0 \quad \text{and} \quad \epsilon^{(d)} > 0 \quad \implies \quad \text{Rotate plots by } \pi/2 \text{ and invert colors in Fig. E.1}$$

Let us focus on the MBZ and the NBZ. Starting from the intersections of the two zones, both can be factored in two sectors of interest,

$$\text{MBZ} = \mathcal{A} \cup \mathcal{B} \quad \text{and} \quad \text{NBZ} = \mathcal{A} \cup \mathcal{C},$$




Figure E.1 | Theoretical representation of mixed-symmetry nematic bands for the normal phase. The white contour represent level lines. In top-left panel, the MBZ and NBZ are depicted. While these plots only account for positive-defined components, in text is described how to map each one on its counterpart with different sign.

being, accordingly to Fig. E.2,

$$\mathcal{A} \equiv \mathcal{A}_1 \cup \mathcal{A}_2, \quad \mathcal{B} \equiv \mathcal{B}_1 \cup \mathcal{B}_2, \quad \mathcal{C} \equiv \mathcal{C}_1 \cup \mathcal{C}_2.$$

These domains satisfy:

$$\mathcal{A} \equiv \left\{ \mathbf{k} \in \text{BZ} \mid \varphi_{\mathbf{k}}^{(s^*)} \geq 0 \wedge \varphi_{\mathbf{k}}^{(d)} \geq 0 \right\} \quad (\text{E.4})$$

$$= \text{MBZ} \cap \text{NBZ},$$

$$\mathcal{B} \equiv \left\{ \mathbf{k} \in \text{BZ} \mid \varphi_{\mathbf{k}}^{(s^*)} \geq 0 \wedge \varphi_{\mathbf{k}}^{(d)} \leq 0 \right\} \quad (\text{E.5})$$

$$= \text{MBZ} \setminus \text{NBZ},$$

$$\mathcal{C} \equiv \left\{ \mathbf{k} \in \text{BZ} \mid \varphi_{\mathbf{k}}^{(s^*)} \leq 0 \wedge \varphi_{\mathbf{k}}^{(d)} \geq 0 \right\} \quad (\text{E.6})$$

$$= \text{NBZ} \setminus \text{MBZ},$$

and will be used later on. Whatever the shape of the mixed band, the following equation always holds

$$|k_x| = \frac{\pi}{2} \quad \text{and} \quad |k_y| = \frac{\pi}{2} \quad \implies \quad \tilde{\epsilon}_{(k_x, k_y)} = 0,$$

since only cosines are involved. This identifies four node points, represented as black dots in Fig. E.2.




Figure E.2 Depiction of the sectors composing the MBZ (Fig. E.2a) and the NBZ (Fig. E.2b), based on the divisions shown in the top-left panel of Fig. E.1. The full green zone composes $\mathcal{A}$, the region where both the s^* -wave and d -wave structure factors are positive defined; its MBZ complementary white counterpart in the left panel gives $\mathcal{B}$, for which the s^* -wave structure factors is positive-definite while the d -wave is negative. Similarly the NBZ complementary white counterpart in the right panel gives $\mathcal{C}$, for which signs are exchanged. The black dots, at the contours intersection, are band nodes.

E.1.2 Bands shrinking in C_4

Consider the simpler situation where C_4 symmetry is imposed. This implies that the renormalised bands are given by

$$\tilde{\epsilon}_{\mathbf{k}} \equiv (u^{(s^*)}V - 2t)(\cos k_x + \cos k_y),$$

with a single self-consistency equation, given by the first row of Tab. 4.2:

$$u^{(s^*)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}).$$

We have the following constraint on $u^{(s^*)}$

$$C_4 \implies 0 < u^{(s^*)} < 2t/V. \quad (\text{E.7})$$

The effective hopping is smaller than the original hopping. We can denominate this net effect “bands shrinking”.

Proof E.1 — Proof of Eq. (E.7): bands shrinking effect in C_4 .

The proof follows from this chain of deductions:

1. For $t > 0$, there is no self-consistent solution admitting $u^{(s^*)} \leq 0$.

Proof. Suppose instead $u^{(s^*)} \leq 0$. For any function $g_{\mathbf{k}}$ defined on the BZ:

$$\sum_{\mathbf{k} \in \text{BZ}} g_{\mathbf{k}} = \sum_{\mathbf{k} \in \text{MBZ}} [g_{\mathbf{k}} + g_{\mathbf{k}+\pi}],$$

where we defined the vector $\boldsymbol{\pi} \equiv (\pi, \pi)$. The self-consistency equation thus reads

$$\begin{aligned} u^{(s^*)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x + \cos k_y) [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) + f(\tilde{\epsilon}_{\mathbf{k}+\boldsymbol{\pi}}; \beta, \tilde{\mu})] \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x + \cos k_y) [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})], \end{aligned} \quad (\text{E.8})$$

having we used the antisymmetry of s^* -wave functions by $\boldsymbol{\pi}$ shifts. Now, if $u^{(s^*)} \leq 0$ the renormalised band $\tilde{\epsilon}_{\mathbf{k}}$ is negative inside the MBZ and positive outside, implying

$$\forall \mathbf{k} \in \text{MBZ} \quad : \quad f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \geq f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}),$$

the equality only being satisfied on the MBZ contour. Then, since inside the MBZ it holds by definition $\cos k_x + \cos k_y \geq 0$, the above equations imply

$$u^{(s^*)} \leq 0 \quad \implies \quad u^{(s^*)} \geq 0.$$

To exclude the possibility of $u^{(s^*)} = 0$, it suffices to notice that from the argument above such a situation only arises when the MBZ contour degenerates to the entirety of the MBZ itself – which is, when the band is flat. But that cannot be the case for $u^{(s^*)} = 0$, as long as $t > 0$. Thus we have proven that

$$u^{(s^*)} > 0.$$

2. For $t > 0$, there is no self-consistent solution admitting $u^{(s^*)} \geq 2t/V$.

Proof. We proceed exactly as did for the previous point, up to Eq. (E.8):

$$u^{(s^*)} = \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x + \cos k_y) [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})].$$

Now, if $u^{(s^*)} \geq 2t/V$ this means the prefactor in front of the renormalised bands is positive. Then $\tilde{\epsilon}_{\mathbf{k}}$ is positive inside the MBZ and negative outside, implying

$$\forall \mathbf{k} \in \text{MBZ} \quad : \quad f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \leq f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}),$$

thus we have

$$u^{(s^*)} \geq 2t/V \quad \implies \quad u^{(s^*)} \leq 0,$$

which is absurd. Thus

$$u^{(s^*)} < 2t/V.$$

Collecting the two points above, we obtain the constraint of Eq. (E.7).

E.1.3 Bands shrinking in $C_2/\mathbb{Z}_2$

Let us extend the arguments above to a weaker symmetry, allowing for


$$C_4 \xrightarrow{\text{SSB}} C_2/\mathbb{Z}_2.$$

We work with the full, anisotropic band of Eq. (4.25). In the normal phase, nematic effects are always absent,

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} = 0,$$

which coincides with Eq. (E.1). Therefore, even if we allow for the breaking of planar rotational symmetry, this does not take place.

Proof E.2 — Proof of Eq. (E.1): non-nematic bands shrinking effect in C_2 .



a | Same-sign mixed bands.

b | Opposite-sign mixed bands.

Figure E.3 | Theoretical example of bands with $u^{(s^*)} > 2t/V$, $u^{(d)} > 0$ (top-left) and $u^{(s^*)} < 0$, $u^{(d)} > 0$ (top right). Two example points are reported, connected by a $\pi/2$ rotation as given in Eqns. (E.15), (E.10).

The proof is organised as a series of sub-proofs. We start by proving the left side of Eq. (E.1), namely that

$$0 < u^{(s^*)} < 2t/V.$$

We use everywhere $t > 0$.

1. *The following implication holds:*

$$u^{(s^*)} \geq 2t/V, \quad u^{(d)} > 0 \quad \implies \quad u^{(s^*)} \leq 0. \quad (\text{E.9})$$

Proof. Let us start by the implication

$$u^{(s^*)} \geq 2t/V > 0, \quad u^{(d)} > 0 \quad \implies \quad \epsilon^{(s^*)} \geq 0, \quad \epsilon^{(d)} > 0.$$

In Fig. E.3a an example band is shown for $\epsilon^{(s^*)} = 1$, $\epsilon^{(d)} = 1/2$. Note that if $u^{(s^*)} = 2t/V$, the renormalised band is purely d -wave since $\epsilon^{(s^*)} = 0$.

Recall Eq. (E.8), and separate in two sums over $\mathcal{A}$ and $\mathcal{B}$:

$$\begin{aligned} u^{(s^*)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \mathcal{A}} (\cos k_x + \cos k_y) [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})] \\ &\quad + \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \mathcal{B}} (\cos k_x + \cos k_y) [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu})]. \end{aligned}$$

Consider the sum over $\mathcal{B}$. Looking at the bottom-left panel of Fig. E.1, such a region contains nodal curves across which bands change sign. Let $\mathcal{R}$ be the clockwise $\pi/2$ rotation: then, any point in $\mathcal{B}$ can be seen as the rotation of a point in $\mathcal{A}$

$$\forall \mathbf{k}' \in \mathcal{B} \quad \exists \quad \mathbf{k} \in \mathcal{A} \mid \mathbf{k}' = \mathcal{R}\mathbf{k}. \quad (\text{E.10})$$

A graphical rendition of this idea is given in Fig. E.3a. Then we can rewrite $u^{(s^*)}$ as a single sum over $\mathcal{A}$,

$$\begin{aligned} u^{(s^*)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \mathcal{A}} (\cos k_x + \cos k_y) \\ &\quad \times [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) + f(\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu})]. \quad (\text{E.11}) \end{aligned}$$

From Eq. (E.4), $\forall \mathbf{k} \in \mathcal{A}$ the functions $\cos k_x \pm \cos k_y$ are positive by construction. By hypothesis $\epsilon^{(s^*)} \geq 0$ and $\epsilon^{(d)} > 0$. Hence:

$$\forall \mathbf{k} \in \mathcal{A} \quad : \quad \begin{aligned} \tilde{\epsilon}_{\mathbf{k}} &= \overbrace{\epsilon^{(s^*)} (\cos k_x + \cos k_y)}^{\geq 0} + \overbrace{\epsilon^{(d)} (\cos k_x - \cos k_y)}^{\geq 0}, \\ \tilde{\epsilon}_{\mathcal{R}\mathbf{k}} &= \underbrace{\epsilon^{(s^*)} (\cos k_x + \cos k_y)}_{\geq 0} - \underbrace{\epsilon^{(d)} (\cos k_x - \cos k_y)}_{\geq 0}. \end{aligned}$$

Summing the two equations above we have:

$$\forall \mathbf{k} \in \mathcal{A} \quad : \quad \tilde{\epsilon}_{\mathbf{k}} + \tilde{\epsilon}_{\mathcal{R}\mathbf{k}} \geq 0,$$

thus $\tilde{\epsilon}_{\mathbf{k}} \geq -\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}$. The Fermi-Dirac distribution is monotonically descendent. Then:

$$\forall \mathbf{k} \in \mathcal{A} \quad : \quad \begin{aligned} f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) &\leq f(-\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu}), \\ f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) &\geq f(\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu}). \end{aligned}$$

This condition, inserted in Eq. (E.11), leads to the absurd of Eq. (E.9).

2. *The following implication holds:*

$$u^{(s^*)} \geq 2t/V > 0, \quad u^{(d)} < 0 \quad \implies \quad u^{(s^*)} \leq 0. \quad (\text{E.12})$$

The proof is identical to the above point, exchanging the roles of $\mathcal{A}$ and $\mathcal{B}$.

3. *The following implication holds:*

$$u^{(s^*)} \geq 2t/V > 0 \quad \implies \quad u^{(d)} = 0. \quad (\text{E.13})$$

The proof follows from combining Eqs. (E.9), (E.12).

4. *It is impossible to have $u^{(s^*)} \geq 2t/V$.*

Proof. From Eq. (E.13), we know that if $u^{(s^*)} \geq 2t/V$ then $u^{(d)} = 0$. But then C_4 is unbroken, and we get back to the discussion of Sec. E.1.2. Then we cannot have $u^{(s^*)} \geq 2t/V$.

5. *The following implication holds:*

$$u^{(s^*)} \leq 0, \quad u^{(d)} > 0 \quad \implies \quad u^{(s^*)} > 0. \quad (\text{E.14})$$

Proof. Let for instance

$$u^{(s^*)} \leq 0, \quad u^{(d)} > 0 \quad \implies \quad \epsilon^{(s^*)} < 0, \quad \epsilon^{(d)} > 0.$$

In Fig. E.3b an example band is shown for $\epsilon^{(s^*)} = -1$, $\epsilon^{(d)} = 1/2$. Proceed as for the first point of this chain. Similarly to Eq. (E.10), Any point in $\mathcal{A}$ can be obtained by rotating some point in $\mathcal{B}$,

$$\forall \mathbf{k}' \in \mathcal{A} \quad \exists \quad \mathbf{k} \in \mathcal{B} \mid \mathbf{k}' = \mathcal{R}\mathbf{k}. \quad (\text{E.15})$$

An equation similar to Eq. (E.11) holds, the sum being performed over $\mathcal{B}$

$$\begin{aligned} u^{(s^*)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \mathcal{B}} (\cos k_x + \cos k_y) \\ &\quad \times [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) + f(\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(-\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu})]. \end{aligned} \quad (\text{E.16})$$

From Eq. (E.5), $\forall \mathbf{k} \in \mathcal{B}$ the form factor $\cos k_x + \cos k_y$ is positive by construction; the form factor $\cos k_x - \cos k_y$ is negative by construction. By hypothesis $\epsilon^{(s^*)} < 0$ and $\epsilon^{(d)} > 0$. Hence:

$$\forall \mathbf{k} \in \mathcal{B} : \quad \begin{aligned} \tilde{\epsilon}_{\mathbf{k}} &= \overbrace{\epsilon^{(s^*)} (\cos k_x + \cos k_y)}^{\leq 0} + \overbrace{\epsilon^{(d)} (\cos k_x - \cos k_y)}^{\leq 0}, \\ \tilde{\epsilon}_{\mathcal{R}\mathbf{k}} &= \overbrace{\epsilon^{(s^*)} (\cos k_x + \cos k_y)}^{\leq 0} - \overbrace{\epsilon^{(d)} (\cos k_x - \cos k_y)}^{\leq 0}. \end{aligned}$$

Summing the two equations above we have:

$$\forall \mathbf{k} \in \mathcal{B} : \quad \tilde{\epsilon}_{\mathbf{k}} + \tilde{\epsilon}_{\mathcal{R}\mathbf{k}} \leq 0,$$

thus $\tilde{\epsilon}_{\mathbf{k}} \leq -\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}$. The Fermi-Dirac distribution is monotonically descendent. Then:

$$\forall \mathbf{k} \in \mathcal{B} : \quad \begin{aligned} f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) &\geq f(-\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu}), \\ f(-\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) &\leq f(\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu}). \end{aligned}$$

This condition, inserted in Eq. (E.11), leads to an absurd:

$$u^{(s^*)} \leq 0, \quad u^{(d)} > 0 \quad \implies \quad u^{(s^*)} > 0.$$

which coincides with Eq. (E.14).

6. *The following implication holds:*

$$u^{(s^*)} \leq 0, \quad u^{(d)} < 0 \quad \implies \quad u^{(s^*)} > 0. \quad (\text{E.17})$$

The proof is identical to the above point, exchanging the roles of $\mathcal{A}$ and $\mathcal{B}$.

7. *The following implication holds:*

$$u^{(s^*)} \leq 0 \quad \implies \quad u^{(d)} = 0. \quad (\text{E.18})$$

The proof follows from combining Eqs. (E.14), (E.17).

8. *It is impossible to have $u^{(s^*)} \leq 0$.*

Proof. From Eq. (E.18), we know that if $u^{(s^*)} \leq 0$ then $u^{(d)} = 0$. But then C_4 is unbroken, and we get back to the discussion of Sec. E.1.2. Then we cannot have $u^{(s^*)} \leq 0$.

From the chain of derivations above, we can conclude self-consistently that also in the nematic structure the constraint of Eq. (E.7),

$$0 < u^{(s^*)} < 2t/V,$$

holds. In the s^* -wave channel, partial breaking of rotational symmetry is not enough to modify the main effect identified under C_4 symmetry, “bands shrinking”. What is missing is the behaviour of the d -wave channel when said effect takes place.

We move on and prove the right side of Eq. (E.1), namely that

$$u^{(d)} = 0.$$

Continuing the enumeration from the above points,

9. *The following implication holds:*

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} > 0 \quad \implies \quad u^{(d)} < 0. \quad (\text{E.19})$$

Proof. Let $0 < u^{(s^*)} < 2t/V$ and $u^{(d)} > 0$. Then $\epsilon^{(d)} > 0$. From Tab. 4.2, the self-consistency equation for $u^{(d)}$ reads

$$\begin{aligned} u^{(d)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{NBZ}} (\cos k_x - \cos k_y) [f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) - f(\tilde{\epsilon}_{\mathcal{R}\mathbf{k}}; \beta, \tilde{\mu})], \end{aligned}$$

where we recognised, as is shown in Fig. E.2b, that $\text{BZ} \setminus \text{NBZ} = \mathcal{R}\text{NBZ}$. It holds:

$$\forall \mathbf{k} \in \text{NBZ} \quad : \quad \begin{aligned} \tilde{\epsilon}_{\mathbf{k}} &= \epsilon^{(s^*)} (\cos k_x + \cos k_y) + \overbrace{\epsilon^{(d)} (\cos k_x - \cos k_y)}^{\geq 0}, \\ \tilde{\epsilon}_{\mathcal{R}\mathbf{k}} &= \epsilon^{(s^*)} (\cos k_x + \cos k_y) - \underbrace{\epsilon^{(d)} (\cos k_x - \cos k_y)}_{\geq 0}, \end{aligned}$$

thus $\tilde{\epsilon}_{\mathcal{R}\mathbf{k}} < \tilde{\epsilon}_{\mathbf{k}}$, and inserting this fact in the self-consistency equation above we get the absurd of Eq. (E.19).

10. *The following implication holds:*

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} < 0 \quad \implies \quad u^{(d)} > 0. \quad (\text{E.20})$$

The proof is identical to the point above.

11. *The following implication holds:*

$$0 < u^{(s^*)} < 2t/V \quad \implies \quad u^{(d)} = 0. \quad (\text{E.21})$$

The proof follows from combining Eqs. (E.19), (E.20).

12. *It is impossible to have $u^{(d)} \neq 0$.* The proof follows from combining Eqs. (E.13), (E.18) and (E.21).

Hence we conclude that Eq. (E.1) holds.

E.2 AF phase

The AF phase divides in two subphases, namely aAF and sAF phases, as reported in Tab. 5.4. The sAF phase is described by five HFPS,

$$\text{sAF HFPS} \quad : \quad m, u^{(s^*)}, u^{(d)}, v^{(s^*)}, v^{(d)},$$

while for the aAF phase

$$\text{aAF HFPS} \quad : \quad m, u^{(s^*)}, u^{(d)}, -iv^{(p_x)}, -iv^{(p_y)}.$$

Self-consistency equations are reported in Eq. (5.68) and Tabs. 5.2, 5.3.

In this section we show that

- Eq. (5.110) holds,

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} = 0. \quad (\text{E.22})$$

Hence also in the AF phase we have a Mott localisation effect;

- Eq. (5.111) holds,

$$\forall \gamma \quad : \quad v^{(\gamma)} = 0. \quad (\text{E.23})$$

Hence the AF gap is not “complexified”.

The discussion is divided in four parts: in Sec. E.2.1, we derive Eq. (E.22) within C_4 symmetry (as in Sec. E.1.2, setting $u^{(d)} = 0$). In Sec. E.2.2 we prove Eq. (E.23) for the sAF phase within C_2 symmetry. In Sec. E.2.3 we do the same for the aAF phase. We conclude, in Sec. E.2.4, by proving Eq. (E.22) within C_2 symmetry.

E.2.1 Bands shrinking in C_4

For the normal phase, we saw that when C_4 symmetry is preserved (Sec. E.1.2) we can conclude immediately that bands renormalisation is actually a shrinking. An identical result holds here: if C_4 symmetry is imposed onto the renormalized bands,

$$\tilde{\epsilon}_{\mathbf{k}\sigma} \equiv (u^{(s^*)}V - 2t)(\cos k_x + \cos k_y).$$

Among the harmonics $u^{(\gamma)}$, just the s^* -wave channel contributes to renormalisation of bands, with a single self-consistency equation

$$u^{(s^*)} = -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (\cos k_x + \cos k_y) \frac{\tilde{\epsilon}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}].$$

We now prove that:

$$0 < u^{(s^*)} < 2t/V. \quad (\text{E.24})$$

This constraint is identical to Eq. (E.7), obtained for the normal phase. Also in AF phase the bare bands are effectively shrunk by $\tilde{H}_V$.

Proof E.3 — Proof of Eq. (E.24).

As we did for the normal phase:

1. For $t > 0$, there is no self-consistent solution admitting $u^{(s^*)} \leq 0$.

Proof. If $t > 0$ and $u^{(s^*)} \leq 0$

$$\forall \mathbf{k} \in \text{MBZ} \quad : \quad \tilde{\epsilon}_{\mathbf{k}} \leq 0,$$

the equality only holding on the MBZ contour. Looking at the self-consistency equation, all other objects inside the sum are positive-defined. This gives

$$u^{(s^*)} \leq 0 \quad \implies \quad u^{(s^*)} \geq 0.$$

But $u^{(s^*)} \neq 0$ as long as $t > 0$, as can be seen by the self-consistency equation. We conclude necessarily

$$u^{(s^*)} > 0,$$

which is absurd.

2. For $t > 0$, there is no self-consistent solution admitting $u^{(s^*)} \geq 2t/V$.

Proof. If $t > 0$ and $u^{(s^*)} \geq 2t/V$

$$\forall \mathbf{k} \in \text{MBZ} \quad : \quad \tilde{\epsilon}_{\mathbf{k}} \geq 0$$

the equality only holding on the MBZ contour. Inverting inequalities from point above, we conclude necessarily

$$u^{(s^*)} < 2t/V.$$

which is absurd.

E.2.2 Selection rules on the magnetic HFPs in sAF phase

We applied HF methods to the EHM and obtained two flavours of antiferromagnetism, each of which involves five HFPs. The scenario appears to be rich. In this section we present an argument

to prove that the actual number of HFPS is much smaller, as many of those are mathematically suppressed.

Let us start from the sAF phase, assuming

$$C_4 \xrightarrow{\text{SSB}} C_2.$$

The self-consistency problem is described by the five HFPS on the first row of Tab. 5.4 and their self-consistency equations of Tabs. 5.2 and 5.3.

Let us define the functional

$$F[f(\mathbf{k})] \equiv \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]}{\tilde{E}_{\mathbf{k}}} f(\mathbf{k}). \quad (\text{E.25})$$

The functional F is linear:

$$F[af(\mathbf{k}) + bg(\mathbf{k})] = aF[f(\mathbf{k})] + bF[g(\mathbf{k})], \quad (\text{E.26})$$

being $a, b \in \mathbb{C}$ and f, g two integrable functions in reciprocal space. Another handy property is the sign-preserving relation for sign-defined functions. Let for instance

$$\forall \mathbf{k} : f(\mathbf{k}) \geq 0 \implies F[f(\mathbf{k})] \geq 0. \quad (\text{E.27})$$

The relation above is implied by the fact that both $\tilde{\text{Th}}_{\mathbf{k}}^{(-)}$ and $\tilde{E}_{\mathbf{k}}$ are positive-defined,

$$\tilde{\text{Th}}_{\mathbf{k}}^{(-)} = \frac{1}{e^{\beta(\tilde{E}_{\mathbf{k}} - \mu)} + 1} - \frac{1}{e^{\beta(-\tilde{E}_{\mathbf{k}} - \mu)} + 1} \geq 0 \quad \text{and} \quad \tilde{E}_{\mathbf{k}} = \sqrt{\tilde{\epsilon}_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}\sigma}|^2} \geq 0,$$

so, being the functional F essentially a weighted integral over MBZ, any positive-defined function has a positive image. An identical statement with inverted inequalities holds for negative-defined function.

Consider now the self-consistency equation for $u^{(s^*)}$: explicitly,

$$\begin{aligned} u^{(s^*)} &= -\frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} (c_x + c_y) \frac{\tilde{\epsilon}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}] \\ &= \frac{[2t - Vu^{(s^*)}]}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]}{\tilde{E}_{\mathbf{k}}} (c_x + c_y)^2 - \frac{Vu^{(d)}}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]}{\tilde{E}_{\mathbf{k}}} (c_x^2 - c_y^2), \end{aligned}$$

having we used the notation $c_\ell \equiv \cos k_\ell$ and substituted the full bare bands

$$\tilde{\epsilon}_{\mathbf{k}} \equiv -\left[2t - u_\sigma^{(s^*)}V\right] (c_x + c_y) + u_\sigma^{(d)}V(c_x - c_y).$$

In terms of the functional F , we can rewrite:

$$\begin{aligned} u^{(s^*)} &= \left[2t - Vu^{(s^*)}\right] \times F[(c_x + c_y)^2] - Vu^{(d)} \times F[c_x^2 - c_y^2] \\ &= \left[2t - Vu^{(s^*)}\right] \times \{F[c_x^2] + 2F[c_x c_y] + F[c_y^2]\} - Vu^{(d)} \times \{F[c_x^2] - F[c_y^2]\}, \end{aligned}$$

having we used the linearity property of Eq. (E.26) in last passage.

Let now:

$$A \equiv F[c_x^2], \quad B \equiv 2F[c_x c_y], \quad C \equiv F[c_y^2], \quad (\text{E.28})$$

which allows us to rewrite the self-consistency equation as:

$$u^{(s^*)} = \left[2t - Vu^{(s^*)}\right] (A + B + C) - Vu^{(d)}(A - C)$$

Repeating an identical procedure on the three self-consistency equations for $u^{(d)}$, $v^{(s^*)}$ and $v^{(d)}$, we get the full system:

$$u^{(s^*)} = \left[2t - Vu^{(s^*)} \right] (A + B + C) - Vu^{(d)}(A - C), \quad (\text{E.29})$$

$$u^{(d)} = \left[2t - Vu^{(s^*)} \right] (A - C) - Vu^{(d)}(A - B + C), \quad (\text{E.30})$$

$$v^{(s^*)} = -Vv^{(s^*)}(A + B + C) - Vv^{(d)}(A - C), \quad (\text{E.31})$$

$$v^{(d)} = -Vv^{(s^*)}(A - C) - Vv^{(d)}(A - B + C). \quad (\text{E.32})$$

These equations define a four-components nonlinear map, whose parameters A, B, C depend on the HFPs themselves, being the functional F dependent on the HFPs. Because of this last point, the four equations are coupled to each other and the map does not decouple in two independent sub-maps. Note that the full map would include another equation for the magnetisation. It holds:

$$v^{(s^*)} = 0 \quad \text{and} \quad v^{(d)} = 0. \quad (\text{E.33})$$

Proof E.4 — Proof of Eq. (E.33): all symmetric complexifiers vanish.

Let us start by applying a linear transform on the map: define the new “sum” and “difference” variables

$$S_u \equiv u^{(s^*)} + u^{(d)}, \quad D_u \equiv u^{(s^*)} - u^{(d)}, \quad S_v \equiv v^{(s^*)} + v^{(d)}, \quad D_v \equiv v^{(s^*)} - v^{(d)},$$

whose matrix representation is

$$\begin{bmatrix} S_u \\ D_u \\ S_v \\ D_v \end{bmatrix} \equiv \begin{bmatrix} 1 & 1 & & \\ 1 & -1 & & \\ & & 1 & 1 \\ & & 1 & -1 \end{bmatrix} \begin{bmatrix} u^{(s^*)} \\ u^{(d)} \\ v^{(s^*)} \\ v^{(d)} \end{bmatrix}.$$

The map transform is obtained by summing and subtracting two couples of equations: Eq. (E.29) and (E.30) (the first two) as well as Eq. (E.31) and (E.32) (the second two). By doing this we get the simplified system:

$$(1 + 2AV)S_u + BVD_u = 2t(2A + B), \quad (\text{E.34})$$

$$(1 + 2CV)D_u + BVS_u = 2t(B + 2C), \quad (\text{E.35})$$

$$(1 + 2AV)S_v + BVD_v = 0, \quad (\text{E.36})$$

$$(1 + 2CV)D_v + BVS_v = 0. \quad (\text{E.37})$$

1. Either $S_v = 0 \wedge D_v = 0$, or $S_v \neq 0 \wedge D_v \neq 0$.

Proof. Let $S_v = 0$. From Eq. (E.37), using $C = F[c_y^2] \geq 0$ implied by Eq. (E.27):

$$\underbrace{(1 + 2CV)}_{>0} D_v = 0 \implies D_v = 0.$$

Identically, let $D_v = 0$. Then applying the same consideration on Eq. (E.36) with $C \leftrightarrow A$, we conclude $S_v = 0$. Either both are non-zero or both are zero.

2. Let $S_v \neq 0 \wedge D_v \neq 0$. Then $B \neq 0$.

Proof. To prove this, consider for example Eq. (E.36): we are working at $V > 0$, thus

$$B = -\frac{1 + 2AV}{V} \frac{S_v}{D_v},$$

which is non-zero because $A = F[c_x^2] \geq 0$, the latter following from Eq. (E.27). We could have used identically Eq. (E.37).

3. Let $S_v \neq 0 \wedge D_v \neq 0$. Hence the following relation holds:

$$(BV)^2 = (1 + 2CV)(1 + 2AV). \quad (\text{E.38})$$

Proof. We multiply Eq. (E.36) by $BV \neq 0$ (see point above),

$$BV(1 + 2AV)S_v + (BV)^2 D_v = 0,$$

and substitute from Eq. (E.37) $BVS_v = -(1 + 2CV)D_v$,

$$[-(1 + 2CV)(1 + 2AV) + (BV)^2] D_v = 0.$$

Since $D_v \neq 0$ by hypothesis, Eq. (E.38) is implied.

4. Let $S_v \neq 0 \wedge D_v \neq 0$. Hence the following relation holds:

$$B + 2C = V(B^2 - 4AC). \quad (\text{E.39})$$

Proof. We multiply Eq. (E.34) by BV on both sides,

$$BV(1 + 2AV)S_u + (BV)^2 D_u = 2tBV(2A + B),$$

and substitute from Eq. (E.35) $BVS_v = -(1 + 2CV)D_v + 2t(B + 2C)$,

$$\begin{aligned} -(1 + 2CV)(1 + 2AV)D_v + 2t(B + 2C)(1 + 2AV) + (BV)^2 D_u &= 2tBV(2A + B) \\ 2t(B + 2C)(1 + 2AV) &= 2tBV(2A + B) \\ B + 2ABV + 2C + 4ACV &= 2ABV + B^2V, \end{aligned}$$

Passing from first to second line we used Eq. (E.38), and in the passage below $t > 0$. Last line implies Eq. (E.39).

5. Let $S_v \neq 0 \wedge D_v \neq 0$. Then $B^2 \neq 4AC$.

Proof. Let $B^2 = 4AC$. Then from Eq. (E.38),

$$(BV)^2 = 1 + 2V(A + C) + 4ACV^2 \implies 1 + 2V(A + C) = 0,$$

which is absurd, being $A \geq 0$, $C \geq 0$ from Eq. (E.27).

6. Using the above point in Eq. (E.39) we get that $V > 0$ satisfies:

$$V = \frac{B + 2C}{B^2 - 4AC}. \quad (\text{E.40})$$

7. Let $S_v \neq 0 \wedge D_v \neq 0$. Then $A = C$.

Proof. We substitute Eq. (E.40) in Eq. (E.38),

$$\begin{aligned} (B^2 - 4AC) \left(\frac{B + 2C}{B^2 - 4AC} \right)^2 - 2 \frac{B + 2C}{B^2 - 4AC} (A + C) - 1 &= 0 \\ (B + 2C)^2 - 2(B + 2C)(A + C) - (B^2 - 4AC) &= 0 \\ B^2 + 4C^2 + 4BC - 2AB - 2BC - 4AC - 4C^2 - B^2 + 4AC &= 0, \end{aligned}$$

Simplifying all recurring terms, we get

$$2B(C - A) = 0.$$

From previous points we know $B \neq 0$. Then $A = C$.

8. *It's impossible to have $S_v \neq 0 \wedge D_v \neq 0$.*

Proof. To prove this, assume the contrary and use the point above to reduce Eq. (E.40) to

$$A = C \implies V = \frac{B + 2C}{B^2 - 4C^2} = \frac{1}{B - 2C}.$$

Inserting the latter in Eq. (E.37),

$$\begin{aligned} \left(1 + \frac{2C}{B - 2C} \right) D_v + \frac{B}{B - 2C} S_v &= 0 \\ \frac{B}{B - 2C} D_v + \frac{B}{B - 2C} S_v &= 0, \end{aligned}$$

which implies

$$D_v + S_v = 0 \implies v^{(s^*)} = 0.$$

Recalling from Tab. 5.3 the explicit self-consistency equation for $v^{(d)}$ and setting $v^{(s^*)} = 0$,

$$v^{(d)} = -V v^{(d)} F[(c_x - c_y)^2],$$

and using the positivity rule of Eq. (E.27) we get

$$V > 0 \quad \text{and} \quad F[(c_x - c_y)^2] \geq 0 \implies v^{(d)} = 0,$$

because no finite positive number can equal any finite negative number, and *viceversa*. Thus

$$S_v \neq 0 \quad \text{and} \quad D_v \neq 0 \implies v^{(s^*)} = v^{(d)} = 0 \implies S_v = D_v = 0.$$

Absurd.

9. Combining the above point with our first point, we get that $S_v = 0 \wedge D_v = 0$.

Eq. (E.33) is a consequence of the last point.

We explored any possibility for S_v, D_v : no stable solution with finite values of the HFPs $v^{(s^*)}$, $v^{(d)}$ exists, thus both must be zero and can be eliminated from the algorithm.

To gain full knowledge on the map fixed point's shape, we should analyse similarly the behaviour of the self-consistency equations for $u^{(s^*)}$, $u^{(d)}$. This is done later in Sec. E.2.4. For now, it suffices to note that – limitedly to the sAF phase – gap complexification is not present. This effect is not a consequence of HF derivation, which remains self-coherent and general; it is instead of the peculiar form of the non-linear map defined by the set of self-consistency equations.

E.2.3 Selection rules on the magnetic HFPS in aAF phase

Let us discuss the aAF phase. It is described by the five HFPS on the second row of Tab. 5.4, as well as the self-consistency equations of Tabs. 5.2 and 5.3. We continue using the functional F as defined in Eq. (E.25). We also define A, B, C as in Eq. (E.28). Let $s_\ell \equiv \sin k_\ell$, and

$$P \equiv F[s_x^2], \quad Q \equiv 2F[s_x s_y], \quad R \equiv F[s_y^2]. \quad (\text{E.41})$$

Using these definitions, four of five self-consistency equations are given by

$$u^{(s^*)} = \left[2t - Vu^{(s^*)} \right] (A + B + C) - Vu^{(d)}(A - C), \quad (\text{E.42})$$

$$u^{(d)} = \left[2t - Vu^{(s^*)} \right] (A - C) - Vu^{(d)}(A - B + C), \quad (\text{E.43})$$

$$v^{(p_x)} = -Vv^{(p_x)}P - Vv^{(p_y)}Q, \quad (\text{E.44})$$

$$v^{(p_y)} = -Vv^{(p_x)}Q - Vv^{(p_y)}R. \quad (\text{E.45})$$

We have:

$$v^{(p_x)} = 0 \quad \text{and} \quad v^{(p_y)} = 0. \quad (\text{E.46})$$

Proof E.5 — Proof of Eq. (E.46): all antisymmetric complexifiers vanish.

Let us write the equation for Q , given in Eq. (E.41), explicitly:

$$\begin{aligned} Q &= F[s_x s_y] \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{MBZ}} \frac{\tilde{\text{T}}\text{h}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]}{\tilde{E}_{\mathbf{k}}} \sin k_x \sin k_y \\ &= \frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\tilde{\text{T}}\text{h}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]}{\tilde{E}_{\mathbf{k}}} \sin k_x \sin k_y \\ &= \frac{1}{2L_x L_y} \sum_{k_x = -\pi}^{+\pi} \sin k_x \sum_{k_y = -\pi}^{+\pi} \frac{\tilde{\text{T}}\text{h}_{(k_x, k_y)}^{(-)}[\beta, \tilde{\mu}]}{\tilde{E}_{(k_x, k_y)}} \sin k_y. \end{aligned}$$

Passing from second to third line, we used MBZ periodicity. The k_y sum at last line is null: even if using antiperiodic harmonics for $\text{Im}\{\tilde{\Delta}_{\mathbf{k}}\}$, we did not break inversion symmetry neither in x nor y direction. Bogoliubov bands $\tilde{E}_{\mathbf{k}}$ and thermal factors must show said symmetry. The product of a symmetric function ($\tilde{\text{T}}\text{h}/\tilde{E}$) with $\sin k_y$ integrates to zero. Then

$$Q = 0.$$

Using this result in Eq. (E.44), we get

$$v^{(p_x)} = -Vv^{(p_x)}F[s_x^2].$$

Using the positivity rule of Eq. (E.27) we obtain

$$V > 0 \quad \text{and} \quad F[s_x^2] \geq 0 \quad \implies \quad v^{(p_x)} = 0.$$

Identically, from Eq. (E.45) we get $v^{(p_y)} = 0$. This proves Eq. (E.46).

This ends this short section and shows that also for the aAF phase all gap complexifiers actually vanish. Eqs. (E.33) and (E.46), combined, give Eq. (E.23). The latter is coherent with the constraint of Eq. (??).




Figure E.4 | Sketch of the Fermi-Dirac distribution $f(x)$ and the pseudo-distribution $f'(x)$ described in text for the normal phase. These example data have been plotted arbitrarily at $mU = \Delta = 0.2$ for $\mu = 0.0, 0.3, 0.6$.

E.2.4 Bands shrinking in $C_2/\mathbb{Z}_2$

A consequence of the two sections above is that the gap function is purely real, non-renormalized and constant across all reciprocal space,

$$\tilde{\Delta}_{\mathbf{k}} = mU \quad \implies \quad \tilde{E}_{\mathbf{k}} = \sqrt{\tilde{\epsilon}_{\mathbf{k}}^2 + (mU)^2}.$$

Bogoliubov bands maintain renormalisation through the $u^{(\gamma)}$ HFPs channel. For the AF phase, Eq. (E.22) holds,

$$0 < u^{(s^*)} < 2t/V \quad \text{and} \quad u^{(d)} = 0.$$

Starting from a rather complicated scenario, we were able to reduce all the AF HFPs down to the pair

$$m, u^{(s^*)}.$$

Perhaps not-so-excitingly, the real net effect of the non-local extension of the HM is, once again, the shrinking effect on bands.

Proof E.6 — Proof of Eq. (E.22): non-nematic bands shrinking effect in C_2 .

While the proof given in Sec. E.1.3 was long and quite elaborated, this one is shorter. Let us start from recalling the normal phase context:

$$\begin{aligned} \text{Normal phase:} \quad u^{(s^*)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}), \\ u^{(d)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) f(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}), \end{aligned}$$

while, using π periodicity, in the present context we can re-write all equations doubling the integration domain,

$$\begin{aligned} \text{AF phase:} \quad u^{(s^*)} &= -\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x + \cos k_y) \frac{\tilde{\epsilon}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}], \\ u^{(d)} &= -\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} (\cos k_x - \cos k_y) \frac{\tilde{\epsilon}_{\mathbf{k}}}{\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}]. \end{aligned}$$

Let us define the pseudo distribution

$$f'(\tilde{\epsilon}_{\mathbf{k}}; \beta, \tilde{\mu}) \equiv -\frac{\tilde{\epsilon}_{\mathbf{k}}}{2\tilde{E}_{\mathbf{k}}} \tilde{\text{Th}}_{\mathbf{k}}^{(-)}[\beta, \tilde{\mu}],$$

which, substituted in the self-consistency equations, makes them very similar to the normal ones, up to a $f \rightarrow f'$ exchange. We plot $f(x)$ and $f'(x)$ in Fig. E.4 for various choices of the chemical potential.

The fact that $u^{(d)} = 0$ follows from the simple consideration that in Sec. E.1.3 (when showing the same for the normal phase) we never used the specific form of f ; instead, we only made use of it being monotonically descendent as a function of $\epsilon_{\mathbf{k}}$. If f' also is a monotonically descendent function of $\epsilon_{\mathbf{k}}$, we can re-use the entire derivation of Sec. E.1.3. Let us show that using a somewhat lighter notation,

$$\epsilon_{\mathbf{k}} \rightarrow x \quad \text{and} \quad mU \rightarrow \Delta,$$

and omitting parametric arguments. The function f' is given by the product

$$f'(x) = g(x)h(x),$$

with

$$g(x) = -\frac{x}{\sqrt{x^2 + \Delta^2}},$$

$$h(x) = \left(\frac{1}{e^{\beta(\sqrt{x^2 + \Delta^2} - \mu)} + 1} - \frac{1}{e^{\beta(-\sqrt{x^2 + \Delta^2} - \mu)} + 1} \right).$$

The following points are true:

1. The function is antisymmetric about $x = 0$, because $g(x)$ is antisymmetric, while $h(x)$ is symmetric.
2. $g(x)$ is monotonically descendent. Its derivative is given by

$$\begin{aligned} \frac{dg}{dx} &= -\frac{\sqrt{x^2 + \Delta^2} - x^2/\sqrt{x^2 + \Delta^2}}{x^2 + \Delta^2} \\ &= -\frac{\Delta^2}{(x^2 + \Delta^2)^{3/2}} < 0. \end{aligned}$$

3. The product $g(x)h(x)$ is monotonically descendent at positive finite doping. To prove this, we can compute the derivative of f' ,

$$\frac{df'}{dx} = \frac{dg}{dx}h(x) + g(x)\frac{dh}{dx}.$$

We have $h(x) > 0$, and as we said $dg/dx < 0$, which makes the first term negative. The second is negative too: let

$$E(x) \equiv \sqrt{x^2 + \Delta^2} \quad \Rightarrow \quad \frac{dE}{dx} = \frac{x}{E(x)} = -g(x).$$

Then:

$$\begin{aligned} \frac{dh}{dx} &= \frac{d}{dx} \left[\frac{1}{e^{\beta(-E(x) - \mu)} + 1} - \frac{1}{e^{\beta(E(x) - \mu)} + 1} \right] \\ &= \beta \frac{dE}{dx} \underbrace{\left[\frac{e^{\beta(-E(x) - \mu)}}{[e^{\beta(-E(x) - \mu)} + 1]^2} + \frac{e^{\beta(E(x) - \mu)}}{[e^{\beta(E(x) - \mu)} + 1]^2} \right]}_{k(x)}. \end{aligned}$$

The term in square brackets is positive, $k(x) > 0$. Taking all together, we have

$$g(x) \frac{dh}{dx} = -\beta g^2(x) k(x) < 0 \quad \implies \quad \frac{df'}{dx} < 0.$$

Hence f' decreases monotonously.

These points end the proof: as anticipated, f' is a monotonically descendent function of $\epsilon_{\mathbf{k}}$, a feature that allows us to re-use the entire derivation of Sec. E.1.3, and conclude that also for the AF phase

$$u^{(d)} = 0,$$

while $u^{(s^*)}$ remains constrained in the limits of Eq. (E.24).

Appendix F

Mixing optimization in HM model

This short appendix is meant to give a practical example on a topic we often referenced to in text: how optimisation of g , the mixing parameter defined in Sec. 2.3.2 and given operatively in Eq. (2.25) through

$$\mathbf{v}_{i+1} \equiv g\mathbf{A}(\mathbf{v}_i) + (1 - g)\mathbf{v}_i,$$

can be relevant in order to ensure convergence of an iterative HF scheme as presented in Sec. 2.3.

F.1 Map analysis for s -wave superconductivity in HM

As we discussed in Sec. ??, conventional repulsive HM cannot exhibit BCS s -wave superconductivity within a MFT general scheme. This is proven at the end of Sec. ??, where we end up with the self-consistency equation

$$w^{(s)} = -\frac{U}{U_c[\beta]} w^{(s)}, \quad (\text{F.1})$$

where $U_c[\beta]$ is defined as in Eq. (6.45),

$$U_c[\beta] \equiv \left[\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{\text{Th}_{\mathbf{k}}^{(-)}[\beta]}{E_{\mathbf{k}}} \right]^{-1}. \quad (\text{F.2})$$

Since by construction $U_c[\beta] > 0$ when $U > 0$, Eq. (F.1) can only be solved by $w^{(s)} = 0$. Now, our aim is the following: given any initialiser $w_0^{(s)}$, our iterative HF algorithm should give

$$w_0^{(s)} \xrightarrow{\text{Alg}} \dots \xrightarrow{\text{Alg}} \sim 0.$$

We ask: is this algorithmic evolution possible at any model parametrisation?

F.1.1 Constraints for algorithmic evolution

Consider the nonlinear map of Eq. (F.1) as “seen by the algorithm”: at step i , we compute the left side by using the value of $w^{(s)}$ at step $i - 1$ (also for computing $U_c[\beta]$). Let us drop the superscript (s) for the sake of readability. Seen this way, we are effectively moving along the numeric succession

$$w_{i+1} = -g \left(\frac{U}{U_c[\beta]} \right)_i w_i + (1 - g)w_i \quad (\text{F.3})$$

the subscript indicating iteration. In previous equation we made use of Eq. (2.25) to generate the $i + 1$ -th step from step i . Let now:

$$U_s \equiv \left(\frac{1}{g} - 1 \right) U_c,$$

$$U_h \equiv \left(\frac{2}{g} - 1 \right) U_c.$$

The subscript indicate a “soft boundary” (first) and an “hard boundary” (second). If $w_i > 0$:

1. It always holds $w_{i+1} \leq w_i$. If instead $w_{i+1} > w_i$, from Eq. (F.3) we have

$$\begin{aligned} -g \left(\frac{U}{U_c[\beta]} \right)_i w_i + (1-g)w_i &> w_i \\ -g \left(\frac{U}{U_c[\beta]} \right)_i w_i - gw_i &> 0, \end{aligned}$$

which is impossible because a sum of negative terms cannot be positive.

2. For $0 < U < U_s$, we have $0 < w_{i+1} < w_i$. Indeed w_{i+1} is greater than zero when:

$$\begin{aligned} -g \left(\frac{U}{U_c} \right)_i w_i + (1-g)w_i &> 0 \\ \frac{1-g}{g} &> \frac{U}{U_c} \\ U_s &> U. \end{aligned}$$

3. For $U_s < U < U_h$, we have $-w_i < w_{i+1} < 0$. From point above, we know $U > U_s$ implies $w_{i+1} < 0$ by simply inverting inequalities. Moreover,

$$\begin{aligned} -g \left(\frac{U}{U_c} \right)_i w_i + (1-g)w_i &> -w_i \\ \frac{2-g}{g} &> \frac{U}{U_c} \\ U_h &> U \end{aligned}$$

4. For $U > U_h$, we have $w_{i+1} < -w_i$. This is proven by inverting inequalities in last point.

If we know U_c , for any given g we can predict the algorithmic image of a given step.

Now, consider U_c as a function of the gap Δ ,

$$U_c(\Delta) = \left[\frac{1}{2L_x L_y} \sum_{\mathbf{k} \in \text{BZ}} \frac{1}{\sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2}} \tanh\left(\frac{\beta}{2} \sqrt{\xi_{\mathbf{k}}^2 + |\Delta|^2}\right) \right]^{-1}$$

[To be continued...]

Bibliography

- [1] Alexander Wietek et al. “Stripes, Antiferromagnetism, and the Pseudogap in the Doped Hubbard Model at Finite Temperature”. en. In: *Physical Review X* 11.3 (July 2021), p. 031007. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.11.031007](https://doi.org/10.1103/PhysRevX.11.031007). URL: <https://link.aps.org/doi/10.1103/PhysRevX.11.031007> (visited on 12/12/2025).
- [2] Mingpu Qin et al. “Absence of Superconductivity in the Pure Two-Dimensional Hubbard Model”. In: *Physical Review X* 10.3 (July 2020), p. 031016. DOI: [10.1103/PhysRevX.10.031016](https://doi.org/10.1103/PhysRevX.10.031016). URL: <https://link.aps.org/doi/10.1103/PhysRevX.10.031016> (visited on 03/21/2026).
- [3] Jan Zaanen and Olle Gunnarsson. “Charged magnetic domain lines and the magnetism of high- T_c oxides”. In: *Physical Review B* 40.10 (Oct. 1989), pp. 7391–7394. DOI: [10.1103/PhysRevB.40.7391](https://doi.org/10.1103/PhysRevB.40.7391). URL: <https://link.aps.org/doi/10.1103/PhysRevB.40.7391> (visited on 03/21/2026).
- [4] Masaru Kato et al. “Soliton Lattice Modulation of Incommensurate Spin Density Wave in Two Dimensional Hubbard Model -A Mean Field Study-”. In: *Journal of the Physical Society of Japan* 59.3 (Mar. 1990), pp. 1047–1058. ISSN: 0031-9015. DOI: [10.1143/JPSJ.59.1047](https://doi.org/10.1143/JPSJ.59.1047). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.59.1047> (visited on 03/21/2026).
- [5] B. Keimer et al. “From quantum matter to high-temperature superconductivity in copper oxides”. en. In: *Nature* 518.7538 (Feb. 2015), pp. 179–186. ISSN: 0028-0836, 1476-4687. DOI: [10.1038/nature14165](https://doi.org/10.1038/nature14165). URL: <https://www.nature.com/articles/nature14165> (visited on 03/15/2025).
- [6] Avinash Singh and Zlatko Tešanović. “Collective excitations in a doped antiferromagnet”. en. In: *Physical Review B* 41.1 (Jan. 1990), pp. 614–631. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.41.614](https://doi.org/10.1103/PhysRevB.41.614). URL: <https://link.aps.org/doi/10.1103/PhysRevB.41.614> (visited on 12/12/2025).
- [7] Avinash Singh, Zlatko Tešanović, and Ju H. Kim. “Doped Hubbard antiferromagnet: Instability and effective interactions”. en. In: *Physical Review B* 44.14 (Oct. 1991), pp. 7757–7759. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.44.7757](https://doi.org/10.1103/PhysRevB.44.7757). URL: <https://link.aps.org/doi/10.1103/PhysRevB.44.7757> (visited on 12/12/2025).
- [8] Giuseppe Grosso and Giuseppe Pastori Parravicini. *Solid State Physics*. Second Edition. Academic Press, 2014.
- [9] J. G. Bednorz and K. A. Muller. “Possible high- T_c superconductivity in the Ba-La-Cu-O system”. en. In: *Zeitschrift für Physik B Condensed Matter* 64.2 (June 1986), pp. 189–193. ISSN: 1431-584X. DOI: [10.1007/BF01303701](https://doi.org/10.1007/BF01303701). URL: <https://doi.org/10.1007/BF01303701> (visited on 04/03/2026).
- [10] Liangzi Deng et al. “Ambient-pressure 151 K superconductivity in HgBa₂Ca₂Cu₃O₈ via pressure quench”. In: *Proceedings of the National Academy of Sciences* 123.11 (Mar. 2026), e2536178123. DOI: [10.1073/pnas.2536178123](https://doi.org/10.1073/pnas.2536178123). URL: <https://www.pnas.org/doi/10.1073/pnas.2536178123> (visited on 04/03/2026).
- [11] D. J. Scalapino. “A common thread: The pairing interaction for unconventional superconductors”. en. In: *Reviews of Modern Physics* 84.4 (Oct. 2012), pp. 1383–1417. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.84.1383](https://doi.org/10.1103/RevModPhys.84.1383). URL: <https://link.aps.org/doi/10.1103/RevModPhys.84.1383> (visited on 04/03/2026).

- [12] Hari Padma et al. “Beyond-Hubbard pairing in a cuprate ladder”. en. In: *Physical Review X* 15.2 (May 2025), p. 021049. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.15.021049](https://doi.org/10.1103/PhysRevX.15.021049). URL: <https://link.aps.org/doi/10.1103/PhysRevX.15.021049> (visited on 04/03/2026).
- [13] Antoine Georges et al. “Dynamical mean-field theory of strongly correlated fermion systems and the limit of infinite dimensions”. en. In: *Reviews of Modern Physics* 68.1 (Jan. 1996), pp. 13–125. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.68.13](https://doi.org/10.1103/RevModPhys.68.13). URL: <https://link.aps.org/doi/10.1103/RevModPhys.68.13> (visited on 03/15/2025).
- [14] J. E. Hirsch. “Two-dimensional Hubbard model: Numerical simulation study”. In: *Phys. Rev. B* 31 (7 Apr. 1985), pp. 4403–4419. DOI: [10.1103/PhysRevB.31.4403](https://doi.org/10.1103/PhysRevB.31.4403). URL: <https://link.aps.org/doi/10.1103/PhysRevB.31.4403>.
- [15] Robin Scholle et al. “Comprehensive mean-field analysis of magnetic and charge orders in the two-dimensional Hubbard model”. In: *Phys. Rev. B* 108 (3 July 2023), p. 035139. DOI: [10.1103/PhysRevB.108.035139](https://doi.org/10.1103/PhysRevB.108.035139). URL: <https://link.aps.org/doi/10.1103/PhysRevB.108.035139>.
- [16] Zhangkai Cao et al. “Dominant p -wave pairing induced by nearest-neighbor attraction in the square-lattice extended Hubbard model”. In: *Phys. Rev. B* 111 (2 Jan. 2025), p. 024509. DOI: [10.1103/PhysRevB.111.024509](https://doi.org/10.1103/PhysRevB.111.024509). URL: <https://link.aps.org/doi/10.1103/PhysRevB.111.024509>.
- [17] Daniel P. Arovas et al. “The Hubbard Model”. In: *Annual Review of Condensed Matter Physics* 13. Volume 13, 2022 (2022), pp. 239–274. ISSN: 1947-5462. DOI: <https://doi.org/10.1146/annurev-conmatphys-031620-102024>. URL: <https://www.annualreviews.org/content/journals/10.1146/annurev-conmatphys-031620-102024>.
- [18] Kazuyoshi Yoshimi, Takeo Kato, and Hideaki Maebashi. “Enhanced Spin Susceptibility toward the Charge-Ordering Transition in a Two-Dimensional Extended Hubbard Model”. In: *Journal of the Physical Society of Japan* 78.10 (Oct. 2009), p. 104002. ISSN: 0031-9015. DOI: [10.1143/JPSJ.78.104002](https://doi.org/10.1143/JPSJ.78.104002). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.78.104002> (visited on 04/17/2026).
- [19] Ashvin Vishwanath. “Dirac Nodes and Quantized Thermal Hall Effect in the Mixed State of d-wave Superconductors”. In: *Physical Review B* 66.6 (Aug. 2002). arXiv:cond-mat/0104213, p. 064504. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.66.064504](https://doi.org/10.1103/PhysRevB.66.064504). URL: <http://arxiv.org/abs/cond-mat/0104213> (visited on 01/05/2026).
- [20] P. Senarath Yapa et al. “Mixed-symmetry superconductivity and the energy gap”. en. In: *Physica C: Superconductivity and its Applications* 633 (June 2025), p. 1354719. ISSN: 09214534. DOI: [10.1016/j.physc.2025.1354719](https://doi.org/10.1016/j.physc.2025.1354719). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0921453425000723> (visited on 01/06/2026).
- [21] Joel Hutchinson and Frank Marsiglio. “Mixed temperature-dependent order parameters in the extended Hubbard model”. en. In: *Journal of Physics: Condensed Matter* 33.6 (Feb. 2021), p. 065603. ISSN: 0953-8984, 1361-648X. DOI: [10.1088/1361-648X/abc801](https://doi.org/10.1088/1361-648X/abc801). URL: <https://iopscience.iop.org/article/10.1088/1361-648X/abc801> (visited on 01/06/2026).
- [22] C. Zhou and H. J. Schulz. “Density of states and tunneling spectra in two-dimensional d-wave superconductors”. In: *Physical Review B* 45.13 (Apr. 1992). Publisher: American Physical Society, pp. 7397–7401. DOI: [10.1103/PhysRevB.45.7397](https://doi.org/10.1103/PhysRevB.45.7397). URL: <https://link.aps.org/doi/10.1103/PhysRevB.45.7397> (visited on 01/06/2026).
- [23] James F. Annett. “Symmetry of the order parameter for high-temperature superconductivity”. In: *Advances in Physics* 39.2 (Apr. 1990). Publisher: Taylor & Francis eprint: <https://doi.org/10.1080/00018739000101481>, pp. 83–126. ISSN: 0001-8732. DOI: [10.1080/00018739000101481](https://doi.org/10.1080/00018739000101481). URL: <https://doi.org/10.1080/00018739000101481> (visited on 01/06/2026).

- [24] R. Micnas, J. Ranninger, and S. Robaszkiewicz. “Superconductivity in narrow-band systems with local nonretarded attractive interactions”. In: *Reviews of Modern Physics* 62.1 (Jan. 1990). Publisher: American Physical Society, pp. 113–171. DOI: [10.1103/RevModPhys.62.113](https://doi.org/10.1103/RevModPhys.62.113). URL: <https://link.aps.org/doi/10.1103/RevModPhys.62.113> (visited on 01/06/2026).
- [25] Manfred Sgrist and Kazuo Ueda. “Phenomenological theory of unconventional superconductivity”. In: *Reviews of Modern Physics* 63.2 (Apr. 1991). Publisher: American Physical Society, pp. 239–311. DOI: [10.1103/RevModPhys.63.239](https://doi.org/10.1103/RevModPhys.63.239). URL: <https://link.aps.org/doi/10.1103/RevModPhys.63.239> (visited on 01/07/2026).
- [26] Anna E. Böhrer et al. “Nematicity and nematic fluctuations in iron-based superconductors”. en. In: *Nature Physics* 18.12 (Dec. 2022), pp. 1412–1419. ISSN: 1745-2481. DOI: [10.1038/s41567-022-01833-3](https://doi.org/10.1038/s41567-022-01833-3). URL: <https://www.nature.com/articles/s41567-022-01833-3>.
- [27] Eugene Kogan and Godfrey Gumbs. “Green’s Functions and DOS for Some 2D Lattices”. en. In: *Graphene* 10.01 (2021), pp. 1–12. ISSN: 2169-3439, 2169-3471. DOI: [10.4236/graphene.2021.101001](https://doi.org/10.4236/graphene.2021.101001). URL: <https://www.scirp.org/journal/paperinformation?paperid=104591> (visited on 03/26/2026).
- [28] Charles L. Cleveland and Rodrigo Medina A. “Obtaining a Heisenberg Hamiltonian from the Hubbard model”. en. In: *American Journal of Physics* 44.1 (Jan. 1976), pp. 44–46. ISSN: 0002-9505, 1943-2909. DOI: [10.1119/1.10537](https://doi.org/10.1119/1.10537). URL: <https://pubs.aip.org/ajp/article/44/1/44/1050180/Obtaining-a-Heisenberg-Hamiltonian-from-the> (visited on 05/08/2026).
- [29] Jan Zaanen. “Watching Rush Hour in the World of Electrons”. In: *Science* 315.5817 (Mar. 2007), pp. 1372–1373. DOI: [10.1126/science.1139805](https://doi.org/10.1126/science.1139805). URL: <https://www.science.org/doi/10.1126/science.1139805> (visited on 04/03/2026).
- [30] Patrick A. Lee, Naoto Nagaosa, and Xiao-Gang Wen. “Doping a Mott insulator: Physics of high-temperature superconductivity”. en. In: *Reviews of Modern Physics* 78.1 (Jan. 2006), pp. 17–85. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.78.17](https://doi.org/10.1103/RevModPhys.78.17). URL: <https://link.aps.org/doi/10.1103/RevModPhys.78.17> (visited on 05/07/2026).
- [31] J. Brad Marston and Ian Affleck. “Large- n limit of the Hubbard-Heisenberg model”. en. In: *Physical Review B* 39.16 (June 1989), pp. 11538–11558. ISSN: 0163-1829. DOI: [10.1103/PhysRevB.39.11538](https://doi.org/10.1103/PhysRevB.39.11538). URL: <https://link.aps.org/doi/10.1103/PhysRevB.39.11538> (visited on 05/07/2026).
- [32] Mi Jiang. “Enhancing d -wave superconductivity with nearest-neighbor attraction in the extended Hubbard model”. en. In: *Physical Review B* 105.2 (Jan. 2022), p. 024510. ISSN: 2469-9950, 2469-9969. DOI: [10.1103/PhysRevB.105.024510](https://doi.org/10.1103/PhysRevB.105.024510). URL: <https://link.aps.org/doi/10.1103/PhysRevB.105.024510> (visited on 05/08/2026).
- [33] F. C. Zhang et al. “A renormalised Hamiltonian approach to a resonant valence bond wavefunction”. en. In: *Superconductor Science and Technology* 1.1 (June 1988), p. 36. ISSN: 0953-2048. DOI: [10.1088/0953-2048/1/1/009](https://doi.org/10.1088/0953-2048/1/1/009). URL: <https://iopscience.iop.org/article/10.1088/0953-2048/1/1/009> (visited on 05/11/2026).
- [34] J. Hubbard. “Electron Correlations in Narrow Energy Bands”. en. In: *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* 276.1365 (1963), pp. 238–257. URL: <http://www.jstor.org/stable/2414761>.
- [35] P. W. Anderson. “Resonating valence bonds: A new kind of insulator?” In: *Materials Research Bulletin* 8.2 (Feb. 1973), pp. 153–160. ISSN: 0025-5408. DOI: [10.1016/0025-5408\(73\)90167-0](https://doi.org/10.1016/0025-5408(73)90167-0). URL: <https://www.sciencedirect.com/science/article/pii/0025540873901670> (visited on 05/11/2026).
- [36] P. Fazekas and P. W. Anderson. “On the ground state properties of the anisotropic triangular antiferromagnet”. en. In: *Philosophical Magazine* 30.2 (Aug. 1974), pp. 423–440. ISSN: 0031-8086. DOI: [10.1080/14786439808206568](https://doi.org/10.1080/14786439808206568). URL: <http://www.tandfonline.com/doi/abs/10.1080/14786439808206568> (visited on 05/11/2026).

- [37] M. T. Béal-Monod, C. Bourbonnais, and V. J. Emery. “Possible superconductivity in nearly antiferromagnetic itinerant fermion systems”. en. In: *Physical Review B* 34.11 (Dec. 1986), pp. 7716–7720. ISSN: 0163-1829. DOI: [10.1103/PhysRevB.34.7716](https://doi.org/10.1103/PhysRevB.34.7716). URL: <https://link.aps.org/doi/10.1103/PhysRevB.34.7716> (visited on 05/11/2026).
- [38] K. Miyake, S. Schmitt-Rink, and C. M. Varma. “Spin-fluctuation-mediated even-parity pairing in heavy-fermion superconductors”. en. In: *Physical Review B* 34.9 (Nov. 1986), pp. 6554–6556. ISSN: 0163-1829. DOI: [10.1103/PhysRevB.34.6554](https://doi.org/10.1103/PhysRevB.34.6554). URL: <https://link.aps.org/doi/10.1103/PhysRevB.34.6554> (visited on 05/11/2026).
- [39] D. J. Scalapino, E. Loh, and J. E. Hirsch. “ d -wave pairing near a spin-density-wave instability”. In: *Physical Review B* 34.11 (Dec. 1986), pp. 8190–8192. DOI: [10.1103/PhysRevB.34.8190](https://doi.org/10.1103/PhysRevB.34.8190). URL: <https://link.aps.org/doi/10.1103/PhysRevB.34.8190> (visited on 05/11/2026).
- [40] Gabriel Kotliar and Dieter Vollhardt. “Strongly Correlated Materials: Insights From Dynamical Mean-Field Theory”. In: *Physics Today* 57 (Mar. 2004). ADS Bibcode: 2004PhT....57c..53K, pp. 53–59. ISSN: 0031-9228. DOI: [10.1063/1.1712502](https://ui.adsabs.harvard.edu/abs/2004PhT....57c..53K). URL: <https://ui.adsabs.harvard.edu/abs/2004PhT....57c..53K> (visited on 05/11/2026).
- [41] Eva Pavarini et al., eds. *Many-body physics: from Kondo to Hubbard: lecture notes of the Autumn School on Correlated Electrons 2015: at Forschungszentrum Jülich, 21-25 September 2015*. en. Band 5. Meeting Name: Autumn School on Correlated Electrons. Forschungszentrum Jülich, 2015. ISBN: 978-3-95806-074-6.
- [42] Piers Coleman. *Introduction to Many-Body Physics*. Cambridge University Press, 2015.
- [43] Gabriele Giuliani and Giovanni Vignale. *Quantum Theory of the Electron Liquid*. Cambridge University Press, 2005.
- [44] Michele Fabrizio. *A Course in Quantum Many-Body Theory*. Springer, 2022.
- [45] Eva Pavarini et al., eds. *Correlated electrons: from models to materials: lecture notes of the Autumn School Correlated Electrons 2012: at Forschungszentrum Jülich, 3-7 September 2012*. eng. Schriften des Forschungszentrums Jülich. Reihe Modeling and Simulation Band 2. Meeting Name: Autumn School Correlated Electrons. Jülich: Forschungszentrum Jülich, Zentralbibliothek, Verl, 2012. ISBN: 978-3-89336-796-2.