

Numerical analysis of superconducting phases in the extended Hubbard model with non-local pairing

University of Pisa, a.y. 2025-2026

Alessandro Gori*

Thesis for the Master's degree in Physics

Abstract

[To be continued. . .]

- Appendix on s -wave SC ghost simulations on HM
- Appendix on triplet pairing in MFT self-consistent scheme
- Set ComputerModern font in all CairoMakie figures

Contents

1	Mean-field theory discussion of the EHM	3
1.1	The EHM symmetries	3
1.1.1	Lattice symmetries	3
1.1.2	Spin and charge symmetries	4
1.1.3	Particle-hole symmetry at half-filling	7
1.2	MFT on Hubbard lattices	9
1.2.1	MFT general procedure, self-consistency and Wick's theorem	9
1.2.2	MFT treatment of the pure Hubbard model	10
1.3	The non-local attraction $\hat{H}_V$ as a source of symmetry-breaking interactions	10
1.3.1	Real space description	11
1.3.2	Fourier transform of the model	12
1.4	Square lattice spatial symmetry channels	14
	Bibliography	19

Draft: April 6, 2026

*a.gori23@studenti.unipi.it / nepero27178@github.com

List of symbols and abbreviations

AF	Antiferromagnetic
BCS	Bardeen-Cooper-Schrieffer (theory)
BZ	Brillouin zone
β	Inverse temperature
$\Delta^{(\gamma)}$	γ -wave component of the gap function
DoF	Degree of freedom
EHM	Extended Hubbard model
EV(s)	Expectation value(s)
FS	Fermi surface
HF	Hartree-Fock
HFP(s)	Hartree-Fock parameter(s)
HM	Hubbard model
L_ℓ	Number of lattice sites in direction ℓ
iHF	Iterative Hartree-Fock
LRT	Linear Response Theory
MBZ	Magnetic Brillouin zone
MFT	Mean-Field Theory
NBZ	Nematic Brillouin zone
NN(s)	Nearest neighbor(s)
RWC(s)	Relevant Wick's contraction(s)
RMP(s)	Renormalized model parameter(s)
SC	Superconductor
SDW	Spin-density wave
SSB	Spontaneous symmetry breaking
T_c	Critical temperature

Introduction

This thesis project is about my favorite ice cream flavor. [To be continued...]

Chapter 1

Mean-field theory discussion of the EHM

The general idea of this project is to apply HF methods to the EHM, derive analytically the approximated quadratic hamiltonian of Eq. (??) and then apply the iHF procedure sketched in Sec. ?? to extract the self-consistent values of the various HFPs. One of the advantages of using iHF methods is that we can impose rather simply the symmetries of the HF approximated wavefunction.

This is most useful when dealing with competing phases. A phase transition is basically described by some spontaneous symmetry breaking (SSB) and the rising to a non-zero value of an order parameter. We will deal with three phases:

1. the normal phase, where no symmetry is broken;
2. the antiferromagnetic phase, where we allow for spin density to be non-uniform across sites;
3. the superconducting phase, where we do not conserve particle number.

Each phase is discussed in a separate chapter. Many other “phases” are possible, based on which symmetries are respected and which are not.

For each phase, we will extract the best approximation to the true ground state wavefunction. In computational terms, we will find three solutions to the general self-consistency problem of Eq. (??). In the end we want to compare the free energy of these three solutions, assessing which one is the most stable. Any HF approximation, being by construction not the true ground state, overestimates energy. Thus, comparing energies of our solutions we get a hint of the features of the true ground state.

1.1 The EHM symmetries

Let us start by enumerating the spatial symmetries of the EHM on the square lattice. We start with those related to the spatial structure of the lattice, and then move to the ones more specific of Hubbard Physics.

1.1.1 Lattice symmetries

We study the EHM on the planar square lattice, with one orbital per site. The hamiltonian of Eq. (??) respects three fundamental lattice symmetries:

1. Discrete translations symmetry in both the x and y directions. We refer to the groups of one-site translations as $T_1^{(x)}$, $T_1^{(y)}$. It holds:

$$T_n^{(\ell)} \supset T_{kn}^{(\ell)} \quad k \in \mathbb{N} \quad (1.1)$$

A shift in direction ℓ by kn sites can be decomposed in k shift by n sites. Invariance by n -sites shifts, thus, implies invariance also for kn -sites shifts.

2. Discrete $\pi/2$ rotations symmetry. This symmetry is described by the four-fold cyclic point group C_4 , containing rotations by angle $\pi/2, \pi, 3\pi/2$ and identity. A similar property as for the translations holds for cyclic point groups,

$$C_n \supset C_{n/k} \quad k, n/k \in \mathbb{N} \quad (1.2)$$

Invariance for rotations of angle $2\pi/n$ implies invariance for rotations of angle $k \times 2\pi/n$.

3. Inversion symmetry. We define I_2 as the inversion group in two-dimensional space, containing the reflection operator and identity. This point may appear redundant, because

$$I_2 \equiv C_2 \quad (1.3)$$

In 2D space, to rotate by π around the origin and to mirror the position of a point are the same operation.

[To be continued...]

1.1.2 Spin and charge symmetries

The conventional HM shows a global $U(2)$ symmetry, extended to $SU(2) \times SU(2)/\mathbb{Z}_2 = SO(4)$ symmetry at half-filling on a bipartite lattice [9], such as the square lattice. The same symmetries are present in the EHM. Let us break down these statements.

Gauge symmetry. The EHM is symmetric under the global transformation

$$\hat{c}_{i\sigma} \rightarrow e^{-i\eta} \hat{c}_{i\sigma} \quad \hat{c}_{i\sigma}^\dagger \rightarrow e^{i\eta} \hat{c}_{i\sigma}^\dagger \quad \eta \in \mathbb{R} \quad (1.4)$$

To prove this, it suffices to note that

$$\hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rightarrow e^{i\eta} \hat{c}_{i\sigma}^\dagger e^{-i\eta} \hat{c}_{j\sigma'} = \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}$$

As a consequence, any product of fermionic operators with an equal number of annihilations and creations is symmetric. Hence the kinetic part $\hat{H}_t$ and the interactions $\hat{H}_U, \hat{H}_V$ are invariant. As a consequence, the total number of particles (charge)

$$\hat{N} = \sum_{i\sigma} \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma}$$

is a good quantum number. The symmetry is described by the transformation of Eq. (1.4), which is a pure phase shift by η . The relative group is $U^c(1)$, with a superscript c indicating “charge”.

Global spin symmetry The EHM is described by three operators: NN hopping, independent of σ ; local repulsion between opposite-spin densities; non-local attraction, independent of σ . Intuitively, the model is invariant under global spin rotations. Let us prove this rigorously.

Let the local spinor:

$$\hat{\Psi}_i \equiv \begin{bmatrix} \hat{c}_{i\uparrow} \\ \hat{c}_{i\downarrow} \end{bmatrix}$$

and the Pauli matrices

$$\tau^x \equiv \begin{bmatrix} & 1 \\ 1 & \end{bmatrix} \quad \tau^y \equiv \begin{bmatrix} & -i \\ i & \end{bmatrix} \quad \tau^z \equiv \begin{bmatrix} 1 & \\ & -1 \end{bmatrix}$$

We define the local spin operator as:

$$\hat{S}_i^\alpha \equiv \frac{1}{2} \hat{\Psi}_i^\dagger \tau^\alpha \hat{\Psi}_i \quad [\hat{S}_j^\alpha, \hat{S}_j^\beta] = i \sum_\gamma \epsilon_{\alpha\beta\gamma} \hat{S}_j^\gamma$$

with $\epsilon_{\alpha\beta\gamma}$ the Levi-Civita tensor. It can be checked by explicit calculation that the $\mathfrak{su}(2)$ spin algebra is actually respected. For example, consider $\alpha = x, \beta = y$:

$$\begin{aligned}
[\hat{S}_j^x, \hat{S}_j^y] &= \frac{1}{4} \left[\hat{\Psi}_j^\dagger \begin{bmatrix} & 1 \\ 1 & \end{bmatrix} \hat{\Psi}_j, \hat{\Psi}_j^\dagger \begin{bmatrix} & -i \\ i & \end{bmatrix} \hat{\Psi}_j \right] \\
&= \frac{1}{4} \left[\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} + \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow}, -i \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} + i \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \right] \\
&= \frac{i}{4} \left[\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow}, \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \right] - \frac{i}{4} \left[\hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow}, \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} \right] \\
&= \frac{i}{2} \left[\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow}, \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \right] \\
&= \frac{i}{2} \left(\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} - \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow} \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow} \right) \\
&= \frac{i}{2} \left(-\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \hat{c}_{j\uparrow} + \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\uparrow} + \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\uparrow}^\dagger \hat{c}_{j\uparrow} \hat{c}_{j\downarrow} - \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \right) \\
&= \frac{i}{2} \left(\hat{c}_{j\uparrow}^\dagger \hat{c}_{j\uparrow} - \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \right) \\
&= i \hat{S}_j^z
\end{aligned}$$

We used Fermi anticommutation rules, $\{\hat{c}_{i\sigma}, \hat{c}_{j\sigma'}^\dagger\} = \delta_{ij} \delta_{\sigma\sigma'}$. Identical derivations can be carried out for all other components.

Let the raising and lowering operators:

$$\hat{S}_j^+ \equiv \frac{\hat{S}_j^x + i \hat{S}_j^y}{2} \quad \hat{S}_j^- \equiv \frac{\hat{S}_j^x - i \hat{S}_j^y}{2}$$

and the global spin operators

$$\hat{S}^\alpha \equiv \sum_{i \in \mathcal{L}} \hat{S}_i^\alpha \quad [\hat{S}^\alpha, \hat{S}^\beta] = i \sum_\gamma \epsilon_{\alpha\beta\gamma} \hat{S}^\gamma$$

being $\mathcal{L}$ the lattice. Explicitly,

$$\hat{S}^z \equiv \frac{1}{2} \sum_{i \in \mathcal{L}} (\hat{n}_{i\uparrow} - \hat{n}_{i\downarrow}) \quad \hat{S}^+ \equiv \frac{1}{2} \sum_{i \in \mathcal{L}} \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow} \quad \hat{S}^- \equiv (\hat{S}^+)^\dagger$$

The EHM hamiltonian commutes with the global spin operator. For the kinetic part $\hat{H}_t$ and the local repulsion $\hat{H}_U$ this is a well known result [pavarini2015many, 9],

$$[\hat{H}_t + \hat{H}_U, \hat{\mathbf{S}}] = 0$$

For the non-local attraction, since it depends solely on total densities on neighbouring sites, spin-rotational invariance must hold. Anyway, let us give a short formal proof following this chain of results:

1. The non-local attraction can be written as:

$$\begin{aligned}
\hat{H}_V &= V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'} \\
&= V \sum_{\langle ij \rangle} \hat{\Psi}_i^\dagger \hat{\Psi}_i \hat{\Psi}_j^\dagger \hat{\Psi}_j
\end{aligned}$$

having we used

$$\hat{\Psi}_i^\dagger \hat{\Psi}_i = \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} + \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow}$$

2. Let now $\hat{R}(\boldsymbol{\theta})$ be any global spin rotation,

$$\hat{R}(\boldsymbol{\theta}) \equiv \exp(-i \boldsymbol{\theta} \cdot \hat{\mathbf{S}}) \quad \text{where} \quad \boldsymbol{\theta} = (\theta_x, \theta_y, \theta_z)$$

Spinors transform as:

$$\hat{R}(\boldsymbol{\theta})\hat{\Psi}_i\hat{R}(\boldsymbol{\theta})^\dagger = \mathcal{R}(\boldsymbol{\theta})\hat{\Psi}_i$$

with $\mathcal{R}(\boldsymbol{\theta}) \in \text{SU}(2)$ the unitary matrix associated to said rotation,

$$\mathcal{R}(\boldsymbol{\theta}) = \exp\left(-\frac{i}{2}\boldsymbol{\theta} \cdot \boldsymbol{\tau}\right) \quad \text{where} \quad \boldsymbol{\tau} = (\tau_x, \tau_y, \tau_z)$$

Let us prove this. Let $\hat{A} = i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}$ e $\hat{B} = \hat{\Psi}_i$. The following identity, a consequence of Baker-Campbell-Housdorff formula, holds:

$$e^{\hat{A}}\hat{B}e^{-\hat{A}} = \hat{B} + [\hat{A}, \hat{B}] + \frac{1}{2!}[\hat{A}, [\hat{A}, \hat{B}]] + \frac{1}{3!}[\hat{A}, [\hat{A}, [\hat{A}, \hat{B}]]] + \dots$$

The first order of the expansion gives:

$$\begin{aligned} [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, \hat{\Psi}_i] &= i \sum_{\alpha} \theta_{\alpha} [\hat{S}^{\alpha}, \hat{\Psi}_i] \\ &= -\frac{i}{2} \sum_{\alpha} \theta_{\alpha} \tau^{\alpha} \hat{\Psi}_i \\ &= -\frac{i}{2} \boldsymbol{\theta} \cdot \boldsymbol{\tau} \hat{\Psi}_i \end{aligned}$$

having we omitted some purely algebraic passages. The second order:

$$\begin{aligned} [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, \hat{\Psi}_i]] &= \left[i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, -\frac{i}{2}(\boldsymbol{\theta} \cdot \boldsymbol{\tau}) \hat{\Psi}_i \right] \\ &= -\frac{i}{2} \boldsymbol{\theta} \cdot \boldsymbol{\tau} [i\boldsymbol{\theta} \cdot \hat{\mathbf{S}}, \hat{\Psi}_i] \\ &= \frac{1}{4}(\boldsymbol{\theta} \cdot \boldsymbol{\tau})^2 \hat{\Psi}_i \end{aligned}$$

and so on. Collecting all terms,

$$\begin{aligned} \hat{R}(\boldsymbol{\theta})\hat{\Psi}_i\hat{R}(\boldsymbol{\theta})^\dagger &= \hat{\Psi}_i - \frac{i}{2}\boldsymbol{\theta} \cdot \boldsymbol{\tau} \hat{\Psi}_i + \frac{1}{2! \times 4}(\boldsymbol{\theta} \cdot \boldsymbol{\tau})^2 \hat{\Psi}_i + \dots \\ &= \sum_{k=1}^{\infty} \frac{1}{k!} \left(-\frac{i}{2}\boldsymbol{\theta} \cdot \boldsymbol{\tau}\right)^k \hat{\Psi}_i \\ &= \mathcal{R}(\boldsymbol{\theta})\hat{\Psi}_i \end{aligned}$$

which ends the sub-proof.

3. Then, the group acts on $\hat{H}_V$ as

$$\begin{aligned} \hat{R}\hat{H}_V\hat{R}^\dagger &= V \sum_{\langle ij \rangle} \left(\hat{\Psi}_i^\dagger \mathcal{R}^\dagger \mathcal{R} \hat{\Psi}_i \right) \left(\Psi_j^\dagger \mathcal{R}^\dagger \mathcal{R} \hat{\Psi}_j \right) \\ &= V \sum_{\langle ij \rangle} \left(\hat{\Psi}_i^\dagger \hat{\Psi}_i \right) \left(\hat{\Psi}_j^\dagger \hat{\Psi}_j \right) \\ &= \hat{H}_V \end{aligned}$$

making the non-local attraction spin-rotationally symmetric.

The EHM is symmetric under global spin rotations. Note that $\hat{H}_V$ is actually symmetric under local spin rotations, meaning that we could rotate the spin at each site by a different amount and get the same interaction. The same holds for $\hat{H}_U$. It is the kinetic sector that makes the EHM only globally $\text{SU}(2)$ symmetric.

Phase	SSBs
AF	$T_1^{(\ell)} \rightarrow T_2^{(\ell)}$ $SU(2) \rightarrow U^z(1)$
SC	$U^c(1) \rightarrow 0$

Table 1.1 | SSB interpretation of the AF and SC phases. In the last line, the 0 on the SSB right-hand side indicates a completely disrupted symmetry. **MOVE ME**

1.1.3 Particle-hole symmetry at half-filling

FROM HERE...

The half-filled EHM is of particular interest due to its particle-hole symmetry (PHS), which is immediately disrupted at infinitesimal doping. Let us prove that. Consider the generic unitary operation

$$P : \hat{c}_{i\sigma} \rightarrow e^{i\eta_{i\sigma}} \hat{h}_{i\sigma}^\dagger$$

being in general η_i a phase dependent on the fermionic quantum state $(i\sigma)$. Inserting the above transformation in the EHM, we get

$$P\{\hat{H}\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma})} \hat{h}_{i\sigma} \hat{h}_{j\sigma}^\dagger + U \sum_i \hat{h}_{i\uparrow} \hat{h}_{i\downarrow} \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\uparrow}^\dagger - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger$$

For the two interactions $\hat{H}_U, \hat{H}_V$ the phase factors cancelled out. We aim to find the conditions under which the above hamiltonian is mapped onto the EHM of (??),

$$P\{\hat{H}\} \stackrel{?}{=} -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma} + U \sum_i \hat{h}_{i\uparrow}^\dagger \hat{h}_{i\downarrow}^\dagger \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}$$

Before starting the derivation, which is an extension of the argument of Arovas et al. [9] about the conventional HM, let us discuss some properties of the hole operators. Due to the presence of the phase factors in the PHS operative definition, the Fermi anticommutation rules hold regardlessly of the phase factor:

$$\begin{aligned} \{\hat{h}_{i\sigma}, \hat{h}_{j\sigma'}^\dagger\} &= e^{i(\eta_{i\sigma} - \eta_{j\sigma'})} \{\hat{c}_{i\sigma}^\dagger, \hat{c}_{j\sigma'}\} \\ &= e^{i(\eta_{i\sigma} - \eta_{j\sigma'})} \delta_{ij} \delta_{\sigma\sigma'} \\ &= \delta_{ij} \delta_{\sigma\sigma'} \end{aligned}$$

and evidently:

$$\hat{h}_{i\sigma} \hat{h}_{i\sigma}^\dagger = \hat{n}_{i\sigma}$$

being $\hat{n}_{i\sigma}$ the fermionic number operator on site i with spin σ . It follows, for quartic interactions

$$\begin{aligned} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma}^\dagger &= -\hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \\ &= \hat{h}_{i\sigma} \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= -\hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} - \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger \delta_{ij} \delta_{\sigma\sigma'} \\ &= -\hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{i\sigma} \hat{h}_{j\sigma'} - \hat{h}_{i\sigma}^\dagger \hat{h}_{i\sigma} + \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^\dagger - \left(\hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \right) \delta_{ij} \delta_{\sigma\sigma'} \\ &= \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'}^\dagger \hat{h}_{j\sigma'} \hat{h}_{i\sigma} + (\hat{n}_{i\sigma} + \hat{n}_{j\sigma'}) - 1 - \left(\hat{h}_{i\sigma} \hat{h}_{j\sigma'}^\dagger - \hat{h}_{i\sigma}^\dagger \hat{h}_{j\sigma'} \right) \delta_{ij} \delta_{\sigma\sigma'} \end{aligned}$$

The first term of the last passage is mirrored with respect to the starting term. Now, let us deal with the kinetic, local and non-local interactions separately.

Kinetic term. Consider the kinetic term,

$$P\{\hat{H}_t\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma})} \hat{h}_{i\sigma} \hat{h}_{j\sigma}^{\dagger}$$

It suffices to anticommute the two fermionic operators, and exchange site labels $i \leftrightarrow j$, to get

$$P\{\hat{H}_t\} = -t \sum_{\langle ij \rangle} \sum_{\sigma} e^{i(\eta_{i\sigma} + \eta_{j\sigma} + \pi)} \hat{h}_{i\sigma}^{\dagger} \hat{h}_{j\sigma}$$

A necessary condition on the unitary operation P is the following,

$$\eta_{i\sigma} + \eta_{j\sigma} = (2k - 1)\pi \quad \text{with} \quad k \in \mathbb{Z}$$

valid for NN sites.

Local repulsion. Substitute in $\hat{H}_U$ the above derivation,

$$\begin{aligned} P\{\hat{H}_U\} &= U \sum_i \hat{h}_{i\uparrow} \hat{h}_{i\downarrow} \hat{h}_{i\downarrow}^{\dagger} \hat{h}_{i\uparrow}^{\dagger} \\ &= U \sum_i \hat{h}_{i\uparrow}^{\dagger} \hat{h}_{i\downarrow}^{\dagger} \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} + U \sum_i (\hat{n}_{i\uparrow} + \hat{n}_{i\downarrow}) - U \sum_i (1) \\ &= U \sum_i \hat{h}_{i\uparrow}^{\dagger} \hat{h}_{i\downarrow}^{\dagger} \hat{h}_{i\downarrow} \hat{h}_{i\uparrow} + U(\hat{N} - L_x L_y) \end{aligned}$$

Thus, being for the half-filled state

$$\langle \hat{N} \rangle = L_x L_y$$

preserves PHS in the local sector.

Non-local attraction. Proceed identically for $\hat{H}_V$,

$$\begin{aligned} P\{\hat{H}_V\} &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma} \hat{h}_{j\sigma'} \hat{h}_{j\sigma'}^{\dagger} \hat{h}_{i\sigma}^{\dagger} \\ &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^{\dagger} \hat{h}_{j\sigma'}^{\dagger} \hat{h}_{j\sigma'} \hat{h}_{i\sigma} - V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} (\hat{n}_{i\sigma} + \hat{n}_{j\sigma'}) + V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} (1) \\ &= -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{h}_{i\sigma}^{\dagger} \hat{h}_{j\sigma'}^{\dagger} \hat{h}_{j\sigma'} \hat{h}_{i\sigma} - 2zV(\hat{N} - L_x L_y) \end{aligned}$$

with $z = 4$ the lattice coordination number. In the last passage, we used that on the square lattice

$$\begin{aligned} \sum_{\langle ij \rangle} (1) &= \sum_{i \in \mathcal{S}/2} \sum_{j \in \text{NN}(i)} (1) \\ &= \frac{L_x L_y}{2} \times z \end{aligned} \tag{1.5}$$

which will turn out quite often. As before, an half-filled phase preserves PHS in the non-local sector. In last passage we used the fact that summing over NNs means overcounting each site $z/2$ times, and spin degeneracy.

No other condition on phases is derived, thus we conclude the half-filled EHM preserves PHS if the two sublattices $\mathcal{S}_a$, $\mathcal{S}_b$ the square lattice can be divided into, being a (trivial) bipartite lattice, differ by a π phase difference under PHS transform. In other words, if P is defined by the unitary operation

$$P : \begin{cases} \hat{c}_{i\sigma} \rightarrow \hat{h}_{i\sigma}^{\dagger} & \text{for } i \in \mathcal{S}_a \\ \hat{c}_{i\sigma} \rightarrow -\hat{h}_{i\sigma}^{\dagger} & \text{for } i \in \mathcal{S}_b \end{cases}$$

and the system is half-filled, PHS is respected. Any other phase choice, as long as it alternates by π on the two sublattices, is good enough. To summarize, we report in Tab. 1.2 all identified symmetries.

Symmetry	Operations
$T_1^{(x)} \otimes T_1^{(y)}$	One-site discrete translations separately in both directions
C_4	Four-fold point group (real $\pi/2$ angle rotations)
$U^c(1)$	Charge-conserving global phase shifts
$SU(2)$	Three-dimensional rotations in spin space
PHS	Particle-hole exchange (only at half filling)
$U^z(1)$	Reduced two-dimensional rotations in spin space

Table 1.2 | Model symmetries and relative group operations. Note that $U^c(1)$ represents the charge conserving group, while $U^z(1)$ represents the subgroup obtained by breaking the $SU(2)$ symmetry with a vector order parameter and preserving its amplitude. Since the latter is “derived” from $SU(2)$, we separate it from the four main symmetries of the model.

1.2 MFT on Hubbard lattices

This section serves as groundwork for everything that follows. We here define the MFT scheme as well as the iterative HF computational scheme we use to estimate parametrically the relevant physical quantities of the various phases. We start by recalling the general features of MFT and then move to the specificity of the EHM, apply Wick’s theorem and finally move to algorithmic aspects.

1.2.1 MFT general procedure, self-consistency and Wick’s theorem

The Mean Field Theory approach to any fermionic quartic operator is given by the following procedure: given the hamiltonian made of the quadratic operators $\hat{A}$ and $\hat{B}$

$$\hat{H} \equiv \hat{A}\hat{B}$$

we define the fluctuations

$$\delta\hat{A} \equiv \hat{A} - \langle\hat{A}\rangle \quad \delta\hat{B} \equiv \hat{B} - \langle\hat{B}\rangle$$

thus the following chain holds:

$$\begin{aligned} \hat{A}\hat{B} &= \langle\hat{A}\rangle\langle\hat{B}\rangle + \delta\hat{A}\langle\hat{B}\rangle + \langle\hat{A}\rangle\delta\hat{B} + \delta\hat{A}\delta\hat{B} \\ &= -\langle\hat{A}\rangle\langle\hat{B}\rangle + \hat{A}\langle\hat{B}\rangle + \langle\hat{A}\rangle\hat{B} + \underbrace{\delta\hat{A}\delta\hat{B}}_{\mathcal{O}(\delta)^2} \end{aligned}$$

which implies

$$\langle\hat{H}\rangle = \langle\hat{A}\rangle\langle\hat{B}\rangle + \langle\delta\hat{A}\delta\hat{B}\rangle$$

Thus, if $|\langle\hat{A}\rangle\langle\hat{B}\rangle| \gg |\langle\delta\hat{A}\delta\hat{B}\rangle|$, in a statistical sense we can neglect the $\mathcal{O}(\delta)^2$ part of the hamiltonian and approximate

$$\hat{H} \simeq \hat{A}\langle\hat{B}\rangle + \langle\hat{A}\rangle\hat{B} - \langle\hat{A}\rangle\langle\hat{B}\rangle \quad \implies \quad \langle\hat{H}\rangle \simeq \langle\hat{A}\rangle\langle\hat{B}\rangle \quad (1.6)$$

Self consistency of MFT procedure is thus embodied by the condition

$$\left| \frac{\langle\delta\hat{A}\delta\hat{B}\rangle}{\langle\hat{A}\rangle\langle\hat{B}\rangle} \right| \ll 1 \quad \implies \quad \left| \frac{\langle\hat{A}\hat{B}\rangle}{\langle\hat{A}\rangle\langle\hat{B}\rangle} - 1 \right| \ll 1 \quad (1.7)$$

which essentially states that the least the fluctuations, the better the approximation. This rule of thumb is necessary in order to interpret the physical relevance of numerical MFT results. We will come back to this in the end, for now assuming the validity of MFT approximation and deriving the mathematical results.

Term	Normal phase	Antiferromagnet	Superconductor
Hartree			
Fock	Prohibited	Prohibited	Prohibited
Cooper	Prohibited	Prohibited	

Table 1.3 | Summary of Wick’s terms selection rules and relevant contributions to the effective hamiltonian for each phase limitedly to $\hat{H}_U$.

1.2.2 MFT treatment of the pure Hubbard model

The pure Hubbard model, $\hat{H}_t + \hat{H}_U$, offers a simple framework for MFT analysis of the AF phase: this is described in detail in App. ???. The key passage is there given by the MFT approximation

$$\hat{n}_{i\uparrow}\hat{n}_{i\downarrow} \simeq \hat{n}_{i\uparrow}\langle\hat{n}_{i\downarrow}\rangle + \langle\hat{n}_{i\uparrow}\rangle\hat{n}_{i\downarrow} + (\text{constants}) \quad (1.8)$$

from which the AF structure is simply recovered. In terms of the discussion of the above paragraph, we have $\hat{A} \equiv \hat{n}_{i\uparrow}$ and $\hat{B} \equiv \hat{n}_{i\downarrow}$. However the “full” operator is given by

$$\hat{n}_{i\uparrow}\hat{n}_{i\downarrow} = \hat{c}_{i\uparrow}^\dagger \hat{c}_{j\downarrow}^\dagger \hat{c}_{j\downarrow} \hat{c}_{i\uparrow}$$

whose terms we can contract in three different ways following Wick’s Theorem:

$$\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \hat{c}_{i\uparrow}\rangle \simeq \underbrace{\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger\rangle\langle\hat{c}_{i\downarrow} \hat{c}_{i\uparrow}\rangle}_{\text{Cooper}} - \underbrace{\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}\rangle\langle\hat{c}_{i\downarrow}^\dagger \hat{c}_{i\uparrow}\rangle}_{\text{Fock}} + \underbrace{\langle\hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow}\rangle\langle\hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow}\rangle}_{\text{Hartree}} \quad (1.9)$$

where the $-$ sign in the Fock term comes from fermionic anticommutation rules.

As a first approximation, the theorem is assumed to hold (which, in a BCS-like fashion, is equivalent to assuming for the ground-state to be a coherent state). The AF ground state reduces translational invariance and rotational symmetry, as explained in Tab. 1.1. Thus, of the three terms above, when considering the pure Hubbard model in AF phase:

- Cooper fluctuations are suppressed, because they break charge conservation;
- Similarly the Fock term is null as well because if $i = j$ and $\sigma' = \bar{\sigma}$ (as is for the local interaction, which contains the operator $\hat{n}_{i\uparrow}\hat{n}_{i\downarrow}$) the expectation values involved are describing a process breaking the survivor $U^z(1)$ spin symmetry.

Thus, correctly, the Wick’s decomposition of pure HM only involves Hartree-terms of Eq. (1.9), getting us back to (1.8). The distinction between the normal phase and the AF phase is given by whether the Hartree fluctuations break or not translational symmetry: AF is indeed a specific SDW ordering which weakens the latter. In general the three parts of the decomposition (Hartree, Fock, Cooper) need to be considered altogether: this is what is done in the next chapters. What said here extends naturally to the EHM: the relevant contributions for each phase given by the local repulsion $\hat{U}$ in the EHM are listed in Tab. 1.3.

1.3 The non-local attraction $\hat{H}_V$ as a source of symmetry-breaking interactions

Let us now extend the discussion to the EHM and consider the NN non-local attraction,

$$\hat{H}_V \equiv -V \sum_{\langle\mathbf{r}\mathbf{r}'\rangle} \sum_{\sigma\sigma'} \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}'\sigma'} \quad (1.10)$$

where we changed notation, and replaced the site index i with its 2D position on the lattice

$$i \rightarrow \mathbf{r} = (x, y)$$

Let also the lattice versors,

$$\underline{\mathbf{x}} \equiv (1, 0) \quad \underline{\mathbf{y}} \equiv (0, 1)$$




Figure 1.1 | Schematic representation of the four NNs of a given site i for a planar square lattice.

as in Fig. 1.1.

Evidently the hamiltonian can be decomposed in various spin terms,

$$\begin{aligned}\hat{H}_V &= \sum_{\sigma\sigma'} \hat{H}_V^{\sigma\sigma'} \\ &= \underbrace{\hat{H}_V^{\uparrow\uparrow} + \hat{H}_V^{\downarrow\downarrow}}_{\text{Same-spin}} + \underbrace{\hat{H}_V^{\uparrow\downarrow} + \hat{H}_V^{\downarrow\uparrow}}_{\text{Opposite-spin}}\end{aligned}\quad (1.11)$$

The role of said terms is crucial in establishing the pairing channel of the dominant physical phase. As an example, the s.s. terms can be seen as a possible source of triplet pairing superconductivity. Next section are devoted to analyze said terms and derive a simple analytical result in reciprocal space.

1.3.1 Real space description

Let us start from the real space form of Eq. (1.11). To carry out a summation over nearest neighbors $\langle \mathbf{r}\mathbf{r}' \rangle$ of a square lattice means precisely to sum over all links of the lattice. Then we can identify the generic opposite-spin (o.s.) term $\hat{H}_V^{\sigma\bar{\sigma}}$ as the one collecting the σ operators of sublattice $\mathcal{S}_a$ and $\bar{\sigma}$ operators of sublattice $\mathcal{S}_b$. The o.s. non-local interactions can be written as a sum of terms over just one of the two sublattices $\mathcal{S}_a$ and $\mathcal{S}_b$. Define the site hamiltonian

$$\hat{h}_V^{(\mathbf{r})} \equiv -V \sum_{\delta} (\hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}+\delta\downarrow} + \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}-\delta\downarrow}) \quad \text{with } \delta \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\} \quad (1.12)$$

Then, it follows,

$$\begin{aligned}\hat{H}_V^{(\text{o.s.})} &= \sum_{\mathbf{r} \in \mathcal{S}_a} \overbrace{\hat{h}_V^{(\mathbf{r})}}^{\hat{H}_V^{\uparrow\downarrow}} + \sum_{\mathbf{r} \in \mathcal{S}_b} \overbrace{\hat{h}_V^{(\mathbf{r})}}^{\hat{H}_V^{\downarrow\uparrow}} \\ &= \sum_{\mathbf{r} \in \mathcal{S}} \hat{h}_V^{(\mathbf{r})}\end{aligned}\quad (1.13)$$

Here the notation of Fig. ?? is used. For each site $\mathbf{r}$ in a given sublattice, the nearest neighbors sites are four – all in the other sublattice. Similarly, the same-spin (s.s.) hamiltonian decomposes as

$$\hat{H}_V^{(\text{s.s.})} = -V \sum_{\mathbf{r} \in \mathcal{S}_a} \sum_{\delta, \sigma} (\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}+\delta\sigma} + \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r}-\delta\sigma}) \quad \text{with } \delta \in \{\underline{\mathbf{x}}, \underline{\mathbf{y}}\}$$

Opposite-spin terms. Consider first the o.s. terms of Eq. (1.12): take e.g. the NNs term $\hat{n}_{i\uparrow} \hat{n}_{\mathbf{r}+\delta\downarrow}$. As in Eq. (1.9), taking Wick's contractions,

$$\begin{aligned}\langle \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}+\delta\downarrow} \rangle &= \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\downarrow}^\dagger \hat{c}_{\mathbf{r}+\delta\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle \\ &\simeq \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\downarrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}+\delta\downarrow} \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Cooper}} - \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\downarrow} \rangle \langle \hat{c}_{\mathbf{r}+\delta\downarrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}+\delta\downarrow}^\dagger \hat{c}_{\mathbf{r}+\delta\downarrow} \rangle}_{\text{Hartree}}\end{aligned}$$

Identical decompositions are given for all others NNs. Of the three terms above:

Sector	Term	Normal phase	Antiferromagnet	Superconductor
o.s.	Hartree			
	Fock	Prohibited	Prohibited	Prohibited
	Cooper	Prohibited	Prohibited	
s.s.	Hartree			
	Fock			
	Cooper	Prohibited	Prohibited	

Table 1.4 | Summary of Wick's terms selection rules and relevant contributions to the effective hamiltonian for each phase limitedly to $\hat{H}_V$, assuming preservation of $U^z(1)$ symmetry.

- The Cooper term breaks $U^c(1)$ charge symmetry, allowing for supeconducting instabilities;
- The Fock term breaks $U^z(1)$ symmetry, because it accounts for a site hop *plus* spin flip process;
- The Hartree term can reduce both translational invariance and spin rotational symmetry.

As a general rule of thumb, throughout this text we will discard the o.s. Fock term everywhere, preserving a robust $U^z(1)$ symmetry. For the normal state as well as for the AF instability only the Hartree term is to be accounted; for the SC phase also the Cooper term contributes. Note that for superconducting instabilities, due to superexchange mechanism (as explained in App. ??) the o.s. term accounts for singlet pairing as well as zero-projection triplet pairing. Which channel is preferred, is a matter of thermodynamic advantage.

Same-spin terms. Consider then the same-spin terms: take e.g. the term $\hat{n}_{\mathbf{r}\uparrow}\hat{n}_{\mathbf{r}+\delta\uparrow}$. As above,

$$\begin{aligned}
\langle \hat{n}_{\mathbf{r}\uparrow}\hat{n}_{\mathbf{r}+\delta\uparrow} \rangle &= \langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow} \hat{c}_{\mathbf{r}\uparrow} \rangle \\
&= \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \rangle \langle \hat{c}_{\mathbf{r}+\delta\uparrow} \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Cooper}} - \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow} \rangle \langle \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{\mathbf{r}\uparrow}^\dagger \hat{c}_{\mathbf{r}\uparrow} \rangle \langle \hat{c}_{\mathbf{r}+\delta\uparrow}^\dagger \hat{c}_{\mathbf{r}+\delta\uparrow} \rangle}_{\text{Hartree}}
\end{aligned}$$

Identical consideration as in the above paragraph hold for each term. The only difference with the o.s. terms is given by the Fock term: since the spin-flip process is absent, now the Fock fluctuations actually contribute as an effective NN hopping term. In other words, this term does not break $U^z(1)$ symmetry and thus is perfectly legitimate in each phase. Tab. 1.4 summarizes the selection rules hereby presented.

1.3.2 Fourier transform of the model

It is useful to derive analytically the reciprocal-space form of the three hamiltonian parts. By the sign $\stackrel{\mathcal{F}}{=}$, we indicate a passage where a Fourier-transform is performed. $\mathcal{F}$ -transform is obtained according to the convention Fourier transform it according to the convention

$$\hat{c}_{\mathbf{r}\sigma} \stackrel{\mathcal{F}}{=} \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{-i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma} \quad (1.14)$$

Kinetic part. The kinetic part is the easiest to transform. Apply the definition,

$$\begin{aligned}
-t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r}'\sigma} + \text{h.c.}] &\stackrel{\mathcal{F}}{=} -t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{i\mathbf{k} \cdot \mathbf{r}} \hat{c}_{\mathbf{k}\sigma}^\dagger \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k}' \in \text{BZ}} e^{-i\mathbf{k}' \cdot \mathbf{r}'} \hat{c}_{\mathbf{k}'\sigma} + \text{h.c.} \\
&= -2t \sum_{\mathbf{k}, \mathbf{k}'} \sum_{\sigma} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}'\sigma} \frac{1}{L_x L_y} \sum_{\mathbf{r} \in S/2} e^{i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}} \left[e^{ik'_x} + e^{-ik'_x} + e^{ik'_y} + e^{-ik'_y} \right] \\
&= \sum_{\mathbf{k} \in \text{BZ}} \sum_{\sigma} \epsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma} \quad (1.15)
\end{aligned}$$

where the bare bands were defined,

$$\epsilon_{\mathbf{k}} \equiv -2t (\cos k_x + \cos k_y) \quad (1.16)$$

Non-local attraction. Consider a generic bond, say, the one connecting NNs $\mathbf{r}$ and $\mathbf{r} \pm \boldsymbol{\delta}$. Then:

$$\begin{aligned} \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r} \pm \boldsymbol{\delta} \sigma'} &= \hat{c}_{\mathbf{r}\sigma}^\dagger \hat{c}_{\mathbf{r} \pm \boldsymbol{\delta} \sigma'}^\dagger \hat{c}_{\mathbf{r} \pm \boldsymbol{\delta} \sigma'} \hat{c}_{\mathbf{r}\sigma} \\ &\stackrel{\mathcal{F}}{=} \frac{1}{(L_x L_y)^2} \sum_{\mathbf{k}_1, \dots, \mathbf{k}_4} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{r}} e^{\pm i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \boldsymbol{\delta}} \hat{c}_{\mathbf{k}_1 \sigma}^\dagger \hat{c}_{\mathbf{k}_2 \sigma'}^\dagger \hat{c}_{\mathbf{k}_3 \sigma'} \hat{c}_{\mathbf{k}_4 \sigma} \end{aligned}$$

Then, the interaction at site $\mathbf{r}$, spin σ with its NNs at spin σ' – indicated as $(\mathbf{r}\sigma\sigma')$ – is given by

$$\begin{aligned} (\mathbf{r}\sigma\sigma') &= -V \sum_{\boldsymbol{\delta} \in \{\mathbf{x}, \mathbf{y}\}} [\hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r} + \boldsymbol{\delta} \sigma'} + \hat{n}_{\mathbf{r}\sigma} \hat{n}_{\mathbf{r} - \boldsymbol{\delta} \sigma'}] \\ &\stackrel{\mathcal{F}}{=} -\frac{V}{(L_x L_y)^2} \sum_{\boldsymbol{\delta}} \sum_{\mathbf{k}_1, \dots, \mathbf{k}_4} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{r}} \\ &\quad \times \left(e^{i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \boldsymbol{\delta}} + e^{-i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \boldsymbol{\delta}} \right) \hat{c}_{\mathbf{k}_1 \sigma}^\dagger \hat{c}_{\mathbf{k}_2 \sigma'}^\dagger \hat{c}_{\mathbf{k}_3 \sigma'} \hat{c}_{\mathbf{k}_4 \sigma} \\ &= -\frac{2V}{(L_x L_y)^2} \sum_{\boldsymbol{\delta}} \sum_{\mathbf{k}_1, \dots, \mathbf{k}_4} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{r}} \cos[(\mathbf{k}_2 - \mathbf{k}_3) \cdot \boldsymbol{\delta}] \hat{c}_{\mathbf{k}_1 \sigma}^\dagger \hat{c}_{\mathbf{k}_2 \sigma'}^\dagger \hat{c}_{\mathbf{k}_3 \sigma'} \hat{c}_{\mathbf{k}_4 \sigma} \end{aligned}$$

The full non-local interaction is given by summing over all sites of $\mathcal{S}/2$ (which is, one sublattice of $\mathcal{S}$). This gives back momentum conservation,

$$\frac{1}{L_x L_y} \sum_{\mathbf{r} \in \mathcal{S}/2} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{r}} = \frac{1}{2} \delta_{\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4}$$

Let $\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4 = \mathbf{K}$, and define $\mathbf{k}, \mathbf{k}'$ such that

$$\mathbf{k}_1 \equiv \mathbf{K} + \mathbf{k} \quad \mathbf{k}_2 \equiv \mathbf{K} - \mathbf{k} \quad \mathbf{k}_3 \equiv \mathbf{K} - \mathbf{k}' \quad \mathbf{k}_4 \equiv \mathbf{K} + \mathbf{k}' \quad \boldsymbol{\delta} \mathbf{k} \equiv \mathbf{k} - \mathbf{k}'$$

Sums over these variables must be intended as over the Brillouin Zone (BZ). Hence:

$$\begin{aligned} \hat{H}_V &= \sum_{\mathbf{r} \in \mathcal{S}_a} \sum_{\sigma\sigma'} (\mathbf{r}\sigma\sigma') \\ &\stackrel{\mathcal{F}}{=} -\frac{V}{L_x L_y} \sum_{\sigma\sigma'} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \cos(\boldsymbol{\delta} \mathbf{k} \cdot \boldsymbol{\delta}) \hat{c}_{\mathbf{K} + \mathbf{k} \sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k} \sigma'}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}' \sigma'} \hat{c}_{\mathbf{K} + \mathbf{k}' \sigma} \\ &= -\frac{V}{L_x L_y} \sum_{\sigma\sigma'} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K} + \mathbf{k} \sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k} \sigma'}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}' \sigma'} \hat{c}_{\mathbf{K} + \mathbf{k}' \sigma} \end{aligned} \quad (1.17)$$

which decomposes in s.s. and o.s. channels as:

$$\begin{aligned} \hat{H}_V &\stackrel{\mathcal{F}}{=} \overbrace{-\frac{V}{L_x L_y} \sum_{\sigma} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K} + \mathbf{k} \sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k} \sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}' \sigma} \hat{c}_{\mathbf{K} + \mathbf{k}' \sigma}}^{\text{s.s.}} \\ &\quad - \underbrace{\frac{V}{L_x L_y} \sum_{\sigma} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K} + \mathbf{k} \sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k} \bar{\sigma}}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}' \bar{\sigma}} \hat{c}_{\mathbf{K} + \mathbf{k}' \sigma}}_{\text{o.s.}} \end{aligned}$$

The o.s. part can be simplified: since

$$\hat{c}_{\mathbf{K} + \mathbf{k} \sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k} \bar{\sigma}}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}' \bar{\sigma}} \hat{c}_{\mathbf{K} + \mathbf{k}' \sigma} = \hat{c}_{\mathbf{K} - \mathbf{k} \bar{\sigma}}^\dagger \hat{c}_{\mathbf{K} + \mathbf{k} \sigma}^\dagger \hat{c}_{\mathbf{K} + \mathbf{k}' \sigma} \hat{c}_{\mathbf{K} - \mathbf{k}' \bar{\sigma}}$$

due to fermionic anti-commutation rules, and since the composite transform

$$\mathbf{k} \rightarrow -\mathbf{k} \quad \mathbf{k}' \rightarrow -\mathbf{k}'$$

leaves the form factor $\cos(\delta k_x) + \cos(\delta k_y)$ invariant, we reduce the spin sum to two times the $\sigma = \uparrow$ part,

$$\hat{H}_V^{(\text{o.s.})} = -\frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \quad (1.18)$$

whiel the explicit s.s. sector is given immediately by:

$$\begin{aligned} \hat{H}_V^{(\text{s.s.})} = & -\frac{V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \\ & \times \left[\hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\uparrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} + \hat{c}_{\mathbf{K}+\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\downarrow} \right] \end{aligned} \quad (1.19)$$

The result here obtained will be used various times in next chapters. Note that different Wick contraction schemes will be referred to as:

$$\overbrace{c_{\mathbf{K}+\mathbf{k}\sigma}^\dagger c_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger c_{\mathbf{K}-\mathbf{k}'\sigma'} c_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{Cooper contraction}} \quad (1.20)$$

$$\overbrace{c_{\mathbf{K}+\mathbf{k}\sigma}^\dagger c_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger c_{\mathbf{K}-\mathbf{k}'\sigma'} c_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{Fock contraction}} \quad (1.21)$$

$$\overbrace{c_{\mathbf{K}+\mathbf{k}\sigma}^\dagger c_{\mathbf{K}-\mathbf{k}\sigma'}^\dagger c_{\mathbf{K}-\mathbf{k}'\sigma'} c_{\mathbf{K}+\mathbf{k}'\sigma}}^{\text{Hartree contraction}} \quad (1.22)$$

When contracting in the Fock channel, a minus sign must be included.

Local repulsion. A somewhat identical argument can be carried out for the local interaction,

$$\hat{H}_U = U \sum_{\mathbf{r} \in \mathcal{S}} \hat{n}_{\mathbf{r}\uparrow} \hat{n}_{\mathbf{r}\downarrow}$$

The only difference here is made by the fact that the interaction is repulsive on-site, thus the final structure factor in the Fourier transform disappears as well as the minus sign,

$$\hat{H}_U \stackrel{\mathcal{F}}{=} \frac{U}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \quad (1.23)$$

When contracting these fermionic operators, identical considerations as for the non-local counterpart (1.17) hold. Note the similarity of this Fourier transform with the o.s. sector of the non-local attraction, given in Eq. (1.17). With this, we conclude the general introduction to the model and pass to the setup of the Hartree-Fock (HF) algorithm.

1.4 Square lattice spatial symmetry channels

In next sections we introduce spatial symmetry structures and discuss their relation with point groups. We will always refer to various spatial structures as “symmetry channels”, throughout the system is able to establish a given order parameter defined in space.

Looking at $\hat{H}$, the main difference between its two-body interactions $\hat{H}_U$ and $\hat{H}_V$ of course relies in locality (first) and non-locality (second). Both operators in principle are able to break C_4 symmetry, as will later be discussed extensively.

For the 2D square lattice, various spatial symmetries are possible. Let i be a specific site, and consider a generic complex function g_{ij} whose domain is given by the site itself and its four NNs (the domain concides with the points of Fig. 1.1). The function can be written in the C_4 basis

$$g_{ij} = \sum_{\gamma} g^{(\gamma)} \varphi_{ij}^{(\gamma)}$$

where $g^{(\gamma)} \in \mathbb{C}$ are the g_{ij} symmetry channels coefficients, while $\varphi_{ij}^{(\gamma)}$ are the form factors listed in Tab. 1.5 and plotted in Fig. 1.3, a simple orthonormal rearrangement of the harmonics basis.

When dealing with phases which respect translational symmetry, or at most reduce it to a weaker symmetry by shrinking the Brillouin Zone, it's useful to work in reciprocal space, as discussed in Sec. 1.3. As said, a particular phase can show specific spatial symmetries, and so will the




Figure 1.2 Form factors at different topologies, as listed in Tab. 1.5. In figures five sites are represented: the hub site i and its four NN. Solid lines represent non-zero values for φ_{δ} , while dashed lines represent vanishing factors.

Structure	Form factor	Graph
s-wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(s)} = \delta_{\mathbf{r}\mathbf{r}'}$	Fig. 1.2a
Extended s-wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(s^*)} = \frac{1}{2} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 1.2b
p_x -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(p_x)} = \frac{1}{\sqrt{2}} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}}]$	Fig. 1.2c
p_y -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(p_y)} = \frac{1}{\sqrt{2}} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 1.2d
$d_{x^2-y^2}$ -wave	$\varphi_{\mathbf{r}\mathbf{r}'}^{(d)} = \frac{1}{2} [\delta_{\mathbf{r}'=\mathbf{r}+\underline{x}} + \delta_{\mathbf{r}'=\mathbf{r}-\underline{x}} - \delta_{\mathbf{r}'=\mathbf{r}+\underline{y}} - \delta_{\mathbf{r}'=\mathbf{r}-\underline{y}}]$	Fig. 1.2e

Table 1.5 Form factors for a function defined on a square lattice, with a prefactor included for normalization. We use lattice vector notation here. The last column indicates the graph representation of each contribution given in Fig. 1.2. Subscript $x^2 - y^2$ is omitted for notational clarity.

Hartree-Fock parameters. The key point to notice is that self-consistency equations involve sums (or integrals) to be computed over fermionic states, which can be heavily simplified by employing said symmetries. Consider the form factors of Tab. 1.5, and their Fourier transforms of Tab. 1.6. They satisfy the relations:

$$\frac{1}{L_x L_y} \sum_{ij} \varphi_{ij}^{(\gamma)} \varphi_{ij}^{(\gamma)*} = \delta_{\gamma\gamma'} \quad \frac{1}{L_x L_y} \sum_{\mathbf{k}} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)*} = \delta_{\gamma\gamma'}$$

If these symmetries are developed by the HFPs, it's of no use to integrate over the entire BZ, since different regions linked by periodicity of the form factors will lead to redundant calculations.

Structure	Structure factor	Graph
s -wave	$\varphi_{\mathbf{k}}^{(s)} = 1$	Fig. 1.2a
Extended s -wave	$\varphi_{\mathbf{k}}^{(s^*)} = \cos k_x + \cos k_y$	Fig. 1.2b
p_x -wave	$\varphi_{\mathbf{k}}^{(p_x)} = i\sqrt{2} \sin k_x$	Fig. 1.2c
p_y -wave	$\varphi_{\mathbf{k}}^{(p_y)} = i\sqrt{2} \sin k_y$	Fig. 1.2d
$d_{x^2-y^2}$ -wave	$\varphi_{\mathbf{k}}^{(d)} = \cos k_x - \cos k_y$	Fig. 1.2e

Table 1.6 | Structure factors derived from the correlation structures of Tab. 1.5. The functions hereby defined are orthonormal, and are plotted in Fig- 1.3.




Figure 1.3 | Representation of the various structure factors listed in Tab. 1.6 on the BZ. For the s^* and $d_{x^2-y^2}$ -wave factors, the real part is plotted; for the p_ℓ factors, the imaginary part is plotted.

Finally, a very useful detail is given by the following argument. Consider the equation

$$\cos(\delta k_x) + \cos(\delta k_y) = \cos k_x \cos k'_x + \sin k_x \sin k'_x + \cos k_y \cos k'_y + \sin k_y \sin k'_y$$

For the sake of readability, the notations

$$c_\ell \equiv \cos k_\ell \quad s_\ell \equiv \sin k_\ell \quad c'_\ell \equiv \cos k'_\ell \quad s'_\ell \equiv \sin k'_\ell$$

are used. Group the four terms above,

$$\underbrace{(c_x c'_x + c_y c'_y)}_{\text{Symmetric}} + \underbrace{(s_x s'_x + s_y s'_y)}_{\text{Anti-symmetric}} \quad (1.24)$$

The first two exhibit inversion symmetry for both arguments $\mathbf{k}, \mathbf{k}'$; the second two exhibit anti-symmetry. Decoupling the symmetric part, we obtain

$$c_x c'_x + c_y c'_y = \frac{1}{2}(c_x + c_y)(c'_x + c'_y) + \frac{1}{2}(c_x - c_y)(c'_x - c'_y)$$

which finally gives:

$$\begin{aligned} \cos(\delta k_x) + \cos(\delta k_y) &= \frac{1}{2}(c_x + c_y)(c'_x + c'_y) && (s^*\text{-wave}) \\ &+ s_x s'_x && (p_x\text{-wave}) \\ &+ s_y s'_y && (p_y\text{-wave}) \\ &+ \frac{1}{2}(c_x - c_y)(c'_x - c'_y) && (d_{x^2-y^2}\text{-wave}) \end{aligned}$$

or, in a more compact form

$$\cos(\delta k_x) + \cos(\delta k_y) = \frac{1}{2} \sum_{\gamma} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} \quad \text{for } \gamma \in \{s^*, p_x, p_y, d_{x^2-y^2}\} \quad (1.25)$$

$$= \frac{1}{2} \sum_{\gamma} \varphi_{\mathbf{k}}^{(\gamma)*} \varphi_{\mathbf{k}'}^{(\gamma)} \quad (1.26)$$

This decomposition will become particularly handy in later calculations.

Bibliography

- [1] J. G. Bednorz and K. A. Muller. “Possible high- T_c superconductivity in the Ba-La-Cu-O system”. en. In: *Zeitschrift für Physik B Condensed Matter* 64.2 (June 1986), pp. 189–193. ISSN: 1431-584X. DOI: [10.1007/BF01303701](https://doi.org/10.1007/BF01303701). URL: <https://doi.org/10.1007/BF01303701> (visited on 04/03/2026).
- [2] Liangzi Deng et al. “Ambient-pressure 151 K superconductivity in HgBa₂Ca₂Cu₃O₈ via pressure quench”. In: *Proceedings of the National Academy of Sciences* 123.11 (Mar. 2026), e2536178123. DOI: [10.1073/pnas.2536178123](https://www.pnas.org/doi/10.1073/pnas.2536178123). URL: <https://www.pnas.org/doi/10.1073/pnas.2536178123> (visited on 04/03/2026).
- [3] B. Keimer et al. “From quantum matter to high-temperature superconductivity in copper oxides”. en. In: *Nature* 518.7538 (Feb. 2015), pp. 179–186. ISSN: 0028-0836, 1476-4687. DOI: [10.1038/nature14165](https://www.nature.com/articles/nature14165). URL: <https://www.nature.com/articles/nature14165> (visited on 03/15/2025).
- [4] D. J. Scalapino. “A common thread: The pairing interaction for unconventional superconductors”. en. In: *Reviews of Modern Physics* 84.4 (Oct. 2012), pp. 1383–1417. ISSN: 0034-6861, 1539-0756. DOI: [10.1103/RevModPhys.84.1383](https://link.aps.org/doi/10.1103/RevModPhys.84.1383). URL: <https://link.aps.org/doi/10.1103/RevModPhys.84.1383> (visited on 04/03/2026).
- [5] Hari Padma et al. “Beyond-Hubbard pairing in a cuprate ladder”. en. In: *Physical Review X* 15.2 (May 2025), p. 021049. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.15.021049](https://link.aps.org/doi/10.1103/PhysRevX.15.021049). URL: <https://link.aps.org/doi/10.1103/PhysRevX.15.021049> (visited on 04/03/2026).
- [6] J. E. Hirsch. “Two-dimensional Hubbard model: Numerical simulation study”. In: *Phys. Rev. B* 31 (7 Apr. 1985), pp. 4403–4419. DOI: [10.1103/PhysRevB.31.4403](https://link.aps.org/doi/10.1103/PhysRevB.31.4403). URL: <https://link.aps.org/doi/10.1103/PhysRevB.31.4403>.
- [7] Robin Scholle et al. “Comprehensive mean-field analysis of magnetic and charge orders in the two-dimensional Hubbard model”. In: *Phys. Rev. B* 108 (3 July 2023), p. 035139. DOI: [10.1103/PhysRevB.108.035139](https://link.aps.org/doi/10.1103/PhysRevB.108.035139). URL: <https://link.aps.org/doi/10.1103/PhysRevB.108.035139>.
- [8] Zhangkai Cao et al. “Dominant p -wave pairing induced by nearest-neighbor attraction in the square-lattice extended Hubbard model”. In: *Phys. Rev. B* 111 (2 Jan. 2025), p. 024509. DOI: [10.1103/PhysRevB.111.024509](https://link.aps.org/doi/10.1103/PhysRevB.111.024509). URL: <https://link.aps.org/doi/10.1103/PhysRevB.111.024509>.
- [9] Daniel P. Arovas et al. “The Hubbard Model”. In: *Annual Review of Condensed Matter Physics* 13. Volume 13, 2022 (2022), pp. 239–274. ISSN: 1947-5462. DOI: <https://doi.org/10.1146/annurev-conmatphys-031620-102024>. URL: <https://www.annualreviews.org/content/journals/10.1146/annurev-conmatphys-031620-102024>.
- [10] Avinash Singh and Zlatko Tešanović. “Collective excitations in a doped antiferromagnet”. en. In: *Physical Review B* 41.1 (Jan. 1990), pp. 614–631. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.41.614](https://link.aps.org/doi/10.1103/PhysRevB.41.614). URL: <https://link.aps.org/doi/10.1103/PhysRevB.41.614> (visited on 12/12/2025).
- [11] Alexander Wietek et al. “Stripes, Antiferromagnetism, and the Pseudogap in the Doped Hubbard Model at Finite Temperature”. en. In: *Physical Review X* 11.3 (July 2021), p. 031007. ISSN: 2160-3308. DOI: [10.1103/PhysRevX.11.031007](https://link.aps.org/doi/10.1103/PhysRevX.11.031007). URL: <https://link.aps.org/doi/10.1103/PhysRevX.11.031007> (visited on 12/12/2025).

- [12] Ashvin Vishwanath. “Dirac Nodes and Quantized Thermal Hall Effect in the Mixed State of d-wave Superconductors”. In: *Physical Review B* 66.6 (Aug. 2002). arXiv:cond-mat/0104213, p. 064504. ISSN: 0163-1829, 1095-3795. DOI: [10.1103/PhysRevB.66.064504](https://doi.org/10.1103/PhysRevB.66.064504). URL: <http://arxiv.org/abs/cond-mat/0104213> (visited on 01/05/2026).
- [13] P. Senarath Yapa et al. “Mixed-symmetry superconductivity and the energy gap”. en. In: *Physica C: Superconductivity and its Applications* 633 (June 2025), p. 1354719. ISSN: 09214534. DOI: [10.1016/j.physc.2025.1354719](https://doi.org/10.1016/j.physc.2025.1354719). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0921453425000723> (visited on 01/06/2026).
- [14] Joel Hutchinson and Frank Marsiglio. “Mixed temperature-dependent order parameters in the extended Hubbard model”. en. In: *Journal of Physics: Condensed Matter* 33.6 (Feb. 2021), p. 065603. ISSN: 0953-8984, 1361-648X. DOI: [10.1088/1361-648X/abc801](https://doi.org/10.1088/1361-648X/abc801). URL: <https://iopscience.iop.org/article/10.1088/1361-648X/abc801> (visited on 01/06/2026).
- [15] C. Zhou and H. J. Schulz. “Density of states and tunneling spectra in two-dimensional d-wave superconductors”. In: *Physical Review B* 45.13 (Apr. 1992). Publisher: American Physical Society, pp. 7397–7401. DOI: [10.1103/PhysRevB.45.7397](https://doi.org/10.1103/PhysRevB.45.7397). URL: <https://link.aps.org/doi/10.1103/PhysRevB.45.7397> (visited on 01/06/2026).
- [16] James F. Annett. “Symmetry of the order parameter for high-temperature superconductivity”. In: *Advances in Physics* 39.2 (Apr. 1990). Publisher: Taylor & Francis _eprint: <https://doi.org/10.1080/00018739000101481>, pp. 83–126. ISSN: 0001-8732. DOI: [10.1080/00018739000101481](https://doi.org/10.1080/00018739000101481). URL: <https://doi.org/10.1080/00018739000101481> (visited on 01/06/2026).
- [17] R. Micnas, J. Ranninger, and S. Robaszkiewicz. “Superconductivity in narrow-band systems with local nonretarded attractive interactions”. In: *Reviews of Modern Physics* 62.1 (Jan. 1990). Publisher: American Physical Society, pp. 113–171. DOI: [10.1103/RevModPhys.62.113](https://doi.org/10.1103/RevModPhys.62.113). URL: <https://link.aps.org/doi/10.1103/RevModPhys.62.113> (visited on 01/06/2026).
- [18] Manfred Sigrist and Kazuo Ueda. “Phenomenological theory of unconventional superconductivity”. In: *Reviews of Modern Physics* 63.2 (Apr. 1991). Publisher: American Physical Society, pp. 239–311. DOI: [10.1103/RevModPhys.63.239](https://doi.org/10.1103/RevModPhys.63.239). URL: <https://link.aps.org/doi/10.1103/RevModPhys.63.239> (visited on 01/07/2026).
- [19] Anna E. Böhrer et al. “Nematicity and nematic fluctuations in iron-based superconductors”. en. In: *Nature Physics* 18.12 (Dec. 2022), pp. 1412–1419. ISSN: 1745-2481. DOI: [10.1038/s41567-022-01833-3](https://doi.org/10.1038/s41567-022-01833-3). URL: <https://www.nature.com/articles/s41567-022-01833-3>.
- [20] Simons Collaboration on the Many-Electron Problem et al. “Absence of Superconductivity in the Pure Two-Dimensional Hubbard Model”. In: *Physical Review X* 10.3 (July 2020), p. 031016. DOI: [10.1103/PhysRevX.10.031016](https://doi.org/10.1103/PhysRevX.10.031016). URL: <https://link.aps.org/doi/10.1103/PhysRevX.10.031016> (visited on 03/21/2026).
- [21] Jan Zaanen and Olle Gunnarsson. “Charged magnetic domain lines and the magnetism of high- T_c oxides”. In: *Physical Review B* 40.10 (Oct. 1989), pp. 7391–7394. DOI: [10.1103/PhysRevB.40.7391](https://doi.org/10.1103/PhysRevB.40.7391). URL: <https://link.aps.org/doi/10.1103/PhysRevB.40.7391> (visited on 03/21/2026).
- [22] Masaru Kato et al. “Soliton Lattice Modulation of Incommensurate Spin Density Wave in Two Dimensional Hubbard Model -A Mean Field Study-”. In: *Journal of the Physical Society of Japan* 59.3 (Mar. 1990), pp. 1047–1058. ISSN: 0031-9015. DOI: [10.1143/JPSJ.59.1047](https://doi.org/10.1143/JPSJ.59.1047). URL: <https://journals.jps.jp/doi/10.1143/JPSJ.59.1047> (visited on 03/21/2026).
- [23] Eugene Kogan and Godfrey Gumbs. “Green’s Functions and DOS for Some 2D Lattices”. en. In: *Graphene* 10.01 (2021), pp. 1–12. ISSN: 2169-3439, 2169-3471. DOI: [10.4236/graphene.2021.101001](https://doi.org/10.4236/graphene.2021.101001). URL: <https://www.scirp.org/journal/paperinformation?paperid=104591> (visited on 03/26/2026).
- [24] Piers Coleman. *Introduction to Many-Body Physics*. Cambridge University Press, 2015.

- [25] Giuseppe Grosso and Giuseppe Pastori Parravicini. *Solid State Physics*. Second Edition. Academic Press, 2014.
- [26] Gabriele Giuliani and Giovanni Vignale. *Quantum Theory of the Electron Liquid*. Cambridge University Press, 2005.
- [27] Michele Fabrizio. *A Course in Quantum Many-Body Theory*. Springer, 2022.