

# Numerical analysis of superconducting phases in the extended Hubbard model with non-local pairing

University of Pisa, a.y. 2025-2026

Alessandro Gori\*

Thesis for the Master's degree in Physics

## Abstract

[To be continued...]

## Contents

<b>1</b>	<b>Mean-field theory analysis of the EHM</b>	<b>1</b>
1.1	Mean-Field theory real space description . . . . .	1
1.1.1	The non-local term as a source of symmetry-breaking interactions . . . . .	2
1.1.2	Reciprocal-space transform of the non-local interaction . . . . .	4
1.2	Anti-Ferromagnetic instability . . . . .	5
1.2.1	Hartree renormalization of chemical potential and gap . . . . .	5
1.2.2	Fock renormalization of the hopping amplitude . . . . .	7
1.3	Superconducting instability . . . . .	8
1.3.1	Mean-field treatment of the non-local term . . . . .	9
1.3.2	Mean-field treatment of the local term . . . . .	9
1.3.3	Topological correlations . . . . .	10
1.4	Mean-Field theory reciprocal space description . . . . .	11
1.4.1	Kinetic term . . . . .	11
1.4.2	Non-local attraction . . . . .	11
1.4.3	Local interaction and gap function . . . . .	13
1.4.4	Nambu formalism and Bogoliubov transform . . . . .	14
1.4.5	A short comment on self-consistency . . . . .	17
1.5	Results of the HF algorithmtm . . . . .	17
	<b>Bibliography</b>	<b>19</b>

*Draft: September 26, 2025*

---

\*a.gori23@studenti.unipi.it / nepero27178@github.com

# Chapter 1

## Mean-field theory analysis of the EHM

This chapter is devoted to develop a Mean Field Theory (MFT) approximation of the Extended Hubbard model (EHM) of Eq. (??),

$$\hat{H} = \underbrace{-t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma}}_{\hat{H}_t} + \underbrace{U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow}}_{\hat{H}_U} - \underbrace{V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'}}_{\hat{H}_V}$$

Mean Field Theory (MFT) is a widely used and simple theoretical tool, often sufficient to describe the leading orders in phase transition phenomena of Many-Body Physics. Here MFT is employed to discuss both the effects of the non-local term  $\hat{H}_V$  onto the AF phase, as well as the insurgence of anisotropic superconductivity – following the path of Bardeen-Cooper-Schrieffer (BCS) theory in describing conventional *s*-wave superconductivity. As will be thoroughly described, the lattice spatial structure directly influences the topology of the gap function, giving rise to anisotropic pairing. Sec. 1.1 studies the non-local attraction  $\hat{H}_V$  in real-space, describing how such interaction can contribute to the hamiltonian as a symmetry-breaking term in given channels. In the following sections, we move to specific channels and study theoretically and numerically the effect of non-local attraction.

### 1.1 Mean-Field theory real space description

The general aim is to study the phase diagram of the model by comparing ground-state energies of different phases. The phases we consider here for the EHM are the Anti-Ferromagnetic ordering (AF), given by a non-uniform distribution of charge in each spin sector, and the Superconducting phase, described by a uniformly distributed charge allowing for Cooper pairing instabilities. For the EHM, three symmetries are “brekable”:

1. (Crystal) translational invariance. By breaking explicitly this symmetry, the obtained state must show a Charge-Density Wave (CDW) ordering;
2. U(1) charge conservation. By breaking this symmetry, we allow for Cooper fluctuations originating superconductivity;
3. SU(2) spin conservation. By breaking this symmetry, we take in all the processes not conserving spin (such as site hop *plus* spin flip).

As a general rule, SU(2) symmetry will be preserved while charge and space symmetries will be alternatively broken, in order to generate a pure AF phase or a pure superconducting phase. “To preserve one symmetry” means to set to zero all operators which break said symmetry, due to selection rules.

App. ?? describes in detail the MFT treatment of the pure Hubbard model,  $\hat{H}_t + \hat{H}_U$ ; the key passage is there given by the approximation

$$\hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \simeq \hat{n}_{i\uparrow} \langle \hat{n}_{i\downarrow} \rangle + \langle \hat{n}_{i\uparrow} \rangle \hat{n}_{i\downarrow} + (\text{constants}) \quad (1.1)$$




Figure 1.1 | Schematic representation of the four NNs of a given site  $i$  for a planar square lattice.

from which the AF structure is simply recovered. However, to perform the above approximation coherently, we are implementing Wick's Theorem on the generic term:

$$\langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle \simeq \underbrace{\langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'}^\dagger \rangle \langle \hat{c}_{j\sigma'} \hat{c}_{i\sigma} \rangle}_{\text{Cooper}} - \underbrace{\langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma'} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{i\sigma} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma} \rangle \langle \hat{c}_{j\sigma'}^\dagger \hat{c}_{j\sigma'} \rangle}_{\text{Hartree}} \quad (1.2)$$

As a first approximation, the theorem is assumed to hold (which, in a BCS-like fashion, is equivalent to assuming for the ground-state to be a coherent state). Of the three terms above, the AF phase description for the pure Hubbard Model only allows for the Hartree term to be non-zero. Cooper fluctuations are suppressed, because only the translational invariance is broken, and the Fock term is null as well because if  $i = j$  and  $\sigma' = \bar{\sigma}$  (as is for the local interaction, which contains the operator  $\hat{n}_{i\uparrow} \hat{n}_{i\downarrow}$ ) the expectation values involved are describing a process breaking SU(2) spin symmetry. Thus, correctly, the Wick's decomposition of Eq. (1.1) only involves Hartree-terms of Eq. (1.2). In general, however, the three terms need to be considered altogether: this is what is done in the next section.

### 1.1.1 The non-local term as a source of symmetry-breaking interactions

Consider now the NN non-local term,

$$\hat{H}_V \equiv -V \sum_{\langle ij \rangle} \sum_{\sigma\sigma'} \hat{n}_{i\sigma} \hat{n}_{j\sigma'} \quad (1.3)$$

Evidently the hamiltonian can be decomposed in various spin terms,

$$\begin{aligned} \hat{H}_V &= \sum_{\sigma\sigma'} \hat{H}_V^{\sigma\sigma'} \\ &= \underbrace{\hat{H}_V^{\uparrow\uparrow} + \hat{H}_V^{\downarrow\downarrow}}_{\text{Same-spin}} + \underbrace{\hat{H}_V^{\uparrow\downarrow} + \hat{H}_V^{\downarrow\uparrow}}_{\text{Opposite-spin}} \end{aligned}$$

Evidently, to carry out a summation over nearest neighbors  $\langle ij \rangle$  of a square lattice means precisely to sum over all links of the lattice. Then we can identify the generic opposite-spin (o.s.) term  $\hat{H}_V^{\sigma\bar{\sigma}}$  as the one collecting the  $\sigma$  operators of sublattice  $\mathcal{S}_a$  and  $\bar{\sigma}$  operators of sublattice  $\mathcal{S}_b$ . The o.s. non-local interactions can be written as a sum of terms over just one of the two sublattices  $\mathcal{S}_a$  and  $\mathcal{S}_b$ , oppositely polarized in the AF configuration (see Fig. ??)

$$\begin{aligned} \hat{H}_V^{(\text{o.s.})} &= \overbrace{\sum_{i \in \mathcal{S}_a} \hat{h}_V^{(i)}}^{\hat{H}_V^{\uparrow\downarrow}} + \overbrace{\sum_{i \in \mathcal{S}_b} \hat{h}_V^{(i)}}^{\hat{H}_V^{\downarrow\uparrow}} \\ &= \sum_{i \in \mathcal{S}} \hat{h}_V^{(i)} \end{aligned} \quad \hat{h}_V^{(i)} = -V \sum_{\ell=x,y} (\hat{n}_{i\uparrow} \hat{n}_{i+\delta_\ell\downarrow} + \hat{n}_{i\downarrow} \hat{n}_{i-\delta_\ell\downarrow})$$

Here the notation of Fig. ?? is used. The two-dimensional lattice is regular-square. For each site  $i$  in a given sublattice, the nearest neighbors sites are four – all in the other sublattice. The notation used is  $i \pm \delta_x$ ,  $i \pm \delta_y$  as in Fig. 1.1. Similarly, the same-spin (s.s.) hamiltonian decomposes as

$$\hat{H}_V^{(\text{s.s.})} = -V \sum_{i \in \mathcal{S}_a} \sum_{\ell=x,y} \sum_{\sigma} (\hat{n}_{i\sigma} \hat{n}_{i+\delta_\ell\sigma} + \hat{n}_{i\sigma} \hat{n}_{i-\delta_\ell\sigma})$$

Note here the summation only on one sublattice. The non-local interaction contribution to energy, as a function of the  $T = 0$  full hamiltonian ground-state<sup>1</sup>  $|\Psi\rangle$ , is given by

$$\begin{aligned} E_V[\Psi] &= \langle \Psi | \hat{H}_V | \Psi \rangle \\ &= -V \sum_{\langle ij \rangle} \sum_{\sigma \sigma'} \langle \hat{n}_{i\sigma} \hat{n}_{j\sigma'} \rangle \\ &= -V \underbrace{\sum_{\langle ij \rangle} \sum_{\sigma} \langle \hat{n}_{i\sigma} \hat{n}_{j\sigma} \rangle}_{\text{s.s.}} - V \underbrace{\sum_{\langle ij \rangle} \sum_{\sigma} \langle \hat{n}_{i\sigma} \hat{n}_{j\bar{\sigma}} \rangle}_{\text{o.s.}} \end{aligned}$$

Shorthand notation has been used:  $\langle \Psi | \cdot | \Psi \rangle = \langle \cdot \rangle$ . The ground-state must realize the condition

$$\frac{\delta}{\delta \langle \Psi |} E[\Psi] = 0$$

being  $E[\Psi]$  the total energy (made up of the three terms of couplings  $t$ ,  $U$  and  $V$ ). [\[Expand derivation?\]](#) The functional derivative must be carried out in a variational fashion including a Lagrange multiplier, the latter accounting for state-norm conservation, as is done normally in deriving the Hartree-Fock approximation for the eigenenergies of the electron liquid [4, 5].

**Opposite-spin terms.** Consider first the o.s. terms: take e.g. the term  $\hat{n}_{i\uparrow} \hat{n}_{i+\delta_x\downarrow}$ . As in Eq. (1.2), Wick's Theorem states that, if the expectation value is performed onto a coherent state,

$$\begin{aligned} \langle \hat{n}_{i\uparrow} \hat{n}_{i+\delta_x\downarrow} \rangle &= \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i+\delta_x\downarrow}^\dagger \hat{c}_{i+\delta_x\downarrow} \hat{c}_{i\uparrow} \rangle \\ &= \underbrace{\langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i+\delta_x\downarrow}^\dagger \rangle \langle \hat{c}_{i+\delta_x\downarrow} \hat{c}_{i\uparrow} \rangle}_{\text{Cooper}} - \underbrace{\langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i+\delta_x\downarrow} \rangle \langle \hat{c}_{i+\delta_x\downarrow}^\dagger \hat{c}_{i\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle \langle \hat{c}_{i+\delta_x\downarrow}^\dagger \hat{c}_{i+\delta_x\downarrow} \rangle}_{\text{Hartree}} \end{aligned}$$

Identical decompositions are given for all others NNs. Of the three terms above:

- The Cooper term breaks U(1) charge symmetry, allowing for superconducting instabilities;
- The Fock term breaks the SU(2) symmetry, because it accounts for a site hop *plus* spin flip process;
- The Hartree term breaks translational invariance, because the mean-field to interact with is given by the local density;

Then, to look for AF instability only the Hartree term is to be accounted; instead, in superconducting instability only the Cooper term contributes. The Fock term we always assume to be suppressed, preserving SU(2) symmetry. Note that for superconducting instabilities, due to superexchange mechanism (as explained in App. ??) the o.s. term account for singlet pairing as well as zero-spin triplet pairing. Which channel is preferred, is a matter of thermodynamic advantage.

**Same-spin terms.** Consider then the same-spin terms: take e.g. the term  $\hat{n}_{i\uparrow} \hat{n}_{i+\delta_x\uparrow}$ . As above,

$$\begin{aligned} \langle \hat{n}_{i\uparrow} \hat{n}_{i+\delta_x\uparrow} \rangle &= \langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i+\delta_x\uparrow}^\dagger \hat{c}_{i+\delta_x\uparrow} \hat{c}_{i\uparrow} \rangle \\ &= \underbrace{\langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i+\delta_x\uparrow}^\dagger \rangle \langle \hat{c}_{i+\delta_x\uparrow} \hat{c}_{i\uparrow} \rangle}_{\text{Cooper}} - \underbrace{\langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i+\delta_x\uparrow} \rangle \langle \hat{c}_{i+\delta_x\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle}_{\text{Fock}} + \underbrace{\langle \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \rangle \langle \hat{c}_{i+\delta_x\uparrow}^\dagger \hat{c}_{i+\delta_x\uparrow} \rangle}_{\text{Hartree}} \end{aligned}$$

Identical consideration as in the above paragraph hold for each term. The only difference with the o.s. terms is given by the Fock term: since the spin-flip process is absent, now the Fock fluctuations actually contribute to the AF phase as an effective NN hopping term [\[Unclear: does the renormalization happen also for the Cooper phase?\]](#). As a final remark, notice that the superconducting instabilities of the s.s. terms account only for triplet pairing. The only possible superconducting ordering established by the means of these terms is odd in real space. Then *s*-wave and *d*-wave superconductivity cannot establish in this channel; *p<sub>ℓ</sub>*-wave superconductivity, instead, can.

<sup>1</sup>Extensions to finite temperatures is simple: minimization must be carried out on free energy, while expectation values must be taken in a thermodynamic fashion.

### 1.1.2 Reciprocal-space transform of the non-local interaction

It is useful to derive analytically the reciprocal-space form of the non-local attraction. Consider a generic bond, say, the one connecting sites  $j$  and  $j \pm \delta_\ell$  (variable  $i$  is here referred to as the imaginary unit to avoid confusion).  $\mathbf{x}_j$  is the 2D notation for the position of site  $j$ , while  $\delta_\ell$  is the 2D notation for the lattice spacing previously indicated as  $\delta_\ell$ . Fourier transform it according to the convention

$$\hat{c}_{j\sigma} = \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{-i\mathbf{k} \cdot \mathbf{x}_j} \hat{c}_{\mathbf{k}\sigma}$$

Then:

$$\begin{aligned} \hat{n}_{j\sigma} \hat{n}_{j \pm \delta_\ell \sigma'} &= \hat{c}_{j\sigma}^\dagger \hat{c}_{j \pm \delta_\ell \sigma'}^\dagger \hat{c}_{j \pm \delta_\ell \sigma'} \hat{c}_{j\sigma} \\ &= \frac{1}{(L_x L_y)^2} \sum_{\nu=1}^4 \sum_{\mathbf{k}_\nu \in \text{BZ}} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} e^{\pm i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_\ell} \hat{c}_{\mathbf{k}_1 \sigma}^\dagger \hat{c}_{\mathbf{k}_2 \sigma'}^\dagger \hat{c}_{\mathbf{k}_3 \sigma'} \hat{c}_{\mathbf{k}_4 \sigma} \end{aligned}$$

Then, the interaction at site  $j$ , spin  $\sigma$  with its NNs at spin  $\sigma'$  – indicated as  $(j\sigma\sigma')$  – is given by

$$\begin{aligned} (j\sigma\sigma') &= -V \sum_{\ell=x,y} \sum_{\delta=\pm\delta_\ell} \hat{n}_{j\sigma} \hat{n}_{j \pm \delta_\ell \sigma'} \\ &= -\frac{V}{(L_x L_y)^2} \sum_{\ell=x,y} \sum_{\nu=1}^4 \sum_{\mathbf{k}_\nu \in \text{BZ}} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} \\ &\quad \times \left( e^{i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_\ell} + e^{-i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_\ell} \right) \hat{c}_{\mathbf{k}_1 \sigma}^\dagger \hat{c}_{\mathbf{k}_2 \sigma'}^\dagger \hat{c}_{\mathbf{k}_3 \sigma'} \hat{c}_{\mathbf{k}_4 \sigma} \\ &= -\frac{2V}{(L_x L_y)^2} \sum_{\ell=x,y} \sum_{\nu=1}^4 \sum_{\mathbf{k}_\nu \in \text{BZ}} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} \cos[(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_\ell] \hat{c}_{\mathbf{k}_1 \sigma}^\dagger \hat{c}_{\mathbf{k}_2 \sigma'}^\dagger \hat{c}_{\mathbf{k}_3 \sigma'} \hat{c}_{\mathbf{k}_4 \sigma} \end{aligned}$$

The full non-local interaction is given by summing over all sites of  $\mathcal{S}_a$  (which is, half the sites of  $\mathcal{S}$ ). This gives back momentum conservation,

$$\frac{1}{L_x L_y} \sum_{j \in \mathcal{S}_a} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} = \frac{1}{2} \delta_{\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4}$$

Let  $\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4 = \mathbf{K}$ , and define  $\mathbf{k}, \mathbf{k}'$  such that

$$\mathbf{k}_1 \equiv \mathbf{K} + \mathbf{k} \quad \mathbf{k}_2 \equiv \mathbf{K} - \mathbf{k} \quad \mathbf{k}_3 \equiv \mathbf{K} - \mathbf{k}' \quad \mathbf{k}_4 \equiv \mathbf{K} + \mathbf{k}' \quad \delta \mathbf{k} \equiv \mathbf{k} - \mathbf{k}'$$

Sums over these variables must be intended as over the Brillouin Zone (BZ). Then, finally

$$\begin{aligned} \hat{H}_V &= \sum_{j \in \mathcal{S}_a} \sum_{\sigma\sigma'} (j\sigma\sigma') \\ &= -\frac{V}{L_x L_y} \sum_{\ell=x,y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \cos(\delta \mathbf{k} \cdot \delta_\ell) \hat{c}_{\mathbf{K} + \mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}'\sigma'} \hat{c}_{\mathbf{K} + \mathbf{k}'\sigma} \\ &= -\frac{V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x) + \cos(\delta k_y)] \hat{c}_{\mathbf{K} + \mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}'\sigma'} \hat{c}_{\mathbf{K} + \mathbf{k}'\sigma} \end{aligned} \quad (1.4)$$

Different Wick contraction schemes lead to different results:

$$\overbrace{\hat{c}_{\mathbf{K} + \mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}'\sigma'} \hat{c}_{\mathbf{K} + \mathbf{k}'\sigma}}^{\text{Cooper contraction}} \quad (1.5)$$

$$\overbrace{\hat{c}_{\mathbf{K} + \mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}'\sigma'} \hat{c}_{\mathbf{K} + \mathbf{k}'\sigma}}^{\text{Fock contraction}} \quad (1.6)$$

$$\overbrace{\hat{c}_{\mathbf{K} + \mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}\sigma'}^\dagger \hat{c}_{\mathbf{K} - \mathbf{k}'\sigma'} \hat{c}_{\mathbf{K} + \mathbf{k}'\sigma}}^{\text{Hartree contraction}} \quad (1.7)$$

[To be continued...]

## 1.2 Anti-Ferromagnetic instability

In this section, the effect of the non-local interaction on the antiferromagnetic phase is discussed. The MFT derivation for the “standard” Hubbard Model is discussed in App. ???. Recalling the main results, the Anti-Ferromagnetic phase specified by the Ansatz (??) (which is explicitly breaking translational invariance in each spin sector, while preserving SU(2) and U(1) symmetries) reduces the hamiltonian to the form of Eq. (??)

$$\hat{H}_t + \hat{H}_U \stackrel{\text{MFT}}{\simeq} -t \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \hat{c}_{\mathbf{r}\sigma}^{\dagger} \hat{c}_{\mathbf{r}'\sigma} + nU \sum_{\mathbf{r}} [\hat{n}_{\mathbf{r}\uparrow} + \hat{n}_{\mathbf{r}\downarrow}] - mU \sum_{\mathbf{r}} (-1)^{x+y} [\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}]$$

In reciprocal space, the hamiltonian decomposes as in Eq. (??),

$$\hat{H}_t + \hat{H}_U \stackrel{\text{MFT}}{\simeq} \sum_{\mathbf{k} \in \text{MBZ}} \sum_{\sigma} \hat{\Psi}_{\mathbf{k}\sigma}^{\dagger} h_{\mathbf{k}\sigma} \hat{\Psi}_{\mathbf{k}\sigma} \quad \text{being} \quad h_{\mathbf{k}\sigma} \equiv \begin{bmatrix} \epsilon_{\mathbf{k}} & -\Delta_{\sigma} \\ -\Delta_{\sigma} & -\epsilon_{\mathbf{k}} \end{bmatrix}$$

and  $\Delta_{\uparrow} = mU$ ,  $\Delta_{\downarrow} = -mU$ . Nambu spinorial formulation is used,

$$\hat{\Psi}_{\mathbf{k}\sigma} \equiv \begin{bmatrix} \hat{c}_{\mathbf{k}\sigma} \\ \hat{c}_{\mathbf{k}+\pi\sigma} \end{bmatrix}$$

and the free electrons energy is simply the tight binding energy

$$\epsilon_{\mathbf{k}} = -2t [\cos(k_x) + \cos(k_y)]$$

which is spin-invariant. The MFT description of the model reduces to a gas of free “ $\gamma$ -fermions”, described by the Nambu spinor of Eq. (??),

$$\hat{\Gamma}_{\mathbf{k}\sigma} = W_{\mathbf{k}\sigma} \hat{\Psi}_{\mathbf{k}\sigma} = \begin{bmatrix} \hat{\gamma}_{\mathbf{k}\sigma}^{(-)} \\ \hat{\gamma}_{\mathbf{k}\sigma}^{(+)} \end{bmatrix}$$

where

$$W_{\mathbf{k}\sigma} = \begin{bmatrix} -\sin \theta_{\mathbf{k}\sigma} & -\cos \theta_{\mathbf{k}\sigma} \\ \cos \theta_{\mathbf{k}\sigma} & -\sin \theta_{\mathbf{k}\sigma} \end{bmatrix} \quad \text{and} \quad \sin 2\theta_{\mathbf{k}\sigma} \equiv \frac{\Delta_{\sigma}}{E_{\mathbf{k}}}$$

These fermions populate the two bands  $\pm E_{\mathbf{k}} = \sqrt{\epsilon_{\mathbf{k}}^2 + \Delta^2}$ .

Consider now the non-local interaction  $\hat{H}_V$ : since only translational invariance is broken in the AF phase, the only relevant contributions coming from Wick’s decomposition are Hartree terms and the same-spin Fock term. The net effect obtained by including this interaction, as I will explain, is a renormalization of the various quantities,

$$\epsilon_{\mathbf{k}} \rightarrow \tilde{\epsilon}_{\mathbf{k}\sigma} \quad E_{\mathbf{k}} \rightarrow \tilde{E}_{\mathbf{k}\sigma} \quad \Delta_{\sigma} \rightarrow \tilde{\Delta}_{\mathbf{k}\sigma} \quad \theta_{\mathbf{k}\sigma} \rightarrow \tilde{\theta}_{\mathbf{k}\sigma}$$

The band energies renormalization is simply

$$\tilde{E}_{\mathbf{k}\sigma} \equiv \sqrt{\tilde{\epsilon}_{\mathbf{k}\sigma}^2 + |\tilde{\Delta}_{\mathbf{k}\sigma}|^2}$$

The physical behavior is the same as for the pure Hubbard model. In next sections the different contributions to renormalization are treated.

### 1.2.1 Hartree renormalization of chemical potential and gap

The same-spin and opposite-spin non-local Hartree terms are

$$\underbrace{-V \sum_{\langle ij \rangle} \sum_{\sigma} [\langle \hat{n}_{i\sigma} \rangle \hat{n}_{j\sigma} + \hat{n}_{i\sigma} \langle \hat{n}_{j\sigma} \rangle]}_{\text{s.s.}} - \underbrace{V \sum_{\langle ij \rangle} \sum_{\sigma} [\langle \hat{n}_{i\sigma} \rangle \hat{n}_{j\bar{\sigma}} + \hat{n}_{i\sigma} \langle \hat{n}_{j\bar{\sigma}} \rangle]}_{\text{o.s.}}$$

Let  $i \rightarrow \mathbf{r} = (x, y)$  and  $j \rightarrow \mathbf{r}' = (x', y')$ . Then, using the Ansatz of Eq. ??, summarized as

$$\langle \hat{n}_{\mathbf{r}\sigma} \rangle = n - (-1)^{x+y+\delta_{\sigma=\uparrow}} m$$

we get

$$\begin{aligned} & \overbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{n}_{\mathbf{r}'\sigma} + \hat{n}_{\mathbf{r}\sigma}] + mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{\delta_{\sigma=\uparrow}} \left[ (-1)^{x'+y'} \hat{n}_{\mathbf{r}'\sigma} + (-1)^{x+y} \hat{n}_{\mathbf{r}\sigma} \right]}^{\text{s.s.}} \\ & \underbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{n}_{\mathbf{r}'\bar{\sigma}} + \hat{n}_{\mathbf{r}\sigma}] + mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} \left[ (-1)^{x'+y'+\delta_{\bar{\sigma}=\uparrow}} \hat{n}_{\mathbf{r}'\bar{\sigma}} + (-1)^{x+y+\delta_{\sigma=\uparrow}} \hat{n}_{\mathbf{r}\sigma} \right]}_{\text{o.s.}} \end{aligned}$$

For a square lattice, if  $\mathbf{r} = (x, y)$  and  $\mathbf{r}' = (x', y')$  are NNs evidently

$$(-1)^{x'+y'} = (-1)^{x+y+1}$$

Moreover,

$$(-1)^{\delta_{\bar{\sigma}=\uparrow}} = (-1)^{\delta_{\sigma=\uparrow}+1}$$

We obtain

$$\begin{aligned} & \overbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\sigma}] + mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{x+y+\delta_{\sigma=\uparrow}} [\hat{n}_{\mathbf{r}\sigma} - \hat{n}_{\mathbf{r}'\sigma}]}^{\text{s.s.}} \\ & \underbrace{-nV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\bar{\sigma}}] + mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{x+y+\delta_{\sigma=\uparrow}} [\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\sigma}]}_{\text{o.s.}} \quad (1.8) \end{aligned}$$

Now, since

$$\sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\sigma}] = \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} [\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\bar{\sigma}}] = z\hat{N}$$

with  $z = 4$  the lattice coordination factor, indicating the number of NNs per site, then the first and third terms of Expr. (1.8) contribute to a pure chemical potential shift. The renormalized chemical potential is:

$$\tilde{\mu} \equiv \mu + 2znV \quad (1.9)$$

The second and fourth terms of Expr. (1.8) are to be reduced to a renormalization of the gap function. Explicitly,

$$\begin{aligned} & mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{x+y+\delta_{\sigma=\uparrow}} [\hat{n}_{\mathbf{r}\sigma} - \hat{n}_{\mathbf{r}'\sigma}] + mV \sum_{\langle \mathbf{r}\mathbf{r}' \rangle} \sum_{\sigma} (-1)^{x+y+\delta_{\sigma=\uparrow}} [\hat{n}_{\mathbf{r}\sigma} + \hat{n}_{\mathbf{r}'\sigma}] \\ & = -2zmV \sum_{\mathbf{r}} (-1)^{x+y} [\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}] \quad (1.10) \end{aligned}$$

Consider now the last term of the pure Hubbard model under MFT approximations of Eq. (??),

$$-mU \sum_{\mathbf{r}} (-1)^{x+y} [\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow}] \quad (\text{Local gap})$$

Expr. (1.10) is formally identical, thus we obtain a contribution to the renormalization of the AF gap,

$$\Delta \rightarrow \Delta + 2zmV + (\text{s.s. contribution}) \quad (1.11)$$

This, together with Eq. (1.9), concludes the non-local Hartree reparametrization of the hamiltonian. Next section is devoted to analyzing the effect of the Fock term.

### 1.2.2 Fock renormalization of the hopping amplitude

From Wick's decomposition of  $\hat{H}_V$ , the only allowed Fock term comes from the same-spin part due to SU(2) symmetry selection rules. Said hamiltonian term is

$$V \sum_{\langle ij \rangle} \sum_{\sigma} \left[ \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} \rangle \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} + \text{h.c.} \right] \quad (1.12)$$

(note the + sign in front of it). A bond-wise hopping amplitude can be defined,

$$\tilde{t}_{ij\sigma} \equiv t - V \langle \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} \rangle$$

Then:

$$\hat{H}_t + V \sum_{\langle ij \rangle} \sum_{\sigma} \left[ \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} \rangle \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} + \text{h.c.} \right] = - \sum_{\langle ij \rangle} \sum_{\sigma} \left[ \tilde{t}_{ij\sigma} \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} + \text{h.c.} \right]$$

In reciprocal space, the effective hopping must be transformed as well. Consider the Fourier Transform given in Eq. (1.4), applied to Eq. (1.12),

$$\begin{aligned} V \sum_{\langle ij \rangle} \sum_{\sigma} \left[ \langle \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} \rangle \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} + \text{h.c.} \right] \\ = \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \sum_{\sigma} [\cos(\delta k_x) + \cos(\delta k_y)] \langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}'\sigma} \end{aligned} \quad (1.13)$$

where the 2 prefactor comes from recognizing that the h.c. generates an identical contribution to the full sum.

In order to proceed, it is now necessary to understand how the AF phase is realized in reciprocal space. As is exposed in App. ??, to impose an AF Ansatz of the form

$$\langle \hat{n}_{\mathbf{r}\sigma} \rangle = n - (-1)^{x+y+\delta_{\sigma=\uparrow}} m \implies \langle \hat{n}_{\mathbf{k}\sigma} \rangle = n \delta_{\mathbf{k}=\mathbf{0}} - \underbrace{(-1)^{\delta_{\sigma=\uparrow}} m}_{m_\sigma} \delta_{\mathbf{k}=\boldsymbol{\pi}} \quad (1.14)$$

leads to an AF ground-state of free fermions at temperature  $\beta$  described by the Nambu spinor of Eq. (??). All parameters are renormalized, thus we must account for renormalized band energies  $\pm \tilde{E}_{\mathbf{k}\sigma}$  as well. The ground-state is realized by simply populating the two bands  $\pm \tilde{E}_{\mathbf{k}\sigma}$  as

$$\bigotimes_{\mathbf{k} \in \text{MBZ}} \bigotimes_{\sigma} \left[ \left( \hat{\gamma}_{\mathbf{k}\sigma}^{(-)} \right)^\dagger f(-\tilde{E}_{\mathbf{k}}; \beta, \mu) + \left( \hat{\gamma}_{\mathbf{k}\sigma}^{(+)} \right)^\dagger f(\tilde{E}_{\mathbf{k}}; \beta, \mu) \right] |\Omega\rangle$$

The  $\hat{\gamma}$  operators are normalized superpositions of two  $\hat{c}$  operators at points in reciprocal space separated by a  $\boldsymbol{\pi}$  shift. It follows that the above state is ultimately a superposition of many-body pure states, each of which has either the  $\mathbf{k}\sigma$  state occupied *or* the  $\mathbf{k} + \boldsymbol{\pi}\sigma$  state for each  $\mathbf{k} \in \text{MBZ}$ ,  $\sigma \in \{\uparrow, \downarrow\}$ . It follows that, when computing generically  $\langle \hat{c}_{\mathbf{k}_1\sigma}^\dagger \hat{c}_{\mathbf{k}_2\sigma} \rangle$ , such expectation value can be non-zero if and only if  $\mathbf{k}_1 = \mathbf{k}_2 + n\boldsymbol{\pi}$ , being  $n \in \mathbb{Z}$ . Going back to Eq. (1.13), this implies only two contributions are non-zero:


$$\mathbf{k} = -\mathbf{k}' \quad \text{or} \quad \mathbf{k} + \boldsymbol{\pi} = -\mathbf{k}'$$

Then Eq. (1.13) is reduced to:

$$\begin{aligned} \frac{2V}{L_x L_y} \sum_{\mathbf{K}, \mathbf{k}} \sum_{\sigma} [\cos(2k_x) + \cos(2k_y)] \\ \left[ \underbrace{\langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}}_{\text{Diagonal terms}} - \underbrace{\langle \hat{c}_{\mathbf{K}+\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}+\mathbf{k}+\boldsymbol{\pi}\sigma} \rangle \hat{c}_{\mathbf{K}-\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}-\boldsymbol{\pi}\sigma}}_{\text{Off-diagonal terms}} \right] \end{aligned} \quad (1.15)$$

Now, the above equation presents *diagonal* and *off-diagonal* terms. Let me discuss them separately.





**Figure 1.2** Representation of the vectors involved in the diagonal terms of Eq. (1.15). In Fig. 1.2a the vectors are generic, and density operator at position  $\mathbf{K} - \mathbf{k}$  (dashed line) interacts with the mean field at position  $\mathbf{K} + \mathbf{k}$  (solid line) of the BZ (light gray area). The red dots at  $\mathbf{0}$  and  $\pi$  represents the only non-zero reciprocal Fourier factors of the averaged density for the AF phase. In Fig. 1.2b, starting from a generic point  $\mathbf{q} \in \text{BZ}$ , a construction of the kind of Fig. 1.2a always exists connecting  $\mathbf{q}$  to  $\pi$ , and analogously to  $\mathbf{0}$ .

**Diagonal terms.** The diagonal terms of Eq. (1.15) are simple density interaction with the mean density field. Consider Fig. 1.2a: density at vector  $\mathbf{q} \equiv \mathbf{K} - \mathbf{k}$  interacts with the mean density at vector  $\mathbf{K} + \mathbf{k}$ . Consider 1.2a: from Eq. (1.14), the only non-zero components of the averaged density in AF phase are at reciprocal points  $\mathbf{0}, \pi$  (in Figure indicated in red). To obtain a nonzero contribution to the hamiltonian, it must hold

$$\mathbf{K} + \mathbf{k} = \mathbf{0} \quad \text{or} \quad \mathbf{K} + \mathbf{k} = \pi$$

Changing perspective, taking any point  $\mathbf{q} \in \text{BZ}$  as in Fig. 1.2b, it can always be connected to each interesting points by exactly one construction. Then, when performing the sum  $\sum_{\mathbf{K}, \mathbf{k}}$  of Eq. (1.15), we get one contribution from the  $\mathbf{0}$  point (where the density amplitude is  $n$ ) and one for each corner of the BZ (where the density amplitude is  $-m_\sigma$ ), because of the reciprocal space tiling. The diagonal part reduces to:

$$\frac{2V}{L_x L_y} \sum_{\mathbf{k}\sigma} [\cos(2k_x) + \cos(2k_y)] \left[ \underbrace{(n - 4m_\sigma) \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{\mathbf{k}\sigma}}_{\text{Diagonal terms}} + (\text{Off-diagonal terms}) \right] \quad (1.16)$$

The tight-binding spectrum is effectively renormalized:

$$\tilde{\epsilon}_{\mathbf{k}\sigma} \equiv \epsilon_{\mathbf{k}} + \frac{2V}{L_x L_y} [\cos(2k_x) + \cos(2k_y)] (n - 4m_\sigma)$$

[I am pretty sure the size denominator should not be there...]

### 1.3 Superconducting instability

This section is devoted to studying the superconducting phase of the system. The only symmetry we assume to break is the U(1) charge symmetry, thus allowing for superconducting fluctuations. The symmetry structure of the pairing mechanism determines the contributing Cooper fluctuations: for  $s$ -wave and  $d$ -wave superconductivity, only the o.s. Cooper term contributes; for  $p_\ell$ -wave

superconductivity, the s.s. term contributes as well [Include hopping renormalization?]. In the following sections, a derivation containing both Cooper terms is proposed.

[To be continued: separate singlet and triplet pairing channels, and describe triplet channel by itself by the means of four-components Nambu spinors. Use selection rules to set  $\Delta^{(p_\ell)} = 0$  in the singlet channel, in order to justify results obtained by a pure space-even simulation containing just the o.s. terms.]

### 1.3.1 Mean-field treatment of the non-local term

This approach leads to the conclusion that the (coherent) ground-state of the system must be an eigenstate of the mean-field effective hamiltonian:

$$\begin{aligned} \hat{H}^{(e)} = & -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_{i \in \mathcal{S}} \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \\ & - V \sum_{i \in \mathcal{S}} \sum_{\ell=x,y} \sum_{\delta=\pm\delta_{\ell}} \left[ \langle \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{i+\delta\downarrow}^{\dagger} \rangle \hat{c}_{i+\delta\downarrow} \hat{c}_{i\uparrow} + \text{h.c.} \right] \end{aligned} \quad (1.17)$$

Note that I am here summing over  $i \in \mathcal{S}$ : this is the same as considering both the  $\uparrow\downarrow$  *plus* the  $\downarrow\uparrow$  terms of the o.s. hamiltonian involved in even-wave pairing. The pairing correlation function is defined across each bond as the pairing expectation

$$g_{ij} \equiv \langle \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{j\downarrow}^{\dagger} \rangle$$

The effective hamiltonian reads:

$$\hat{H}^{(e)} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_{i \in \mathcal{S}} \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} - V \sum_{\langle ij \rangle} \left[ g_{ij} \hat{c}_{j\downarrow} \hat{c}_{i\uparrow} + g_{ij}^* \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{j\downarrow}^{\dagger} \right] \quad (1.18)$$

As in standard BCS theory, this hamiltonian – being quadratic in the electronic operators – can be diagonalized via a Bogoliubov rotation. Superconducting pairing can arise both from the local  $U$  term and from the non-local  $V$  term. In next sections it is assumed the  $V$  term generates dominant superconductivity via its weak non-local pairing.

### 1.3.2 Mean-field treatment of the local term

The mean-field description of the local (on-site)  $U$  interaction is given in detail in App. ??, along with a simple numerical analysis of the insurgence of antiferromagnetic ordering in a Hartree-Fock approximation scheme. Here the Cooper pairing is likewise assumed to dominate. Performing an analysis analogous to the one carried out in last section, we get the decoupling

$$U \sum_{i \in \mathcal{S}} \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \simeq U \sum_{i \in \mathcal{S}} \left[ f_i \hat{c}_{i\downarrow} \hat{c}_{i\uparrow} + f_i^* \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{i\downarrow}^{\dagger} \right]$$

being

$$f_i \equiv \langle \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{i\downarrow}^{\dagger} \rangle$$


Collect  $f$  and  $g$  in the unique function of two variables:

$$C(i, j) = \begin{cases} f_i & \text{if } i = j \\ g_{ij} & \text{if } |i - j| = 1 \\ (\dots) & \text{otherwise} \end{cases}$$

which expresses the generic correlator  $\langle \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{j\downarrow}^{\dagger} \rangle$ . The correlator for  $|i - j| > 1$  is left unexpressed, and supposed to be subdominant. The decoupled hamiltonian, apart from pure energy shifts and suppressed terms, is given by

$$\begin{aligned} \hat{H}^{(e)} = & -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_i \left[ f_i \hat{c}_{i\downarrow} \hat{c}_{i\uparrow} + f_i^* \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{i\downarrow}^{\dagger} \right] \\ & - V \sum_{\langle ij \rangle} \left[ g_{ij} \hat{c}_{j\downarrow} \hat{c}_{i\uparrow} + g_{ij}^* \hat{c}_{i\uparrow}^{\dagger} \hat{c}_{j\downarrow}^{\dagger} \right] \end{aligned} \quad (1.19)$$

[To be continued...]



**Figure 1.3** | Form factors at different topologies, as listed in Tab. 1.1. In figures five sites are represented: the hub site  $i$  and its four NN. Solid lines represent non-zero values for  $\varphi_\delta$ , while dashed lines represent vanishing factors.

Structure	Form factor	Graph
$s$ -wave	$\varphi_{ij}^{(s)} = \delta_{ij}$	Fig. 1.3a
Extended $s$ -wave	$\varphi_{ij}^{(s^*)} = \delta_{j=i+\delta_x} + \delta_{j=i-\delta_x} + \delta_{j=i+\delta_y} + \delta_{j=i-\delta_y}$	Fig. 1.3b
$p_x$ -wave	$\varphi_{ij}^{(p_x)} = \delta_{j=i+\delta_x} - \delta_{j=i-\delta_x}$	Fig. 1.3c
$p_y$ -wave	$\varphi_{ij}^{(p_y)} = \delta_{j=i+\delta_y} - \delta_{j=i-\delta_y}$	Fig. 1.3d
$d_{x^2-y^2}$ -wave	$\varphi_{ij}^{(d)} = \delta_{j=i+\delta_x} + \delta_{j=i-\delta_x} - \delta_{j=i+\delta_y} - \delta_{j=i-\delta_y}$	Fig. 1.3e

**Table 1.1** | First four spatial structures for the correlation function  $C(i, j)$ . In the middle column, all spatial dependence is included in the  $\delta$ s, while  $f^s, g^{(\gamma)} \in \mathbb{C}$ . The last column indicates the graph representation of each contribution given in Fig. ???. Subscript  $x^2 - y^2$  is omitted for notational clarity.

### 1.3.3 Topological correlations

Topology plays an important role in establishing SC, giving rise to anisotropic pairing as well as real space structures for the Cooper pairs. The correlator  $g_{ij}$  is a function of position, specifically of its variables difference  $\delta \equiv \mathbf{x}_j - \mathbf{x}_i$ . Over the square lattice with NN interaction, the latter can assume four values:  $\delta = \pm\delta_x, \pm\delta_y$ . For a function of space defined over the four rim sites  $\mathbf{x}_i \pm \delta_\ell$  of Fig. 1.1, various symmetry structures can be defined under the planar rotations group  $\text{SO}(2)$ . In other words, the function  $g_\delta$  can be decomposed in planar harmonics (which are simply the sine-cosine basis). Equivalently, given two NN sites  $i, j$

$$g_{ij} = \sum_{\gamma} g^{(\gamma)} \varphi_{ij}^{(\gamma)}$$

where  $g^{(\gamma)}$  are the  $g_{ij}$  symmetries-decomposition coefficients while  $\varphi_{ij}^{(\gamma)}$  are the form factors listed in Tab. 1.1, a simple orthonormal rearrangement of the harmonics basis.

SC is established with a given symmetry – which means, symmetry breaking in the phase transition proceeds in a specific channel. Conventional BCS superconductivity arises from the only possible spatial structure of the local pairing,  $s$ -wave – here appearing as a local term (Fig. 1.3a)

and extended on a non-local term (Fig. 1.3b). Cuprates exhibit a tendency towards  $d_{x^2-y^2}$  SC, while other materials towards  $p$ -wave types – eventually with some chirality, as is the case for  $p_x \pm ip_y$  SCs. To establish SC under a certain symmetry  $\gamma$  means that Cooper pairs acquire said symmetry – which implies, for correlations,  $g^{(\gamma')} = g^{(\gamma)}\delta_{\gamma\gamma'}$ . and  $g_{ij} \propto \varphi_{ij}^{(\gamma)}$ .

## 1.4 Mean-Field theory reciprocal space description

In this BCS-like approach, a self-consistent equation for the gap function must be retrieved in order to further investigate the model and extract the conditions for the formation of a superconducting phase with a given pairing topology. In order to do so, let me take a step back and perform explicitly the Fourier-transform of the various terms of Eq. ??.

### 1.4.1 Kinetic term

The kinetic part is trivial to transform. The followed convention is

$$\hat{c}_{j\sigma} = \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{-i\mathbf{k} \cdot \mathbf{x}_j} \hat{c}_{\mathbf{k}\sigma}$$

Calculation is carried out in App. ?.?. Let

$$\epsilon_{\mathbf{k}} \equiv -2t [\cos(k_x \delta_x) + \cos(k_y \delta_y)]$$

then we have

$$\begin{aligned} -t \sum_{\langle ij \rangle} \sum_{\sigma} \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} &= \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}} \hat{c}_{\mathbf{k}\sigma}^{\dagger} \hat{c}_{\mathbf{k}\sigma} \\ &= \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} \left[ \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} + \hat{c}_{\mathbf{k}\downarrow}^{\dagger} \hat{c}_{\mathbf{k}\downarrow} \right] \\ &= \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} \left[ \hat{c}_{\mathbf{k}\uparrow}^{\dagger} \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow}^{\dagger} \hat{c}_{-\mathbf{k}\downarrow} \right] \end{aligned}$$

In last passage I used fermionic anti-commutation rules and reversed the sign of the mute variable. This will become useful later.

### 1.4.2 Non-local attraction

Consider a generic bond, say, the one connecting sites  $j$  and  $j \pm \delta_{\ell}$  (variable  $i$  is here referred to as the imaginary unit to avoid confusion).  $\mathbf{x}_j$  is the 2D notation for the position of site  $j$ , while  $\delta_{\ell}$  is the 2D notation for the lattice spacing previously indicated as  $\delta_{\ell}$ . Fourier transform it according to the convention

$$\hat{c}_{j\sigma} = \frac{1}{\sqrt{L_x L_y}} \sum_{\mathbf{k} \in \text{BZ}} e^{-i\mathbf{k} \cdot \mathbf{x}_j} \hat{c}_{\mathbf{k}\sigma}$$

Then:

$$\begin{aligned} \hat{n}_{j\uparrow} \hat{n}_{j \pm \delta_{\ell} \downarrow} &= \hat{c}_{j\uparrow}^{\dagger} \hat{c}_{j \pm \delta_{\ell} \downarrow}^{\dagger} \hat{c}_{j \pm \delta_{\ell} \downarrow} \hat{c}_{j\uparrow} \\ &= \frac{1}{(L_x L_y)^2} \sum_{\nu=1}^4 \sum_{\mathbf{k}_{\nu} \in \text{BZ}} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} e^{\pm i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_{\ell}} \hat{c}_{\mathbf{k}_1 \uparrow}^{\dagger} \hat{c}_{\mathbf{k}_2 \downarrow}^{\dagger} \hat{c}_{\mathbf{k}_3 \downarrow} \hat{c}_{\mathbf{k}_4 \uparrow} \end{aligned}$$

It follows,

$$\begin{aligned} \hat{h}_V^{(j)} &= -\frac{V}{(L_x L_y)^2} \sum_{\ell=x,y} \sum_{\nu=1}^4 \sum_{\mathbf{k}_{\nu} \in \text{BZ}} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} \\ &\quad \times \left( e^{i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_{\ell}} + e^{-i(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_{\ell}} \right) \hat{c}_{\mathbf{k}_1 \uparrow}^{\dagger} \hat{c}_{\mathbf{k}_2 \downarrow}^{\dagger} \hat{c}_{\mathbf{k}_3 \downarrow} \hat{c}_{\mathbf{k}_4 \uparrow} \\ &= -\frac{2V}{(L_x L_y)^2} \sum_{\ell=x,y} \sum_{\nu=1}^4 \sum_{\mathbf{k}_{\nu} \in \text{BZ}} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} \cos[(\mathbf{k}_2 - \mathbf{k}_3) \cdot \delta_{\ell}] \hat{c}_{\mathbf{k}_1 \uparrow}^{\dagger} \hat{c}_{\mathbf{k}_2 \downarrow}^{\dagger} \hat{c}_{\mathbf{k}_3 \downarrow} \hat{c}_{\mathbf{k}_4 \uparrow} \end{aligned}$$

The full non-local interaction is given by summing over all sites of  $\mathcal{S}$ . This gives back momentum conservation,

$$\frac{1}{L_x L_y} \sum_{j \in \mathcal{S}} e^{i[(\mathbf{k}_1 + \mathbf{k}_2) - (\mathbf{k}_3 + \mathbf{k}_4)] \cdot \mathbf{x}_j} = \delta_{\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4}$$

Let  $\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4 = \mathbf{K}$ , and define  $\mathbf{k}, \mathbf{k}'$  such that

$$\mathbf{k}_1 \equiv \mathbf{K} + \mathbf{k} \quad \mathbf{k}_2 \equiv \mathbf{K} - \mathbf{k} \quad \mathbf{k}_3 \equiv \mathbf{K} - \mathbf{k}' \quad \mathbf{k}_4 \equiv \mathbf{K} + \mathbf{k}' \quad \delta \mathbf{k} \equiv \mathbf{k} - \mathbf{k}'$$

Sums over these variables must be intended as over the Brillouin Zone (BZ). Then, finally

$$\begin{aligned} \hat{H}_V &\simeq \sum_{j \in \mathcal{S}} \hat{h}_V^{(j)} \\ &= -\frac{2V}{L_x L_y} \sum_{\ell=x,y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} \cos(\delta \mathbf{k} \cdot \boldsymbol{\delta}_\ell) \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \\ &= -\frac{2V}{L_x L_y} \sum_{\ell=x,y} \sum_{\mathbf{K}, \mathbf{k}, \mathbf{k}'} [\cos(\delta k_x \delta_\ell) + \cos(\delta k_y \delta_\ell)] \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \end{aligned}$$

Eventually, in the second passage the prefactor 2 can be absorbed by reintroducing the spin DoF<sup>2</sup>.

Taking in the mean-field approximation (with Cooper pair symmetry breaking), we get

$$\hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \simeq \langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} + \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \langle \hat{c}_{\mathbf{K}-\mathbf{k}'\downarrow} \hat{c}_{\mathbf{K}+\mathbf{k}'\uparrow} \rangle + \dots$$

Take e.g.  $\langle \hat{c}_{\mathbf{K}+\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{K}-\mathbf{k}\downarrow}^\dagger \rangle$ : the only non-zero contribution can come from the  $\mathbf{K} = \mathbf{0}$  term, as will be discussed self-consistently in Sec. 1.4.5. Then finally:

$$\hat{H}_V \simeq - \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} \left[ \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \hat{\phi}_{\mathbf{k}'} + \langle \hat{\phi}_{\mathbf{k}'} \rangle \hat{\phi}_{\mathbf{k}}^\dagger \right]$$

having I defined the pairing operator

$$\hat{\phi}_{\mathbf{k}} \equiv \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{\mathbf{k}\uparrow} \quad \hat{\phi}_{\mathbf{k}}^\dagger \equiv \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow}^\dagger$$

and the two-body potential

$$V_{\mathbf{k}\mathbf{k}'} = \frac{2V}{L_x L_y} [\cos(\delta k_x \delta_x) + \cos(\delta k_y \delta_y)]$$

Now, consider the term

$$\cos(\delta k_x) + \cos(\delta k_y) = \cos k_x \cos k'_x + \sin k_x \sin k'_x + \cos k_y \cos k'_y + \sin k_y \sin k'_y$$

For the sake of readability, the notations

$$c_\ell \equiv \cos k_\ell \quad s_\ell \equiv \sin k_\ell \quad c'_\ell \equiv \cos k'_\ell \quad s'_\ell \equiv \sin k'_\ell$$

are used. Group the four terms above,

$$\underbrace{(c_x c'_x + c_y c'_y)}_{\text{Symmetric}} + \underbrace{(s_x s'_x + s_y s'_y)}_{\text{Anti-symmetric}} \quad (1.20)$$

The first two exhibit inversion symmetry for both arguments  $\mathbf{k}, \mathbf{k}'$ ; the second two exhibit anti-symmetry. Decoupling the symmetric part,

$$c_x c'_x + c_y c'_y = \frac{1}{2}(c_x + c_y)(c'_x + c'_y) + \frac{1}{2}(c_x - c_y)(c'_x - c'_y)$$

<sup>2</sup>Justification can be given in two ways: either commuting appropriately the  $\hat{c}$  operators, or by carrying out the previous space sums independently over the two sublattices.

Structure	Structure factor	Graph
$s$ -wave	$\varphi_{\mathbf{k}}^{(s)} = 1$	Fig. 1.3a
Extended $s$ -wave	$\varphi_{\mathbf{k}}^{(s^*)} = \cos k_x + \cos k_y$	Fig. 1.3b
$p_x$ -wave	$\varphi_{\mathbf{k}}^{(p_x)} = i\sqrt{2} \sin k_x$	Fig. 1.3c
$p_y$ -wave	$\varphi_{\mathbf{k}}^{(p_y)} = i\sqrt{2} \sin k_y$	Fig. 1.3d
$d_{x^2-y^2}$ -wave	$\varphi_{\mathbf{k}}^{(d)} = \cos k_x - \cos k_y$	Fig. 1.3e

**Table 1.2** | Structure factors derived from the correlation structures of Tab. ???. The functions hereby defined are orthonormal, and define the various components of the non-local topological effective potential.

which finally gives:

$$\begin{aligned}
\cos(\delta k_x) + \cos(\delta k_y) &= \frac{1}{2}(c_x + c_y)(c'_x + c'_y) && (s^*\text{-wave}) \\
&+ s_x s'_x && (p_x\text{-wave}) \\
&+ s_y s'_y && (p_y\text{-wave}) \\
&+ \frac{1}{2}(c_x - c_y)(c'_x - c'_y) && (d_{x^2-y^2}\text{-wave})
\end{aligned}$$

In other words, the two-body potential decomposes as

$$\begin{aligned}
V_{\mathbf{k}\mathbf{k}'} &= \sum_{\gamma} V^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} \quad \text{where } \gamma = s^*, p_x, p_y, d_{x^2-y^2} \\
&= \frac{V}{L_x L_y} \sum_{\gamma} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*}
\end{aligned}$$

being  $\varphi_{\mathbf{k}}^{(\gamma)}$  the reciprocal-space expressions for the form factors of Tab. 1.1, listed explicitly in Tab. 1.2, and  $V_{\mathbf{k}\mathbf{k}'}^{(\gamma)}$  the symmetry-resolved components of the non-local attraction. Then the two-body potential has been decomposed in its planar symmetry components, each of which will naturally couple only to identically structured parameters in the full hamiltonian.

Define now the non-local gap function

$$\mathcal{V}_{\mathbf{k}} \equiv \sum_{\mathbf{k}'} V_{\mathbf{k}\mathbf{k}'} \langle \hat{\phi}_{\mathbf{k}'}^{\dagger} \rangle \quad (1.21)$$

one gets immediately

$$\hat{H}_V \simeq - \sum_{\mathbf{k}} \left[ \mathcal{V}_{\mathbf{k}} \hat{\phi}_{\mathbf{k}} + \mathcal{V}_{\mathbf{k}}^* \hat{\phi}_{\mathbf{k}}^{\dagger} \right] \quad (1.22)$$

To assume symmetry is broken in a specific symmetry channel  $\gamma$  means precisely to assume  $g_{ij} \propto \varphi_{ij}^{(\gamma)}$ , which in turn implies  $\langle \hat{\phi}_{\mathbf{k}} \rangle \propto \varphi_{\mathbf{k}}^{(\gamma)}$ . Of course, in Eq. (1.21) only the  $\gamma$  component of the potential survives, implying the gap function acquires the same symmetry,

$$\begin{aligned}
\mathcal{V}_{\mathbf{k}} &\propto \sum_{\mathbf{k}'} \frac{V}{L_x L_y} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} \varphi_{\mathbf{k}'}^{(\gamma)} \\
&\propto \varphi_{\mathbf{k}}^{(\gamma)}
\end{aligned}$$

where I used orthonormality of the  $\varphi_{\mathbf{k}}^{(\gamma)}$  functions.

### 1.4.3 Local interaction and gap function

A very similar argument can be carried out for the local  $U$  term. Without delving in too many details, the local gap  $\mathcal{U}_{\mathbf{k}}$  is given by

$$\mathcal{U}_{\mathbf{k}} \equiv \frac{U}{2L_x L_y} \sum_{\mathbf{k}} \langle \hat{\phi}_{\mathbf{k}} \rangle \quad (1.23)$$

evidently independent of  $\mathbf{k}$ , correctly. Identical considerations as in the above section hold for the local gap. The local part of the hamiltonian then gets

$$\hat{H}_U \simeq \sum_{\mathbf{k}} \left[ \mathcal{U}_{\mathbf{k}} \hat{\phi}_{\mathbf{k}} + \mathcal{U}_{\mathbf{k}}^* \hat{\phi}_{\mathbf{k}}^\dagger \right] \quad (1.24)$$

and the full gap function is simply

$$\Delta_{\mathbf{k}} \equiv \mathcal{V}_{\mathbf{k}} - \mathcal{U}_{\mathbf{k}} \quad (1.25)$$

Notice here that the only possible topology here is  $s$ -wave; define trivially the  $s$ -wave component of the total two-body interaction,

$$V^{(s)} = -\frac{U}{2L_x L_y}$$

Then the full effective interaction is collected in

$$\begin{aligned} \hat{H}_U + \hat{H}_V &\simeq - \sum_{\gamma} \sum_{\mathbf{k}, \mathbf{k}'} V^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)} \varphi_{\mathbf{k}'}^{(\gamma)*} \left[ \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \hat{\phi}_{\mathbf{k}'} + \langle \hat{\phi}_{\mathbf{k}} \rangle \hat{\phi}_{\mathbf{k}'}^\dagger \right] \\ &= - \sum_{\mathbf{k}} \left[ \Delta_{\mathbf{k}} \hat{\phi}_{\mathbf{k}} + \Delta_{\mathbf{k}}^* \hat{\phi}_{\mathbf{k}}^\dagger \right] \end{aligned}$$

The full self-consistency equation is given by

$$\Delta_{\mathbf{k}} \equiv \sum_{\mathbf{k}'} \left[ V^{(s)} + V_{\mathbf{k}\mathbf{k}'} \right] \langle \hat{\phi}_{\mathbf{k}'}^\dagger \rangle \quad (1.26)$$

The gap function decomposes in symmetry channels as well,

$$\Delta_{\mathbf{k}} = \sum_{\gamma} \Delta^{(\gamma)} \varphi_{\mathbf{k}}^{(\gamma)}$$

If SC arises in a specific symmetry channel,  $\Delta_{\mathbf{k}}$  will show the same symmetry. It follows, due to orthonormality and using Eq. (1.26),

$$\begin{aligned} \Delta^{(\gamma)} &= \frac{1}{L_x L_y} \sum_{\mathbf{k}} \varphi_{\mathbf{k}}^{(\gamma)*} \Delta_{\mathbf{k}} \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k}} \varphi_{\mathbf{k}}^{(\gamma)*} \sum_{\mathbf{k}'} \left[ V^{(s)} + V_{\mathbf{k}\mathbf{k}'} \right] \langle \hat{\phi}_{\mathbf{k}'}^\dagger \rangle \\ &= \frac{1}{L_x L_y} \sum_{\mathbf{k}} \varphi_{\mathbf{k}}^{(\gamma)*} \sum_{\mathbf{k}' \gamma'} V^{(\gamma')} \varphi_{\mathbf{k}}^{(\gamma')} \varphi_{\mathbf{k}'}^{(\gamma')*} \langle \hat{\phi}_{\mathbf{k}'}^\dagger \rangle \\ &= V^{(\gamma)} \sum_{\mathbf{k}} \varphi_{\mathbf{k}}^{(\gamma)*} \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle \end{aligned} \quad (1.27)$$

This result provides a set of self-consistency equations for each symmetry channel, listed in Tab. 1.3. Notice that to reconstruct self-consistently the full  $s$ -wave phase transition, the actual gap function is given by

$$\Delta^{(s)} + \Delta^{(s^*)} (c_x + c_y)$$

The  $s$ -wave transition is the only one equipped of both the local and the non-local parts. Within this structure, we are finally able to move to Nambu formalism.

#### 1.4.4 Nambu formalism and Bogoliubov transform

Define the Nambu spinor<sup>3</sup> as in BCS

$$\hat{\Psi}_{\mathbf{k}} \equiv \begin{bmatrix} \hat{c}_{\mathbf{k}\uparrow} \\ \hat{c}_{-\mathbf{k}\downarrow}^\dagger \end{bmatrix}$$

---

<sup>3</sup>Notice that the spinor is here differently defined with respect to App. ??, where because of the HF prevalence in mean-field decoupling the spinor components were homogeneously fermions creations or destructions.

Structure	Self-consistency equation	Graph
$s$ -wave	$\Delta^{(s)} = -\frac{U}{2L_x L_y} \sum_{\mathbf{k}} \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle$	Fig. 1.3a
Extended $s$ -wave	$\Delta^{(s^*)} = \frac{V}{L_x L_y} \sum_{\mathbf{k}} (c_x + c_y) \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle$	Fig. 1.3b
$p_x$ -wave	$\Delta^{(p_x)} = -i\sqrt{2} \frac{V}{L_x L_y} \sum_{\mathbf{k}} s_x \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle$	Fig. 1.3c
$p_y$ -wave	$\Delta^{(p_y)} = -i\sqrt{2} \frac{V}{L_x L_y} \sum_{\mathbf{k}} s_y \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle$	Fig. 1.3d
$d_{x^2-y^2}$ -wave	$\Delta^{(d)} = \frac{V}{L_x L_y} \sum_{\mathbf{k}} (c_x - c_y) \langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle$	Fig. 1.3e

**Table 1.3** Symmetry resolved self-consistency equations for the MFT parameters  $\Delta^{(\gamma)}$ , based on Eq. (1.26) and (1.27). By computing  $\langle \hat{\phi}_{\mathbf{k}}^\dagger \rangle$ , it is possible to reconstruct the various components of the gap function.

Evidently,

$$\phi_{\mathbf{k}} = \hat{\Psi}_{\mathbf{k}}^\dagger \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \hat{\Psi}_{\mathbf{k}} \quad \phi_{\mathbf{k}}^\dagger = \hat{\Psi}_{\mathbf{k}}^\dagger \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \hat{\Psi}_{\mathbf{k}} \quad (1.28)$$

The full hamiltonian is then given by:

$$\hat{H} = \sum_{\mathbf{k}} \hat{\Psi}_{\mathbf{k}} h_{\mathbf{k}} \hat{\Psi}_{\mathbf{k}} \quad h_{\mathbf{k}} \equiv \begin{bmatrix} \epsilon_{\mathbf{k}} & -\Delta_{\mathbf{k}}^* \\ -\Delta_{\mathbf{k}} & -\epsilon_{\mathbf{k}} \end{bmatrix} \quad (1.29)$$

Let  $\tau^\alpha$  for  $\alpha = x, y, z$  be the Pauli matrices. Define:

$$\hat{s}_{\mathbf{k}}^\alpha \equiv \hat{\Psi}_{\mathbf{k}}^\dagger \tau^\alpha \hat{\Psi}_{\mathbf{k}} \quad \text{for } \alpha = x, y, z$$

As can be shown easily, these operators realize spin-1/2 algebra.  $\hat{H}$  represents an ensemble of  $L_x L_y$  independent spins subject to pseudo-magnetic fields. Note that, differently from App. ?? where the chemical potential is inserted later (because in Nambu formalism it accounts for a diagonal term) here the chemical potential is part of the  $z$  component of the pseudo-magnetic field, since

$$\begin{aligned} \hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{-\mathbf{k}\downarrow} &= \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} + \hat{c}_{-\mathbf{k}\downarrow}^\dagger \hat{c}_{-\mathbf{k}\downarrow} \\ &= \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{\mathbf{k}\uparrow} - \hat{c}_{-\mathbf{k}\downarrow} \hat{c}_{-\mathbf{k}\downarrow}^\dagger + \mathbb{I} \\ &= \hat{\Psi}_{\mathbf{k}}^\dagger \tau^z \hat{\Psi}_{\mathbf{k}} + \mathbb{I} \end{aligned} \quad (1.30)$$

and then it follows

$$\begin{aligned} -\mu \hat{N} &= -\mu \sum_{\mathbf{k} \in \text{BZ}} [\hat{n}_{\mathbf{k}\uparrow} + \hat{n}_{-\mathbf{k}\downarrow}] \\ &= -\mu \sum_{\mathbf{k} \in \text{BZ}} \hat{\Psi}_{\mathbf{k}}^\dagger \tau^z \hat{\Psi}_{\mathbf{k}} - \mu L_x L_y \end{aligned}$$

Then, adding a term  $-\mu \hat{N}$  to  $\hat{H}$ , apart from an irrelevant total energy increase, changes the pseudo-field whose explicit form becomes

$$\mathbf{b}_{\mathbf{k}} \equiv \begin{bmatrix} -\text{Re}\{\Delta_{\mathbf{k}}\} \\ -\text{Im}\{\Delta_{\mathbf{k}}\} \\ \epsilon_{\mathbf{k}} - \mu \end{bmatrix} \quad (1.31)$$

This hamiltonian behaves as an ensemble of spins in local magnetic fields precisely as in Eq. (??),

$$\hat{H} - \mu \hat{N} = \sum_{\mathbf{k} \in \text{BZ}} \mathbf{b}_{\mathbf{k}} \cdot \hat{\mathbf{s}}_{\mathbf{k}} \quad \text{where} \quad \hat{\mathbf{s}}_{\mathbf{k}\sigma} = \begin{bmatrix} \hat{s}_{\mathbf{k}}^x \\ \hat{s}_{\mathbf{k}}^y \\ \hat{s}_{\mathbf{k}}^z \end{bmatrix} \quad (1.32)$$



Proceed as in App. ?? and diagonalize via a rotation,

$$d_{\mathbf{k}} \equiv \begin{bmatrix} -E_{\mathbf{k}} & \\ & E_{\mathbf{k}} \end{bmatrix} \quad \text{being} \quad E_{\mathbf{k}} \equiv \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}$$

and  $\xi_{\mathbf{k}} \equiv \epsilon_{\mathbf{k}} - \mu$ . Given the pseudoangles

$$\tan(2\theta_{\mathbf{k}}) \equiv \frac{|\Delta_{\mathbf{k}}|}{\epsilon_{\mathbf{k}}} \quad \tan(2\zeta_{\mathbf{k}}) \equiv \frac{\text{Im}\{\Delta_{\mathbf{k}}\}}{\text{Re}\{\Delta_{\mathbf{k}}\}}$$

the general diagonalizer will be an orthogonal rotation matrix

$$\begin{aligned} W_{\mathbf{k}} &= e^{i(\theta_{\mathbf{k}} - \frac{\pi}{2})\tau^y} e^{i\zeta_{\mathbf{k}}\tau^z} \\ &= \begin{bmatrix} -\sin\theta_{\mathbf{k}} & -\cos\theta_{\mathbf{k}} \\ \cos\theta_{\mathbf{k}} & -\sin\theta_{\mathbf{k}} \end{bmatrix} \begin{bmatrix} e^{i\zeta_{\mathbf{k}}} & \\ & e^{-i\zeta_{\mathbf{k}}} \end{bmatrix} \\ &= \begin{bmatrix} -\sin\theta_{\mathbf{k}}e^{i\zeta_{\mathbf{k}}} & -\cos\theta_{\mathbf{k}}e^{-i\zeta_{\mathbf{k}}} \\ \cos\theta_{\mathbf{k}}e^{i\zeta_{\mathbf{k}}} & -\sin\theta_{\mathbf{k}}e^{-i\zeta_{\mathbf{k}}} \end{bmatrix} \end{aligned} \quad (1.33)$$

given by a rotation of angle  $\zeta_{\mathbf{k}}$  around the  $z$  axis, to align the  $x$  axis with the field projection onto the  $xy$  plane, followed by a rotation around the  $y$  axis to anti-align with the pseudo-field. The MFT-BCS solution is given by a degenerate Fermi gas at ground state, whose quasi-particles occupy two bands  $\pm E_{\mathbf{k}}$  and their fermionic operators are given by

$$\hat{\gamma}_{\mathbf{k}}^{(-)} \equiv [W_{\mathbf{k}}\hat{\Psi}_{\mathbf{k}}]_1 \quad \hat{\gamma}_{\mathbf{k}}^{(+)} \equiv [W_{\mathbf{k}}\hat{\Psi}_{\mathbf{k}}]_2$$

The diagonalization operators are given by

$$\hat{\Gamma}_{\mathbf{k}} \equiv W_{\mathbf{k}}\hat{\Psi}_{\mathbf{k}} \quad \text{where} \quad \hat{\Gamma}_{\mathbf{k}} = \begin{bmatrix} \hat{\gamma}_{\mathbf{k}}^{(-)} \\ \hat{\gamma}_{\mathbf{k}}^{(+)} \end{bmatrix}$$

then, using Eq. (??),

$$\langle [\hat{\Psi}_{\mathbf{k}}^\dagger]_i [\hat{\Psi}_{\mathbf{k}}]_j \rangle = [W_{\mathbf{k}}]_{1i} [W_{\mathbf{k}}^\dagger]_{j1} f(-E_{\mathbf{k}}; \beta, 0) + [W_{\mathbf{k}}]_{2i} [W_{\mathbf{k}}^\dagger]_{j2} f(E_{\mathbf{k}}; \beta, 0)$$

where in the Fermi-Dirac function chemical potential was set to zero, because it already was included in the diagonalized hamiltonian. Recalling Eq. (??), it follows

$$\langle \phi_{\mathbf{k}}^\dagger \rangle = [W_{\mathbf{k}}]_{11} [W_{\mathbf{k}}^\dagger]_{21} f(-E_{\mathbf{k}}; \beta, 0) + [W_{\mathbf{k}}]_{21} [W_{\mathbf{k}}^\dagger]_{22} f(E_{\mathbf{k}}; \beta, 0) \quad (1.34)$$

$$= \frac{1}{2} \sin(2\theta_{\mathbf{k}}) e^{i2\zeta_{\mathbf{k}}} \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right) \quad (1.35)$$

The last passage has been obtained by computing the matrix element from the explicit form of  $W_{\mathbf{k}}$  of Eq. (1.33) and by the simple relation

$$\begin{aligned} \frac{1}{e^{-x} + 1} - \frac{1}{e^x + 1} &= \frac{e^x - 1}{e^x + 1} \\ &= \tanh\left(\frac{x}{2}\right) \end{aligned}$$

Eqns. (1.34), (1.35) give us both the algorithmic formula (first row) and its theoretical counterpart (second row) to compute the order parameters in the HF approach at each point in  $k$ -space ( $k_x, k_y$ ). We can finally derive the BCS self-consistency equation

$$\Delta_{\mathbf{k}} \equiv \frac{1}{2} \sum_{\mathbf{k}'} [V^{(s)} + V_{\mathbf{k}\mathbf{k}'}] \frac{|\Delta_{\mathbf{k}}|}{\sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}} e^{i\text{Im}\{\Delta_{\mathbf{k}}\}/\text{Re}\{\Delta_{\mathbf{k}}\}} \tanh\left(\frac{\beta}{2} \sqrt{\xi_{\mathbf{k}}^2 + |\Delta_{\mathbf{k}}|^2}\right) \quad (1.36)$$

The whole point of the HF algorithm is to find an iterative solution for each symmetry channel, using the self-consistency equation projection of Tab. 1.3.

Notice that the  $z$  component of the spin operators is related to density: using Eq. (??),

$$\langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^z \hat{\Psi}_{\mathbf{k}} \rangle = \langle [\hat{\Psi}_{\mathbf{k}}^\dagger]_1 [\hat{\Psi}_{\mathbf{k}}]_1 \rangle - \langle [\hat{\Psi}_{\mathbf{k}}^\dagger]_2 [\hat{\Psi}_{\mathbf{k}}]_2 \rangle$$

I proceed in as done previously, and from Eq. (1.30),

$$\begin{aligned} \langle \hat{n}_{\mathbf{k}\uparrow} \rangle + \langle \hat{n}_{-\mathbf{k}\downarrow} \rangle &= 1 + \langle \hat{\Psi}_{\mathbf{k}}^\dagger \tau^z \hat{\Psi}_{\mathbf{k}} \rangle \\ &= 1 + \left( |[W_{\mathbf{k}}]_{11}|^2 - |[W_{\mathbf{k}}]_{12}|^2 \right) f(-E_{\mathbf{k}}; \beta, 0) \\ &\quad + \left( |[W_{\mathbf{k}}]_{21}|^2 - |[W_{\mathbf{k}}]_{22}|^2 \right) f(E_{\mathbf{k}}; \beta, 0) \end{aligned} \quad (1.37)$$

$$= 1 - \cos(2\theta_{\mathbf{k}}) \tanh\left(\frac{\beta E_{\mathbf{k}}}{2}\right) \quad (1.38)$$

The expectation value for the density is needed in order to extract the optimal chemical potential  $\mu$  for the target density we aim to simulate at the given parametrization. This is numerically obtained by using Eq. (1.37) directly on the diagonalization matrix of  $h_{\mathbf{k}}$ .

#### 1.4.5 A short comment on self-consistency

The Bogoliubov fermions in spinor representation satisfy obviously  $\hat{\Psi}_{\mathbf{k}} = W_{\mathbf{k}}^\dagger \hat{\Gamma}_{\mathbf{k}}$ . Consider e.g.

$$\langle \hat{c}_{\mathbf{k}\sigma}^\dagger \hat{c}_{-\mathbf{k}\sigma}^\dagger \rangle$$

which is a spin-symmetric anomalous Cooper pair. For simplicity, take  $\sigma = \uparrow$ . Expand:

$$\begin{aligned} \langle \hat{c}_{\mathbf{k}\uparrow}^\dagger \hat{c}_{-\mathbf{k}\uparrow}^\dagger \rangle &= \langle [\hat{\Psi}_{\mathbf{k}}^\dagger]_1 [\hat{\Psi}_{-\mathbf{k}}^\dagger]_1 \rangle \\ &= \langle [W_{\mathbf{k}} \hat{\Gamma}_{\mathbf{k}}^\dagger]_1 [W_{-\mathbf{k}} \hat{\Gamma}_{-\mathbf{k}}^\dagger]_1 \rangle \end{aligned}$$

This expectation value is taken over the ground-state, the latter being the vacuum of  $\Gamma$  fermions. Evidently the above expectation cannot assume non-zero values. Obviously the same holds for  $\sigma = \downarrow$ , and this argument explains why the Ferromagnetic terms of the hamiltonian decomposition do not contribute to Cooper instability. An identical argument, with the exchange

$$(\sigma, \sigma) \rightarrow (\uparrow, \downarrow) \quad \text{and} \quad (\mathbf{k}, -\mathbf{k}) \rightarrow (\mathbf{K} + \mathbf{k}, \mathbf{K} - \mathbf{k}) \quad \text{with} \quad \mathbf{K} \neq \mathbf{0}$$

justifies why in Sec. 1.4.4 the only relevant contribution was given by  $\mathbf{K} = \mathbf{0}$ . In the next sections, the results of the self-consistent HF algorithm are exposed.

### 1.5 Results of the HF algorithm

[To be continued...]



# Bibliography

- [1] Zhangkai Cao et al. *p-wave superconductivity induced by nearest-neighbor attraction in the square-lattice extended Hubbard model*. en. arXiv:2408.01113 [cond-mat]. Jan. 2025. DOI: [10.48550/arXiv.2408.01113](https://doi.org/10.48550/arXiv.2408.01113). URL: <http://arxiv.org/abs/2408.01113> (visited on 03/15/2025).
- [2] Piers Coleman. *Introduction to Many-Body Physics*. Cambridge University Press, 2015.
- [3] Michele Fabrizio. *A Course in Quantum Many-Body Theory*. Springer, 2022.
- [4] Gabriele Giuliani and Giovanni Vignale. *Quantum Theory of the Electron Liquid*. Cambridge University Press, 2005.
- [5] Giuseppe Grosso and Giuseppe Pastori Parravicini. *Solid State Physics*. Second Edition. Academic Press, 2014.
- [6] J. E. Hirsch. “Two-dimensional Hubbard model: Numerical simulation study”. In: *Phys. Rev. B* 31 (7 Apr. 1985), pp. 4403–4419. DOI: [10.1103/PhysRevB.31.4403](https://doi.org/10.1103/PhysRevB.31.4403). URL: <https://link.aps.org/doi/10.1103/PhysRevB.31.4403>.
- [7] Robin Scholle et al. “Comprehensive mean-field analysis of magnetic and charge orders in the two-dimensional Hubbard model”. In: *Phys. Rev. B* 108 (3 July 2023), p. 035139. DOI: [10.1103/PhysRevB.108.035139](https://doi.org/10.1103/PhysRevB.108.035139). URL: <https://link.aps.org/doi/10.1103/PhysRevB.108.035139>.