

1. (12 pts) The two parts are not related.

- (a) Determine whether the following statement is **true** or **false**, and explain your answer: The set in \mathbb{R}^3 described in cartesian coordinates as $A = \{(x, y, z) \in \mathbb{R}^3 : z^2 = x^2 + y^2\}$ is the same as the set in \mathbb{R}^3 described in spherical coordinates as $B = \{(\rho, \theta, \phi) : \phi = \frac{3\pi}{4}\}$, under the usual convention $\rho \geq 0$, $\theta \in [0, 2\pi)$ and $\phi \in [0, \pi]$.

False, B is the lower half of a cone
but A is a full cone.


- (b) A thin lamina occupies the region

$$D = \{(x, y) : 1 \leq x^2 + y^2 \leq 16 \text{ and } y \geq |x|\}.$$

If the density function ρ at each point (x, y) is inversely proportional to the square of the distance of the point to the origin, find the y coordinate of the center of mass of the lamina (the \bar{y}).

$$\rho(x, y) = \frac{k}{(\sqrt{x^2 + y^2})^2} = \frac{k}{x^2 + y^2} = \frac{k}{r^2} \text{ in polar}$$

$$D = \{(r, \theta) : \frac{\pi}{4} \leq \theta \leq \frac{3\pi}{4}, 1 \leq r \leq 4\}$$



$$M = \iint_D \rho(x, y) dA = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \int_1^4 \frac{k}{r^2} r dr d\theta$$

$$= \frac{\pi}{2} k \ln r \Big|_1^4 = \frac{\pi}{2} k \ln 4$$


$$M_x = \iint_D y \rho(x, y) dA = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \int_1^4 r \sin \theta \frac{k}{r^2} r dr d\theta$$

$$= 3k [\cos \theta]_{\frac{\pi}{4}}^{\frac{3\pi}{4}}$$

$$\text{So } \bar{y} = \frac{3k\sqrt{2}}{\frac{\pi}{2}k \ln 4} = \frac{6\sqrt{2}}{\pi \ln 4}$$

$$= 3k \sqrt{2}$$

2. (8 pts) Let $f(x, y, z) = xy$. Set up but do not evaluate $\iiint_E f(x, y, z) dV$ in cylindrical coordinates, where E is the solid that lies above the sphere $x^2 + y^2 + z^2 = 9$, under the cone $z = -\sqrt{x^2 + y^2}$ and satisfies $y \leq 0$.



$$x^2 + y^2 + z^2 = 9 \Rightarrow$$

$$r^2 + z^2 = 9$$

$$\Rightarrow z = \pm \sqrt{9 - r^2}$$

$$z = -\sqrt{9 - r^2} \text{ bec. we want lower part}$$

$$z = -\sqrt{x^2 + y^2} \Rightarrow z = -\sqrt{r^2} \Rightarrow z = -r$$

$$\text{So } -\sqrt{9 - r^2} \leq z \leq -r$$

$$y \leq 0 \Rightarrow \theta \in [\pi, 2\pi]$$

Find projection:

$$\begin{cases} r^2 + z^2 = 9 \\ z = -r \end{cases} \Rightarrow$$

$$2r^2 = 9 \Rightarrow r = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2}$$


So

$$\iiint_E f dV = \int_{\pi}^{2\pi} \int_0^{\frac{3\sqrt{2}}{2}} \int_{-\sqrt{9-r^2}}^{-r} r \cos \theta r \sin \theta r dz dr d\theta$$

545

3. (16 pts) [You should be able to answer each part regardless of whether you have answered the other one]
Let $f(x, y, z) = z$.

(a) Set up but do not evaluate $\iiint_E f(x, y, z) dV$ in the order $dx dy dz$, where E is the solid in the first octant bounded by the coordinate planes, the circular cylinder $x^2 + y^2 = 4$ and circular cylinder $x^2 + z^2 = 4$. (make sure to involve the given function in your formula!)



With $u_1(y, z) \leq x \leq u_2(y, z)$

(1) $\Rightarrow x = \sqrt{4 - y^2}$

(2) $\Rightarrow x = \sqrt{4 - z^2}$

(3) $x = 0$

(1) $x^2 + y^2 = 4$

(2) $x^2 + z^2 = 4$

(3) $x = 0$

(4) $y = 0$

(5) $z = 0$

Projection.

(1), (2) $\rightarrow y^2 = z^2 \Rightarrow y = \pm z \Rightarrow y = z$ ($y, z \geq 0$)

(4) $y = 0$ (5) $z = 0$

(1), (3) $\Rightarrow y = 2$ (2), (3) $\Rightarrow z = 2$ int. 3.

On D_1 $0 \leq x \leq \sqrt{4 - z^2}$

D_2 $0 \leq x \leq \sqrt{4 - y^2}$

$D_1 = \{(y, z): 0 \leq z \leq 2, 0 \leq y \leq z\}$

$D_2 = \{(y, z): 0 \leq z \leq 2, z \leq y \leq 2\}$

5.55 (b) Evaluate $\iiint_E f(x, y, z) dV$ using cylindrical coordinates.

$0 \leq z \leq \sqrt{4 - x^2} \Rightarrow$

$\Rightarrow 0 \leq z \leq \sqrt{4 - r^2 \cos^2 \theta}$

$0 \leq r \leq 2, 0 \leq \theta \leq \frac{\pi}{2}$

$\int_0^2 \int_0^{\frac{\pi}{2}} \int_0^{\sqrt{4 - r^2 \cos^2 \theta}} z r dz d\theta dr =$

$= \int_0^2 \int_0^{\frac{\pi}{2}} \frac{1}{2} (4 - r^2 \cos^2 \theta) r d\theta dr = \int_0^2 \int_0^{\frac{\pi}{2}} 2r d\theta dr - \frac{1}{2} \int_0^2 \int_0^{\frac{\pi}{2}} r^3 \cos^2 \theta d\theta dr$

$= 4 \cdot \frac{\pi}{2} - \frac{1}{2} \cdot \frac{2^4}{4} \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta = 2\pi - 2 \int_0^{\frac{\pi}{2}} \frac{1 + \cos 2\theta}{2} d\theta = 2\pi - 2 \cdot \frac{\pi}{4}$

So

$\iiint_E f(x, y, z) dV =$

$= \int_0^2 \int_0^{\frac{\pi}{2}} \int_0^{\sqrt{4 - z^2}} z dx dy dz$

$+ \int_0^2 \int_0^{\frac{\pi}{2}} \int_0^{\sqrt{4 - y^2}} z dx dy dz$

4. (12 pts) Let R be the trapezoid in the xy plane defined by the points $(1,1)$, $(2,2)$, $(2,0)$ and $(4,0)$, as in the picture, and you are given the transformation $x = u + v$ and $y = u - v$.

- (a) Compute the Jacobian determinant $\frac{\partial(x,y)}{\partial(u,v)}$.

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} = -2$$

- (b) Find the inverse transformation T^{-1} (that is, $u = u(x,y)$ and $v = v(x,y)$).

$$\begin{cases} x = u + v \\ y = u - v \end{cases} \Rightarrow \begin{cases} x + y = 2u \\ x - y = 2v \end{cases} \Rightarrow \begin{cases} u = \frac{x+y}{2} \\ v = \frac{x-y}{2} \end{cases}$$


- (c) Find the image S of R under T^{-1} in the uv plane (that is, the set $S = T^{-1}(R)$) and draw a picture of it.

$$y = x \Rightarrow 2v = 0 \Rightarrow v = 0$$

$$y = 4 - x \Rightarrow u = 2$$

$$y = 0 \Rightarrow u = v$$

$$y = 2 - x \Rightarrow u = 1$$



- (d) Use your work in the parts (a)-(c) to calculate $\iint_R e^{\frac{x-y}{x+y}} dA$ (you can use the back of the page if you run out of space).

$$\int_1^2 \int_0^u e^{\frac{2v}{2u} \cdot |-2|} dv du =$$

$$= \int_1^2 \left[2 \int_0^u u e^v dv \right] du = \int_1^2 2u(e-1) du$$

$$= 3(e-1)$$

