2 Fourier Series and LTI Systems

- Eigenfunctions Revisited
- 3 Frequency Response of LTI System
 - Continuous Time
 - Discrete Time
 - Periodic Signals
 - Examples using Frequency Response
- 4 Freq Shaping and Filtering
 - Quick Review of Analogue Filters (non-assessable)
 - Key Observation
 - CT Low Pass Filter
 - CT High Pass Filter
 - CT Band Pass Filter
 - DT Low Pass Filter
 - DT High Pass Filter
 - DT Band Pass Filter
 - Moving Average Filter
 - Other Types of Filters



- Previous focus was unit samples $\delta[n]$ and impulses $\delta(t)$ convolution
- Alternative focus is eigenfunctions of LTI systems

Eigenfunction definition:

A signal for which the system output is a (possibly complex) constant times the input is referred to as an eigenfunction of the system, and the amplitude factor is refereed to as the systems eigenvalue.

stralian tional jversity

eigenvalue

eigenvalue

figenfunction

final jversity

Convenor: R. A. Kennedy

Fourier Series and LTI Systems – Eigenfunctions Revisited

- \bullet $e^{j\omega t}$, e^{st} are eigenfunctions of CT LTI systems
- ullet $e^{j\omega n}$, z^n are eigenfunctions of DT LTI systems

- Study of CT LTI systems using $e^{j\omega t}$:
 - Fourier series (FS) periodic signals
 - Fourier transform (FT) general signals
- Study of CT LTI systems using e^{st} :
 - Laplace transform (outside the scope of this course)
- Study of DT LTI systems using $e^{j\omega n}$:
 - Discrete-time Fourier series (DTFS) periodic signals
 - Discrete-time Fourier transform (DTFT) general signals
- Study of DT LTI systems using z^n :
 - z-transform (to be covered in ENGN4537 DT Signal Processing)



Fourier Series and LTI Systems – Eigenfunctions Revisited

$$x(t) = e^{j\omega t} \longrightarrow \text{CT LTI} \quad h(t) \longrightarrow y(t) = ?$$

$$y(t) = x(t) * h(t) = h(t) * x(t)$$

$$= \int_{-\infty}^{\infty} h(\tau)x(t - \tau)d\tau$$

$$= \int_{-\infty}^{\infty} h(\tau)e^{j\omega(t - \tau)}d\tau$$

$$= e^{j\omega t} \int_{-\infty}^{\infty} h(\tau)e^{-j\omega\tau}d\tau$$

Therefore

where
$$H(j\omega)=\int_{-\infty}^{\infty}h(\tau)e^{-j\omega\tau}d\tau$$
 .

Generalisation: If the input to an LTI system is expressed as a linear combination of periodic complex exponentials, the output can also be expressed in this form.



Fourier Series and LTI Systems – Eigenfunctions Revisited

$$x[n] = e^{j\omega n} \longrightarrow \text{DT LTI} \quad h[n] \longrightarrow y[n] = ?$$

$$y[n] = x[n] * h[n] = h[n] * x[n]$$

$$= \sum_{k=-\infty}^{\infty} h[k]x[n-k]$$

$$= \sum_{k=-\infty}^{\infty} h[k]e^{j\omega(n-k)}$$

$$= e^{j\omega n} \sum_{k=-\infty}^{\infty} h[k]e^{-j\omega k}$$

Therefore

$$y[n] = e^{j\omega n} H(e^{j\omega})$$

where $H(e^{j\omega}) = \sum_{k=-\infty}^{\infty} h[k]e^{-j\omega k}$







- 2 Fourier Series and LTI Systems
 - Eigenfunctions Revisited

Frequency Response of LTI System

- Continuous Time
- Discrete Time
- Periodic Signals
- Examples using Frequency Response

4 Freq Shaping and Filtering

- Quick Review of Analogue Filters (non-assessable)
- Key Observation
- CT Low Pass Filter
- CT High Pass Filter
- CT Band Pass Filter
- DT Low Pass Filter
- DT High Pass Filter
- DT Band Pass Filter
- Moving Average Filter
- Other Types of Filters



- 2 Fourier Series and LTI Systems
 - Eigenfunctions Revisited

3 Frequency Response of LTI System

- Continuous Time
- Discrete Time
- Periodic Signals
- Examples using Frequency Response

4 Freq Shaping and Filtering

- Quick Review of Analogue Filters (non-assessable)
- Key Observation
- CT Low Pass Filter
- CT High Pass Filter
- CT Band Pass Filter
- DT Low Pass Filter
- DT High Pass Filter
- DT Band Pass Filter
- Moving Average Filter
- Other Types of Filters



Definition (CT Frequency Response)

The CT Frequency Response is defined by

$$H(j\omega) \triangleq \int_{-\infty}^{\infty} h(t) e^{-j\omega t} dt = H(j\omega)$$

Note that ω need not be multiples of some ω_0 . ω can take any value (still have eigenfunctions).



- 2 Fourier Series and LTI Systems
 - Eigenfunctions Revisited


Frequency Response of LTI System

- Continuous Time
- Discrete Time
- Periodic Signals
- Examples using Frequency Response

4 Freq Shaping and Filtering

- Quick Review of Analogue Filters (non-assessable)
- Key Observation
- CT Low Pass Filter
- CT High Pass Filter
- CT Band Pass Filter
- DT Low Pass Filter
- DT High Pass Filter
- DT Band Pass Filter
- Moving Average Filter
- Other Types of Filters





Definition (DT Frequency Response)

The DT Frequency Response is defined by

$$H(e^{j\omega}) \triangleq \sum_{n=-\infty}^{\infty} h[n] e^{-j\omega n} = H(e^{j\omega})$$

Note that ω need not be multiples of some ω_0 . ω can take any value (still have eigenfunctions).



- 2 Fourier Series and LTI Systems
 - Eigenfunctions Revisited

3 Frequency Response of LTI System

- Continuous Time
- Discrete Time
- Periodic Signals
- Examples using Frequency Response

4 Freq Shaping and Filtering

- Quick Review of Analogue Filters (non-assessable)
- Key Observation
- CT Low Pass Filter
- CT High Pass Filter
- CT Band Pass Filter
- DT Low Pass Filter
- DT High Pass Filter
- DT Band Pass Filter
- Moving Average Filter
- Other Types of Filters



Frequency Response of LTI System – Periodic Signals

Continuous Time

$$\left\langle \begin{array}{c} periodic \\ signal \end{array} \right\rangle \, x(t) \longrightarrow \left\langle \begin{array}{c} h(t) \\ signal \end{array} \right\rangle \, y(t)$$

Fourier Series representation

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t} \longrightarrow h(t) \longrightarrow y(t) = \sum_{k=-\infty}^{\infty} \underbrace{a_k \underbrace{H(jk\omega_0)}_{b_k} e^{jk\omega_0 t}}_{b_k}$$

The Fourier coefficients map like

b like
$$y(t) = \sum_{k=-\infty}^{\infty} b_k e^{jk} w_0 t$$

$$a_k \to b_k \triangleq H(jk\omega_0) a_k = \infty$$

Here the complex gain (eigenvalue) $H(jk\omega_0) \in \mathbb{C}$ scales/filters the periodic signal component at frequency $k\omega_0$, that is, $e^{jk\omega_0t}$ (eigenfunction).



Frequency Response of LTI System – Periodic Signals

Discrete Time

$$\left\langle \begin{array}{c} periodic \\ signal \end{array} \right\rangle x[n] \longrightarrow \left\langle \begin{array}{c} h[n] \\ signal \end{array} \right\rangle y[n]$$

DT

Fourier Series representation

$$x[n] = \sum_{k=0}^{N-1} a_k e^{jk\omega_0 n} \longrightarrow h[n] \longrightarrow y(t) = \sum_{k=0}^{N-1} \underbrace{a_k H(e^{jk\omega_0})}_{b_k} e^{jk\omega_0 n}$$

The Fourier coefficients map like

$$a_k \to b_k \triangleq H(e^{jk\omega_0}) a_k$$

Here complex gain $H(e^{jk\omega_0}) \in \mathbb{C}$ scales/filters the periodic signal component at frequency $k\omega_0$.




- 2 Fourier Series and LTI Systems
 - Eigenfunctions Revisited
- **3** Frequency Response of LTI System
 - Continuous Time
 - Discrete Time
 - Periodic Signals
 - Examples using Frequency Response
- 4 Freq Shaping and Filtering
 - Quick Review of Analogue Filters (non-assessable)
 - Key Observation
 - CT Low Pass Filter
 - CT High Pass Filter
 - CT Band Pass Filter
 - DT Low Pass Filter
 - DT High Pass Filter
 - DT Band Pass Filter
 - Moving Average Filter
 - Other Types of Filters



Frequency Response of LTI System –Given h(t) find $H(j\omega)$

Remember:

- h(t) impulse response $\delta(t) \to h(t)$
- $H(j\omega)$ frequency response LTI system, by definition $H(j\omega)=\int_{-\infty}^{\infty}h(t)e^{-j\omega t}dt$





Frequency Response of LTI System –Given h[n] find $H(e^{j\omega)}$

Remember:

- h[n] impulse response $\delta[n] \to h[n]$
- $H(e^{j\omega})$ frequency response LTI system, by definition $H(e^{j\omega})=\sum_{n=-\infty}^{\infty}h[n]e^{-j\omega n}$
- Example: $h[n] = \alpha^n u[n]$, $-1 < \alpha < 1$ $H(e^{j\omega}) = \sum_{n=-\infty}^{\infty} \chi^n u[n] e^{-j\omega n} = \sum_{n=0}^{\infty} \chi^n e^{-j\omega n}$

X in geometric sone



Frequency Response of LTI System – Finding y(t)

RC circuit with input x(t). We know $h(t) = \frac{1}{RC} e^{-\frac{t}{RC}} u(t)$, what is y(t)?

- Could find y(t) = x(t) * h(t)
- Use Fourier series to solve for y(t) instead
- Output Fourier coefficients given by $b_k = H(jk\omega_0)a_k$
- First need $H(j\omega)$: $H(j\omega) = \int_{-\infty}^{\infty} h(t) e^{-j\omega t} dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-j\omega t} dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} \frac{1}{Rc} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt$ $= \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt = \int_{-\infty}^{\infty} e^{-t} (j\omega + /Rc) dt$ $= \int_$

Frequency Response of LTI System –Finding y(t) Cont.



RC circuit with input x(t). We know $h(t) = \frac{1}{RC}e^{-\frac{t}{RC}}u(t)$, what is y(t)?

- $x(t) = \cos(2\pi 1000t)$ (1kHz) $\leftarrow \omega_0 = 2\pi(000)$
- $R = \frac{1000}{2\pi}$
- $C = 10 \ \mu F$
- Output Fourier coefficients given by $b_k = H(jk\omega_0)a_k$

Need to find
$$H(jk\omega_0)$$
:

$$H(j\omega)$$
, $\omega = k\alpha$

Need to find
$$H(jk\omega_0)$$
: $+|(j\omega)\rangle = |(j\omega)\rangle$



jkwoRC+1 jk 21/000 x 1000 x 10 x 16 +

Need to find FS coefficients x(t), a_k :

$$x(+) = 10^{12} \pi 1000t$$

$$Q_{1} = \frac{1}{2}$$


$$Q_{-1} = \frac{1}{2}$$




Frequency Response of LTI System –Finding y(t) Cont.

RC circuit with input x(t). We know $h(t) = \frac{1}{RC} e^{-\frac{t}{RC}} u(t)$, what is y(t)?

• Output Fourier coefficients given by $b_k = H(jk\omega_0)a_k$








$$=2 \times 0.04975 \quad Cos(wot-1.4711)$$

$$= 0.0995 \cos \left(27110000t - 1.4711\right)$$



Part 2 Slide 51/many Convenor: R. A. Kennedy

CT Example Summary



Frequency Response of LTI System –Given h[n] find $H(e^{j\omega})$

Difference equation: $y[n] - \frac{1}{4}y[n-1] = x[n]$

Sol. 1: Solve difference equation to find h[n], then find $H(e^{j\omega})$

$$y[n] = \frac{1}{4}y[n-1] + x[n]$$

$$h[n] = \frac{1}{4}h[n-1] + \delta[n]$$

$$h[n]$$

$$h[0] = \frac{1}{4}h[-1] + \delta[0] = 0 + 1 = 1$$

 $h[1] = \frac{1}{4}h[0] + \delta[1] = \frac{1}{4} + 0 = \frac{1}{4}$
 $h[2] = \frac{1}{4}h[1] + \delta[2] = \frac{1}{4}(\frac{1}{4})^{2} + 3$

$$h[n] = \left(\frac{1}{4}\right)^n u[n]$$



$$H(ejw) = \sum_{n=-\infty}^{\infty} h[n]e^{-jwn} = \sum_{n=-\infty}^{\infty} (\frac{1}{4})^n u[n]e^{-jwn}$$

$$= \sum_{n=0}^{\infty} (\frac{1}{4})^n u[n]e^{-jwn}$$

$$= \sum_{n=-\infty}^{\infty} (\frac{1}{4})^n u[n]e^{-jwn}$$

$$= \sum_{n=-\infty}^{\infty} (\frac{1}{4})^n u[n]e^{-jwn}$$

Frequency Response of LTI System –Given h[n] find $H(e^{j\omega)}$



Difference equation: $y[n] - \frac{1}{4}y[n-1] = x[n]$

Sol. 2: Avoids recursion

Sol. 2: Avoids recursion
$$H(e^{j\omega})e^{j\omega n} - \frac{1}{4}H(e^{j\omega})e^{j\omega n}e^{-j\omega} = e^{j\omega n}$$

